


ORIGINAL ARTICLE

Update on the role of antiseptics in the management of chronic wounds with critical colonisation and/or biofilm

Paulo J. Alves¹  | Ruben T. Barreto² | Brigitte M. Barrois³ | Luc G. Gryson⁴ | Sylvie Meaume⁵ | Stan J. Monstrey⁶

¹Wounds Research Laboratory, Universidade Católica Portuguesa, Porto, Portugal

²Advanced Wound Diagnosis and Treatment Centre, Funchal, Portugal

³Société Française de l'Escarre, Saint-Saturnin, France

⁴Belgian Defence Military Medical Component, Brussels, Belgium

⁵Department of Geriatrics and Wound Care Unit, Hospital Rothschild, APHP Assistance Publique Hôpitaux de Paris, Sorbonne Université, Paris, France

⁶Department of Plastic Surgery, Ghent University Hospital, Ghent, Belgium

Correspondence

Paulo J. Alves, Universidade Católica Portuguesa, Institute of Health Sciences, Wounds Research Lab, Centre for Interdisciplinary Research in Health (CIIS), Rua Diogo de Botelho, 1327, 4169-005 Porto, Portugal.
Email: pjalves@porto.ucp.pt

Abstract

Biofilms play a major role in delaying chronic wounds from healing. A wound infiltrated with biofilm, or “critically colonised” wound, may become clinically infected if the number of microbes exceeds a critical level. Chronic wound biofilms represent a significant treatment challenge by demonstrating recalcitrance towards antimicrobial agents. However, a “window of opportunity” may exist after wound debridement when biofilms are more susceptible to topical antiseptics. Here, we discuss the role of antiseptics in the management of chronic wounds and biofilm, focusing on povidone-iodine (PVP-I) in comparison with two commonly used antiseptics: polyhexanide (PHMB) and silver. This article is based on the literature reviewed during a focus group meeting on antiseptics in wound care and biofilm management, and on a PubMed search conducted in March 2020. Compared with PHMB and silver, PVP-I has a broader spectrum of antimicrobial activity, potent antibiofilm efficacy, no acquired bacterial resistance or cross-resistance, low cytotoxicity, good tolerability, and an ability to promote wound healing. PVP-I represents a viable therapeutic option in wound care and biofilm management, with the potential to treat biofilm-infiltrated, critically colonised wounds. We propose a practical algorithm to guide the management of chronic, non-healing wounds due to critical colonisation or biofilm, using PVP-I.

KEYWORDS

biguanides, biofilms, povidone-iodine, silver, wound healing

1 | INTRODUCTION

Wound healing is a complicated and tightly regulated process that is essential to restore the normal barrier function of the skin, thereby preventing further damage or infection.^{1,2} The normal wound healing process involves sequential but overlapping phases, including inflammation, proliferation, and remodelling.¹ These phases are

mediated by a range of cell types including fibroblasts, keratinocytes, endothelial cells, and macrophages, the activity of which is carefully coordinated by a range of growth factors, cytokines, and chemokines.³ However, when a wound fails to progress through the normal successive phases of wound healing in an orderly and timely manner, a chronic wound may result.^{1,4} Chronic wounds, comprising vascular leg ulcers (eg, venous and arterial

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2020 The Authors. International Wound Journal published by Medicalhelplines.com Inc (3M) and John Wiley & Sons Ltd.

ulcers), diabetic foot ulcers, and pressure ulcers, represent a significant burden to the individual and to the healthcare system.^{5,6} There are many factors that may be responsible for the delay and/or failure of a chronic wound to heal, including the patient's age, the nutritional status (eg, obesity) of the patient, how oxygenated the wound is, and the presence of either an underlying chronic disease (eg, diabetes) or an immunocompromised condition (eg, cancer).^{7,8} Certain therapies may also hinder wound healing, including chemotherapeutic agents, radiation therapy and long-term use of corticosteroids, and non-steroidal anti-inflammatory drugs.^{9,10}

Perhaps the most important factor that can affect the healing of a chronic wound, and subsequently increase the likelihood the wound may become infected, is the presence of a biofilm.¹¹⁻¹⁴ Biofilms are generally polymicrobial communities, attached either to each other or to a surface that become encased within an extracellular polymeric substance (EPS).^{11,15,16} Most biofilms located within chronic wounds are composed of 10% to 20% microorganisms and 80% to 90% EPS.¹¹ Biofilms cause chronic wounds to become locked in an inflammatory state,^{17,18} and they are thought to be the root cause of approximately 80% of all infections in humans,¹¹ and most medical device-related infections.¹⁹ A recent systematic review and meta-analysis has proposed that the prevalence of biofilms in chronic wounds is 78.2%, although a more conservative estimate suggests that the figure may be 60%.^{13,20} Biofilms in chronic wounds are difficult to visualise macroscopically, and whether located on the surface of a wound or deeper in the wound bed, the identification and diagnosis of biofilms in chronic wounds can be challenging.²¹ Although tissue biopsies are regarded as one of the most reliable methods to detect biofilms in wounds, the procedure can be invasive, painful, and expensive, and usually requires a specifically trained practitioner to be performed.²²⁻²⁴ Furthermore, the distribution of biofilms within chronic wounds is not thought to be uniform,²⁵ so a single biopsy sample taken from just one small area of a wound may not be enough to confirm the presence of biofilm within the wound as a whole. An algorithm to help identify (and treat) suspected biofilms has been developed, which is intended to guide the clinical management of chronic wounds.¹¹ In addition, recent consensus guidelines have been proposed to help clinicians recognise the signs and symptoms of biofilms in chronic, non-healing wounds, thereby optimising patient care.²¹ Nevertheless, there are currently no “gold standard” diagnostic tests to enable clinicians to confirm the presence of biofilms in chronic wounds.^{11,18,21,26}

The “wound infection continuum” conceptualises the relationship between bacteria, wound, and host,¹² and a key phase of this continuum is “critical colonisation”,

Key messages

- chronic wounds with biofilm, or “critically colonised wounds”, are slow to heal and pose a significant burden to the patient and the healthcare system
- biofilms in chronic wounds are very often tolerant and resistant to antibiotics and antiseptics, leading to treatment failure
- a time-dependent “window of opportunity” is thought to exist after wound debridement during which biofilms are increasingly susceptible to treatment, in particular topical antiseptics
- the topical antiseptic PVP-I demonstrates a broad spectrum of antimicrobial activity, potent antibiofilm efficacy, no acquired bacterial resistance or cross-resistance, low cytotoxicity, good tolerability, and an ability to promote wound healing, making it a promising therapeutic option in wound care and biofilm management
- a new algorithm has been proposed for the management of chronic, non-healing wounds due to critical colonisation and/or biofilm, using PVP-I

which marks the proliferation of microbes within the wound and the development of biofilm.⁸ Agreement is yet to be reached on how best to define the term critical colonisation, but Cutting (2003) suggested it refers to a wound that has become compromised by microbes, resulting in delayed healing without causing the classic signs and symptoms traditionally associated with clinical infection.²⁷ A critically colonised—or more generally biofilm infiltrated—wound has the potential to deteriorate to clinical infection once microbes reach a critical level (10^5 colony-forming units per gram tissue); should the numbers of bacteria increase above this level, the likelihood of infection increases as the host's immune system may no longer be able to control the proliferating microbes.^{8,28} In contrast, an infected wound marks the presence of multiplying microbes, which have already overwhelmed the host's immune system, leading to the traditional clinical signs of inflammation, including wound redness, swelling, warmth, pain, and potentially fever.²⁸⁻³⁰ Alternative terms do exist in the literature to describe a wound that is critically colonised, including “sub-clinically infected”, but at present there appears to be no consensus on how best to describe this stage in the wound infection continuum.^{8,12,29} White and Cutting (2008) argued that critical

colonisation is a cause of delayed healing and failure to acknowledge this important clinical step in the wound infection continuum could jeopardise the early diagnosis and treatment of a chronic wound.²⁹ Therefore, if a biofilm-infiltrated chronic wound can be treated before advancing beyond critical colonisation, this should not only improve the status of the wound but also help to avoid any significant personal and economic burden should the wound subsequently become infected.²⁹

To date, therapeutic intervention to eradicate biofilms in chronic wounds has relied principally on the use of conventional antibiotics and antiseptics.¹⁶ However, chronic wound biofilms can be highly tolerant and resistant to antibiotics and antiseptics.^{21,31} A wide range of molecular mechanisms are thought to contribute to the recalcitrance of biofilms towards antimicrobial agents, which may subsequently lead to treatment failure.³¹⁻³⁵ Debridement also represents a viable treatment strategy against biofilms, but it is unable to completely remove all biofilm and so is not recommended for use alone.²¹ Biofilms may reform quickly, even after repeated debridement, yet a time-dependent “window of opportunity” is thought to exist after debridement during which biofilms are more susceptible to treatment, in particular topical antiseptics.^{21,31,36} By delaying biofilm reformation after debridement, topical antiseptics may reduce the risk of infection and subsequent need for antibiotics, helping to minimise the potential for antibiotic resistance to develop.²¹ Indeed, recent consensus guidelines recommend topical antiseptics as first-line therapy in the treatment of stalled (chronic) wounds.²¹

Here, we review the antibiofilm efficacy, safety, and tolerability of topical antiseptics in chronic wound care and biofilm management, focusing specifically on povidone-iodine (PVP-I) in comparison with two commonly used antiseptics in wound care, polyhexamethylene biguanide/polyhexanide (PHMB), and silver. We also propose a new practical clinical guide or algorithm for the treatment of chronic, non-healing wounds due to critical colonisation or biofilm.

2 | METHODS

This narrative review results from a Focus Group meeting on “antiseptics in wound care and biofilm management” held in December 2019. The review is based primarily on the literature reviewed and recommended by the authors during the meeting. Additional English language publications of relevance were identified following literature searches conducted in PubMed in March 2020, using various combinations of the key terms: “antimicrobial”, “biofilm”, “critical colonisation”, “chronic

wound”, “cytotoxicity”, “non-healing wound”, “polyhexamethylene biguanide”, “polyhexanide”, “polihexanide”, “povidone iodine”, “silver”, “silver colloid”, “silver compounds”, “silver ion”, “silver nanoparticles”, “topical antiseptics”, “wound dressing”, and “wound healing”. The key terms in all searches could be combined using Boolean operators such as “OR” or “AND”. No date restrictions in the searches were employed. Only full-text articles identified from the searches that were considered directly relevant were included in this review, and most of these articles were open access. Reference lists of identified papers were also hand-searched to identify any further papers of interest. Reports and academic dissertations were not considered for inclusion in this review.

3 | RESULTS AND DISCUSSION

3.1 | Antimicrobial spectrum of activity

Differences observed in the antimicrobial spectrum of activity of PVP-I, PHMB, and silver may stem from the varying mechanisms of action of each antiseptic. An overview of the antimicrobial activity and mechanisms of action of PVP-I, PHMB, and silver is presented in Table 1.³⁷⁻⁷⁰

3.1.1 | Povidone-iodine

PVP-I is an iodophor or iodine-releasing agent, consisting of a complex of iodine and a neutral polymer base (polyvinylpyrrolidone), which acts as a reservoir of free active iodine.^{37,44,67} PVP-I has a particularly broad antimicrobial spectrum of activity that includes Gram-positive and Gram-negative bacteria (including strains resistant to antiseptics and antibiotics), fungi, protozoa, viruses, bacterial spores, and amoeba.³⁷⁻⁴³ Studies suggest that PVP-I exhibits a rapid onset of activity, with antimicrobial efficacy evident after a contact time of 1 minute, although there were certain study limitations to take into consideration.^{71,72} Some studies have even demonstrated potent antimicrobial efficacy of PVP-I in as little as 15 seconds against certain enveloped viruses (eg, Ebola, Middle East Respiratory Syndrome coronavirus [MERS-CoV], and Severe Acute Respiratory Syndrome coronavirus [SARS-CoV]).⁷³⁻⁷⁵ Of significance, a recent *in vitro* study has shown that PVP-I also provides rapid and potent activity against SARS-CoV-2, the virus responsible for the ongoing coronavirus disease 2019 (COVID-19) pandemic.⁷⁶ In the study, all topical and oral PVP-I products tested provided $\geq 99.99\%$ virucidal activity against SARS-CoV-2 within 30 seconds of contact, suggesting that PVP-I could

TABLE 1 Antimicrobial activity and mechanisms of action for povidone-iodine, polyhexanide, and silver

Characteristic	Povidone-iodine	Polyhexanide	Silver
Spectrum of activity	Broad spectrum of activity against Gram-positive and Gram-negative bacteria, fungi, protozoa, viruses, bacterial spores, and amoeba ³⁷⁻⁴³	Antimicrobial efficacy on Gram-negative bacteria, Gram-positive bacteria, and <i>Candida albicans</i> ⁵¹	Bactericidal activity against Gram-negative and Gram-positive bacteria, fungi, and viruses ⁵⁶⁻⁶²
Bacterial resistance	No reports of bacterial resistance ^{44,45}	No reports of bacterial resistance ⁵¹⁻⁵³	Gram-negative bacteria, including <i>E coli</i> , <i>P aeruginosa</i> , <i>K pneumoniae</i> , and <i>E cloacae</i> , may all express resistance ⁶³⁻⁶⁶
Cross-resistance	No reports of cross-resistance to antibiotics or other antiseptics ^{40,46,47}	Prolonged exposure of MRSA to a low concentration of PHMB in vitro associated with reduced susceptibility, not only to PHMB, but also to daptomycin ⁵⁴	No cross-resistance reported ⁴⁹
Mechanism of action	Iodine oxidises fatty acids, amino acids, and nucleic acids, leading to destabilisation of cell membranes, deactivation of cytosolic enzymes, and disruption of internal metabolic pathways ⁴⁸⁻⁵⁰	Disrupts cell membranes, increasing membrane permeability via interaction with acidic, negatively charged membrane phospholipids, and inhibits internal metabolic processes ^{54,55}	Ionic form reacts with the thiol groups in enzymes and proteins, adversely affecting enzyme function, cell replication, and energy generation in a non-specific manner ^{56,67-69} Silver ions also interfere with electron transport and/or membrane ion-exchange systems ^{69,70}

Abbreviations: MRSA, methicillin-resistant *Staphylococcus aureus*; PHMB, polyhexanide.

form a valuable part of future COVID-19 infection control strategies.⁷⁶ There have been no reports of bacterial resistance to iodine despite more than 150 years of use, possibly due to the multiple mechanisms of action exhibited by free iodine.^{44,45,48,77} In addition, no evidence of cross-resistance to antibiotics or other antiseptics has been observed with PVP-I use in a wide range of Gram-positive and Gram-negative bacterial species.^{40,46,47}

3.1.2 | Polyhexanide

PHMB is a biguanide, a strong base, which is highly positively charged at physiological pH.⁵¹ Findings from in vitro studies have demonstrated efficacy of PHMB on Gram-negative bacteria, Gram-positive bacteria, and *Candida albicans*,⁵¹ but it is without rapid onset of antimicrobial activity.⁷² There have been no reports of bacteria acquiring resistance to PHMB,⁵¹⁻⁵³ possibly a consequence of its non-specific mechanisms of action.⁵⁵ Although recent reviews suggest no evidence of cross-resistance to antibiotics in Gram-positive and Gram-negative bacteria after low-level exposure to PHMB,^{46,47} reduced susceptibility of methicillin-resistant *Staphylococcus aureus* (MRSA) to daptomycin and other cell wall-targeting antibiotics has been reported.⁵⁴

3.1.3 | Silver-containing products

Silver-containing products used in wound care (eg, silver salts, colloidal silver, and more recently, silver nanoparticles) require the release of positively charged silver ions in order to exhibit antimicrobial activity.^{68,78,79} The conversion process of inert metal to active ionised form is thought to be facilitated by interaction of the silver contained in wound dressings with aqueous media (eg, wound exudate).⁸⁰ The rate of onset of antimicrobial efficacy of certain higher silver release formulations has been demonstrated within 30 minutes of contact with clinically relevant bacteria.⁸¹ Silver has demonstrated bactericidal activity against Gram-negative and Gram-positive bacteria, and may also target fungi and viruses.⁵⁶⁻⁶² Bacterial resistance to silver has been documented.⁶³⁻⁶⁶ Evidence suggests silver resistance is encoded on plasmids.⁸² Plasmid-encoded resistance to silver is of particular concern in the clinical setting as plasmid transfer between bacteria within polymicrobial-infected chronic wounds may confer silver resistance across multiple bacterial species.⁸³ Furthermore, plasmid-mediated metallic salt resistance has been shown to be associated with coresistance to antibiotics.⁸⁴ No cross-resistance to antibiotics has been reported to date.⁴⁹

3.2 | Antibiofilm efficacy

3.2.1 | Povidone-iodine

The World Union of World Healing Societies 2016 Position Document recognises iodine as a suitable antimicrobial agent to manage biofilms,¹⁸ and numerous studies have been conducted to investigate the antibiofilm efficacy of PVP-I. Sub-inhibitory concentrations of PVP-I (0.17%, 0.35%, and 0.7% w/v) inhibited biofilm development by *Staphylococcus epidermidis* and *S aureus*, two of the most prevalent bacterial species found within chronic wounds.^{17,85} Using an in vitro model of chronic wound biofilms, Hill et al (2010) demonstrated that mixed *Pseudomonas* and *Staphylococcus* biofilms were disrupted by PVP-I 1% (w/v) solution; in contrast, these mixed biofilms were unaffected by treatment with ciprofloxacin and flucloxacillin.⁸⁶ Furthermore, in the same study, mature 7-day mixed biofilms were completely destroyed by PVP-I-containing dressing.⁸⁶ PVP-I at low doses (0.25% w/w) completely eradicated established biofilms of multi-drug resistant *S aureus*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa* and *C albicans* in vitro.⁸⁷ In a complex biofilm model, PVP-I 10% solution destroyed both early (90 minutes) and mature (48 hours) biofilms formed by *Candida auris*, an emerging multi-drug resistant yeast.^{88,89} The antibiofilm efficacy of PVP-I is rapid in onset in vitro, with total eradication of mature 3-day biofilms of *S aureus* and *P aeruginosa* achieved after only 15 minutes' exposure.⁹⁰ Using a basally perfused biofilm model, PVP-I 10% w/v demonstrated greater effectiveness against a biofilm community consisting of *P aeruginosa*, *Streptococcus pyogenes*, MRSA, and *Bacteriodes fragilis* compared with PHMB 0.5% v/v and silver acetate 0.05% w/v.⁹¹ In the study, PVP-I (57%) achieved the largest reduction in average bacterial count over time vs PHMB (44%) and silver acetate (27%).⁹¹ In a further study, iodine-containing wound dressings demonstrated greater antimicrobial efficacy against mature biofilms of *P aeruginosa* and *S aureus* over a 24-hour period compared with silver-based dressings using an in vitro static diffusion model.⁹²

3.2.2 | Polyhexanide

PHMB is the most used antiseptic to treat critically colonised and locally infected acute and chronic wounds.⁹³ Investigations into the antibiofilm efficacy of PHMB have demonstrated that wound irrigation by PHMB was effective against MRSA 3- and 6-day biofilms in a porcine wound model.⁹⁴ Application of a PHMB-containing biocellulose dressing to non-healing locally

infected and/or critically colonised wounds provided good efficacy against existing biofilm in 10 (63%) adult patients, thereby facilitating wound healing.⁹⁵ In addition, comparable efficacy of PHMB to chlorhexidine (CHG) was demonstrated against *P aeruginosa* biofilm grown in routinely used microtitre plates and on silicone materials in vitro in artificial wound fluid.⁹⁶ However, although PHMB was equally as effective as saline solution in reducing the bacterial load in venous leg ulcers, neither treatment was able to eliminate biofilm from the wound tissue.⁹⁷ Furthermore, conflicting evidence exists regarding the activity of PHMB against *S aureus* biofilms.^{98,99}

3.2.3 | Silver-containing products

Silver has been positioned among first-line options for the treatment of wound infections and is recommended for use in biofilm treatment.^{12,18} Silver sulfadiazine 5 to 10 µg/mL eradicated mature *P aeruginosa* biofilm in vitro,¹⁰⁰ and colloidal silver 100 and 150 µL almost completely eliminated *S aureus* biofilm in vitro 24 hours after treatment.¹⁰¹ Application of a silver-containing wound dressing to a 24-hour biofilm composed of either *P aeruginosa*, *Enterobacter cloacae*, *S aureus*, or a mixed bacterial community, resulted in total bacterial killing after 48 hours' exposure.¹⁰² Antibiofilm efficacy of silver ions against *S epidermidis* is thought to be mediated by the binding of silver ions to proteins and polysaccharides within the EPS, leading to breakdown of the EPS and destabilisation of the overall biofilm structure.¹⁰³ Silver nanoparticles are a relatively new addition to the range of available antimicrobial treatment options.¹⁰⁴ Interaction of silver nanoparticles with an aqueous environment (eg, an exuding wound) promotes oxidation of the nanoparticles and subsequent release of antimicrobial silver ions.¹⁰⁴ A concentration-dependent effect on *P aeruginosa* biofilm development and architecture has been demonstrated, with complete inhibition of biofilm growth achieved with a high concentration (18 µg/mL) of silver nanoparticles.¹⁰⁵ Potent antibiofilm efficacy of silver nanoparticles has been shown against *C auris* by inhibition of biofilm formation and by the disruption and distortion of pre-formed biofilms.¹⁰⁶ Silver nanoparticles have demonstrated potent inhibition of biofilm formation by *Escherichia coli*, *P aeruginosa*, and *Serratia proteamaculans*.¹⁰⁷ A 7-day treatment in vitro study by Kostenko et al (2010) concluded that the efficacy of silver against biofilms formed by MRSA, *P aeruginosa*, and *E coli* was determined by the type of silver species and base materials used in the wound dressings.¹⁰⁸

3.3 | Cytotoxicity and tolerability

A pre-requisite of any wound dressing is to shield the wound, offering not only a physical protective barrier but also a first line of defence against antimicrobial invasion of the wound. Intimate contact between the wound dressing and those cells that are key to the wound healing process is inevitable, so it is vital that a wound dressing demonstrates good cell compatibility.¹⁰⁹ Cytotoxic effects of a wound dressing would reduce the viability, proliferation, and migration of cells involved in the wound healing process, and so decrease the healing rate.¹⁰⁹ Studies have shown different levels of cytotoxicity among PVP-I, PHMB, and silver-containing products (Table 2).¹¹⁰⁻¹²⁹ Cytotoxicity data, in particular data derived from in vitro studies, must be interpreted with caution as any cytotoxic effects observed in cultured cell types can be magnified and may not be truly reflective of the in vivo or clinical setting.^{49,130,131}

3.3.1 | Povidone-iodine

A cytotoxicity assay conducted in cultured murine fibroblasts showed that PVP-I had the lowest cytotoxicity compared with PHMB, silver nitrate, and silver sulfadiazine.¹³² A further in vitro cytotoxicity test using murine fibroblasts demonstrated that PVP-I was less cytotoxic than a variety of antiseptics, including PHMB.¹³³ Furthermore, of the antiseptics tested, PVP-I was unique in provoking a revitalisation of fibroblasts, which may be pivotal to improved wound healing and better tissue tolerance with PVP-I.¹³³ In an in vitro study to investigate the cytotoxic effect of commonly used antiseptics on human fibroblasts and mesenchymal stromal cells, PVP-I was the only antiseptic to show remaining cell viability at the minimal bactericidal concentration (MBC; 1.32 g/L); in the same study, PHMB was 100% cytotoxic at the commercially available concentration of 0.04%, which was below the estimated MBC.¹³⁰ In vitro

TABLE 2 In vitro cytotoxicity of povidone-iodine, polyhexanide and silver-containing products against cell types instrumental to the wound healing process

Antiseptic	Fibroblasts	Keratinocytes	Endothelial cells
Povidone-iodine	Decrease in human fibroblast cell survival, cell migration, and cell viability ^{110,111}	Decrease in human keratinocyte cell viability ¹¹¹	Cytotoxic damage in bovine corneal endothelial cells ¹²⁴
Polyhexanide	Complete cell destruction of human skin fibroblasts ¹¹² Time-dependent cytotoxicity in human dermal fibroblasts ¹¹³ High cytotoxicity in murine fibroblasts after up to 72 hours of incubation ¹¹⁴	Concentration- and time-dependent cytotoxicity in human dermal keratinocytes ¹¹³ High cytotoxicity in human keratinocytes after up to 72 hours of incubation ¹¹⁴	Large reduction in human endothelial cell number and viability ¹²⁵
Silver-containing products	Cytotoxic effects of silver-based dressings, silver nitrate, and silver ions in fibroblasts ¹¹⁵⁻¹¹⁷ Significant change in cell morphology and decrease in cell proliferation and collagen synthesis of human diabetic fibroblasts (silver-containing dressings) ¹¹⁸ Inhibition of human dermal fibroblast proliferation with silver-dependent cell loss (silver nitrate) ¹¹⁹ Time- and concentration-dependent cytotoxicity in human fibroblasts (silver nitrate) ¹²⁰ Dose- and time-dependent cytotoxicity in human periodontal fibroblasts (silver nanoparticles) ¹²¹	Cytotoxic effects of silver-based dressings, silver nitrate, and silver ions in keratinocytes ¹¹⁵⁻¹¹⁷ Inhibition of human keratinocyte growth (nanocrystalline silver dressing) ¹²² Decrease in human epidermal keratinocyte viability, metabolism, and proliferatory and migratory potential (silver nanoparticles) ¹²³	Formation of reactive oxygen species and induction of apoptosis in human umbilical vein endothelial cells (silver nanoparticles) ^{126,127} Dose-dependent inhibition of the proliferation of rat vascular endothelial cells (silver nanoparticles) ¹²⁸ Increase in human umbilical vein endothelial cell membrane permeability (silver nanoparticles) ^{127,129}

evidence does exist to suggest that PVP-I may also be cytotoxic towards certain cell types involved in the wound healing process, including fibroblasts, keratinocytes, and endothelial cells.^{110,111,124} However, Bigliardi et al (2017) commented that PVP-I is well tolerated in human studies when used in appropriate concentrations, and that pronounced cytotoxicity with PVP-I has only been observed in certain *in vitro* studies.¹³⁴ Indeed, the densities of dendrocytes and microvessels in chronic leg ulcers were higher following 6 weeks of PVP-I treatment compared with silver sulfadiazine and CHG, with no evidence of dendrocytotoxicity, which may be considered a sign of *in vivo* cytotoxicity.¹³⁵ In contrast, silver sulfadiazine and CHG promoted changes in the superficial microvasculature and induced dendrocytotoxicity.¹³⁵ Overall, PVP-I has a good tolerability profile.⁴⁹ The prevalence of allergic contact dermatitis caused by PVP-I has been estimated to be approximately 0.4%, with reports of anaphylaxis exceptionally rare.¹³⁶

3.3.2 | Polyhexanide

There are reports to suggest that PHMB is cytotoxic towards cells that are crucial to the wound healing process. A range of concentrations of PHMB (0.005%–1.0% v/v) demonstrated high cytotoxicity against cultured human keratinocytes and murine fibroblasts after 24 and 72 hours of incubation *in vitro*.¹¹⁴ In an *in vitro* cytotoxicity test using cultured human skin fibroblasts, PHMB was extremely cytotoxic and appeared to induce complete cell destruction in the majority of cells.¹¹² When used *in vitro* at or below concentrations commonly employed in human wound care, PHMB (0.01%, 0.04%, and 0.1%) demonstrated both time- and concentration-dependent cytotoxicity in cultured human keratinocytes and osteoblasts, but only time-dependent cytotoxicity in cultured fibroblasts.¹¹³ Furthermore, exposure of cultured human endothelial cells to PHMB (0.0006%–0.01%) resulted in a large reduction in cell number and viability.¹²⁵ The overall tolerability profile of PHMB is good, with report of contact dermatitis found to be rare.¹³⁶ However, a recent case report identified PHMB as an emerging allergen, which may have induced an anaphylactic reaction.¹³⁷

3.3.3 | Silver-containing products

Studies have indicated cytotoxic effects of silver-based dressings and silver nitrate in both cultured fibroblasts and keratinocytes,¹¹⁵ with silver-based dressings also shown to cause a significant delay in re-epithelialisation.¹¹⁶ In addition, silver-based dressings

have been shown to significantly alter the cell morphology and decrease cell proliferation and collagen synthesis of cultured human diabetic fibroblasts *in vitro* compared with silver-free dressings, suggesting the use of such dressings should be closely monitored when treating diabetic wounds.¹¹⁸ Although the cytotoxic effects of silver nanoparticles are thought to be dependent on a number of factors, including nanoparticle size, shape, concentration, and aggregation, it remains to be determined if the cytotoxicity is due to inherent properties of the nanoparticles themselves or due to the release of ionic silver following oxidation.^{138–140} Indeed, ionic silver has been found to be significantly more cytotoxic *in vitro* in human dermal fibroblasts and epidermal keratinocytes compared with silver nanoparticles.¹¹⁷ Additional *in vitro* studies provide further evidence of the cytotoxicity of silver-based products on fibroblasts, keratinocytes, and endothelial cells.^{119–123,126–129} Nevertheless, silver has a good tolerability profile, and although argyria can result from the long-term use of silver-based products, any discoloration of the skin is not thought to result in pathological tissue damage or to be in any way a danger to life.¹⁴¹

3.4 | Wound management

In wound management, it is important to not only consider the antimicrobial efficacy and potential cytotoxicity of antiseptics, but also to be aware of the way in which antiseptics may impact the complex cellular and extracellular mechanisms involved in the wound healing process.¹⁴² One of the many properties an ideal antiseptic should possess is its ability to facilitate wound healing.⁴⁹

3.4.1 | Povidone-iodine

PVP-I enhanced wound healing via increased expression of transforming growth factor- β (TGF- β), neovascularisation, and re-epithelialisation in a rat acute skin wound model.¹⁴³ *In vitro* evidence suggests that PVP-I may facilitate wound healing by exerting an anti-inflammatory effect, scavenging superoxide anions, and inhibiting the production of reactive oxygen species by human polymorphonuclear neutrophils.¹⁴⁴ Indeed, treatment of venous leg ulcers with PVP-I in combination with hydrocolloid dressing reduced bacteria-related inflammation, vasculitis, and phagocytic infiltration of the ulcers compared with hydrocolloid dressing alone, resulting in an improved ulcer healing rate.¹⁴⁵ In a further study, the healing rate of chronic leg ulcers was significantly increased by PVP-I vs controls, reducing the

time to healing by 2 to 9 weeks.¹³⁵ Recently, a phase IV prospective study conducted in 106 adult patients showed that PVP-I foam dressing achieved a shorter epithelialisation time compared with hydrocellular foam dressing and conventional petrolatum gauze when used as a split-thickness skin graft donor site dressing.¹⁴⁶ Further evidence supports the role of PVP-I in wound healing when investigated within in vivo human studies.^{147,148} Compared with silver-based foam dressings or control gauze, PVP-I 3% foam dressing was the most effective dressing in wound healing by promoting neovascularisation, re-epithelialisation, and collagen deposition in an in vivo rat wound model.¹⁴⁹ Similarly, PVP-I 10% solution promoted rapid neovascularisation more effectively than silver nitrate solution in an in vivo mouse wound model.¹⁵⁰

3.4.2 | Polyhexanide

Evidence to date suggests that PHMB may also be beneficial for wound healing. In a recent systematic review, it was concluded that PHMB may promote the healing of chronic wounds, reduce the bacterial load, eradicate MRSA, and lessen wound-related pain.¹⁵¹ A preclinical study using a mouse wound model demonstrated that PHMB had more beneficial effects on the microcirculation, angiogenesis, and epithelialisation compared with chitosan.¹⁵² In a further study, PHMB had a more positive effect upon the blood flow of intact human skin in vivo than octenidine, suggesting value in the treatment of critically perfused wounds, such as burns.¹⁴² Comparison of a PHMB-containing dressing with a silver-based dressing in patients with critically colonised and locally infected wounds demonstrated that the PHMB-containing dressing was significantly faster and more effective at removing the critical bacterial load over a 28-day period.¹⁵³ In addition, PHMB has been shown to protect keratinocytes from bacterial damage by *S aureus* and re-establish normal cell proliferation in vitro in a dose-dependent manner.¹⁵⁴ Nevertheless, a recent in vitro study has suggested that PHMB may exert pro-inflammatory effects, including increased cytokine secretion and nuclear factor kappa B activation,¹⁵⁵ both of which would hinder the wound healing process.

3.4.3 | Silver-containing products

A Cochrane meta-analysis concluded that there is insufficient evidence to confirm whether silver-containing products can promote wound healing.¹⁵⁶ Furthermore, the heterogeneous nature of the evidence regarding the

effectiveness of silver-based treatments in wound care is thought to have hindered the development of treatment guidelines.^{157,158} As such, authors of a recent qualitative literature analysis proposed that silver-containing wound dressings should be chosen with care if the wound healing process is not to be impeded.¹⁵⁷ Nevertheless, there is evidence to suggest that silver can have beneficial effects on the healing of chronic wounds, including those wounds showing signs of critical colonisation.^{159,160} For example, Duan et al demonstrated that a sub-cytotoxic concentration of silver ions may promote the proliferation of human skin keratinocytes in vitro.¹⁶¹ Additionally, silver nanoparticles decreased the generation of pro-inflammatory cytokines by human keratinocytes and fibroblasts in an in vitro wound healing model.¹⁶²

3.5 | Algorithm for the treatment of chronic, non-healing wounds due to critical colonisation or biofilm

Of the three antiseptics discussed in this review, PVP-I has particular characteristics that are ideal for the treatment of chronic, non-healing wounds due to critical colonisation and/or biofilm, namely its potent antibiofilm efficacy, broad spectrum of antimicrobial activity, wound healing properties, and rapidity of action. Therefore, we have proposed a new practical clinical guide or algorithm to remove biofilm and manage critically colonised wounds, using PVP-I (Figure 1).

When biofilm presence within a chronic wound is strongly suspected, clinicians should adopt an early intervention plan to remove the biofilm as soon as possible and reduce the risk of infection.^{21,23} Wound bed preparation is vital if successful wound healing is to occur, as described in the TIMERS (Tissue, Inflammation/infection, Moisture imbalance, Epithelial edge advancement, Repair/regeneration, and Social factors) framework to guide wound care (Figure 2).^{23,163} According to the new algorithm proposed here, intensive mechanical washing or cleansing of the wound with either soap or PVP-I scrub should help to prepare the wound bed by removing debris and biofilm from the wound. This should preferably be performed without causing any additional trauma to the wound.²⁶ Ideally, cleansing of the wound should be performed with each change in dressing.¹² This is particularly the case if the likelihood of biofilm is high, but the unique characteristics of each particular wound bed should determine the frequency of dressing changes. Following wound cleansing, debridement of the wound can help to disrupt any remaining biofilm, remove necrotic tissue, and stimulate wound healing.^{12,30,163} Debridement may be achieved using a number of different methods,




FIGURE 1 A proposed new algorithm for the treatment of chronic, non-healing wounds due to critical colonisation and/or biofilm.

*Secondary dressing is to be used to keep the primary dressing securely in place




FIGURE 2 TIMERS framework for the management of chronic, non-healing wounds. (Data reproduced with permission of MA Healthcare Limited, from Atkin et al, 2019; permission conveyed through Copyright Clearance Center, Inc.).²³ NPWT, negative pressure wound therapy

including surgical, mechanical, and chemical techniques.¹² The wound can then be disinfected using gauze impregnated with PVP-I dermic solution. Given the rapid onset of action of PVP-I, employing a contact time of at least 1 minute may be sufficient to eradicate the majority of any microbes remaining in the wound.^{71,72}

Biofilm is not completely removed by debridement and it may quickly regrow within 24 hours, so debridement alone is not an appropriate treatment strategy.^{21,36} Regrowth of biofilm must be controlled according to the status of the wound, in particular the amount of exudate the wound is producing. A balanced, moist wound


environment is seen as critical for wound healing.²³ Not enough exudate or excessive production of exudate will hinder the wound healing process.²³ Dry wounds are devoid of exudate, which hinders the activity of tissue-repairing cells (Figure 3).¹⁶³ With low and moderate exuding wounds, the wound bed and surrounding skin become increasingly wet. The excessive amount of exudate produced by a highly exuding wound may result in maceration of the surrounding skin.³⁰ Therefore, it is important to manage moisture levels by selecting the correct dressing.²³ Numerous types of dressing are available to both protect the wound and promote healing (Table 3).^{1,164} Choosing the most appropriate option from the extensive range of dressings available can be a difficult treatment decision, but it should ultimately be tailored to the characteristics of each wound and to the

patient.^{23,30} Dressings that manage the exudate and encourage a balanced wound environment are crucial for improved patient outcomes.¹⁶⁵ The new algorithm proposes that low to moderate exuding wounds can be treated with PVP-I gel and PVP-I tulle and covered with a secondary dressing. Highly exuding wounds may be treated with PVP-I gel applied beneath an absorbent dressing. Until signs of improvement in the wound bed surface are evident, dressings should be changed daily and regularly checked for discoloration, as any change in colour can indicate reapplication of PVP-I is needed in order to maintain its clinical efficacy.¹⁶⁶ Regular monitoring of the healing status of the wound is required according to specific criteria, including assessment of the size and depth of the wound, and the amount and type of exudate.³⁰ If there is an improvement in wound healing,



FIGURE 3 Different types of exuding wound. Images were kindly provided by the authors P.J.A. and S.M., with permission

TABLE 3 Different types of wound dressing currently available for wound management and their potential clinical uses

Level of wound exudate	Type of dressing	Properties	Clinical use
Low  High	Hydrogels	Rehydrates dry wounds, easily removed/changed, may cause overhydration, removes necrotic tissue	Dry wounds
	Films	Occlusive, retains moisture, only for non-exudative wounds	Superficial wounds with very limited exudate
	Gauze	Inexpensive, drying, may cause further injury upon changing	Clean and dry wounds with low exudate levels
	Hydrocolloids	Long-lasting, keep moist environment, not for use on wounds with high exudate levels due to impermeable nature, occlusive, not for infected wounds	Low-to-moderate drainage wounds
	Foams	Moderately absorbent, insulating, for use on moderately exuding wounds, minimal trauma during dressing changes	Low-to-moderate drainage wounds
	Gelling fibres	Highly absorbent	Moderate-to-highly exuding wounds
	Polyacrylate polymers	Highly absorbent	Highly exuding wounds
	Alginates	Highly absorbent, haemostatic	Infected and non-infected wounds with large amount of exudate

Note: Data obtained from Han and Ceilley, and Shi et al.^{1,164}

treatment may revert to the standard of care (Figure 2).²³ Reassessment of the wound should be made on a weekly basis and, if necessary, the procedure outlined in the algorithm can be restarted.

4 | CONCLUSIONS

Recent evidence suggests that the majority of chronic wounds have biofilms which can hinder wound healing and result in ineffective treatment, burdening both the patient and the healthcare system. PVP-I demonstrates potent efficacy against biofilms formed by a variety of microbes found to be prevalent within chronic wounds, including *S aureus*, *S epidermidis*, and *P aeruginosa*. Given how diverse the microbial community can be within chronic wounds, the broader spectrum of antimicrobial activity of PVP-I should be advantageous vs the more limited spectrum of antimicrobial activity shown by PHMB and silver-containing products. PVP-I also fulfils all of the other requirements of an ideal antiseptic for chronic wound care, including a lack of acquired bacterial resistance or cross-resistance, wound healing properties, low cytotoxicity, and good tolerability. Collectively, these characteristics of PVP-I suggest that it represents a highly viable therapeutic option in wound care and biofilm management, with the potential to be particularly

effective during the critically colonised, biofilm-infiltrated stage of the wound infection continuum. The proposed new algorithm utilising PVP-I should help to guide clinicians in the treatment of patients with chronic, non-healing wounds, which prove particularly unresponsive to treatment.

ACKNOWLEDGEMENTS

Medical writing assistance in the preparation of this manuscript was provided by Chris Cammack (CircleScience, an Ashfield Company, part of UDG Healthcare plc) and funded by Meda Pharma S.p.A., a Viatrix company. All authors were speakers at the Focus Group meeting, funded by Viatrix.

CONFLICT OF INTEREST

P.J.A. has served as a consultant for Viatrix. R.T.B. has served as a consultant for Viatrix. B.M.B. has served as a consultant for Viatrix. L.G.G. has served as a consultant for Viatrix. S.M. has no conflicts of interest. S.J.M. has received honoraria as a speaker and has been a member of advisory boards for Viatrix.

AUTHOR CONTRIBUTIONS

All authors contributed to developing the algorithm, analysis and interpretation of information presented, drafting and revising the article, gave final approval of

the version to be published, and agree to be held accountable for all aspects of the work.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article

ORCID

Paulo J. Alves  <https://orcid.org/0000-0002-6348-3316>

REFERENCES

- Han G, Ceilley R. Chronic wound healing: a review of current management and treatments. *Adv Ther*. 2017;34(3):599-610.
- Sorg H, Tilkorn DJ, Hager S, Hauser J, Mirastschijski U. Skin wound healing: an update on the current knowledge and concepts. *Eur Surg Res*. 2017;58(1-2):81-94.
- Barrientos S, Stojadinovic O, Golinko MS, Brem H, Tomic-Canic M. Growth factors and cytokines in wound healing. *Wound Repair Regen*. 2008;16(5):585-601.
- Frykberg RG, Banks J. Challenges in the treatment of chronic wounds. *Adv Wound Care (New Rochelle)*. 2015;4(9):560-582.
- Nunan R, Harding KG, Martin P. Clinical challenges of chronic wounds: searching for an optimal animal model to recapitulate their complexity. *Dis Model Mech*. 2014;7(11):1205-1213.
- Olsson M, Jarbrink K, Divakar U, et al. The humanistic and economic burden of chronic wounds: a systematic review. *Wound Repair Regen*. 2019;27(1):114-125.
- Guo S, Dipietro LA. Factors affecting wound healing. *J Dent Res*. 2010;89(3):219-229.
- Percival SL. Importance of biofilm formation in surgical infection. *Br J Surg*. 2017;104(2):e85-e94.
- Beitz JM. Pharmacologic impact (aka "breaking bad") of medications on wound healing and wound development: a literature-based overview. *Ostomy Wound Manage*. 2017;63(3):18-35.
- Jacobson LK, Johnson MB, Dedhia RD, Niknam-Bienia S, Wong AK. Impaired wound healing after radiation therapy: a systematic review of pathogenesis and treatment. *JPRAS Open*. 2017;13:92-105.
- Percival SL, Vuotto C, Donelli G, Lipsky BA. Biofilms and wounds: an identification algorithm and potential treatment options. *Adv Wound Care (New Rochelle)*. 2015;4(7):389-397.
- International Wound Infection Institute. Wound infection in clinical practice. *Wounds International*. London: IWII; 2016.
- Malone M, Bjarnsholt T, McBain AJ, et al. The prevalence of biofilms in chronic wounds: a systematic review and meta-analysis of published data. *J Wound Care*. 2017;26(1):20-25.
- Ganesh K, Sinha M, Mathew-Steiner SS, Das A, Roy S, Sen CK. Chronic wound biofilm model. *Adv Wound Care (New Rochelle)*. 2015;4(7):382-388.
- Wolcott RD, Hanson JD, Rees EJ, et al. Analysis of the chronic wound microbiota of 2,963 patients by 16S rDNA pyrosequencing. *Wound Repair Regen*. 2016;24(1):163-174.
- Kadam S, Shai S, Shahane A, Kaushik KS. Recent advances in non-conventional antimicrobial approaches for chronic wound biofilms: have we found the 'Chink in the armor'? *Bio-medicine*. 2019;7(2):35.
- Thomson CH. Biofilms: do they affect wound healing? *Int Wound J*. 2011;8(1):63-67.
- World Union of Wound Healing Societies. Florence congress, position document. Management of biofilm. *Wounds International*. London: WUWHS; 2016.
- Khatoon Z, McTiernan CD, Suuronen EJ, Mah TF, Alarcon EI. Bacterial biofilm formation on implantable devices and approaches to its treatment and prevention. *Heliyon*. 2018;4(12):e01067.
- James GA, Swogger E, Wolcott R, et al. Biofilms in chronic wounds. *Wound Repair Regen*. 2008;16(1):37-44.
- Schultz G, Bjarnsholt T, James GA, et al. Consensus guidelines for the identification and treatment of biofilms in chronic nonhealing wounds. *Wound Repair Regen*. 2017;25(5):744-757.
- Smith ME, Robinowitz N, Chaulk P, Johnson K. Comparison of chronic wound culture techniques: swab versus curetted tissue for microbial recovery. *Br J Community Nurs*. 2014;19(Suppl(9 0)):S22-S26.
- Atkin L, Bucko Z, Conde Montero E, et al. Implementing TIMERS: the race against hard-to-heal wounds. *J Wound Care*. 2019;28(3):S1-S49.
- Høiby N, Bjarnsholt T, Moser C, et al. ESCMID guideline for the diagnosis and treatment of biofilm infections 2014. *Clin Microbiol Infect*. 2015;21(Suppl 1):S1-S25.
- Fazli M, Bjarnsholt T, Kirketerp-Møller K, et al. Nonrandom distribution of *Pseudomonas aeruginosa* and *Staphylococcus aureus* in chronic wounds. *J Clin Microbiol*. 2009;47(12):4084-4089.
- Swanson T, Keast D, Cooper R, et al. Ten top tips: identification of wound infection in a chronic wound. *Wounds Int*. 2015;6(2):22-27.
- Cutting KF. Wound healing, bacteria and topical therapies. *EWMA*. 2003;3:17-19.
- Kingsley A. A proactive approach to wound infection. *Nurs Stand*. 2001;15(30):50-54. 56, 58.
- White R, Cutting K. Critical colonisation of chronic wounds: microbial mechanisms. *Wounds UK*. 2008;4:70-78.
- Gupta S, Andersen C, Black J, et al. Management of chronic wounds: diagnosis, preparation, treatment, and follow-up. *Wounds*. 2017;29(9):S19-S36.
- Omar A, Wright JB, Schultz G, Burrell R, Nadworny P. Microbial biofilms and chronic wounds. *Microorganisms*. 2017;5(1):9.
- Stewart PS, Costerton JW. Antibiotic resistance of bacteria in biofilms. *Lancet*. 2001;358(9276):135-138.
- Hall CW, Mah TF. Molecular mechanisms of biofilm-based antibiotic resistance and tolerance in pathogenic bacteria. *FEMS Microbiol Rev*. 2017;41(3):276-301.
- Lebeaux D, Ghigo JM, Beloin C. Biofilm-related infections: bridging the gap between clinical management and fundamental aspects of recalcitrance toward antibiotics. *Microbiol Mol Biol Rev*. 2014;78(3):510-543.
- Percival SL, Hill KE, Malic S, Thomas DW, Williams DW. Antimicrobial tolerance and the significance of persister cells in recalcitrant chronic wound biofilms. *Wound Repair Regen*. 2011;19(1):1-9.
- Wolcott RD, Rumbaugh KP, James G, et al. Biofilm maturity studies indicate sharp debridement opens a time-dependent therapeutic window. *J Wound Care*. 2010;19(8):320-328.

37. Lachapelle J-M, Castel O, Casado AF, et al. Antiseptics in the era of bacterial resistance: a focus on povidone iodine. *Clin Pract.* 2013;10(5):579-592.
38. Kawana R, Kitamura T, Nakagomi O, et al. Inactivation of human viruses by povidone-iodine in comparison with other antiseptics. *Dermatology.* 1997;195(Suppl 2):29-35.
39. Wutzler P, Sauerbrei A, Klocking R, Brogmann B, Reimer K. Virucidal activity and cytotoxicity of the liposomal formulation of povidone-iodine. *Antiviral Res.* 2002;54(2):89-97.
40. Kunisada T, Yamada K, Oda S, Hara O. Investigation on the efficacy of povidone-iodine against antiseptic-resistant species. *Dermatology.* 1997;195(Suppl 2):14-18.
41. Fleischer W, Reimer K. Povidone-iodine in antiseptics—state of the art. *Dermatology.* 1997;195(Suppl 2):3-9.
42. Lacey RW, Catto A. Action of povidone-iodine against methicillin-sensitive and -resistant cultures of *Staphylococcus aureus*. *Postgrad Med J.* 1993;69(Suppl 3):S78-S83.
43. Gatti S, Cevini C, Bruno A, Penso G, Rama P, Scaglia M. In vitro effectiveness of povidone-iodine on *Acanthamoeba* isolates from human cornea. *Antimicrob Agents Chemother.* 1998;42(9):2232-2234.
44. Durani P, Leaper D. Povidone-iodine: use in hand disinfection, skin preparation and antiseptic irrigation. *Int Wound J.* 2008;5(3):376-387.
45. Cooper RA. Iodine revisited. *Int Wound J.* 2007;4(2):124-137.
46. Kampf G. Biocidal agents used for disinfection can enhance antibiotic resistance in gram-negative species. *Antibiotics (Basel).* 2018;7(4):110.
47. Kampf G. Antibiotic resistance can be enhanced in gram-positive species by some biocidal agents used for disinfection. *Antibiotics (Basel).* 2019;8(1):13.
48. Eggers M. Infectious disease management and control with povidone iodine. *Infect Dis Ther.* 2019;8(4):581-593.
49. Bigliardi PL, Alsagoff SAL, El-Kafrawi HY, Pyon JK, Wa CTC, Villa MA. Povidone iodine in wound healing: a review of current concepts and practices. *Int J Surg.* 2017;44:260-268.
50. Kanagalingam J, Feliciano R, Hah JH, Labib H, Le TA, Lin JC. Practical use of povidone-iodine antiseptic in the maintenance of oral health and in the prevention and treatment of common oropharyngeal infections. *Int J Clin Pract.* 2015;69(11):1247-1256.
51. Fjeld H, Lingaas E. Polyhexanide - safety and efficacy as an antiseptic. *Tidsskr nor Laegeforen.* 2016;136(8):707-711.
52. Moore K, Gray D. Using PHMB antimicrobial to prevent wound infection. *Wounds UK.* 2007;3:96-102.
53. Gilliver S. PHMB: a well-tolerated antiseptic with no reported toxic effects. *J Wound Care.* 2009;S9-S14.
54. Renzoni A, Von Dach E, Landelle C, et al. Impact of exposure of methicillin-resistant *Staphylococcus aureus* to polyhexanide in vitro and in vivo. *Antimicrob Agents Chemother.* 2017;61(10):e00272-e00217.
55. Hubner NO, Kramer A. Review on the efficacy, safety and clinical applications of polyhexanide, a modern wound antiseptic. *Skin Pharmacol Physiol.* 2010;23:17-27.
56. Jung WK, Koo HC, Kim KW, Shin S, Kim SH, Park YH. Antibacterial activity and mechanism of action of the silver ion in *Staphylococcus aureus* and *Escherichia coli*. *Appl Environ Microbiol.* 2008;74(7):2171-2178.
57. Vila Dominguez A, Ayerbe Algaba R, Miro Canturri A, Rodriguez Villodres A, Smani Y. Antibacterial activity of colloidal silver against gram-negative and gram-positive bacteria. *Antibiotics (Basel).* 2020;9(1):36.
58. Kang J, Dietz MJ, Hughes K, Xing M, Li B. Silver nanoparticles present high intracellular and extracellular killing against *Staphylococcus aureus*. *J Antimicrob Chemother.* 2019;74(6):1578-1585.
59. Liao S, Zhang Y, Pan X, et al. Antibacterial activity and mechanism of silver nanoparticles against multidrug-resistant *Pseudomonas aeruginosa*. *Int J Nanomedicine.* 2019;14:1469-1487.
60. Lu L, Sun RW, Chen R, et al. Silver nanoparticles inhibit hepatitis B virus replication. *Antivir Ther.* 2008;13(2):253-262.
61. Mallmann EJ, Cunha FA, Castro BN, Maciel AM, Menezes EA, Fechine PB. Antifungal activity of silver nanoparticles obtained by green synthesis. *Rev Inst Med Trop Sao Paulo.* 2015;57(2):165-167.
62. Xiang DX, Chen Q, Pang L, Zheng CL. Inhibitory effects of silver nanoparticles on H1N1 influenza A virus in vitro. *J Virol Methods.* 2011;178(1-2):137-142.
63. Randall CP, Gupta A, Jackson N, Busse D, O'Neill AJ. Silver resistance in Gram-negative bacteria: a dissection of endogenous and exogenous mechanisms. *J Antimicrob Chemother.* 2015;70(4):1037-1046.
64. Finley PJ, Norton R, Austin C, Mitchell A, Zank S, Durham P. Unprecedented silver resistance in clinically isolated Enterobacteriaceae: major implications for burn and wound management. *Antimicrob Agents Chemother.* 2015;59(8):4734-4741.
65. Panacek A, Kvitek L, Smekalova M, et al. Bacterial resistance to silver nanoparticles and how to overcome it. *Nat Nanotechnol.* 2018;13(1):65-71.
66. Sutterlin S, Tano E, Bergsten A, Tallberg AB, Melhus A. Effects of silver-based wound dressings on the bacterial flora in chronic leg ulcers and its susceptibility in vitro to silver. *Acta Derm Venereol.* 2012;92(1):34-39.
67. McDonnell G, Russell AD. Antiseptics and disinfectants: activity, action, and resistance. *Clin Microbiol Rev.* 1999;12(1):147-179. [published correction appears in Clin Microbiol Rev 2001 Jan;14(1):227].
68. Kramer A, Dissemmond J, Kim S, et al. Consensus on wound antiseptics: update 2018. *Skin Pharmacol Physiol.* 2018;31(1):28-58.
69. Muller M. Bacterial silver resistance gained by cooperative interspecies redox behavior. *Antimicrob Agents Chemother.* 2018;62(8):e00672-e00618.
70. Ketikidis I, Banti CN, Kourkoumelis N, et al. Conjugation of penicillin-G with silver(I) ions expands its antimicrobial activity against Gram negative bacteria. *Antibiotics (Basel).* 2020;9(1):25.
71. Deshmukh N, Kramer JW, Kjellberg SI. A comparison of 5-minute povidone-iodine scrub and 1-minute povidone-iodine scrub followed by alcohol foam. *Mil Med.* 1998;163(3):145-147.
72. Koburger T, Hubner NO, Braun M, Siebert J, Kramer A. Standardized comparison of antiseptic efficacy of triclosan, PVP-iodine, octenidine dihydrochloride, polyhexanide and chlorhexidine digluconate. *J Antimicrob Chemother.* 2010;65(8):1712-1719.

73. Eggers M, Eickmann M, Kowalski K, Zorn J, Reimer K. Povidone-iodine hand wash and hand rub products demonstrated excellent in vitro virucidal efficacy against Ebola virus and modified vaccinia virus Ankara, the new European test virus for enveloped viruses. *BMC Infect Dis*. 2015;15:375.
74. Eggers M, Eickmann M, Zorn J. Rapid and effective virucidal activity of povidone-iodine products against Middle East respiratory syndrome coronavirus (MERS-CoV) and modified vaccinia virus Ankara (MVA). *Infect Dis Ther*. 2015;4(4):491-501.
75. Eggers M, Koburger-Janssen T, Eickmann M, Zorn J. In vitro bactericidal and virucidal efficacy of povidone-iodine gargle/mouthwash against respiratory and oral tract pathogens. *Infect Dis Ther*. 2018;7(2):249-259.
76. Anderson DE, Sivalingam V, Kang AEZ, et al. Povidone-iodine demonstrates rapid in-vitro virucidal activity against SARS-CoV-2, the virus causing COVID-19 disease. *Infect Dis Ther*. 2020;9(3):669-675. <https://doi.org/10.1007/s40121-020-00316-3>.
77. Sibbald RG, Leaper D, Queen D. Iodine made easy. *Wounds Int*. 2011;2:2.
78. Pavlik V, Sojka M, Mazurova M, Velebny V. Dual role of iodine, silver, chlorhexidine and octenidine as antimicrobial and antiprotease agents. *PLoS One*. 2019;14(1):e0211055.
79. Percival SL, Finnegan S, Donelli G, Vuotto C, Rimmer S, Lipsky BA. Antiseptics for treating infected wounds: efficacy on biofilms and effect of pH. *Crit Rev Microbiol*. 2016;42(2):293-309.
80. Melaiye A, Youngs WJ. Silver and its application as an antimicrobial agent. *Expert Opin Ther Patents*. 2005;15(2):125-130.
81. Yin HQ, Langford R, Burrell RE. Comparative evaluation of the antimicrobial activity of ACTICOAT antimicrobial barrier dressing. *J Burn Care Rehabil*. 1999;20(3):195-200.
82. Davis IJ, Richards H, Mullany P. Isolation of silver- and antibiotic-resistant *Enterobacter cloacae* from teeth. *Oral Microbiol Immunol*. 2005;20(3):191-194.
83. Hosny AEM, Rasmy SA, Aboul-Magd DS, Kashef MT, El-Bazza ZE. The increasing threat of silver-resistance in clinical isolates from wounds and burns. *Infect Drug Resist*. 2019;12:1985-2001.
84. McHugh GL, Moellering RC, Hopkins CC, Swartz MN. *Salmonella typhimurium* resistant to silver nitrate, chloramphenicol, and ampicillin. *Lancet*. 1975;1(7901):235-240.
85. Oduwale KO, Glynn AA, Molony DC, et al. Anti-biofilm activity of sub-inhibitory povidone-iodine concentrations against *Staphylococcus epidermidis* and *Staphylococcus aureus*. *J Orthop Res*. 2010;28(9):1252-1256.
86. Hill KE, Malic S, McKee R, et al. An in vitro model of chronic wound biofilms to test wound dressings and assess antimicrobial susceptibilities. *J Antimicrob Chemother*. 2010;65(6):1195-1206.
87. Capriotti K, Pelletier J, Barone S, Capriotti J. Efficacy of dilute povidone-iodine against multi-drug resistant bacterial biofilms, fungal biofilms and fungal spores. *Clin Res Dermatol Open Access*. 2018;5(1):1-5.
88. Kean R, McCloud E, Townsend EM, et al. The comparative efficacy of antiseptics against *Candida auris* biofilms. *Int J Antimicrob Agents*. 2018;52(5):673-677.
89. Forsberg K, Woodworth K, Walters M, et al. *Candida auris*: the recent emergence of a multidrug-resistant fungal pathogen. *Med Mycol*. 2019;57(1):1-12.
90. Johani K, Malone M, Jensen SO, et al. Evaluation of short exposure times of antimicrobial wound solutions against microbial biofilms: from in vitro to in vivo. *J Antimicrob Chemother*. 2018;73(2):494-502.
91. Oates A, Lindsay S, Mistry H, Ortega F, McBain AJ. Modelling antiseptics using defined populations of facultative and anaerobic wound pathogens grown in a basally perfused biofilm model. *Biofouling*. 2018;34(5):507-518.
92. Thorn RM, Austin AJ, Greenman J, Wilkins JP, Davis PJ. In vitro comparison of antimicrobial activity of iodine and silver dressings against biofilms. *J Wound Care*. 2009;18(8):343-346.
93. Dissemmond J, Gerber V, Kramer A, et al. A practice-oriented recommendation for treatment of critically colonised and locally infected wounds using polihexanide. *J Tissue Viability*. 2010;19(3):106-115.
94. Davis SC, Harding A, Gil J, et al. Effectiveness of a polyhexanide irrigation solution on methicillin-resistant *Staphylococcus aureus* biofilms in a porcine wound model. *Int Wound J*. 2017;14(6):937-944.
95. Lenselink E, Andriessen A. A cohort study on the efficacy of a polyhexanide-containing biocellulose dressing in the treatment of biofilms in wounds. *J Wound Care*. 2011;20(11):534, 6-9-539.
96. Hubner NO, Matthes R, Koban I, Randler C, Muller G, Bender C, et al. Efficacy of chlorhexidine, polihexanide and tissue-tolerable plasma against *Pseudomonas aeruginosa* biofilms grown on polystyrene and silicone materials. *Skin Pharmacol Physiol*. 2010;23(Suppl 1):28-34.
97. Borges EL, Frison SS, Honorato-Sampaio K, et al. Effect of polyhexamethylene biguanide solution on bacterial load and biofilm in venous leg ulcers: a randomized controlled trial. *J Wound Ostomy Continence Nurs*. 2018;45(5):425-431.
98. Kamaruzzaman NF, Chong SQY, Edmondson-Brown KM, Ntow-Boahene W, Bardiau M, Good L. Bactericidal and anti-biofilm effects of polyhexamethylene biguanide in models of intracellular and biofilm of *Staphylococcus aureus* isolated from bovine mastitis. *Front Microbiol*. 2017;8:1518.
99. Ueda S, Kuwabara Y. Susceptibility of biofilm *Escherichia coli*, *Salmonella Enteritidis* and *Staphylococcus aureus* to detergents and sanitizers. *Biocontrol Sci*. 2007;12(4):149-153.
100. Bjarnsholt T, Kirketerp-Moller K, Kristiansen S, et al. Silver against *Pseudomonas aeruginosa* biofilms. *Apmis*. 2007;115(8):921-928.
101. Goggin R, Jardeleza C, Wormald PJ, Vreugde S. Colloidal silver: a novel treatment for *Staphylococcus aureus* biofilms? *Int Forum Allergy Rhinol*. 2014;4(3):171-175.
102. Percival SL, Bowler P, Woods EJ. Assessing the effect of an antimicrobial wound dressing on biofilms. *Wound Repair Regen*. 2008;16(1):52-57.
103. Chaw KC, Manimaran M, Tay FE. Role of silver ions in destabilization of intermolecular adhesion forces measured by atomic force microscopy in *Staphylococcus epidermidis* biofilms. *Antimicrob Agents Chemother*. 2005;49(12):4853-4859.
104. Lee SH, Jun BH. Silver nanoparticles: synthesis and application for nanomedicine. *Int J Mol Sci*. 2019;20(4):865.

105. Guo J, Qin S, Wei Y, et al. Silver nanoparticles exert concentration-dependent influences on biofilm development and architecture. *Cell Prolif*. 2019;52(4):e12616.
106. Lara HH, Ixtepan-Turrent L, Jose Yacamán M, Lopez-Ribot J. Inhibition of *Candida auris* biofilm formation on medical and environmental surfaces by silver nanoparticles. *ACS Appl Mater Interfaces*. 2020;12(19):21183-21191. <https://doi.org/10.1021/acsami.9b20708>.
107. Radzig MA, Nadtochenko VA, Koksharova OA, Kiwi J, Lipasova VA, Khmel IA. Antibacterial effects of silver nanoparticles on gram-negative bacteria: influence on the growth and biofilms formation, mechanisms of action. *Colloids Surf B Biointerfaces*. 2013;102:300-306.
108. Kostenko V, Lyczak J, Turner K, Martinuzzi RJ. Impact of silver-containing wound dressings on bacterial biofilm viability and susceptibility to antibiotics during prolonged treatment. *Antimicrob Agents Chemother*. 2010;54(12):5120-5131.
109. Wiegand C, Abel M, Hipler UC, Elsner P. Effect of non-adhering dressings on promotion of fibroblast proliferation and wound healing in vitro. *Sci Rep*. 2019;9(1):4320.
110. Liu JX, Werner JA, Buza JIII, Kirsch T, Zuckerman JD, Virk MS. Povidone-iodine solutions inhibit cell migration and survival of osteoblasts, fibroblasts, and myoblasts. *Spine (Phila Pa 1976)*. 2017;42(23):1757-1762.
111. Hirsch T, Koerber A, Jacobsen F, et al. Evaluation of toxic side effects of clinically used skin antiseptics in vitro. *J Surg Res*. 2010;164(2):344-350.
112. Ahani E, Toliyat T, Mahmoudi RM. Comparing size particle, release study and cytotoxicity activity of PHMB encapsulated in different liposomal formulations: neutral and cationic liposomes. *Bioeng Res*. 2019;1(3):1-6.
113. Yabes JM, White BK, Murray CK, et al. In vitro activity of Manuka Honey and polyhexamethylene biguanide on filamentous fungi and toxicity to human cell lines. *Med Mycol*. 2017;55(3):334-343.
114. Rembe JD, Fromm-Dornieden C, Schafer N, Bohm JK, Stuermer EK. Comparing two polymeric biguanides: chemical distinction, antiseptic efficacy and cytotoxicity of polyaminopropyl biguanide and polyhexamethylene biguanide. *J Med Microbiol*. 2016;65(8):867-876.
115. Poon VK, Burd A. In vitro cytotoxicity of silver: implication for clinical wound care. *Burns*. 2004;30(2):140-147.
116. Burd A, Kwok CH, Hung SC, et al. A comparative study of the cytotoxicity of silver-based dressings in monolayer cell, tissue explant, and animal models. *Wound Repair Regen*. 2007;15(1):94-104.
117. Galandakova A, Frankova J, Ambrozova N, et al. Effects of silver nanoparticles on human dermal fibroblasts and epidermal keratinocytes. *Hum Exp Toxicol*. 2016;35(9):946-957.
118. Zou SB, Yoon WY, Han SK, Jeong SH, Cui ZJ, Kim WK. Cytotoxicity of silver dressings on diabetic fibroblasts. *Int Wound J*. 2013;10(3):306-312.
119. Hidalgo E, Dominguez C. Study of cytotoxicity mechanisms of silver nitrate in human dermal fibroblasts. *Toxicol Lett*. 1998;98(3):169-179.
120. Hidalgo E, Bartolome R, Barroso C, Moreno A, Dominguez C. Silver nitrate: antimicrobial activity related to cytotoxicity in cultured human fibroblasts. *Skin Pharmacol Appl Skin Physiol*. 1998;11(3):140-151.
121. Hernandez-Sierra JF, Galicia-Cruz O, Angelica SA, Ruiz F, Pierdant-Perez M, Pozos-Guillen AJ. In vitro cytotoxicity of silver nanoparticles on human periodontal fibroblasts. *J Clin Pediatr Dent*. 2011;36(1):37-41.
122. Lam PK, Chan ES, Ho WS, Liew CT. In vitro cytotoxicity testing of a nanocrystalline silver dressing (Acticoat) on cultured keratinocytes. *Br J Biomed Sci*. 2004;61(3):125-127.
123. Szmyd R, Goralczyk AG, Skalniak L, et al. Effect of silver nanoparticles on human primary keratinocytes. *Biol Chem*. 2013;394(1):113-123.
124. Naor J, Savion N, Blumenthal M, Assia EI. Corneal endothelial cytotoxicity of diluted povidone-iodine. *J Cataract Refract Surg*. 2001;27(6):941-947.
125. Ince A, Schutze N, Hendrich C, Jakob F, Eulert J, Lohr JF. Effect of polyhexanide and gentamycin on human osteoblasts and endothelial cells. *Swiss Med Wkly*. 2007;137(9-10):139-145.
126. Khan I, Bahuguna A, Krishnan M, et al. The effect of biogenic manufactured silver nanoparticles on human endothelial cells and zebrafish model. *Sci Total Environ*. 2019;679:365-377.
127. Guo H, Zhang J, Boudreau M, et al. Intravenous administration of silver nanoparticles causes organ toxicity through intracellular ROS-related loss of inter-endothelial junction. *Part Fibre Toxicol*. 2016;13:21.
128. Cui J, Zhang YD. Effects of different doses of nano silver on vascular endothelial cell proliferation in vitro. *Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi*. 2011;27(6):697-699.
129. Sun X, Shi J, Zou X, Wang C, Yang Y, Zhang H. Silver nanoparticles interact with the cell membrane and increase endothelial permeability by promoting VE-cadherin internalization. *J Hazard Mater*. 2016;317:570-578.
130. van Meurs SJ, Gawlitta D, Heemstra KA, Poolman RW, Vogely HC, Kruyt MC. Selection of an optimal antiseptic solution for intraoperative irrigation: an in vitro study. *J Bone Joint Surg Am*. 2014;96(4):285-291.
131. Leaper DJ, Durani P. Topical antimicrobial therapy of chronic wounds healing by secondary intention using iodine products. *Int Wound J*. 2008;5(2):361-368.
132. Muller G, Kramer A. Biocompatibility index of antiseptic agents by parallel assessment of antimicrobial activity and cellular cytotoxicity. *J Antimicrob Chemother*. 2008;61(6):1281-1287.
133. Muller G, Kramer A. Comparative study of in vitro cytotoxicity of povidone-iodine in solution, in ointment or in a liposomal formulation (Repithel) and selected antiseptics. *Dermatology*. 2006;212(Suppl 1):91-93.
134. Bigliardi P, Langer S, Cruz JJ, Kim SW, Nair H, Srisawasdi G. An Asian perspective on povidone iodine in wound healing. *Dermatology*. 2017;233(2-3):223-233.
135. Fumal I, Braham C, Paquet P, Pierard-Franchimont C, Pierard GE. The beneficial toxicity paradox of antimicrobials in leg ulcer healing impaired by a polymicrobial flora: a proof-of-concept study. *Dermatology*. 2002;204(Suppl 1):70-74.
136. Lachapelle JM. A comparison of the irritant and allergenic properties of antiseptics. *Eur J Dermatol*. 2014;24(1):3-9.

137. Schunter JA, Stocker B, Brehler R. A case of severe anaphylaxis to polyhexanide: cross-reactivity between biguanide anti-septics. *Int Arch Allergy Immunol.* 2017;173(4):233-236.
138. Akter M, Sikder MT, Rahman MM, et al. A systematic review on silver nanoparticles-induced cytotoxicity: physicochemical properties and perspectives. *J Adv Res.* 2018;9:1-16.
139. Xiu ZM, Zhang QB, Puppala HL, Colvin VL, Alvarez PJ. Negligible particle-specific antibacterial activity of silver nanoparticles. *Nano Lett.* 2012;12(8):4271-4275.
140. Beer C, Foldbjerg R, Hayashi Y, Sutherland DS, Autrup H. Toxicity of silver nanoparticles - nanoparticle or silver ion? *Toxicol Lett.* 2012;208(3):286-292.
141. Lansdown AB. A pharmacological and toxicological profile of silver as an antimicrobial agent in medical devices. *Adv Pharmacol Sci.* 2010;2010:910686.
142. Rothenberger J, Krauss S, Tschumi C, Rahmanian-Schwarz A, Schaller HE, Held M. The effect of Polyhexanide, Octenidine Dihydrochloride, and tea tree oil as topical antiseptic agents on in vivo microcirculation of the human skin: a noninvasive quantitative analysis. *Wounds.* 2016;28(10):341-346.
143. Wang L, Qin W, Zhou Y, et al. Transforming growth factor beta plays an important role in enhancing wound healing by topical application of povidone-iodine. *Sci Rep.* 2017;7(1):991.
144. Beukelman CJ, van den Berg AJ, Hoekstra MJ, Uhl R, Reimer K, Mueller S. Anti-inflammatory properties of a liposomal hydrogel with povidone-iodine (Repithel) for wound healing in vitro. *Burns.* 2008;34(6):845-855.
145. Pierard-Franchimont C, Paquet P, Arrese JE, Pierard GE. Healing rate and bacterial necrotizing vasculitis in venous leg ulcers. *Dermatology.* 1997;194(4):383-387.
146. Pak CS, Park DH, Oh TS, et al. Comparison of the efficacy and safety of povidone-iodine foam dressing (Betafoam), hydrocellular foam dressing (Allevyn), and petrolatum gauze for split-thickness skin graft donor site dressing. *Int Wound J.* 2019;16(2):379-386.
147. Goldenheim PD. An appraisal of povidone-iodine and wound healing. *Postgrad Med J.* 1993;69(Suppl 3):S97-S105.
148. Niedner R. Cytotoxicity and sensitization of povidone-iodine and other frequently used anti-infective agents. *Dermatology.* 1997;195(Suppl 2):89-92.
149. Lee JW, Song KY. Evaluation of a polyurethane foam dressing impregnated with 3% povidone-iodine (Betafoam) in a rat wound model. *Ann Surg Treat Res.* 2018;94(1):1-7.
150. Kjolseth D, Frank JM, Barker JH, et al. Comparison of the effects of commonly used wound agents on epithelialization and neovascularization. *J Am Coll Surg.* 1994;179(3):305-312.
151. To E, Dyck R, Gerber S, Kadavil S, Woo KY. The effectiveness of topical polyhexamethylene biguanide (PHMB) agents for the treatment of chronic wounds: a systematic review. *Surg Technol Int.* 2016;29:45-51.
152. Goertz O, Lauer H, Hirscht T, et al. Evaluation of angiogenesis, epithelialisation and microcirculation after application of polyhexanide, chitosan and sodium chloride in rodents. *Int Wound J.* 2016;13(6):1161-1167.
153. Eberlein T, Haemmerle G, Signer M, Gruber Moesenbacher U, Traber J, Mittlboeck M, et al. Comparison of PHMB-containing dressing and silver dressings in patients with critically colonised or locally infected wounds. *J Wound Care.* 2012;21(1):12-20.
154. Wiegand C, Abel M, Ruth P, Hipler UC. HaCaT keratinocytes in co-culture with *Staphylococcus aureus* can be protected from bacterial damage by polyhexanide. *Wound Repair Regen.* 2009;17(5):730-738.
155. Kim HR, Shin DY, Chung KH. In vitro inflammatory effects of polyhexamethylene biguanide through NF-kappaB activation in A549 cells. *Toxicol in Vitro.* 2017;38:1-7.
156. Storm-Versloot MN, Vos CG, Ubbink DT, Vermeulen H. Topical silver for preventing wound infection. *Cochrane Database Syst Rev.* 2010;3:CD006478.
157. Khansa I, Schoenbrunner AR, Kraft CT, Janis JE. Silver in wound care-friend or foe?: a comprehensive review. *Plast Reconstr Surg Glob Open.* 2019;7(8):e2390.
158. Rodriguez-Arguello J, Lienhard K, Patel P, et al. A scoping review of the use of silver-impregnated dressings for the treatment of chronic wounds. *Ostomy Wound Manage.* 2018;64(3):14-31.
159. Lo SF, Chang CJ, Hu WY, Hayter M, Chang YT. The effectiveness of silver-releasing dressings in the management of non-healing chronic wounds: a meta-analysis. *J Clin Nurs.* 2009;18(5):716-728.
160. Woo KY, Coutts PM, Sibbald RG. A randomized controlled trial to evaluate an antimicrobial dressing with silver alginate powder for the management of chronic wounds exhibiting signs of critical colonization. *Adv Skin Wound Care.* 2012;25(11):503-508.
161. Duan X, Peng D, Zhang Y, et al. Sub-cytotoxic concentrations of ionic silver promote the proliferation of human keratinocytes by inducing the production of reactive oxygen species. *Front Med.* 2018;12(3):289-300.
162. Frankova J, Pivodova V, Vagnerova H, Juranova J, Ulrichova J. Effects of silver nanoparticles on primary cell cultures of fibroblasts and keratinocytes in a wound-healing model. *J Appl Biomater Funct Mater.* 2016;14(2):e137-e142.
163. Harries RL, Bosanquet DC, Harding KG. Wound bed preparation: TIME for an update. *Int Wound J.* 2016;13(Suppl 3):8-14.
164. Shi C, Wang C, Liu H, et al. Selection of appropriate wound dressing for various wounds. *Front Bioeng Biotechnol.* 2020;8:182.
165. Bishop SM, Walker M, Rogers AA, Chen WY. Importance of moisture balance at the wound-dressing interface. *J Wound Care.* 2003;12(4):125-128.
166. Dedo DD, Alonso WA, Ogura JH. Povidone-iodine: an adjunct in the treatment of wound infections, dehiscences, and fistulas in head and neck surgery. *Trans Sect Otolaryngol Am Acad Ophthalmol Otolaryngol.* 1977;84(1):68-74.

How to cite this article: Alves PJ, Barreto RT, Barrois BM, Gryson LG, Meaume S, Monstrey SJ. Update on the role of antiseptics in the management of chronic wounds with critical colonisation and/or biofilm. *Int Wound J.* 2021;18:342-358. <https://doi.org/10.1111/iwj.13537>