



FEMS Microbiology Letters, 366, 2019, fnz071

doi: [10.1093/femsle/fnz071](https://doi.org/10.1093/femsle/fnz071)

Advance Access Publication Date: 10 April 2019

Research Letter

RESEARCH LETTER – Biotechnology & Synthetic Biology

Aspergillus niger citrate exporter revealed by comparison of two alternative citrate producing conditions

Dorett I Odoni^{1,*,\dagger,\ddagger,\#}, Marta Vazquez-Vilar^{1,\ddagger,\S}, Merlijn P van Gaal^{1,\¶}, Tom Schonewille¹, Vitor AP Martins dos Santos¹, Juan Antonio Tamayo-Ramos^{1,2}, Maria Suarez-Diez^{1,\ddagger} and Peter J Schaap¹

¹Laboratory of Systems and Synthetic Biology, Wageningen University & Research, Stippeneng 4, 6708 WE Wageningen, The Netherlands and ²International Research Center in Critical Raw Materials-ICCRAM, Advanced Materials, Nuclear Technology and Applied Bio/Nanotechnology, University of Burgos, Plaza Misael Bañuelos s/n, 09001 Burgos, Spain

*Corresponding author: Im Neuenheimer Feld 280, 69120 Heidelberg, Germany. Tel: +49 6221 42 3611; E-mail: dorettwur@gmail.com

One sentence summary: A comparative transcriptomics approach to disentangle *Aspergillus niger* citrate production is combined with other bioinformatics approaches to pinpoint a citrate exporter, which is validated by heterologous expression in *Saccharomyces cerevisiae*.

^{\dagger}These authors contributed equally to this work.

^{\ddagger}Current address: Bioinformatics and Omics Data Analytics (BODA), German Cancer Research Center (DKFZ), Heidelberg, Germany

^{\S}Current address: Instituto de Biología Molecular y Celular de Plantas (IBMCP), Consejo Superior de Investigaciones Científicas (CSIC), Universidad Politécnica de Valencia, Valencia, Spain.

^{\¶}Current address: Laboratory of Microbiology, Wageningen University & Research, Wageningen, The Netherlands

Editor: Matthias Steiger

^{\#}Dorett I Odoni, <http://orcid.org/0000-0001-9085-3899>

Juan Antonio Tamayo-Ramos, <http://orcid.org/0000-0002-7071-002X>

Vitor AP Martins dos Santos, <http://orcid.org/0000-0002-2352-9017>

^{\ddagger}Maria Suarez-Diez, <http://orcid.org/0000-0001-5845-146X>

Peter J Schaap, <http://orcid.org/0000-0002-4346-6084>

ABSTRACT

Currently, there is no consensus regarding the mechanism underlying *Aspergillus niger* citrate biosynthesis and secretion. We hypothesise that depending on the experimental setup, extracellular citrate accumulation can have fundamentally different underlying transcriptomic landscapes. We show that varying the amount and type of supplement of an arginine auxotrophic *A. niger* strain results in transcriptional down-regulation of citrate metabolising enzymes in the condition in which more citrate is accumulated extracellularly. This contrasts with the transcriptional adaptations when increased citrate production is triggered by iron limitation. By combining gene expression data obtained from these two very distinct experimental setups with hidden Markov models and transporter homology approaches, we were able to compile a shortlist of the most likely citrate transporter candidates. Two candidates (An17g01710 and An09g06720m.01) were

Received: 10 January 2019; Accepted: 5 April 2019

© FEMS 2019. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

heterologously expressed in the yeast *Saccharomyces cerevisiae*, and one of the resultant mutants showed the ability to secrete citrate. Our findings provide steps in untangling the complex interplay of different mechanisms underlying *A. niger* citrate accumulation, and we demonstrate how a comparative transcriptomics approach complemented with further bioinformatics analyses can be used to pinpoint a fungal citrate exporter.

Keywords: *Aspergillus niger*; citrate; transport; transcriptomics; homology; MDR

INTRODUCTION

Aspergillus niger citrate production is a notorious example of a process that requires a unique combination of unusual nutrient and environmental conditions (Karaffa and Kubicek 2003). Carbon excess relative to iron, zinc, copper, manganese, phosphorus, magnesium, potassium and nitrogen reportedly all leads to an increase in *A. niger* citrate secretion in the form of either absolute yield or relative productivity (Chesters and Rolinson 1951; Karaffa and Kubicek 2003). The result of this is that although *A. niger* citrate production has been subject to study since Curries fundamental breakthroughs regarding *A. niger* citrate fermentation more than 100 years ago (Currie 1917), it is still not fully understood.

One problem might be that there are multiple factors at play, each independently or in combination influencing citrate secretion, and studies, therefore, contradict each other depending on which aspect or time point of citrate accumulation was investigated (Karaffa and Kubicek 2003). Citrate secretion under conditions of high glucose concentrations in the medium is generally accepted to be the result of overflow metabolism, i.e. excess flux through glycolysis and ultimately the TCA cycle, leading to the 'undesired' (from the perspective of the fungus) accumulation of citrate (Legiša and Matthey 2007). Nevertheless, the viewpoint that citrate is solely an overflow metabolite is changing, and *A. niger* citrate secretion might also be regarded as a response to environmental conditions such as competition (Andersen et al. 2011), or low iron bioavailability (Odoni et al. 2017). Further understanding of the various aspects that lead to increased citrate production in *A. niger* can provide tools for metabolic engineering approaches to further control and modulate citrate production.

An example tool for the control and modulation of citrate production can be found at transporter level (Karaffa and Kubicek 2003). It has been shown that overexpressing or introducing specific transporters for the product of interest can often overcome product limitation (van der Straat and de Graaff 2014). Dynamic models of metabolism have highlighted the citrate exporter as one of the proteins whose overexpression could lead to increased citrate production rates (Alvarez-Vasquez, González-Alcón and Torres 2000). The citrate transport system in *A. niger* has been described (Netik et al. 1997), and a comprehensive list of putative citrate transporter candidates in *A. niger* has been compiled using a transcriptomics approach (Yin et al. 2017), but thus far, gene expression analyses failed to pinpoint a definite *A. niger* citrate exporter.

Here, we explore how combining multiple gene expression datasets with other bioinformatic approaches can narrow down the list of putative citrate exporter candidates. For this, we worked with well-defined *A. niger* *argB* knock-out mutants. The $\Delta argB$ mutation induces an arginine auxotrophy that can be overcome by media supplementation with either arginine or citrulline (Lenouvel, van de Vondervoort and Visser 2002). We studied the impact of supplement type and amount on citrate production and performed transcriptome analysis to pinpoint the

metabolic adaptations associated with the higher citrate producing condition. We compared the results of this study with previous data regarding changes in citrate production triggered by low iron concentrations in the medium (Odoni et al. 2017). Expression data were combined with hidden Markov models (HMMs) and homology approaches to compile a shortlist of the most promising citrate transporter candidates. We validated our approach by heterologously expressing two selected citrate exporter candidates in *Saccharomyces cerevisiae*, and show that one of these yeast transformant strains indeed accumulated a measurable amount of extracellular citrate.

MATERIALS AND METHODS

Strains and media

Aspergillus niger strains NW305 (*cspA*, *goxC17*, $\Delta argB$) (Ruijter et al. 2003), and NW186 (*cspA1*, *goxC17*, *prtF28*, $\Delta argB$) - a derivative of NW185 (Ruijter, Van De Vondervoort and Visser 1999) with restored uridine prototrophy (Odoni et al. 2017) - were used for identification of putative citrate exporter candidates. Strains were maintained on complete medium (CM) as described in Odoni et al. (2017). *Aspergillus niger* spores were harvested with Saline-Tween (0.9% NaCl and 0.001% Tween80).

Escherichia coli DH5 α was used for standard gene cloning purposes. *Escherichia coli* was cultured in Luria broth (LB) medium (5 g L⁻¹ yeast extract, 10 g L⁻¹ peptone, 10 g L⁻¹ NaCl) supplemented 100 mg L⁻¹ ampicillin when necessary.

Saccharomyces cerevisiae strain CENPK2-1D (MAT α ; his3D1; leu2-3.112; ura3-52; trp1-289; MAL2-8c; SUC2) was used for validating the citrate transporter candidates. Preparation of CENPK2-1D yeast electro-competent cells and transformation was performed as described in Suga and Hatakeyama (2003). Yeast-transformed cells were selected in synthetic media (SD) with 2% (w/v) dextrose, 0.67% (w/v) Yeast Nitrogen Base without amino acids (BD), 0.14% (w/v) Yeast Synthetic Drop-out Medium supplement without uracil, tryptophan, histidine and leucine (Sigma-Aldrich), 0.0076% (w/v) histidine, 0.0076% (w/v) tryptophan and 0.038% (w/v) leucine. For *S. cerevisiae* transformation, the cell suspension was mixed with plasmid or linear DNA, transferred to a pre-chilled cuvette (0.2 cm Gene Pulse, Bio-Rad, Uden, The Netherlands), pulsed at 2.5 kV, 25 μ F, 200 Ω using Gene Pulser Xcell (Bio-Rad) and plated on SD agar-plates.

Transporter identification

Experimental setup for *A. niger* RNA seq analysis

For pre-growth of *A. niger* NW186 and NW305, a total of 1×10^6 spores mL⁻¹ were inoculated in 1 L Erlenmeyer shake flasks containing 200 mL medium with the following composition: 1.2 g L⁻¹ NaNO₃, 0.5 g L⁻¹ KH₂PO₄, 0.2 g L⁻¹ MgSO₄·7H₂O, 40 μ L L⁻¹ Vishniac solution, and supplemented with 50 g L⁻¹ (~250 mM) glucose as carbon source, and either 1.1 mM (0.2 g L⁻¹) arginine or 5 mM (0.88 g L⁻¹) citrulline. After 24 h of pre-growth, 11 g of *A. niger* mycelium was transferred to controlled fermentors, containing the same medium as for the pre-growth.

The supernatants were collected for determination of extracellular metabolite concentrations, especially carbon sources and organic acids, by high-performance liquid chromatography (HPLC) as described previously (Odoni et al. 2017).

RNA isolation, sequencing and RNA seq data processing and analysis

Total RNA extraction, RNA sequencing and RNA seq data processing and statistical analysis of the supplement experiment were performed as described in Odoni et al. (2017). The aligned .bam files were submitted to the European Nucleotide Archive (ENA) under the accession number PRJEB24704.

Shortlisting putative citrate transporter candidates

To assemble a list of likely citrate transporter candidates, the RNA seq analysis was complemented with a HMM approach and a homology based approach. HMMs were built as described (Sloothaak et al. 2015) from two sets of proteins ('citrate transport' and 'GO:00 15137') downloaded from the UniProt database (Bateman et al. 2015). The genomes used for the homology approach were downloaded from the JGI database (Nordberg et al. 2014): *Aspergillus kawachii* (Futagami et al. 2011), *Aspergillus nidulans* (Galagan et al. 2005; Arnaud et al. 2012), *Aspergillus flavus* (Arnaud et al. 2012), *Aspergillus fumigatus* (Nierman et al. 2005; Fedorova et al. 2008; Joardar et al. 2012), *Aspergillus terreus* (Arnaud et al. 2012), *Yarrowia lipolytica* (Dujon et al. 2004) and *S. cerevisiae* (Dujon et al. 2004). Proteins with transmembrane helix structures (from the *A. niger* ATCC 1015 *in silico* proteome) were identified using the stand-alone TMHMM 2.0 software package (Sonnhammer, Von Heijne and Krogh 1998; Krogh et al. 2001). Protein localisation was predicted using the stand-alone protComp software (www.softberry.com). The resulting shortlists for each approach can be found in Supplementary Data 4 (Supporting Information).

Transporter validation

The coding sequences of two putative citrate transporter genes, with ATCC 1015 protein IDs 1165828 (*A. niger* CBS 513.88: An17g01710, *A. niger* NRRL3: 06350) and 212337 (*A. niger* CBS 513.88: An09g06720m.01, *A. niger* NRRL3: 00550) were amplified from *A. niger* N402 genomic DNA. The amplified fragments were introduced into a derivative of the pYES-plasmid, whereby the GAL1 inducible promoter was replaced by the CUP1 inducible promoter (Mascorro-Gallardo, Covarrubias and Gaxiola 1996). The plasmid was PCR-amplified with the Q5 polymerase following the manufacturer's protocol.

The coding sequence of the citrate exporter candidate 1165828 (An17g01710) was cloned using Golden Gate. The Golden Gate reaction mixture was prepared as follows: 400 U of T4 DNA ligase (NEB), 10 U of BsmBI (NEB), 1.5 μ L of BSA (1 mg·mL⁻¹), 1.5 μ L of T4 DNA ligase buffer (NEB), ~40 fmol of each PCR product and water to bring the volume up to 15 μ L. The reaction mixture was incubated in a thermocycler according to the following program: 37°C for 10 min prior to 25 cycles of digestion-ligation (37°C for 3 min, 16°C for 4 min) followed by a final digestion step (55°C for 10 min) and a heat inactivation step (80°C for 10 min). *Escherichia coli* DH5 α competent cells were transformed with 1 μ L of the digestion-ligation reaction, and transformants were selected on ampicillin plates. Plasmid extraction was performed using the GeneJET Plasmid Miniprep kit (Thermo Fisher Scientific, Waltham, Massachusetts/US) following the manufacturer's instructions. CENPK2-1D competent cells were transformed with a sequence-verified plasmid.

The coding sequence of the citrate exporter candidate 212337 (An09g06720m.01) was cloned using the yeast homologous recombination assembly strategy. For the in-yeast assembly, CENPK2-1D cells were directly transformed with 100 ng of the PCR-amplified pYES-CUP1 plasmid and an equimolar amount of insert, and plated on selective medium. Positive transformants were picked and grown overnight in liquid medium. Plasmids were extracted using the GeneJET Plasmid Miniprep kit following the manufacturer's instructions with minor modifications. Lysis was performed by transferring the cells to Lysing MatrixC 2 mL tubes (MP Biomedicals, Santa Ana, California/US) and homogenising them for 40 s with a FastPrep-24 from MP Biomedicals. *Escherichia coli* DH5 α competent cells were transformed with the extracted plasmids for propagation, and the plasmids were then again extracted using the GeneJET Plasmid Miniprep kit following the manufacturer's instructions, and sequenced for insert verification.

For the growth experiments, yeast transformant strains were pre-grown in 15 mL SD medium supplemented with 0.0076% uracil to allow growth of the parent strain. A total of 300 μ L of the pre-growth culture were inoculated in 100 mL shake flasks containing 20 mL SD medium described above, containing either 20 g·L⁻¹ glucose or 20.44 g·L⁻¹ glycerol. For CUP1 promoter-induction, CuSO₄ (final concentration 1 mM) was added after 4 h of growth.

RESULTS

Aspergillus niger biomass and citrate yields under different culture conditions

We used the two *A. niger* N402 derivative strains NW305 and NW186. N402, as well as wildtype *A. niger*, has the means to produce major quantities of gluconate, oxalate and citrate, depending on the external condition, but requires pH <3 to avoid excess oxalate production in favour of increased citrate production (Ruijter, Van De Vondervoort and Visser 1999). However, both NW186 and NW305 have a mutation in glucose oxidase (*goxC17*), impairing the natural *A. niger* gluconate production. In addition, NW186 has a mutation in oxaloacetate acetylhydrolase (*prtF28*), impairing oxalate production in this strain (Ruijter, Van De Vondervoort and Visser 1999). Thus, while NW305 can produce both oxalate and citrate, but not gluconate, NW186 can produce neither gluconate nor oxalate, but retains its capacity to produce citrate, and does so even at higher pH levels (Ruijter, Van De Vondervoort and Visser 1999). In addition, NW305 and NW186 are ornithine transcarbamylase (*argB*) knock-out mutants, making them arginine auxotrophic (Lenouvel, van de Vondervoort and Visser 2002). Supplementation of the medium with either arginine or citrulline can restore growth of the Δ *argB* mutants, but we found that addition of excess (5 mM) citrulline increases total citrate production when compared to addition of (1.1 mM; 'standard' condition) arginine. We compared gene expression data from the experiment described above, hereafter referred to as supplement experiment, to data from one of our previous experimental setups, described in Odoni et al. (2017), in which increased citrate secretion was induced by iron limitation (hereafter referred to as iron experiment).

Citrate production yields of the considered conditions are shown in Table 1. In the supplement experiment, total citrate yield is almost doubled in NW186 + Fe.c compared to NW186 + Fe.a. In fact, and not surprisingly, excess citrulline

Table 1. Final biomass and citrate yields of *A. niger* NW305 and NW186 grown with varying iron concentrations and/or different supplements in the medium.

| Experiment | Strain | Fe | Suppl ¹ | Final biomass (g) | Citrate yield | |
|------------|--------|----|--------------------|--------------------------|-----------------------------|-------------------------------|
| | | | | | (g·g ⁻¹ biomass) | (g·g ⁻¹ substrate) |
| Iron | NW305 | ++ | a | 0.35 ± 0.02 | 0.67 ± 0.07 | 0.02 ± 0.002 |
| Iron | NW305 | - | a | 0.19 ± 0.02 | 4.92 ± 0.35 | 0.11 ± 0.002 |
| Iron | NW186 | - | a | 0.22 ± 1e ⁻⁰³ | 21.88 ± 1.51 | 0.46 ± 0.02 |
| Supplement | NW186 | + | a | 1.87 ± 0.04 | 1.80 ± 0.14 | 0.17 ± 0.03 |
| Supplement | NW186 | + | c | 3.96 ± 0.09 | 2.98 ± 0.35 | 0.31 ± 0.03 |

¹Suppl: supplement, with a = arginine, c = citrulline




Figure 1. Differences in total citrate yields and production rates on glucose as carbon source. *Aspergillus niger* strains NW186 and NW305 were either grown without addition of iron (circles), or varying amounts of Fe(II)SO₄ (triangles = 1 g·L⁻¹, squares = 10 g·L⁻¹) and supplemented with either arginine (filled symbols) or excess citrulline (empty symbols). Sample points for RNA extraction (t = 48 h) are marked in orange. Measurement points were taken once every 24 h and show the average of two biological replicates. Note that glucose consumption, rather than time, is plotted on the x-axis.

compared to the standard amount of arginine has an overall stimulating effect on metabolism; it leads to increased glucose consumption (Fig. 1), doubled final biomass (Table 1), and increased CO₂ production (337.79 ± 15.86 mM in NW186 + Fe.c and 265.51 ± 9.28 mM in NW186 + Fe.a). On the other hand, not adding iron to the culture medium increases the citrate per glucose production rate, but reduced glucose consumption is observed (Fig. 1), and total biomass production remains limited (Table 1).

General transcriptome analysis

Transcriptomic data analysis of the iron experiment has already been described in Odoni *et al.* (2017). Here, we performed transcriptome analysis on the supplement experiment, i.e. NW186 supplemented with either 1.1 mM arginine or 5 mM citrulline. RNA for RNA sequencing (RNA seq) was extracted after 48 h of growth. The annotated genome of *A. niger* ATCC 1015 (Andersen *et al.* 2011) was used as reference to map the RNA seq reads (Table 2). Genes with count per million (CPM) ≥ 1 were considered expressed (Supplementary Data 1, Supporting Information). The supplement change from arginine to excess citrulline induces

major transcriptional adaptations, and over 20% of the annotated genes are differentially expressed (Table 2; Supplementary Data 2, Supporting Information).

From the 466 enzyme commission (EC) numbers that have been included in KEGG maps of metabolism (Kanehisa and Goto 2000; Kanehisa *et al.* 2015), and can be found in the annotated genome of *A. niger* ATCC 1015, 37% show differential expression. Pathway enrichment analysis shows prevalence of differentially expressed genes in pathways associated to biomass formation, such as starch and sucrose metabolism, and pathways related to amino acid biosynthesis, such as phenylalanine, tyrosine and tryptophan biosynthesis and metabolism, and valine, leucine and isoleucine biosynthesis (Supplementary Data 3, Supporting Information). There is also an enrichment in differentially expressed genes related to fatty acid biosynthesis, and synthesis and degradation of ketone bodies pathways, upon addition of excess citrulline instead of arginine (Supplementary Data 3, Supporting Information). Note that, although we observed higher glucose consumption in NW186 + Fe.c, glycolysis as a pathway did not show enrichment of differentially expressed genes. Interestingly, the enzymes converting citrulline to arginine also showed no differential expression in the supplement experiment (Fig. 2; Supplementary Data 3, Supporting Information). Most of the pathways enriched in the arginine/citrulline supplement experiment (Supplementary Data 3, Supporting Information) were also enriched in the iron experiment (Odoni *et al.* 2017). Phenylalanine, tyrosine and tryptophan biosynthesis, and fatty acid biosynthesis pathways, showed similar behaviours in both experiments, and most of the enzymes in these pathways are down-regulated in the condition with higher citrate production.

Analysis of the expression of enzymes involved in citrate metabolism revealed quite a contrasting transcriptomic landscape in the two experimental setups (Fig. 2). While citrate biosynthesis genes are up-regulated in the condition corresponding to increased extracellular citrate secretion in the iron experiment, there is down-regulation of citrate metabolising enzymes in the condition corresponding to increased extracellular citrate accumulation in the arginine/citrulline supplement experiment. We also found that alternative oxidase (AOX, or non-electrogenic ubiquinol oxidase, EC 1.10.3.11) was up-regulated in the condition corresponding to the highest citrate secretion in both experimental setups (Fig. 2; Supplementary Data 2, Supporting Information).

Aspergillus niger citrate exporter identification

To identify the elusive citrate exporter, we aimed to combine the gene expression data obtained from the two experimental setups with complementary bioinformatic approaches.

For one, we constructed two HMMs from multiple sequence alignments of biochemically characterised citrate transporters obtained from the UniProt database (Bateman et al. 2015). The sequences selected to build the models matched the terms 'citrate transport' and 'GO:0015137: citrate transmembrane transporter activity', and differences in the selected sets likely reflect differences in protein annotation in UniProt. The HMMs were used to identify and score new citrate transporter candidates in the ATCC 1015 *in silico* proteome. In addition, we identified all (predicted) plasma-membrane proteins from the ATCC 1015 *in silico* proteome and performed protein BLAST against genomes of other sequenced *Aspergilli* or yeast that are known to either produce (*A. kawachii*, *Y. lipolytica*) or not produce (*A. flavus*, *A. terreus*, *S. cerevisiae*) citrate. Finally, we also included 10 genes up- and downstream of the *A. niger* citrate synthase gene (*citA*) as putative citrate transporter candidates. We thereby followed the notion that it is not uncommon that fungal transporters are encoded by genes in close genomic proximity to the genes encoding biosynthesis of their transport target (van der Straat and de Graaff 2014).

The lists of putative citrate exporter candidates obtained from each of these steps were then filtered and ranked based on the following selection criteria (from basic to more specific importance, with putative candidate transporters not fulfilling these criteria dropping out at each step):

- (1) Average nucleotide coverage (avNtCov) ≥ 1 in all conditions.
- (2) avNtCov ≥ 50 in NW186 -Fe.a (highest citrate production rate) and NW186 +Fe.c (highest overall citrate yield).
- (3) Gene expression up-regulated in NW186 -Fe.a compared to NW305 -Fe.a.
- (4) Gene expression up-regulated in NW186 + Fe.c compared to NW186 +Fe.a.

These four points were considered the minimum requirement, and proteins had to adhere to these criteria to be considered for the HMM and proximity approaches (Supplementary Data 4, Supporting Information; sheets 'HMM_approach' and 'proximity_to_citA' approach). For a more stringent selection, applied to the proteins found based on homology in other filamentous fungi and yeasts, the following additional criteria had to be met:

- (5) Encoded protein product contains ≥ 1 predicted transmembrane helix domain (tmhmm).
- (6) *Aspergillus niger* protein is more similar (higher NCBI Blast '% identity') to *A. kawachii* homologue than *A. flavus* and *A. terreus* homologues.
- (7) *Aspergillus niger* protein is more similar to *Y. lipolytica* homologue than to *S. cerevisiae* homologue.
- (8) Gene expression up-regulated in NW305 -Fe.a compared to NW305 ++Fe.a. (Note that this was not considered a minimal requirement, as NW305 will shift to oxalate production under iron limited conditions, and thus the direct inference on citrate production is not as clear cut as between NW186 and NW305, see also Odoni et al. (2017)).

The final shortlists of each approach can be found in Supplementary Data 4 (Supporting Information; sheets 'HMM_approach', 'homology_approach' and 'proximity_to_citA'). The genes that passed the selection criteria were then ordered from high to low based on their log₂ fold-changes (log₂FC values) in NW186 -Fe.a vs NW305 -Fe.a (Supplementary Data 4, Supporting Information, sheet 'all_sorted'; Table 3), as we consider this the biologically most relevant condition for *A.*




Figure 3. Citrate secretion in *S. cerevisiae* transformed with CexA. Solid line = grown on glucose, dashed line = grown on glycerol, filled symbols = induced and empty symbols = non-induced. Note that there was no measurable extracellular citrate accumulation in the untransformed parent strain. Measurements indicate the average of two biological replicates.

niger citrate export (Odoni et al. 2017). As a result of our filtering and prioritisation criteria, and considering the predicted protein localisation, the protein with the *A. niger* ATCC 1015 protein ID 1165828 (*A. niger* CBS 513.88: An17g01710, *A. niger* NRRL3: 06350) is the top most likely citrate exporter candidate (Table 3). In addition, we choose another protein, with protein ID 212337 (*A. niger* CBS 513.88: An09g06720m.01, *A. niger* NRRL3: 00550), as a second putative citrate exporter candidate. This protein was chosen based on being the only predicted plasma-membrane transporter candidate in the close vicinity of the *A. niger* citrate synthase gene (*citA*), although the lower expression levels suggest lesser importance as actual citrate exporter.

In vivo validation of selected citrate exporter candidates in *S. cerevisiae*

Saccharomyces cerevisiae was transformed with plasmids containing either one of the putative citrate exporter candidate genes under the control of a copper inducible promoter (CUP1). In the first experiment, we added CuSO₄ after 4 h of growth to the medium of all strains to induce expression of the transporter proteins, and found that neither the untransformed control strain, nor the strain transformed with the protein 212337 (An09g06720m.01) encoding gene secreted any citrate. However, under these exact same conditions, the strain transformed with the protein 1165828 (An17g01710) encoding gene accumulated a small amount of extracellular citrate after induction of gene expression with CuSO₄, indicating that this could be a citrate exporter. To further verify our initial results, we grew *S. cerevisiae* transformed with the promising citrate exporter gene with either glucose or glycerol in the medium, and either did or did not induce expression with CuSO₄. Confirming our initial observations, we found accumulation of citrate in the extracellular medium in *S. cerevisiae* when heterologous gene expression was induced (Fig. 3). Again, no citrate accumulation was observed in the extracellular medium of the untransformed control strain, irrespective of whether or not we added CuSO₄ to the medium, excluding the possibility that the observed effect was merely a result of intracellularly accumulated citrate due

Table 3. Shortlist of average nucleotide coverage and log₂ fold-changes of putative A. niger citrate exporter candidates found by the different approaches mentioned (Supplementary Data 4, Supporting Information; sheet 'all.sorted').

| proteinid | tmhmm | NW305 -Fe.a | NW186 -Fe.a | log2FC | FDR | NW186 +Fe.a | NW186 +Fe.c | log2FC | FDR | NW305 -Fe.a | NW305 ++Fe.a | log2FC | FDR |
|-----------|-------|----------------|----------------|--------|------|----------------|-----------------|--------|------|----------------|----------------|--------|------|
| 1165828 | 12 | 14.61 ± 0.99 | 139.89 ± 22.45 | 3.42 | 0.00 | 34.03 ± 1.52 | 350.11 ± 111.78 | 3.30 | 0.00 | 14.61 ± 0.99 | 4.73 ± 0.46 | -1.75 | 0.00 |
| 1186388 | 14 | 22.65 ± 1.00 | 64.92 ± 2.47 | 1.67 | 0.00 | 70.22 ± 5.08 | 250.45 ± 19.01 | 1.78 | 0.00 | 22.65 ± 1.00 | 6.90 ± 0.82 | -1.88 | 0.00 |
| 212337 | 4 | 52.79 ± 7.20 | 123.91 ± 0.68 | 1.44 | 0.00 | 34.70 ± 1.91 | 116.80 ± 16.22 | 1.66 | 0.00 | 52.79 ± 7.20 | 41.65 ± 2.17 | -0.44 | 0.48 |
| 1187835 | 12 | 24.73 ± 0.33 | 53.79 ± 6.51 | 1.26 | 0.00 | 87.03 ± 8.32 | 109.85 ± 46.96 | 0.27 | 0.89 | 24.73 ± 0.33 | 9.74 ± 1.07 | -1.51 | 0.00 |
| 1155853 | 5 | 62.05 ± 1.00 | 132.17 ± 18.17 | 1.23 | 0.00 | 61.63 ± 2.29 | 108.63 ± 3.11 | 0.77 | 0.05 | 62.05 ± 1.00 | 61.59 ± 6.45 | -0.16 | 0.85 |
| 1105147 | 14 | 202.52 ± 38.35 | 407.77 ± 8.54 | 1.15 | 0.01 | 34.77 ± 2.66 | 377.25 ± 25.34 | 3.42 | 0.00 | 202.52 ± 38.35 | 2.88 ± 0.05 | -6.32 | 0.00 |
| 1120210 | 7 | 76.98 ± 1.19 | 110.49 ± 2.48 | 0.66 | 0.28 | 55.93 ± 0.80 | 65.12 ± 9.50 | 0.16 | 0.93 | 76.98 ± 1.19 | 73.30 ± 2.75 | -0.24 | 0.75 |
| 1178251 | 5 | 1.77 ± 0.11 | 2.25 ± 0.34 | 0.62 | 0.55 | 1.34 ± 0.12 | 1.07 ± 0.15 | -0.37 | 0.76 | 1.77 ± 0.11 | 2.69 ± 0.11 | 0.51 | 0.44 |
| 1137850 | 0 | 88.34 ± 5.03 | 118.68 ± 1.26 | 0.58 | 0.43 | 51.68 ± 4.67 | 219.35 ± 35.84 | 2.03 | 0.00 | 88.34 ± 5.03 | 26.41 ± 1.27 | -1.91 | 0.00 |
| 1126790 | 1 | 37.23 ± 1.28 | 47.80 ± 1.56 | 0.53 | 0.52 | 54.66 ± 1.84 | 26.97 ± 3.30 | -1.07 | 0.00 | 37.23 ± 1.28 | 55.51 ± 1.16 | 0.44 | 0.46 |
| 1141368 | 0 | 283.79 ± 3.09 | 367.47 ± 40.06 | 0.52 | 0.55 | 350.33 ± 11.21 | 420.75 ± 115.76 | 0.20 | 0.91 | 283.79 ± 3.09 | 254.87 ± 14.52 | -0.33 | 0.63 |
| 1164596 | 4 | 196.19 ± 1.18 | 249.19 ± 20.70 | 0.50 | 0.59 | 75.44 ± 5.94 | 113.96 ± 9.73 | 0.53 | 0.41 | 196.19 ± 1.18 | 148.70 ± 9.82 | -0.56 | 0.28 |
| 1146389 | 1 | 108.54 ± 4.88 | 138.31 ± 19.36 | 0.50 | 0.59 | 116.63 ± 2.51 | 186.06 ± 4.83 | 0.63 | 0.21 | 108.54 ± 4.88 | 108.48 ± 6.72 | -0.15 | 0.86 |
| 1082759 | 1 | 87.33 ± 5.64 | 99.05 ± 0.51 | 0.34 | 0.81 | 64.14 ± 0.23 | 68.81 ± 11.91 | 0.05 | 0.98 | 87.33 ± 5.64 | 58.61 ± 6.89 | -0.72 | 0.11 |
| 1125366 | 4 | 132.02 ± 17.77 | 146.93 ± 8.62 | 0.31 | 0.84 | 37.41 ± 1.48 | 109.57 ± 0.12 | 1.50 | 0.00 | 132.02 ± 17.77 | 75.85 ± 6.71 | -0.92 | 0.02 |
| 1019516 | 2 | 144.19 ± 12.89 | 161.84 ± 26.21 | 0.31 | 0.85 | 192.89 ± 0.71 | 218.35 ± 9.92 | 0.12 | 0.95 | 144.19 ± 12.89 | 135.13 ± 13.09 | -0.23 | 0.76 |
| 1104917 | 0 | 188.49 ± 34.04 | 207.27 ± 13.58 | 0.30 | 0.86 | 273.96 ± 4.26 | 290.30 ± 37.78 | 0.02 | 0.99 | 188.49 ± 34.04 | 106.56 ± 2.73 | -0.97 | 0.02 |
| 1121445 | 11 | 57.61 ± 1.74 | 63.01 ± 2.36 | 0.28 | 0.87 | 72.90 ± 0.47 | 104.82 ± 12.52 | 0.46 | 0.54 | 57.61 ± 1.74 | 37.15 ± 4.55 | -0.80 | 0.05 |
| 40744 | 3 | 34.08 ± 1.51 | 37.29 ± 3.56 | 0.27 | 0.88 | 132.37 ± 11.64 | 40.86 ± 6.76 | -1.76 | 0.00 | 34.08 ± 1.51 | 82.38 ± 0.63 | 1.11 | 0.00 |
| 1178940 | 0 | 194.27 ± 13.70 | 213.06 ± 21.16 | 0.27 | 0.88 | 664.99 ± 5.60 | 612.42 ± 39.79 | -0.16 | 0.93 | 194.27 ± 13.70 | 393.00 ± 37.13 | 0.85 | 0.05 |
| 1113466 | 2 | 338.24 ± 10.36 | 365.49 ± 48.70 | 0.24 | 0.91 | 455.53 ± 13.84 | 401.65 ± 53.28 | -0.24 | 0.88 | 338.24 ± 10.36 | 423.61 ± 7.25 | 0.12 | 0.89 |
| 1156700 | 5 | 98.40 ± 3.12 | 101.76 ± 3.19 | 0.20 | 0.93 | 434.89 ± 16.18 | 257.81 ± 44.29 | -0.80 | 0.07 | 98.40 ± 3.12 | 225.43 ± 20.94 | 1.03 | 0.00 |
| 1165578 | 12 | 110.33 ± 11.86 | 112.62 ± 0.55 | 0.19 | 0.94 | 107.22 ± 2.38 | 158.33 ± 12.11 | 0.51 | 0.44 | 110.33 ± 11.86 | 53.59 ± 0.79 | -1.20 | 0.00 |
| 1120082 | 5 | 146.10 ± 6.70 | 142.51 ± 2.09 | 0.13 | 0.96 | 149.21 ± 1.74 | 172.49 ± 17.22 | 0.16 | 0.93 | 146.10 ± 6.70 | 142.19 ± 0.58 | -0.18 | 0.82 |
| 1121621 | 12 | 509.21 ± 69.19 | 497.06 ± 41.68 | 0.13 | 0.96 | 287.69 ± 2.48 | 366.00 ± 94.79 | 0.29 | 0.82 | 509.21 ± 69.19 | 503.96 ± 74.66 | -0.18 | 0.83 |
| 1147788 | 5 | 82.76 ± 8.14 | 76.08 ± 3.74 | 0.06 | 0.99 | 93.44 ± 3.39 | 136.89 ± 28.41 | 0.51 | 0.50 | 82.76 ± 8.14 | 72.84 ± 1.82 | -0.34 | 0.62 |
| 1092996 | 4 | 108.82 ± 13.42 | 100.63 ± 3.30 | 0.04 | 0.99 | 103.47 ± 3.68 | 121.62 ± 8.56 | 0.18 | 0.92 | 108.82 ± 13.42 | 92.85 ± 0.01 | -0.39 | 0.54 |
| 1163949 | 0 | 29.36 ± 1.78 | 26.80 ± 1.03 | 0.02 | 0.99 | 35.36 ± 1.44 | 32.46 ± 1.63 | -0.17 | 0.93 | 29.36 ± 1.78 | 27.31 ± 0.94 | -0.27 | 0.71 |
| 1141295 | 2 | 197.00 ± 8.52 | 180.50 ± 1.00 | 0.01 | 1.00 | 112.86 ± 4.46 | 136.16 ± 7.77 | 0.20 | 0.91 | 197.00 ± 8.52 | 150.75 ± 3.28 | -0.55 | 0.29 |
| 1103501 | 14 | 140.87 ± 5.80 | 128.08 ± 13.18 | 0.01 | 1.00 | 96.46 ± 6.34 | 206.50 ± 33.35 | 1.05 | 0.00 | 140.87 ± 5.80 | 3.30 ± 0.06 | -5.56 | 0.00 |

Table 3. Continued

| proteinid | tmhmm | NW305 -Fe.a | NW186 -Fe.a | log2FC | FDR | NW186 +Fe.a | NW186 +Fe.c | log2FC | FDR | NW305 -Fe.a | NW305 ++Fe.a | log2FC | FDR |
|-----------|-------|------------------|------------------|--------|------|-----------------|------------------|--------|------|------------------|-----------------|--------|------|
| 1172271 | 1 | 134.47 ± 5.07 | 121.28 ± 1.70 | 0.00 | 1.00 | 66.79 ± 2.94 | 82.65 ± 12.41 | 0.24 | 0.87 | 134.47 ± 5.07 | 107.55 ± 1.94 | -0.47 | 0.42 |
| 1219631 | 0 | 1.92 ± 0.37 | 1.56 ± 0.08 | -0.01 | 1.00 | 4.76 ± 0.47 | 5.85 ± 1.00 | 0.28 | 0.85 | 1.92 ± 0.37 | 1.64 ± 0.37 | -0.30 | 0.72 |
| 177364 | 1 | 2.82 ± 0.18 | 2.61 ± 0.00 | -0.03 | 0.99 | 1.88 ± 0.07 | 1.87 ± 0.32 | 0.01 | 1.00 | 2.82 ± 0.18 | 1.46 ± 0.05 | -1.04 | 0.03 |
| 1137953 | 2 | 40.68 ± 3.74 | 34.00 ± 3.57 | -0.11 | 0.97 | 141.12 ± 2.32 | 63.58 ± 6.84 | -1.21 | 0.00 | 40.68 ± 3.74 | 85.18 ± 6.98 | 0.92 | 0.02 |
| 1117050 | 3 | 40.36 ± 1.55 | 32.08 ± 4.00 | -0.16 | 0.95 | 46.35 ± 3.98 | 53.35 ± 4.46 | 0.15 | 0.94 | 40.36 ± 1.55 | 21.28 ± 0.77 | -1.07 | 0.00 |
| 1141371 | 0 | 1571.19 ± 246.87 | 1249.67 ± 120.07 | -0.17 | 0.95 | 551.49 ± 6.68 | 518.11 ± 5.14 | -0.15 | 0.94 | 1571.19 ± 246.87 | 740.57 ± 40.61 | -1.22 | 0.00 |
| 1147568 | 4 | 7482.49 ± 509.41 | 5782.62 ± 105.11 | -0.22 | 0.92 | 3164.92 ± 14.66 | 3994.04 ± 231.82 | 0.28 | 0.84 | 7482.49 ± 509.41 | 6622.43 ± 20.86 | -0.36 | 0.60 |
| 1181871 | 12 | 49.66 ± 1.89 | 36.63 ± 0.94 | -0.29 | 0.87 | 37.12 ± 0.83 | 30.34 ± 6.81 | -0.36 | 0.74 | 49.66 ± 1.89 | 35.43 ± 0.48 | -0.65 | 0.16 |
| 1209512 | 0 | 50.80 ± 5.25 | 36.89 ± 3.37 | -0.30 | 0.85 | 89.37 ± 2.45 | 60.30 ± 0.80 | -0.61 | 0.25 | 50.80 ± 5.25 | 87.27 ± 4.03 | 0.62 | 0.20 |
| 1011311 | 0 | 55.14 ± 2.22 | 38.64 ± 6.57 | -0.36 | 0.79 | 119.93 ± 7.34 | 98.35 ± 0.78 | -0.34 | 0.76 | 55.14 ± 2.22 | 88.47 ± 3.26 | 0.55 | 0.28 |
| 1168387 | 0 | 60.92 ± 2.05 | 40.64 ± 0.63 | -0.45 | 0.66 | 91.23 ± 0.81 | 80.83 ± 10.18 | -0.22 | 0.89 | 60.92 ± 2.05 | 59.55 ± 5.52 | -0.22 | 0.78 |
| 1141800 | 1 | 43.05 ± 0.75 | 25.30 ± 1.81 | -0.61 | 0.36 | 120.51 ± 4.34 | 73.38 ± 12.23 | -0.79 | 0.07 | 43.05 ± 0.75 | 59.09 ± 1.45 | 0.35 | 0.60 |
| 1169809 | 12 | 113.09 ± 7.38 | 58.75 ± 7.85 | -0.79 | 0.14 | 195.35 ± 3.82 | 323.04 ± 63.23 | 0.66 | 0.24 | 113.09 ± 7.38 | 140.31 ± 24.63 | 0.18 | 0.83 |
| 1164227 | 2 | 43.66 ± 2.41 | 19.30 ± 1.11 | -1.01 | 0.01 | 124.93 ± 0.50 | 30.60 ± 1.03 | -2.08 | 0.00 | 43.66 ± 2.41 | 100.11 ± 12.83 | 1.04 | 0.00 |
| 1148364 | 0 | 66.99 ± 4.34 | 29.59 ± 0.47 | -1.02 | 0.01 | 19.77 ± 1.34 | 31.72 ± 2.23 | 0.60 | 0.35 | 66.99 ± 4.34 | 21.58 ± 1.63 | -1.75 | 0.00 |
| 1222809 | 0 | 172.06 ± 39.87 | 70.73 ± 9.77 | -1.10 | 0.05 | 52.51 ± 0.89 | 74.55 ± 15.51 | 0.44 | 0.61 | 172.06 ± 39.87 | 14.50 ± 1.37 | -3.74 | 0.00 |
| 1115499 | 0 | 41.79 ± 0.54 | 14.21 ± 0.30 | -1.40 | 0.00 | 25.87 ± 1.86 | 19.65 ± 1.58 | -0.44 | 0.59 | 41.79 ± 0.54 | 30.57 ± 3.99 | -0.60 | 0.24 |
| 1146086 | 1 | 73.65 ± 8.51 | 23.91 ± 0.59 | -1.47 | 0.00 | 61.61 ± 1.55 | 63.36 ± 9.68 | 0.00 | 1.00 | 73.65 ± 8.51 | 26.66 ± 0.81 | -1.62 | 0.00 |
| 37369 | 0 | 0.03 ± 0.02 | - ± - | #N/A | #N/A | 0.03 ± 0.02 | 0.02 ± 0.01 | #N/A | #N/A | 0.03 ± 0.02 | - ± - | #N/A | #N/A |
| 1178960 | 0 | 0.23 ± 0.05 | 0.46 ± 0.02 | #N/A | #N/A | 0.71 ± 0.07 | 0.50 ± 0.00 | #N/A | #N/A | 0.23 ± 0.05 | 0.73 ± 0.11 | #N/A | #N/A |

to different media conditions. These data reconfirm that the *A. niger* transporter protein identified as top most likely citrate exporter candidate (Table 3) is indeed a citrate exporter. When initially making our findings publicly available (Odoni et al. 2018) we named this newly identified *A. niger* citrate exporter 'CitT' (Citrate Transporter). However, Steiger et al., who independently recognised this transporter as *A. niger* citrate exporter using a homology approach, argue that our name for this exporter can lead to confusion with the *citT* gene in *E. coli*, which encodes a citrate/succinate antiporter (Steiger et al. 2019). Therefore, adhering to the nomenclature given by (Steiger et al. 2019), we refer to this transporter protein as CexA.

DISCUSSION

Combining the expression values from our experimental setups with the complementary *in silico* homology approach ultimately led to the identification of the *A. niger* citrate exporter CexA (Table 3). Notably, the only other successful approach to identify the *A. niger* citrate exporter thus far was also via a homology approach, using an itaconate transporter from *Ustilago Maydis* as template (Steiger et al. 2019). The rationale behind using this template was that itaconate and citrate share a considerable amount of chemical similarity to justify transport via the same system (Steiger et al. 2019). This rationale proved to be successful, and we have also previously advocated the use of sequence similarity to identify new transporters of a given substrate of interest (Sloothaak et al. 2015, 2016). However, these approaches rely on the availability of the appropriate templates for any given substrate, which might not always be the case (Sloothaak et al. 2015).

Here, we used a homology approach, which we specifically conceived for the scenario in which it is not known which transporters are responsible for citrate export in the organisms used for comparison. Our only assumption was that the presence of a transporter performing this function is either more (*A. kawachii*, *Y. lipolytica*) or less (*A. flavus*, *A. terreus*, *S. cerevisiae*) likely. However, these organisms might share a variety of homologous proteins. Thus, sorting the list of putative citrate transporter proteins based on their log₂FC values in the experimental setup in which we hypothesise that differences in citrate secretion are crucial for survival of the fungus was key (Table 3).

Interestingly, CexA has not been listed as putative citrate exporter candidate using a transcriptomics approach in the citrate producing strain H915-1 (Yin et al. 2017), although *cexA* (An17g01710) gene expression was found to be highly induced by sucrose (Yuan et al. 2008). This suggests different transcriptional regulation of citrate secretion under different citrate producing conditions. Indeed, the contrasting transcriptomic landscapes of enzymes involved in citrate metabolism (Fig. 2) imply very distinct molecular mechanisms underlying the phenotype of increased citrate accumulation in the two experimental setups used for this study.

In the supplement experiment, we observed a higher glucose consumption rate when NW186 was supplemented with excess citrulline (Fig. 1), although this was not reflected in up-regulation of glycolysis at transcript level (Supplementary Data 2, Supporting Information). A high glucose consumption rate results in high glycolytic flux that will lead to excess NADH, the turn-over of which might be limited by the capacity of further biosynthetic processes (Karaffa and Kubicek 2003). The observed down-regulation of TCA cycle enzymes downstream of citrate in NW186 +Fe.c compared to NW186 +Fe.a prevents citrate to

be further metabolised, which might be a mechanism to prevent further generation of NADH (Gallmetzer and Burgstaller 2002). This mechanism represents citrate production via overflow metabolism in its most classical form (Legiša and Matthey 2007).

Aspergillus niger strains used in academia and industry are subject to conditions they would not usually encounter in nature, and the surplus of glucose or another carbon source encountered by the fungus would almost always suggest citrate production via overflow metabolism. Given these conditions, it has been reported that alternative oxidase (also non-electrogenic ubiquinol oxidase, or Aox1) is required for efficient *A. niger* citrate production to avoid excess production of ATP (Ruijter, Kubicek and Visser 2002). Aox1 in *A. niger* effectively decouples ATP generation and NADH re-oxidation, as the reduction of molecular oxygen to water by alternative oxidase bypasses proton translocation via the oxidative phosphorylation complexes III and IV, resulting in a lower ATP yield (Joseph-Horne, Hollomon and Wood 2001).

As expected in a condition of metabolic overflow, up-regulation of *aox1* was accompanied with down-regulation of cytochrome-dependent respiratory enzymes when supplementing NW186 with excess citrulline, indicating a switch from cytochrome-dependent respiration to alternative respiration (Karaffa and Kubicek 2003). However, in the iron experiment, the observed up-regulation of *aox1* was not accompanied with down-regulation of cytochrome-dependent respiratory enzymes (Supplementary Data 2, Supporting Information), possibly indicating that no such switch takes place. In a different study, it was found that incubating detached roots of the plant *Poa annua* with citrate increased protein concentration of alternative oxidase without actually increasing the activity of the alternative respiration pathway itself (Millenaar 2002). The authors of the study hypothesised that the chelating properties of citrate might lead to the withdrawal of the Fe in the active centre of alternative oxidase, thereby rendering the protein inactive and evoking increased transcription to compensate for the inactive protein. In a similar line of reasoning, the transcriptional increase of *aox1* observed in the iron experiment might be a futile attempt to counteract high intracellular accumulation of citrate preceding its secretion in NW186 -Fe.a.

Due to the chelating properties of citrate and other organic acids, we have previously discussed up-regulation of *A. niger* citrate biosynthesis and secretion as 'active' response to overcome iron limitation (Odoni et al. 2017). Remarkably, CexA, which seems to be the only active citrate exporter based on the inability of *A. niger cexA* KO mutants to secrete citrate (Steiger et al. 2019), is a major facilitator protein belonging to the subfamily of H⁺-drug antiporters. In plants, these transporters are used for (trace) metal homeostasis (Magalhaes 2010; Remy and Duque 2014), relating these findings back to our notion that a probable biological role for citrate is to act as an *A. niger* iron siderophore. Another conceivable role for citrate is to aid in the hydrolysis of sucrose to glucose and fructose, a process citrate can be used for (Lowe 1937). Again, the need to use citrate to fulfil a biological function extracellularly offers a possible explanation why sucrose is such a good carbon source for triggering *A. niger* citrate secretion (Hossain, Brooks and Maddox 1984), and also why *cexA* (An17g01710) gene expression was highly induced on sucrose (Yuan et al. 2008).

SUPPLEMENTARY DATA

Supplementary data are available at [FEMSLE](https://academic.oup.com/femsle/article/366/7/fnz071/5437674) online.

ACKNOWLEDGEMENTS

We would like to thank Ruben van Heck, Thanaporn Laothanachareon, Naveed Ishaque and Matthias Schlesner for their involvement in shaping this manuscript.

FUNDING

This work was supported by a grant in the framework of the BE-BASIC program F01.011 Transport processes in the production of organic acids by *Aspergillus niger*, and the WUR IPOP Systems Biology program KB-17-003.02.026 Genome-wide metabolic modelling and data integration of organic acid production in filamentous fungi.

Conflict of interest. None declared.

REFERENCES

- Alvarez-Vasquez F, González-Alcón C, Torres NV. Metabolism of citric acid production by *Aspergillus niger*: model definition, steady-state analysis and constrained optimization of citric acid production rate. *Biotechnol Bioeng* 2000;**70**:82–108.
- Andersen MR, Salazar MP, Schaap PJ et al. Comparative genomics of citric-acid-producing *Aspergillus niger* ATCC 1015 versus enzyme-producing CBS 513.88. *Genome Res* 2011;**21**:885–97.
- Arnaud MB, Cerqueira GC, Inglis DO et al. The *Aspergillus* Genome Database (AspGD): recent developments in comprehensive multispecies curation, comparative genomics and community resources. *Nucleic Acids Res* 2012;**40**:D653–9.
- Bateman A, Martin MJ, O'Donovan C et al. UniProt: a hub for protein information. *Nucleic Acids Res* 2015;**43**:D204–12.
- Chesters CG, Rolinson GN. Zinc in the metabolism of a strain of *Aspergillus niger*. *J Gen Microbiol* 1951;**5**:553–8.
- Currie JN. The citric acid fermentation of *Aspergillus niger*. *J Biol Chem* 1917;**31**:15–37.
- Dujon B, Sherman D, Fischer G et al. Genome evolution in yeasts. *Nature* 2004;**430**:35–44.
- Fedorova ND, Khaldi N, Joardar VS et al. Genomic islands in the pathogenic filamentous fungus *Aspergillus fumigatus*. *PLoS Genet* 2008;**4**:e1000046.
- Futagami T, Mori K, Yamashita A et al. Genome announcement: genome sequence of the white Koji mold *Aspergillus kawachii* IFO 4308, used for brewing the Japanese distilled spirit shochu. *Eukaryot Cell* 2011;**10**:1586–7.
- Galagan JE, Calvo SE, Cuomo C et al. Sequencing of *Aspergillus nidulans* and comparative analysis with *A. fumigatus* and *A. oryzae*. *Nature* 2005;**438**:1105–15.
- Gallmetzer M, Burgstaller W. Efflux of organic acids in *Penicillium simplicissimum* is an energy-spilling process, adjusting the catabolic carbon flow to the nutrient supply and the activity of catabolic pathways. *Microbiology* 2002;**148**:1143–9.
- Hossain M, Brooks JD, Maddox IS. The effect of the sugar source on citric acid production by *Aspergillus niger*. *Appl Microbiol Biotechnol* 1984;**19**:393–7.
- Joardar V, Abrams NF, Hostetler J et al. Sequencing of mitochondrial genomes of nine *Aspergillus* and *Penicillium* species identifies mobile introns and accessory genes as main sources of genome size variability. *BMC Genomics* 2012;**13**:698.
- Joseph-Horne T, Hollomon DW, Wood PM. Fungal respiration: a fusion of standard and alternative components. *Biochim Biophys Acta Bioenerg* 2001;**1504**:179–95.
- Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res* 2000;**28**:27–30.
- Kanehisa M, Sato Y, Kawashima M et al. KEGG as a reference resource for gene and protein annotation. *Nucleic Acids Res* 2015;**44**:D457–62.
- Karaffa L, Kubicek CP. *Aspergillus niger* citric acid accumulation: do we understand this well working black box? *Appl Microbiol Biotechnol* 2003;**61**:189–96.
- Krogh A, Larsson B, Von Heijne G et al. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol* 2001;**305**:567–80.
- Legiša M, Mattey M. Changes in primary metabolism leading to citric acid overflow in *Aspergillus niger*. *Biotechnol Lett* 2007;**29**:181–90.
- Lenouvel F, van de Vondervoort PJI, Visser J. Disruption of the *Aspergillus niger argB* gene: a tool for transformation. *Curr Genet* 2002;**41**:425–31.
- Lowe B. Effect of acid upon sugars and hydrolysis of sugars. In: *Experimental Cookery, from the Chemical and Physical Standpoint*. New York: John Wiley & Sons, 1937.
- Magalhaes JV. How a microbial drug transporter became essential for crop cultivation on acid soils: aluminium tolerance conferred by the multidrug and toxic compound extrusion (MATE) family. *Ann Bot* 2010;**106**:199–203.
- Mascorro-Gallardo JO, Covarrubias AA, Gaxiola R. Construction of a CUP1 promoter-based vector to modulate gene expression in *Saccharomyces cerevisiae*. *Gene* 1996;**172**:169–70.
- Millenaar FF. Role of sugars and organic acids in regulating the concentration and activity of the alternative oxidase in *Poa annua* roots. *J Exp Bot* 2002;**53**:1081–8.
- Netik A, Torres N V., Riol JM et al. Uptake and export of citric acid by *Aspergillus niger* is reciprocally regulated by manganese ions. *Biochim Biophys Acta Biomembr* 1997;**1326**:287–94.
- Nierman WC, Pain A, Anderson MJ et al. Genomic sequence of the pathogenic and allergenic filamentous fungus *Aspergillus fumigatus*. *Nature* 2005;**438**:1151–6.
- Nordberg H, Cantor M, Dusheyko S et al. The genome portal of the department of energy Joint genome institute: 2014 updates. *Nucleic Acids Res* 2014;**42**:D26–31.
- Odoni DI, Laothanachareon T, Vazquez-vilar M et al. *Aspergillus niger* citrate exporter revealed by comparison of two alternative citrate producing conditions. *bioRxiv* 2018:1–36.
- Odoni DI, van Gaal MP, Schonewille T et al. *Aspergillus niger* secretes citrate to increase iron bioavailability. *Front Microbiol* 2017;**8**:1424.
- Remy E, Duque P. Beyond cellular detoxification: a plethora of physiological roles for MDR transporter homologs in plants. *Front Physiol* 2014;**5** MAY:201.
- Ruijter GJG, Bax M, Patel H et al. Mannitol is required for stress tolerance in *Aspergillus niger* conidiospores. *Eukaryot Cell* 2003;**2**:690–8.
- Ruijter GJG, Kubicek CP, Visser J. Production of organic acids by fungi. In: *Industrial Applications*. Berlin, Heidelberg: Springer Berlin Heidelberg, 2002, 213–30.
- Ruijter GJG, Van De Vondervoort PJI, Visser J. Oxalic acid production by *Aspergillus niger*: an oxalate-non-producing mutant produces citric acid at pH 5 and in the presence of manganese. *Microbiology* 1999;**145**:2569–76.
- Sloothaak J, Odoni DI, De Graaff LH et al. *Aspergillus niger* membrane-associated proteome analysis for the identification of glucose transporters. *Biotechnol Biofuels* 2015;**8**:150.
- Sloothaak J, Tamayo-Ramos JA, Odoni DI et al. Identification and functional characterization of novel xylose transporters from the cell factories *Aspergillus niger* and *Trichoderma reesei*. *Biotechnol Biofuels* 2016, DOI: 10.1186/s13068-016-0564-4.

- Sonnhammer ELL, Von Heijne G, Krogh A. A hidden Markov model for predicting transmembrane helices in protein sequences. 1998;6:175–82.
- Steiger MG, Rassinger A, Mattanovich D et al. Engineering of the citrate exporter protein enables high citric acid production in *Aspergillus niger*. *Metab Eng* 2019;52:224–31.
- Suga M, Hatakeyama T. High-efficiency electroporation by freezing intact yeast cells with addition of calcium. *Curr Genet* 2003;43:206–11.
- van der Straat L, de Graaff LH. Pathway transfer in fungi: transporters are the key to success. *Bioeng Bugs* 2014;5:335–9.
- Yin X, Shin HD, Li J et al. Comparative genomics and transcriptome analysis of *Aspergillus niger* and metabolic engineering for citrate production. *Sci Rep* 2017;7:41040.
- Yuan XL, Roubos JA, Van Den Hondel CAMJJ et al. Identification of InuR, a new Zn(II)2Cys6 transcriptional activator involved in the regulation of inulinolytic genes in *Aspergillus niger*. *Mol Genet Genomics* 2008;279:11–26.