



Published in final edited form as:

Psychol Med. 2013 August ; 43(8): 1723–1734. doi:10.1017/S0033291712002516.

Remission, continuation, and incidence of eating disorders during early pregnancy: A validation study on a population-based birth cohort

Hunna J. Watson^{1,2,3}, Ann Von Holle⁴, Robert M. Hamer⁴, Cecilie Knoph Berg⁵, Leila Torgersen⁵, Per Magnus⁶, Camilla Stoltenberg^{6,7}, Patrick Sullivan^{4,8}, Ted Reichborn-Kjennerud^{5,9}, and Cynthia M. Bulik^{4,10,*}

¹Centre for Clinical Interventions, Department of Health in Western Australia, Australia

²Eating Disorders Program, Princess Margaret Hospital for Children, Department of Health in Western Australia, Australia

³School of Paediatrics and Child Health, The University of Western Australia, Australia

⁴Department of Psychiatry, University of North Carolina at Chapel Hill, USA

⁵Division of Mental Health, Norwegian Institute of Public Health, Norway

⁶Division of Epidemiology, Norwegian Institute of Public Health, Norway

⁷Department of Public Health and Primary Health Care, University of Bergen, Norway

⁸Department of Genetics, University of North Carolina at Chapel Hill, USA

⁹Department of Psychiatry, University of Oslo, Norway

¹⁰Department of Nutrition, University of North Carolina at Chapel Hill, USA

Abstract

Background—The objective of this study was to validate previously published rates of remission, continuation, and incidence of broadly defined eating disorders during pregnancy. The previous rate modeling was done by our group (Bulik *et al.* 2007) and yielded participants halfway into recruitment of the planned 100,000 pregnancies in the Norwegian Mother and Child (MoBa) Cohort at the Norwegian Institute of Public Health. This study aimed to internally validate the findings with the completed cohort.

Methods—77267 pregnant women enrolled at 17 weeks gestation between 2001 and 2009 were included. Participants were split into a “training” sample ($n=41243$) based on participants in the MoBa version 2 dataset of the original study and a “validation” sample ($n=36024$) comprising individuals in the MoBa version 5 dataset that were not in the original study (Bulik *et al.* 2007). Internal validation of all original rate models involved fitting a calibration model to compare model parameters between the “training” and “validation” samples as well as bootstrap estimates of bias in the entire version 5 dataset.

*Address for correspondence: Professor Cynthia Bulik, Department of Psychiatry, University of North Carolina at Chapel Hill, 101 Manning Drive, CB#7160, Chapel Hill, North Carolina 27599-7160. (cynthia_bulik@med.unc.edu).

Results—Remission, continuation, and incidence estimates from the “training” sample remained stable when evaluated via a split sample validation procedure. Pre-pregnancy prevalence estimates in the “validation” sample were 0.1% for anorexia nervosa, 1.0% for bulimia nervosa (BN), 3.3% for binge eating disorder (BED), and 0.1% for purging disorder (EDNOS-P). In early pregnancy, estimates were 0.2% for BN, 4.8% for BED, and <0.01% for EDNOS-P. Consistent with the original study, incident BN and EDNOS-P during pregnancy were rare. For BED, the adjusted incidence rate in the “validation” sample was 1.17 per 1000 person-weeks. The highest rates were for full or partial remission for BN and EDNOS-P, and continuation for BED.

Conclusions—This study provides evidence of validity of previously estimated rates of remission, continuation, and incidence of eating disorders during pregnancy. Eating disorders during pregnancy were relatively common, occurring in nearly 1 in every 20 women, although almost all were cases of BED. Pregnancy was a window of remission from BN but a window of vulnerability for onset and continuation of BED. Training to detect the signs and symptoms of eating disorders by obstetricians/gynecologists and interventions to enhance pregnancy and neonatal outcomes warrant attention.

Keywords

Eating disorders; epidemiology; incidence; pregnancy; prospective; remission; The Norwegian Mother and Child Cohort Study

Introduction

Eating disorders in pregnancy are poorly understood but warrant attention. Epidemiological data from our group suggest, first, that eating disorders during pregnancy are reasonably common, with prevalence estimates ranging between 0.1% (purging disorder, EDNOS-P) to 4.8% (binge eating disorder, BED) (Bulik *et al.* 2007). Second, pregnancy is a high-risk period for onset of BED, occurring at a rate of 1.1 per 1000 person-weeks (Bulik *et al.* 2007). Eating disorder symptoms during pregnancy are more prevalent among those with a recent or past history of eating disorders (Micali *et al.* 2007b). Over a quarter of pregnant women with eating disorders purge and 11% report dieting for weight loss at 32 weeks (Micali *et al.* 2007b). Offspring of women with eating disorders are at higher risk for birth complications including perinatal mortality, premature birth, low birth weight, and birth defects (Bulik *et al.* 1999, Micali *et al.* 2007a, Sollid *et al.* 2004). Persistence of eating disorders beyond pregnancy may increase child vulnerability through risk factors associated with the expression of illness. Mothers with and without eating disorders self-report different feeding styles, with restrictive feeding styles and infant feeding problems more common among mothers with eating disorders characterized by binge eating (Reba-Harrelson *et al.* 2010). Indeed, mothers with eating disorders express concern about knowing how to feed their children appropriately (Mazzeo *et al.* 2005). Positively, pregnancy appears to be a “window” for remission of bulimia nervosa (BN) (Bulik *et al.* 2007). Recognition of eating disorders before and during pregnancy is a necessary first step to engaging individuals in treatment and parenting-based interventions to improve eating disorder and obstetric outcomes and promote healthy child development.

Research on the course of eating disorders through pregnancy originated in the 1980s and has typically used retrospective reports or small prospective samples with clinical or at-risk participants (e.g., Blais *et al.* 2000, Crow *et al.* 2004, Koubaa *et al.* 2005, Lacey and Smith 1987, Rocco *et al.* 2005, Tiller and Treasure 1998). Although valuable clinical data had accumulated, an epidemiological perspective on the prevalence and course of eating disorders through pregnancy was absent, meaning there were no reliable estimates of the percentage of women who experience, develop, or remit from eating disorders during pregnancy, which is essential to guide research, health planning, and service provision.

To address this, we (Bulik *et al.* 2007) examined data from an ongoing prospective population-based pregnancy cohort study, which was approximately halfway toward the goal of recruiting 100000 pregnancies, the Norwegian Mother and Child Cohort Study (MoBa; Magnus *et al.* 2006). We provided estimates of eating disorder prevalence pre-pregnancy, and rates of incidence, continuation, and remission of eating disorders during pregnancy. Eating disorders, either new or continuing, were common. Six months prior to pregnancy, the prevalence was 0.1% for anorexia nervosa (AN), 0.7% for BN, 3.5% for BED, and 0.1% for EDNOS-P. During pregnancy, estimates were 0.2% (BN), 4.8% (BED), and 0.1% (EDNOS-P). A prominent and somewhat unexpected finding was the relatively high prevalence of BED onset (1.7%) which was more probable among women with lifetime and psychosocial adversities (Berg *et al.* 2011). Full or partial remission during pregnancy was the most common course for BN and EDNOS-P, but BED had a high continuance rate.

The findings of the study by our group (Bulik *et al.* 2007) are significant, yet unconfirmed, due primarily to methodological practicalities that impede progress in the external validation of these findings; notably, the need for a large population sample to ensure adequate statistical power. The first reason why it is important to replicate prevalence and course estimates is that it would be damaging to the community if health planning and policy was misaligned with community need on the basis of unreproducible scientific evidence, and ultimately disparaging to academic enterprise. Second, given the absence of widespread data on the prevalence of eating disorders in pregnancy, all large-scale field data at present have substantial implications for health research.

Hence, the purpose of this study was to internally validate the statistical modeling of incidence, remission and continuation used in the first study. Furthermore, it is important to note that validation of modeling is distinct from other types of validation such as validation of a data collection instrument. Internal validation of a statistical model entails an assessment of the ability of a certain model to accurately predict outcomes. In this particular case, we evaluate the performance of models predicting rates of incidence, remission and incidence across eating disorder subtypes.

We hypothesized that the models would internally validate given evidence of the stability of eating disorder prevalence in Norway (Zachrisson *et al.* 2008). Given the context, we chose a split sample approach with model recalibration to determine if observed estimates in the latter half of the sample were similar to predicted estimates. Outcomes from this analysis can provide evidence towards reproducibility of the original model and its findings – a critical yet rare process (Altman *et al.* 2009). At the time of publication of the original

estimates (Bulik *et al.* 2007), data collection from the cohort sample was incomplete. The MoBa goal of recruitment of more than 100000 pregnancies (recruited from 1999 to 2009) is now completed enabling us to conduct a statistical approach to validation.

Method

Participants

This study is nested within the MoBa study, which is conducted by the Norwegian Institute of Public Health (Magnus *et al.* 2006). The total sample comprised 77267 pregnant women with valid MoBa data (version 5, release May 2010). Participants were split into a “training” sample ($n=41243$) based on participants in the MoBa version 2 dataset (released April 2006) of the original study and a “validation” sample ($n=36024$) comprising individuals in the MoBa version 5 dataset who were not in the original study (Bulik *et al.* 2007) (Fig. 1). The total sample is less than the overall MoBa cohort as inclusion criteria (below) were necessary to enhance internal validity; participant flow is depicted in Figure 1. The split approximately halved the cohort across time creating in essence a temporal validation. It should also be noted that the “training” sample and the “validation” sample are recruited from overlapping but to some extent different parts of the total population of Norway.

The Medical Birth Registry of Norway (MBRN) monitors trends in birth and administrates a complete nationwide registry with consecutive registration of all births with gestational age > 16 weeks since 1967; notification of births to MBRN is compulsory for physicians and midwives. MoBa is a nationwide prospective population-based pregnancy cohort study that recruited pregnant women via postal invitation before an ultrasound appointment in week 17–18 of pregnancy in Norway between 1999 to 2009 (Magnus *et al.* 2006), and 38.5% of invited women consented to participate (<http://www.fhi.no/moba-en>). The cohort now includes 108000 children, 90700 mothers and 71500 fathers. Approval for this research was granted by appropriate regional committees, the Norwegian Data Inspectorate, and the Institutional Review Board of the University of North Carolina at Chapel Hill.

Inclusion criteria for this study were women with a first pregnancy during the study period, singleton birth, and live birth. Exclusion criteria were a missing pregnancy ID precluding data linking, completion of the pilot version of the questionnaire, weight <30 or >300 kg before and during pregnancy, height <1 m, women who returned the MoBa survey after birth, missing responses precluding assessment of eating disorder caseness, and a missing age value.

Measures

The Norwegian Mother and Child Cohort Study (MoBa) Questionnaire 1 [<http://www.fhi.no/dokumenter/1f32a49514.pdf>] contained items on eating disorders and behaviors that were previously used for studies of eating disorders in the Norwegian Institute of Public Health Twin Panel (Harris *et al.* 2002, Reichborn-Kjennerud *et al.* 2003, Reichborn-Kjennerud *et al.* 2004a, Reichborn-Kjennerud *et al.* 2004b). Items were designed to operationalize *DSM-IV* criteria for AN, BN, and eating disorders not otherwise specified (American Psychiatric Association 1994). Questions for binge eating addressed eating an

unusually large amount of food with an accompanying sense of loss of control and respondents were instructed to distinguish between pregnancy-related nausea and vomiting and self-induced vomiting as a compensatory method. Respondents included in this study completed questionnaire 1 at a median of 17.1 weeks gestation (inter-quartile range 15.9–18.9 weeks and range 4.0–42.1 weeks).

Eating disorder classifications—Algorithms in the original study (Bulik *et al.* 2007) constructed from Questionnaire 1 items were used to define eating disorder diagnoses: broadly defined AN, defined as meeting *DSM-IV* criteria for AN (with the exception of amenorrhea and also endorsing a body mass index (BMI) < 19.0 kg/m² at the time of low weight); broadly defined BN (endorsing at least weekly frequency of binge eating and purging and categorized as BN any type, BN purging type, BN non-purging type); broadly defined BED (binge eating at least weekly in the absence of compensatory behaviors); and EDNOS-P (purging at least weekly in the absence of binge eating). Due to practical difficulties in determining low weight in the presence of pregnancy-related weight gain, AN was assessed prior to pregnancy only. BN, BED, and EDNOS-P were assessed for both six months prior to pregnancy (retrospective assessment) and at the time of survey completion. Self-reported weight and height were used to calculate pre-pregnancy BMI and BMI at the time of assessment.

Diagnostic classifications pre-pregnancy comprised the categories of AN, BN purging type, BN non-purging type, BN any type, BED, EDNOS-P, and “missing”. These classifications were also applied during pregnancy, with the exception of AN due to the difficulties noted earlier.

If an individual had a missing response on one criterion but scored positively on all other criteria for a diagnosis, a classification of missing was assigned, otherwise no eating disorder was indicated. The BN any type includes BN purging and non-purging types as well as an additional category of people; this category was assigned when individuals met criteria for BN including endorsing non-purging compensatory behaviors (i.e., fasting and exercise) but had missing values for the purging items (i.e., laxatives and self-induced vomiting).

Definition of remission, incidence, and continuation—Remission described individuals who experienced an eating disorder pre-pregnancy and had no eating disorder during pregnancy. For BN, remission described an absence of both binge eating and compensatory behaviors during pregnancy. Partial remission pertained to individuals with BN pre-pregnancy who reported ongoing binge eating but the absence of compensatory behaviors during pregnancy. Continuation described individuals who presented with the same eating disorder pre- and during pregnancy. Incidence referred to onset of a broadly defined eating disorder when none was present in the six months prior to pregnancy.

Statistical analysis

We used two different methods to internally validate the statistical models used in the original analysis. The first method is model calibration, which provides information regarding accuracy of the predicted rates from the “training” sample. The second method is

the bootstrap estimate of bias, which indicates whether estimates from the entire version 5 dataset differ from the true population estimates. Given that the MoBa dataset does not include the entire population of Norwegian pregnancies between 1999 and 2009, estimate bias in the models is possible. In the original study (Bulik *et al.* 2007) regression models estimated rates of eating disorder remission, continuation, and incidence during pregnancy by diagnosis type, also, measures of association between sociodemographic measures and BED incidence were defined and discussed.

The split samples were used to calculate calibration statistics (Steyerberg 2009, Steyerberg *et al.* 2004). The calibration method is a multi-step procedure estimating the degree to which parameter estimates from a “training” sample predict observed values in the “validation” sample. Regression coefficients are first estimated using the original models with the “training” data. Next, a linear combination of those coefficients and any relevant covariables from the “validation” dataset — the linear predictor — comprise the only covariate in the final calibration model with response values from the “validation” dataset. Considering the specification of generalized linear regression models, an offset is also used with the linear predictor. For an intercept-only calibration, the updated alpha (α :intercept) should approximate zero in the final model thus providing evidence that the outcome value as predicted by the “training” sample is no different from the observed response value from the “validation” sample. For measures of rates, the α :intercept is reported for unadjusted results. Age-adjusted results in the original paper are a product of combinations of model coefficients and not eligible for the calibration procedure. For measures of association between sociodemographic variables and outcomes β_{overall} is reported, which ideally should approximate one in the final model. The unreliability [U]-statistic and Brier score, as described in Steyerberg (2004), indicate a measure of the overall model performance and miscalibration, respectively, in the context of the split sample validation design. The Brier score and U -statistic approximate zero if updated estimates provide good fit. Simulations prior to calibration determined that a sample size of at least 500 was required to obtain at least 80% power to detect a 20% difference in rates from models using the proposed validation method (i.e., calibration α statistically significantly different from zero). BED is the only eating disorder that meets this sample size criterion thus any failure to find a difference would be worthwhile noting for this particular eating disorder only.

A final measure of internal validity, the bootstrap estimate of bias (Efron and Tibshirani 1993, Good 2006), was applied to estimated rates for all eating disorder subtypes. The bootstrap sampling method generates samples with replacement from the original sample. We generated 1000 bootstrap samples and obtained model parameters from each sample. Rate estimates across the 1000 samples were averaged to obtain the mean, and the standard deviation of estimates was used to form the 95% confidence intervals (CIs) of the bootstrapped mean estimate. Estimate bias calculations are described; these are the original estimated mean in the observed sample subtracted from the bootstrap mean. A positive bias indicates the original estimate is under-estimating the true population value, conversely, a negative estimate indicates over-estimation of the parameter. Bias much greater than the standard error of the estimated mean may indicate poor estimation (Efron and Tibshirani 1993).

It should be noted that the models for EDNOS-P did not converge in more than 60% of the samples for the continuation and remission rates because of sparse cell counts. Incidence models frequently did not converge for BN purging and BN non-purging groups (68% and 38%, respectively). It should be noted that lack of convergence in this situation leaves at most 62% of the replications to form the estimated rates in the bootstrap estimate for those groups. The lack of convergence precluded any specification for the aforementioned combination of outcomes and groups in estimates presented in Table 6.

Data analysis was based on version 5 of the quality-assured MoBa data file released for research in 2010. The number of respondents in the “training” sample from MoBa version 5 dataset does not identically match those from the original version 2 dataset (Bulik *et al.* 2007). There were >1900 pregnancy identification numbers not linked to a unique maternal identification number in the version 2 dataset precluding identification of singleton births for these pregnancies; hence, these mothers were not incorporated into the original study sample.

Results

Sample demographics

Table 1 presents the sociodemographic composition of the “training” and “validation” samples. The “validation” sample appears to represent a more advantaged group with close to 10 point higher proportions with >4 years of university education and the two highest income thresholds, and had elevated primiparity (64.4% versus 48.1%) and cohabitation (54.5% versus 47.4%).

Eating disorder prevalence

The prevalence of eating disorders before and during pregnancy, and frequencies of remission, incidence, and continuation are shown in Table 2. In the 6 months before pregnancy, the prevalence ranged from 0.1% for AN to 3.5% for broad BED across both the “training” and “validation” samples. In both samples, the most common course of illness was continuation for BED (training: 62%, validation: 60%, total (data not shown): 61%), remission and/or partial remission for BN (training: 69%, validation: 78%, total: 74%), and remission for EDNOS-P (training: 79%, validation: 80%, total: 79%). Eating disorders during pregnancy were relatively common (occurring in 1 in every 21 women), though these were mainly made up of BED (1 in 23 women).

Remission, continuation, and incidence

Table 3 shows age-adjusted rates of remission, incidence, and continuation, by sample across eating disorder subtypes. The remission rate was highest for EDNOS-P, followed by BED broad, and BN in both the “training” and “validation” samples. The incidence rate was highest for broad BED, at 1.22 (95% CI=1.14–1.31) and 1.17 (95% CI=1.09–1.27) per 1000 person-weeks for the “training” and “validation” samples, respectively.

Validation models

The validation analysis quantified differences between the split samples by comparing the observed rates in the “validation” sample with the predicted rates from the “training” sample (Table 4). A positive alpha estimate in the calibration model indicates that the observed rate in the “validation” sample is higher than the predicted rate from the “training” sample. For example, the alpha for BED remission is 0.04 (95% CI=−0.03–0.12) indicating the observed rate in the “validation” sample is 4 percent higher than the rate in the “training” sample. According to the calibration estimates for BED (see Table 4), the “validation” sample had higher rates of favorable outcomes (remission) and lower rates of unfavorable outcomes (continuation and incidence) relative to the “training” sample. The effects were mixed for BN (lower continuation rates and higher incidence in the “validation” sample) and opposite effects were observed for EDNOS-P. All CIs spanned zero, with one exception for BN continuation, thus inferring no statistically significant differences in predicted and observed rates. Brier scores and *U*-statistics were close to zero, indicating good model performance. Overall, the rate models were well-calibrated and internally valid, as hypothesized (see Table 4).

Characteristics associated with incidence

Exploratory analysis in the original paper (Bulik *et al.* 2007) included measures of association between sociodemographic predictors and BED incidence. These measures were estimated for the “training” and “validation” samples and the differences between those estimates quantified (Table 5). The differences are not inconsequential as they exceed a 20% difference in the “validation” versus the “training” sample for half of the predictors including ‘ever smoke?’ (21% higher), ‘infertility treatment’ (34% lower), ‘minimum combined income’ (38% lower), and ‘total live births’ (60% lower). All 95% CIs for the β_{overall} estimate span one discounting any evidence that estimates from the “validation” sample differ from the “training” sample; there is one exception, which is “total live births”. The “validation” sample indicates a 60% lower estimated association ($\beta_{\text{overall}}=0.40$, 95% CI=−0.06–0.85) with incidence in the “validation” sample than predictions based on the “training” sample. In the “validation” sample the estimates indicate lower incidence for the nulliparous relative to women with 2+ live births, and the strength of that association is about half that of the “training” sample (−0.23 v. −0.41).

Bias estimates of rates from sample

The bootstrap age-adjusted estimates of remission, continuation, and incidence are shown in Table 6 with bias estimates (original estimated mean in the total dataset subtracted from the bootstrap mean). Negative bias suggests that the total MoBa cohort rate estimate is larger in the sample than a rate obtained from the entire population, and vice versa for positive bias estimates. Bias in almost all cases was negative. However, for some groups the standard errors were similar to the size of the bias indicating substantial variability and little evidence to distinguish from a bias estimate of zero. Some exceptions were for remission bias estimates indicating evidence for upward bias in the Moba cohort (i.e., overestimation), for BN purging (−0.29, *se*=0.09) for example. For continuation, mostly negative bias estimates exceed the standard error for BN any type (−0.10, *se*=0.08), BN purging (−0.28, *se*=0.10).

There is no evidence for substantial bias for any incidence estimates as the bias is <0.01 and standard errors exceed the estimated bias.

Discussion

Updating previously published models of remission, continuation, and incidence of eating disorders in pregnancy established by our group (Bulik *et al.* 2007) with a “validation” sample via a parsimonious calibration approach indicates validity of the original predictions. With only one exception, all observed rates in the “validation” sample did not differ from predicted rates in the “training” sample. This was consistent with our expectations, given evidence of stable prevalence of eating disorders in Norwegian adults (Zachrisson *et al.* 2008). Of note is that in spite of changes to the characteristics of the cohort over time (e.g., socioeconomic status, primiparity, cohabitation), the basic findings of the original study were unchanged, providing evidence of generalizability. In estimates of bias, the story is mixed with consistent negative bias suggesting more extreme estimates in the MoBa cohort than what might be found in the population. However, variability around these estimates does not provide conclusive evidence supporting this result. Lastly, validation of the exploratory aims estimating associations between sociodemographic predictors and BED incidence did not reveal significant departure from original estimates, but did indicate a level of variability around those estimates.

While validation was the focus of this study, some general findings are worthy of discussion. The prevalence of broadly defined eating disorders during pregnancy in the “validation” sample was 4.7%, comparable to the prevalence estimate of 4.8% in the approximate first half of the MoBa cohort, as shown in the present and our previous study (Bulik *et al.* 2007). These are likely to underestimate the true population prevalence given that other poorly defined EDNOS presentations could not be captured by the self-report methodology. The observed prevalence in this study is higher than the 0.5% prevalence of self-reported history of eating disorders in a United Kingdom pregnancy cohort ($N=12254$) (Micali *et al.* 2007b) and the point prevalence of 3.8%–4.0% in the Norwegian adult female population (Götestam and Agras 1995, Zachrisson *et al.* 2008); the difference is probably explained by the lowering of binge/purge thresholds from 2 per week to 1 per week in the MoBa studies, the less strict weight criterion for AN in the MoBA studies, and the inclusion of AN and BN only and use of a single-item self-report in Micali *et al.* The lowered binge/purge thresholds are commensurate with those proposed for *DSM-5*. The prevalence of eating disorders in this study, and eating disorder behaviors more generally among pregnant women is alarmingly high; previous research shows that binge eating occurs among 17%–44%, self-induced vomiting for weight control in 1%–2%, and dieting in 3%–37% (Abraham *et al.* 1994, Fairburn and Welch 1990, Soares *et al.* 2009). The morbidity, heightened risk of birth complications, and negative neonatal outcomes associated with eating disorders (Bulik *et al.* 1999, Micali *et al.* 2007a, Sollid *et al.* 2004) make identification of eating pathology imperative.

Fewer than half of obstetricians/gynecologists (ob/gyn) assess eating disorder history, body image concerns, and eating disorder behaviors, despite assessing related constructs of body weight, BMI, exercise, and dietary practices (Leddy *et al.* 2009). Lack of training in

identification of signs and symptoms, a perception that assessment falls outside the scope of practice, and lack of awareness of the consequences of eating disorders in pregnancy may explain this (Leddy *et al.* 2009). Nevertheless, vigilance for potential signs and symptoms can be easily incorporated into routine ob/gyn practice, via screening questions posed to the individual and through assiduity to selected anthropometric, biochemical, dietary intake, and clinical data, such as reproductive history (Nickols-Richardson 2008).

Pregnancy appears to be a vulnerability window for onset of some eating disorders, consistent with the findings of our former study (Bulik *et al.* 2007) and case reports (Tiller and Treasure 1998). As found in Bulik *et al.*, onset cases generally comprised BED, while BN and EDNOS-P onset were rare. Specific physical and psychological factors have a conjectured role in eating disorder onset during pregnancy (Knoph Berg *et al.* 2008, Tiller and Treasure 1998). For BED onset specifically, low maternal education, low combined income, a native language other than the official country language, lifetime adversities, anxiety and depression, low social support, and weight concerns, are putative vulnerability factors (Knoph Berg *et al.* 2011). Given that we are not well-informed about BED prevention and that pregnancy is a risk period for mental illnesses (e.g., depression), attention to broad mental health promoting mechanisms is advisable. Health care providers can help with social support, skills to manage stress, body image issues, and anxiety and depression. Moreover, given documented differences in nutrition during pregnancy in women with BED (Siega-Riz *et al.* 2008), nutritional counseling may play a valuable role in ensuring healthy balanced nutrition throughout pregnancy and the subsequent lactation period.

Seventy-four percent with BN pre-pregnancy met criteria for remission or partial remission in early pregnancy. Improvement in binge-purge behaviors during pregnancy has been noted elsewhere (Crow *et al.* 2004, Lacey and Smith 1987), along with a reduction in general health risk behaviors, such as alcohol, tobacco, and other drug use (Crow *et al.* 2004). Maternal desire for healthy fetal development appears to motivate behavioral change during pregnancy (Lemberg and Phillips 1989). Previous studies have suggested that cognitive symptoms of BN (i.e., body dissatisfaction, weight concern) remain problematic or worsen during pregnancy (Crow *et al.* 2004, Micali *et al.* 2007a), even in the context of decreasing binge-purge, restricting, and health risk behaviors (Crow *et al.* 2004, Lemberg and Phillips 1989) and binge-purge symptoms may return after childbirth (Crow *et al.* 2008). Pregnancy potentially offers a window to neutralize barriers to help-seeking (e.g., shame, ambivalence about treatment) and enhance engagement in treatment.

This study has several limitations. First, low power to detect differences in outcomes for the BN and EDNOS-P groups is a significant limitation. However, given the dearth of data on course of eating disorders during pregnancy, a decision was made to report all relevant information. Second, diagnostic measures involved self-report rather than clinical diagnostic interview, a practical preclusion due to the size of the sample; additionally the measure has not been psychometrically validated, but is based on *DSM* criteria. Third, the diagnostic criteria do not correspond directly to *DSM-IV* and may in fact be closer to *DSM-5*. Fourth, the overall prevalence of broadly defined eating disorders is conservative, given that the assessment of AN during pregnancy is methodologically compromised due to inability to

assess the weight criterion, hence prevalence of AN during pregnancy did not contribute to the overall prevalence estimate. Further, EDNOS generally is a heterogeneous and poorly defined diagnostic category; although some broadly agreed presentations such as BED, EDNOS-P, and subthreshold AN and BN were captured within this study, it was not possible to capture undefined presentations with the self-report method; hence, the observed overall prevalence of eating disorders likely underestimates the true population prevalence. Fifth, there may be selection bias in the recruitment into MoBa. The prevalences of eating disorder and eating disorder subtypes may differ between MoBa participants and the general Norwegian pregnant population, potentially influencing remission, continuation, and incidence rates during pregnancy. Lastly, we make the assumption here that eating disorder rates remain the same over time and we have temporal validation, however, it could be the case that the validation models are spurious and there is a change over time paired with a bad model predictive ability; however, there is no way to confirm this.

The high prevalence of broadly defined eating disorders (primarily BED) among 1 in every 21 pregnant women and association between maternal eating disorders and birth complications (Bulik *et al.* 1999, Micali *et al.* 2007a, Sollid *et al.* 2004) underscore the need for detection and treatment of eating disorders during pregnancy. Physicians, midwives, and health care professionals play an important role in optimizing maternal and birth outcomes, therefore knowledge of the potential serious consequences of eating disorders coupled with identification and management strategies is vital.

Acknowledgments

The Norwegian Mother and Child Cohort Study is supported by the Norwegian Ministry of Health and the Ministry of Education and Research, NIH/NIEHS (contract no NO-ES-75558), NIH/NINDS (grant no.1 UO1 NS 047537-01), and the Norwegian Research Council/FUGE (grant no. 151918/S10). We are grateful to all the participating families in Norway who take part in this ongoing cohort study.

References

- Abraham S, King W, Llewellyn-Jones D. Attitudes to body weight, weight gain and eating behavior in pregnancy. *Journal of Psychosomatic Obstetrics & Gynecology*. 1994; 15:189–95. [PubMed: 7881504]
- Altman DG, Vergouwe Y, Royston P, Moons KG. Prognosis and prognostic research: Validating a prognostic model. *BMJ*. 2009; 338:b605. [PubMed: 19477892]
- American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. American Psychiatric Association; Washington, DC: 1994.
- Berg CK, Torgerson L, Von Holle A, Hamer RM, Bulik C, Reichborn-Kjennerud T. Factors associated with eating disorders in pregnancy. *International Journal of Eating Disorders*. 2011; 44:124–133. [PubMed: 20127938]
- Blais MA, Becker AE, Burwell RA, Flores AT, Nussbaum KM, Greenwood DN, Ekeblad ER, Herzog DB. Pregnancy: Outcome and impact on symptomatology in a cohort of eating-disordered women. *International Journal of Eating Disorders*. 2000; 27:140–9. [PubMed: 10657887]
- Bulik C, Sullivan P, Fear J, Pickering A, Dawn A. Fertility and reproduction in women with anorexia nervosa: a controlled study. *Journal of Clinical Psychiatry*. 1999; 2:130–135. [PubMed: 10084645]
- Bulik CM, Von Holle A, Hamer R, Knoph Berg C, Torgersen L, Magnus P, Stoltenberg C, Siega-Riz AM, Sullivan P, Reichborn-Kjennerud T. Patterns of remission, continuation and incidence of broadly defined eating disorders during early pregnancy in the Norwegian Mother and Child Cohort Study (MoBa). [Corrigendum appears in *Psychological Medicine*, in press]. *Psychological Medicine*. 2007; 37:1109–18. [PubMed: 17493296]

- Crow SJ, Agras WS, Crosby R, Halmi K, Mitchell JE. Eating disorder symptoms in pregnancy: a prospective study. *International Journal of Eating Disorders*. 2008; 41:277–9. [PubMed: 18027861]
- Crow SJ, Keel PK, Thuras P, Mitchell JE. Bulimia symptoms and other risk behaviors during pregnancy in women with bulimia nervosa. *International Journal of Eating Disorders*. 2004; 36:220–3. [PubMed: 15282692]
- Efron, B.; Tibshirani, R. An introduction to the bootstrap. Chapman & Hall/CRC; Boca Raton: 1993.
- Fairburn CG, Welch SL. The impact of pregnancy on eating habits and attitudes to shape and weight. *International Journal of Eating Disorders*. 1990; 9:153–160.
- Good, PI. Resampling methods: a practical guide to data analysis. Birkhäuser; Boston: 2006.
- Gøtestam KG, Agras WS. General population-based epidemiological study of eating disorders in Norway. *International Journal of Eating Disorders*. 1995; 18:119–26. [PubMed: 7581413]
- Harris JR, Magnus P, Tambs K. The Norwegian Institute of Public Health Twin Panel: a description of the sample and program of research. *Twin Research*. 2002; 5:415–23. [PubMed: 12613498]
- Knoph Berg C, Bulik CM, Von Holle A, Torgersen L, Hamer R, Sullivan P, Reichborn-Kjennerud T. Psychosocial factors associated with broadly defined bulimia nervosa during early pregnancy: findings from the Norwegian Mother and Child Cohort Study. *Australian & New Zealand Journal of Psychiatry*. 2008; 42:396–404. [PubMed: 18473258]
- Knoph Berg C, Torgersen L, Von Holle A, Hamer RM, Bulik CM, Reichborn-Kjennerud T. Factors associated with binge eating disorder in pregnancy. *International Journal of Eating Disorders*. 2011; 44:124–33. [PubMed: 20127938]
- Koubaa S, Hallstrom T, Lindholm C, Hirschberg AL. Pregnancy and neonatal outcomes in women with eating disorders.[Erratum appears in *Obstet Gynecol*. 2008 May;111(5):1217 Note: Kouba, Saloua [corrected to Koubaa, Saloua]]. *Obstetrics & Gynecology*. 2005; 105:255–60. [PubMed: 15684148]
- Lacey JH, Smith G. Bulimia nervosa: The impact of pregnancy on mother and baby. *British Journal of Psychiatry*. 1987; 150:777–81. [PubMed: 3651730]
- Leddy MA, Jones C, Morgan MA, Schulkin J. Eating disorders and obstetric-gynecologic care. *Journal of Women's Health*. 2009; 18:1395–401.
- Lemberg R, Phillips J. The impact of pregnancy on anorexia nervosa and bulimia. *International Journal of Eating Disorders*. 1989; 8:285–295.
- Magnus P, Irgens LM, Haug K, Nystad W, Skjaerven R, Stoltenberg C. MoBa Study G. Cohort profile: The Norwegian Mother and Child Cohort Study (MoBa). *International Journal of Epidemiology*. 2006; 35:1146–50. [PubMed: 16926217]
- Mazzeo SE, Zucker NL, Gerke CK, Mitchell KS, Bulik CM. Parenting concerns of women with histories of eating disorders. *International Journal of Eating Disorders*. 2005; 37(Suppl):S77–9. discussion S87–9. [PubMed: 15852326]
- Micali N, Simonoff E, Treasure J. Risk of major adverse perinatal outcomes in women with eating disorders. *British Journal of Psychiatry*. 2007a; 190:255–259. [PubMed: 17329747]
- Micali N, Treasure J, Simonoff E. Eating disorders symptoms in pregnancy: a longitudinal study of women with recent and past eating disorders and obesity. *Journal of Psychosomatic Research*. 2007b; 63:297–303. [PubMed: 17719368]
- Nickols-Richardson, SM. Anorexia nervosa and bulimia nervosa during pregnancy. In: Lammi-Keefe, CJ.; Couch, SC.; Philipson, EH., editors. *Handbook of nutrition and pregnancy*. Humana Press; NJ: 2008. p. 115-134.
- Reba-Harrelson L, Von Holle A, Hamer RM, Torgersen L, Reichborn-Kjennerud T, Bulik CM. Patterns of maternal feeding and child eating associated with eating disorders in the Norwegian Mother and Child Cohort Study (MoBa). *Eating Behaviors*. 2010; 11:54–61. [PubMed: 19962121]
- Reichborn-Kjennerud T, Bulik CM, Kendler KS, Roysamb E, Maes H, Tambs K, Harris JR. Gender differences in binge-eating: A population-based twin study. *Acta Psychiatrica Scandinavica*. 2003; 108:196–202. [PubMed: 12890274]
- Reichborn-Kjennerud T, Bulik CM, Kendler KS, Roysamb E, Tambs K, Torgersen S, Harris JR. Undue influence of weight on self-evaluation: A population-based twin study of gender differences. *International Journal of Eating Disorders*. 2004a; 35:123–32. discussion 133–5. [PubMed: 14994348]

- Reichborn-Kjennerud T, Bulik CM, Tambs K, Harris JR. Genetic and environmental influences on binge eating in the absence of compensatory behaviors: A population-based twin study. *International Journal of Eating Disorders*. 2004b; 36:307–14. [PubMed: 15478129]
- Rocco PL, Orbitello B, Perini L, Pera V, Ciano RP, Balestrieri M. Effects of pregnancy on eating attitudes and disorders: A prospective study. *Journal of Psychosomatic Research*. 2005; 59:175–9. [PubMed: 16198191]
- Siega-Riz AM, Haugen M, Meltzer HM, Von Holle A, Hamer R, Torgersen L, Knopf-Berg C, Reichborn-Kjennerud T, Bulik CM. Nutrient and food group intakes of women with and without bulimia nervosa and binge eating disorder during pregnancy. *American Journal of Clinical Nutrition*. 2008; 87:1346–55. [PubMed: 18469258]
- Soares RM, Nunes MA, Schmidt MI, Giacomello A, Manzolli P, Camey S, Buss C, Drehmer M, Melere C, Hoffman J, Ozcariz S, Manenti CN, Pinheiro AP, Duncan BB. Inappropriate eating behaviors during pregnancy: Prevalence and associated factors among pregnant women attending primary care in southern Brazil. *International Journal of Eating Disorders*. 2009; 42:387–93. [PubMed: 19115363]
- Sollid CP, Wisborg K, Hjort J, Secher NJ. Eating disorder that was diagnosed before pregnancy and pregnancy outcome. *American Journal of Obstetrics & Gynecology*. 2004; 190:206–10. [PubMed: 14749661]
- Steyerberg, EW. *Clinical prediction models: A practical approach to development, validation, and updating (statistics for biology and health)*. Springer; New York: 2009.
- Steyerberg EW, Borsboom GJJM, van Houwelingen HC, Eijkemans MJC, Habbema JDF. Validation and updating of predictive logistic regression models: a study on sample size and shrinkage. *Statistics in Medicine*. 2004; 23:2567–86. [PubMed: 15287085]
- Tiller J, Treasure J. Eating disorders precipitated by pregnancy. *European Eating Disorders Review*. 1998; 6:178–187.
- Wilson GT, Sysko R. Frequency of binge eating episodes in bulimia nervosa and binge eating disorder: Diagnostic considerations. *International Journal of Eating Disorders*. 2009; 42:603–10. [PubMed: 19610014]
- Zachrisson HD, Vedul-Kjelsås E, Göttestam KG, Mykletun A. Time trends in obesity and eating disorders. *International Journal of Eating Disorders*. 2008; 41:673–80. [PubMed: 18537168]




Fig. 1. Participant flow to achieve final analysis sample. MoBa, The Norwegian Mother and Child Cohort Study. *Extrapolated from the reported 38.5% participation rate [<http://www.fhi.no>]. **Criteria not mutually exclusive

Table 1

Sociodemographic characteristics of women in the MoBa “training” and “validation” samples

Characteristic	Training Sample (<i>n</i> =41243)	Validation Sample (<i>n</i> =36024)
Maternal age (yrs), <i>M</i> (<i>SD</i>)	29.9 (4.6)	30.1 (4.7)
Number of previous live births, <i>N</i> (%)		
0	19823 (48.1)	23216 (64.4)
1+	13695 (33.2)	8744 (24.3)
2+	7725 (18.7)	4064 (11.3)
Marital status, <i>N</i> (%)		
Married	20173 (49.3)	14995 (41.9)
Cohabiting	19399 (47.4)	19499 (54.5)
Single	915 (2.2)	789 (2.2)
Divorced/widowed	422 (1.0)	467 (1.3)
Education, <i>N</i> (%)		
< 3 yrs high school	3676 (9.4)	2129 (6.3)
Vocational high school	5666 (14.5)	4045 (11.9)
3-yr high school general studies, junior college	6554 (16.8)	4488 (13.2)
Regional technical college/4-yr university degree	15924 (40.8)	13760 (40.4)
University/technical college, >4 yrs	7217 (18.5)	9618 (28.3)
Combined minimum income, <i>N</i> (%)		
0–\$36K (200000 NOK)	3898 (10.2)	2714 (8.1)
>\$36K (200000 NOK)	21696 (57.0)	13217 (39.4)
>\$89K (500000 NOK)	8208 (21.6)	10523 (31.3)
>\$125K (700000 NOK)	4275 (11.2)	7128 (21.2)
Ever smoked (yes), <i>N</i> (%)	20394 (49.6)	18081 (50.4)
Smoking during pregnancy (yes), <i>N</i> (%)	4429 (10.8)	2266 (6.3)
Ever drank alcohol (yes), <i>N</i> (%)	38924 (95.5)	34488 (96.6)

Table 2
Prevalence and course of illness of broadly defined eating disorders during pregnancy in a Norwegian population-based pregnancy cohort (MoBa)

Eating disorder	Prevalence*		Course of Illness			
	Before pregnancy n (%)	During pregnancy n (%)	Remission n (%)	Partial remission n (%)	Continuation n (%)	Incidence n (%)
Training sample (n=41243)						
AN (broad)	40 (0.1)	-	-	-	-	-
BN (any type)	310 (0.7)	102 (0.2)	84 (36.5)	75 (32.6)	71 (28.9)	21 (0.1)
BN (purging)	128 (0.3)	55 (0.1)	40 (41.7)	17 (17.7)	39 (36.4)	7 (0.0)
BN (non-purging)	113 (0.3)	33 (0.1)	32 (30.5)	54 (51.4)	17 (15.6)	11 (0.0)
BED (broad)	1480 (3.5)	1954 (4.8)	510 (38.3)	-	821 (61.5)	801 (2.1)
EDNOS-P	46 (0.1)	12 (0.03)	22 (78.6)	-	3 (10.0)	5 (0.0)
Any eating disorder	1876 (4.3)	2068 (4.8)	-	-	-	-
Validation sample (n=36024)						
AN (broad)	32 (0.1)	-	-	-	-	-
BN (any type)	362 (1.0)	95 (0.2)	113 (40.6)	104 (37.4)	60 (20.1)	22 (0.1)
BN (purging)	132 (0.3)	41 (0.1)	48 (48.0)	20 (20.0)	29 (26.9)	4 (0.0)
BN (non-purging)	174 (0.5)	34 (0.1)	54 (34.8)	82 (52.9)	17 (10.2)	13 (0.0)
BED (broad)	1218 (3.3)	1691 (4.8)	456 (40.3)	-	674 (59.5)	690 (2.1)
EDNOS-P	46 (0.1)	15 (<0.01)	24 (80.0)	-	4 (12.5)	5 (0.0)
Any eating disorder	1658 (4.4)	1801 (4.7)	-	-	-	-

AN, anorexia nervosa; BN, bulimia nervosa; BED, binge eating disorder; EDNOS-P, eating disorder not otherwise specified-purging disorder. BN any type includes BN purging, BN non-purging, and individuals who could not reliably be categorized as BN purging or non-purging due to missing data.

* Prevalence was determined from the full sample with 81 320 observations prior to exclusions regarding existent status before and during pregnancy required for rate calculations

Table 3

Age-adjusted rates of remission, continuation, and incidence of eating disorders in pregnancy in 1,000 person-weeks (95% CI) in a Norwegian population-based pregnancy cohort (MoBa)

Course	Eating disorder				EDNOS-P ^d
	BN (any type)	BN (purging)	BN (non-purging)	BED (broad)	
	Training sample (n=41,243)				
Remission ^b	19.99 (16.61 – 24.06)	21.73 (16.35 – 28.88)	17.34 (12.79 – 23.50)	21.77 (20.28 – 23.36)	43.79 (35.75 – 53.64)
Partial remission ^c	18.62 (15.35 – 22.58)	10.49 (6.68 – 16.47)	28.09 (22.90 – 34.44)	-	-
Continuation	16.94 (13.86 – 20.70)	22.11 (17.19 – 28.43)	8.27 (5.04 – 13.57)	35.01 (33.52 – 36.56)	5.89 (2.00 – 17.39)
Incidence	0.03 (0.02 – 0.05)	0.01 (0.00 – 0.02)	0.02 (0.01 – 0.03)	1.22 (1.14 – 1.31)	0.01 (0.00 – 0.02)
	Validation sample (n=36,024)				
Remission ^b	22.39 (19.23 – 26.08)	25.89 (20.54 – 32.64)	19.32 (15.43 – 24.20)	22.68 (21.07 – 24.42)	43.83 (36.25 – 53.01)
Partial remission ^c	20.94 (17.82 – 24.61)	10.28 (6.33 – 16.68)	29.47 (25.17 – 34.50)	-	-
Continuation	11.65 (9.25 – 14.66)	15.88 (11.53 – 21.87)	5.86 (3.73 – 9.21)	33.57 (31.96 – 35.27)	7.20 (2.86 – 18.12)
Incidence	0.04 (0.02 – 0.05)	0.01 (0.00 – 0.02)	0.02 (0.01 – 0.04)	1.17 (1.09 – 1.27)	0.01 (0.00 – 0.02)

BN, bulimia nervosa; BED, binge eating disorder; EDNOS-P, eating disorder not otherwise specified-purging disorder.

^aEDNOS-P rates, BN purging incidence, and BN non-purging incidence and continuation calculations except for BED were not age-adjusted due to small sample size.

^bRemission indicates rate of no eating disorder at time of survey completion during pregnancy.

^cPartial remission in BN indicates absence of compensatory behaviors during early pregnancy.

Table 4

Recalibration estimates and performance statistics for the recalibrated rate models of remission, continuation, and incidence of eating disorders during pregnancy in a Norwegian population-based pregnancy cohort (MoBa)

Course	Recalibration type estimate	Performance statistics	
	α :intercept (95% CI)	<i>U</i> -statistic	Brier score
BED			
Remission ^a	0.04 (-0.03 – 0.12)	0.000362	0.000028
Continuation	-0.04 (-0.09 – 0.01)	0.000547	0.000028
Incidence	-0.04 (-0.11 – 0.04)	0.000015	0.000028
BN			
Remission ^a	0.11 (-0.03 – 0.26)	0.002441	0.000028
Partial remission ^b	0.13 (-0.03 – 0.28)	0.002924	0.000028
Continuation	-0.36 (-0.59 – -0.13)	0.014905	0.000028
Incidence	0.14 (-0.29 – 0.56)	0.000006	0.000028
EDNOS-P ^c			
Remission ^a	<0.01 (-0.19 - 0.19)	0.000000	0.000028
Continuation	0.20 (-0.72 – 1.12)	0.002439	0.000028
Incidence	0.14 (-0.74 – 1.01)	0.000001	0.000028

BED, binge eating disorder; BN, bulimia nervosa; EDNOS-P, eating disorder not otherwise specified-purging disorder.

^aRemission indicates rate of no eating disorder at time of survey completion during pregnancy.

^bPartial remission in BN indicates absence of compensatory behaviors during early pregnancy.

^cEDNOS-P and incidence calculations except for BED were not age-adjusted due to small sample size

Table 5

BED incidence rates by sociodemographic characteristics

Predictor	Training sample (n=41243)		Validation Sample (n=36024)		BED incidence $\beta_{overall}$ (95% CI)
	Estimate (se)		Estimate (se)		
Ever smoke? (no)	-0.58 (0.07)		-0.70 (0.08)		1.21 (0.93 – 1.48)
Infertility treatment (no)	0.22 (0.15)		0.15 (0.14)		0.66 (-0.62 – 1.95)
Pregnant before? (no)	-0.32 (0.08)		-0.30 (0.08)		0.94 (0.45 – 1.43)
Marital status					1.81 (0.94 – 2.67)
Married	-0.42 (0.20)		-0.79 (0.19)		-
Cohabiting	-0.42 (0.20)		-0.72 (0.19)		-
Single	Referent		referent		-
Mother's education					1.11 (0.76 – 1.47)
<3-yr high school	0.76 (0.14)		0.92 (0.14)		-
Vocational high school	0.54 (0.13)		0.40 (0.14)		-
3-yr high school general studies, junior college	0.45 (0.13)		0.38 (0.13)		-
Regional technical college/4-yr university degree	0.25 (0.11)		0.10 (0.11)		-
University, technical college, > 4 yrs	Referent		referent		-
Minimum combined income					0.62 (0.10 – 1.13)
0-\$36K (200000 NOK)	0.59 (0.16)		0.33 (0.16)		-
>\$36K (200000 NOK)	0.25 (0.13)		0.26 (0.11)		-
>\$89K (500000 NOK)	0.23 (0.15)		-0.01 (0.12)		-
>\$125K (700000 NOK)	referent		referent		-
Total abortions	-0.43 (0.09)		-0.39 (0.09)		-
Total live births					0.40 (-0.06-0.85)
0	-0.41 (0.09)		-0.23 (0.12)		-
1	-0.12 (0.09)		-0.18 (0.13)		-
2+	referent		referent		-

The left columns show poisson regression parameter estimates indicating the natural log of ratio of incidence rates for the predictor versus the referent in the "training" and "validation" samples. The right column shows calibration estimates for univariate models with "validation" versus "training" sample data predicting BED incidence by sociodemographic characteristics. BED, binge eating disorder.

* At least one significant difference between groups at an α -level of .05.

Table 6

Bootstrap age-adjusted rates of remission, continuation, and incidence of eating disorders during pregnancy in 1,000 person-weeks (95% CI) and bias estimates in a Norwegian population-based pregnancy cohort (MoBa)^a

Course	BN (any type)	BN (purging)	BN (non-purging)	BED (broad)
Remission ^b	21.5 (19.0,24.1)	23.8 (19.8,28.4)	18.7 (15.4,22.2)	22.4 (21.3,23.5)
Partial remission ^c	19.7 (17.5,22.4)	10.6 (7.6,14.3)	28.7 (25.3,32.7)	-
Continuation	13.8 (11.7,16.1)	18.6 (15.1,22.8)	6.5 (4.3,8.8)	34.3 (33.2,35.4)
Incidence	0.033 (0.023,0.043)			1.200 (1.141,1.264)
Bias estimate (bootstrap <i>se</i>)				
Remission ^b	-0.02 (0.06)	-0.29 (0.09)	-0.08 (0.09)	-0.01 (0.03)
Partial remission ^c	-0.24 (0.06)	-0.38 (0.16)	-0.32 (0.06)	-
Continuation	-0.10 (0.08)	-0.28 (0.10)	-0.11 (0.18)	-0.02 (0.02)
Incidence	>-0.01 (0.15)			>-0.01 (0.03)

BN, bulimia nervosa; BED, binge eating disorder; EDNOS-P, eating disorder not otherwise specified-purging disorder.

^aEDNOS-P and incidence calculations for BN (purging) and BN (non-purging) were not included because models did not converge for >30% of all the samples.

^bRemission indicates rate of no eating disorder at time of survey completion during pregnancy.

^cPartial remission in BN indicates absence of compensatory behaviors during early pregnancy.