

1 **Title (max 85 characters):** Exercise-induced salivary hormone
2 responses to high-intensity, self-paced running

3
4 **Submission type:** Original investigation.

5
6 **Authors:** Diogo V. Leal^{1,2*}, Lee Taylor^{3,4,5}, John Hough^{1,6}

7
8
9 **Affiliations:** ¹Institute of Sport and Physical Activity Research,
10 School of Sport and Physical Activity, University of
11 Bedfordshire, Polhill Avenue, Bedford, Bedfordshire, United
12 Kingdom;

13 ²Research Center in Sports Sciences, Health Sciences and
14 Human Development, University Institute of Maia, Maia,
15 Portugal;

16 ³School of Sport, Exercise and Health Sciences, Loughborough
17 University, Loughborough, United Kingdom;

18 ⁴Human Performance Research Centre, University of
19 Technology Sydney (UTS), Australia.

20 ⁵Sport & Exercise Discipline Group, Faculty of Health,
21 University of Technology Sydney (UTS), Australia.

22 ⁶School of Science and Technology, Nottingham Trent
23 University, Nottingham, NG11 8NS, United
24 Kingdom.

25
26 ***Address for Correspondence:**

27 Diogo Vaz Leal
28 University Institute of Maia, Av. Carlos Oliveira Campos, 4475-
29 690 Castelo da Maia, Portugal

30 **Email:** diogo.leal@ismai.pt

31 **Phone:** +351 22 986 60 00

32 **ORCID:** 0000-0002-4046-6820

33

34 **Preferred Running Head:** Salivary steroids responses to the
35 RPE_{TP}

36

37 **Abstract word count:** 250

38 **Text-only word count:** 3452

39 **Number of figures:** 5

40 **Number of tables:** 1

41

42 **Abstract**

43 **Purpose:** Physical overexertion can lead to detrimental
44 overreaching states without sufficient recovery, which may be
45 identifiable by blunted exercise-induced cortisol and
46 testosterone responses. A running test (RPE_{TP}) elicits
47 reproducible plasma cortisol and testosterone elevations (in a
48 healthy state) and may detect blunted hormonal responses when
49 overreached. This current study determines the salivary cortisol
50 and testosterone responses reproducibility to the RPE_{TP}, to
51 provide greater practical validity using saliva compared to the
52 previously utilized blood sampling. Secondly, the relationship
53 between the salivary and plasma responses will be assessed.
54 **Methods:** Twenty-three active, healthy males completed the
55 RPE_{TP} on three occasions. Saliva (N=23) and plasma (N=13)
56 were collected Pre-, Post- and 30 min Post-Exercise. **Results:**
57 Salivary cortisol did not elevate in any RPE_{TP}-trial, and reduced
58 concentrations occurred 30 min Post-Exercise ($P = 0.029$, $\eta^2 =$
59 0.287); trial differences were observed ($P < 0.001$, $\eta^2 = 0.463$).
60 The RPE_{TP} elevated ($P < 0.001$, $\eta^2 = 0.593$) salivary testosterone
61 with no effect of trial ($P = 0.789$, $\eta^2 = 0.022$). Intra-individual
62 variability was 25% in cortisol and 17% in testosterone. ‘Fair’
63 ICCs of 0.46 (cortisol) and 0.40 (testosterone) were found.
64 Salivary and plasma cortisol positively correlated ($R = 0.581$, P
65 $= 0.037$) yet did not for testosterone ($R = 0.345$, $P = 0.248$).
66 **Conclusions:** The reproducibility of salivary testosterone
67 response to the RPE_{TP} is evident and supports its use as a
68 potential tool, subject to further confirmatory work, to detect
69 hormonal dysfunction during overreaching. Salivary cortisol
70 responds inconsistently in a somewhat individualized manner to
71 the RPE_{TP}.

72

73

74

75

76 **Keywords:** testosterone, cortisol, preventive measures, stress,
77 overreaching.

78

79

80 **Introduction**

81

82 Effective physical performance adaptations require an
83 appropriately prescribed and periodized training program.¹
84 **When overreaching occurs, a reduced athletic capacity**
85 **(transiently or otherwise) may be observed, due to imbalanced**
86 **overload and recovery periodisation.**² Appropriate recovery may
87 elicit a “supercompensatory” performance response referred to
88 as functional overreaching (FOR).³ Yet, insufficient recovery
89 from prolonged periods of intensified-training may lead to “non-
90 functional overreaching” (NFOR) requiring weeks/months for
91 full recovery – whilst – unchecked NFOR can progress to
92 overtraining syndrome (OTS) which can, on occasions, demand
93 years for full recovery to occur.⁴ Prevalence of NFOR/OTS
94 during an elite athlete’s career can range from ~35%⁵ to 67%⁶
95 yet little progress has been made regarding objective biomarkers
96 that detect the onset/magnitude of overreaching.³

97

98 Resting cortisol and testosterone concentrations have been
99 proposed as overreaching/OTS markers, as they provide a ratio
100 of catabolic to anabolic activity.³ However, their alterations at
101 rest are inconsistent when comparing pre to post periods of
102 overload.^{7,8} Recently, their acute responses to exercise have
103 shown promise as an indicator of hormonal dysfunction
104 following intensified-training periods.^{9,10} Blunted exercise-
105 induced salivary cortisol and testosterone responses were shown
106 following a 30-min cycling bout, known as the 55/80 [1 min at
107 55% maximal workload (\dot{W}_{max}) and 4 min at 80% \dot{W}_{max}]
108 following an 11-day⁹ and a 10-day¹⁰ intensified-training period,
109 suggesting these exercise-induced salivary hormones are
110 potentially useful biomarkers of overreaching/OTS. Recently, a
111 treadmill-derivative [rating of perceived exertion protocol
112 (RPE_{TP})]¹¹ of the 55/80 cycle¹² was developed and shown to
113 induce reproducible elevations of plasma testosterone but not
114 cortisol.¹¹

115

116 Therefore, this study primarily sought to, in attempt to increase
117 practical validity, determine whether the same RPE_{TP} cortisol
118 and testosterone responses in plasma¹¹ could be replicated in
119 saliva, in healthy (i.e. non-overreached) adult, male individuals.
120 If saliva was to show such validity, the RPE_{TP} could become a
121 more practical tool to detect and subsequently inform
122 practitioner decision-making, regarding any potential hormonal
123 dysregulation associated with overreaching/OTS. Secondly,
124 this study also **intended** to assess the relationship between saliva
125 and blood cortisol and testosterone responses to the RPE_{TP}, albeit
126 in a subsection of previously measured participants. It was
127 hypothesised that (i) salivary testosterone but not cortisol would
128 acutely elevate in response to the RPE_{TP}; (ii) these responses

129 would be reproducible; and (iii) the salivary hormone responses
130 would correlate with their venous surrogates.

131

132 **Methods**

133

134 **Participants**

135

136 This study was completed to expand upon previously published
137 data.¹¹ Twenty-three ‘recreationally-trained’ and ‘trained’
138 (categorised as per¹³ and reflective of performance levels 2 and
139 3) males [age 21 ± 2 years; height 177 ± 6 cm; body mass 76.1
140 ± 13.1 kg; maximal heart rate (HR_{max}) 191 ± 9 beats·min⁻¹;
141 maximum oxygen uptake ($\dot{V}O_{2max}$) 55 ± 6 mL·kg⁻¹·min⁻¹]
142 volunteered to participate in this study. Partial data from thirteen
143 participants from the previously published study¹¹
144 (physiological, plasma cortisol, plasma testosterone and
145 anthropometric data) are included in the present study. The study
146 was conducted in accordance with the 2013 Declaration of
147 Helsinki under ethical approval [University of Bedfordshire
148 Research Ethics Committee (2014ISPAR003)]. Following
149 verbal and written study descriptions participants provided
150 written informed consent.

151

152 **Design**

153

154 The original research¹¹ examined the reproducibility of plasma
155 cortisol and testosterone responses to two novel running
156 protocols. The present study extends this work¹¹ by examining
157 salivary cortisol and testosterone responses to one of these
158 running protocols (RPE_{TP}), given saliva is a more ecologically
159 valid sample compared to plasma.

160

161 Participants undertook 5 main trials within a temperature-
162 controlled laboratory (see Figure 1). On the first visit, a
163 submaximal running test followed by a $\dot{V}O_{2max}$ test was
164 undertaken to physiologically characterise participants. During
165 the subsequent 4 visits (separated by a minimum of 4 to 7 days),
166 3 exercise trials (T1, T2 and T3), and one resting control trial
167 (CTL) were completed. In each exercise trial, participants
168 undertook the RPE_{TP} (Figure 1), which has been detailed
169 previously.¹¹ CTL identified the influence of the circadian
170 rhythm on the hormones measured.^{14,15} All participants woke
171 before 8 AM on the morning of the trial which started at 12 PM
172 for diurnal variation control purposes.¹⁶ A 76-statement
173 recovery-stress questionnaire (RESTQ-76¹⁷) was completed
174 before the start of each exercise bout, as used previously¹¹. No
175 differences in RESTQ-76¹⁷ metrics were observed prior to trial
176 completion, indicating the participants’ were in a similar state of
177 well-being and pre-disposition to exercise prior to all trials,
178 likely not overreached and that any alterations in the hormones

179 examined were not due to pre-trial stress and/or well-being (or
180 variation in said measures pre-trial).

181

182 Abstinence from exercise, caffeine and alcohol intake 24 hours
183 before each main trial was requested, and a standard breakfast
184 chosen by the participant was consumed by 9 AM (repeated prior
185 to each visit). The participants' pre-trial 24 h nutritional intake
186 was determined via a weighed food diary. Nutrition analysis
187 software (Dietplan, Version 6.70.74, Forestfield, West Sussex,
188 UK) was used on the food diaries and mean energy (9851 ± 4182
189 kJ), carbohydrate ($56\% \pm 12\%$), fat ($25\% \pm 13\%$), and protein
190 ($17\% \pm 2\%$) intake were determined. Euhydration was
191 confirmed by a urine osmolality of ≤ 700 mOsm \cdot kg \cdot H $_2$ O $^{-1}$.¹⁸
192 Food consumption was not allowed until the end of each main
193 experimental trial but water was provided *ad libitum* up to 10
194 min before saliva sample collection.

195

196 *** Insert Figure 1 near here ***

197

198 Methodology

199

200 **Submaximal running and $\dot{V}O_{2max}$ tests.** The protocols used for
201 determination of $\dot{V}O_{2max}$ have been detailed and justified in
202 previous research.^{11,19} Briefly, a 4x4-min-stage, incremental
203 treadmill-run submaximal test was completed, to determine the
204 running speed/oxygen consumption ($\dot{V}O_2$) relationship.²⁰ The
205 initial speed was self-selected between 6.5 – 12.0 km \cdot h $^{-1}$ and
206 increased by 1 km \cdot h $^{-1}$ every stage. The speed corresponding to a
207 HR of ~ 150 beats \cdot min $^{-1}$ (range: 8.0 – 14.0 km \cdot h $^{-1}$) on the
208 submaximal test was noted and, after a 20-min recovery period,
209 used on the incline-ramped $\dot{V}O_{2max}$ test.^{11,20} The speed was
210 maintained throughout with a 1% increase in gradient every
211 minute until volitional exhaustion. Expired gas was analysed
212 through a breath-by-breath ergospirometry system (MetaLyzer
213 3B, Cortex, Leipzig, Germany). This protocol determines the
214 velocity at $\dot{V}O_{2max}$ ($v\dot{V}O_{2max}$), from which percentages were
215 used in the original study¹¹. Such inferences were not required
216 for this study as only the self-paced RPE_{TP} protocol was utilized.
217 The participants' $\dot{V}O_{2max}$ was established in accordance with the
218 British Association of Sports and Exercise Sciences' criteria.¹⁹

219

220 **RPE_{TP} and CTL:** Briefly and as described in full previously¹¹,
221 the RPE_{TP} is a self-paced, continuous, 30-min running bout, with
222 alternating blocks of 1 min at 11 (fairly light) and 4 min at 15
223 (hard) on the 6-20 Borg scale²¹. Speed was self-adjusted to
224 maintain exertion in the target range and blinded from the
225 participant to maintain the exertion in the target range. Saliva
226 samples were collected pre-, immediately post-, and 30 min post-
227 exercise in all exercise trials. The CTL followed the same
228 scheme as in Figure 1 for the RPE_{TP}, but no exercise was

229 completed, therefore sample timepoints are referred to as pre-
230 CTL, post-CTL and 30 min post-CTL. Blood samples were also
231 collected in the first 13 participants immediately before saliva
232 sampling. Heart rate (HR) and RPE were measured in the last
233 15s of each stage via short-range radio telemetry (Polar FT1,
234 Polar Electro Oy, Kempele, Finland) and the 6-20 Borg scale,²¹
235 respectively.

236

237 ***Saliva handling and analysis:*** Saliva samples were collected
238 into 7 mL polystyrene sterile containers (Sterilin, Thermo
239 Scientific, Loughborough, UK) by unstimulated passive drool,
240 with eyes closed, head tilted slightly forward and avoiding any
241 orofacial movement.²² Water consumption was not allowed
242 within the 10 min preceding sampling. Minimum collection time
243 was 3 min for each participant to allow for collection of
244 sufficient sample volume (~2 mL). Samples were then
245 centrifuged at 14600 g for 10 min (Espresso Microcentrifuge,
246 Thermo Scientific, Loughborough, UK) and the supernatant was
247 transferred into 1.5 mL aliquots (Eppendorf, Hamburg,
248 Germany) to be stored at -80°C until further analysis.

249

250 Salivary cortisol and testosterone concentrations were
251 determined by using commercially available enzyme-linked
252 immunosorbent assay (ELISA) kits (Salimetrics, PA 16803,
253 USA). All samples were analysed in duplicate and average
254 concentrations were used. The determined mean intra-assay CVs
255 were 4.8% (salivary cortisol) and 4.4% (salivary testosterone).
256 The present analyses resulted in in-lab mean inter-assay CVs of
257 5.1% and 6.8% for salivary cortisol and testosterone,
258 respectively.

259

260 ***Venous plasma handling and analysis:*** All analytical
261 procedures for blood collection, treatment and analysis have
262 been detailed previously.¹¹ Briefly, whole blood samples were
263 collected by venepuncture from the antecubital fossa into
264 tripotassium ethylenediaminetetraacetic acid (K₃EDTA) tubes,
265 centrifuged at 4°C for 10 min (1500g) and the plasma stored at -
266 80°C before further analysis. The ELISA kits (IBL International,
267 Hamburg, Germany) mean intra- and inter-assay CVs were 3.0%
268 and 4.6%, and 3.5% and 5.7% for plasma cortisol and
269 testosterone, respectively. The venous blood sample data was
270 taken from previously published work¹¹ and has been used for
271 correlation with the salivary data presented in this present study
272 only.

273

274 **Statistical Analysis**

275

276 The IBM Statistical Package for Social Sciences® (SPSS)
277 Statistics version 23.0 (SPSS Inc., Chicago, IL) was used for all
278 statistical analysis. The Shapiro-Wilk test and scatter plots were

279 used for verification of normality and homoscedasticity of raw
280 data, respectively. When non-normally distributed (all
281 variables), log transformation to base 10 was completed with
282 subsequent normality rechecked. All data were then deemed
283 normally distributed, except for speed (how this analysis was
284 completed is detailed below). Magnitude of effect was examined
285 using the Cohen's *d* effect sizes (ES)²³, determined by hand as
286 described in Vincent and Weir,²⁴ and labeled using consistent
287 thresholds of < 0.2 trivial, 0.21 – 0.49 small, 0.50 – 0.80
288 moderate, > 0.80 large.^{24,25} The alpha level of significance was
289 set as $P < 0.05$. Data is reported as mean (SD), and all results are
290 presented as raw data to facilitate comprehension. Salivary
291 cortisol and HR_{max} data were collapsed for all correlation
292 analysis.

293

294 *Salivary Hormone and Physiological Data Analysis:* Salivary
295 testosterone, speed and HR data sets were collapsed given there
296 were no significant differences between any trial (excluding
297 CTL) ($P > 0.05$). Salivary cortisol was not collapsed as a trial
298 effect was observed. A two-way repeated measures analysis of
299 variance (ANOVA) was used on the normalised data (salivary
300 hormones and HR), with unchanged significant effects observed.
301 On finding an effect, paired sample *t*-tests were used, and
302 Bonferroni adjustments applied (also used to examine the
303 hormonal responses during CTL), with partial eta squared (η^2)
304 values determining the size of the effect. A non-parametric 2-
305 related sample Wilcoxon test was used for between-trial
306 comparisons for speed.

307

308 *Reproducibility Analysis:* Intra-individual coefficients of
309 variation (CV_i) for all physiological and hormonal
310 measurements were calculated. The intra-individual mean
311 concentrations (\bar{X}_i) and SDs (SD_i) were used to calculate the CV_i
312 using the equation $CV_i = (SD_i/\bar{X}_i)*100$. A two-way model based
313 on the examination of single measures intraclass correlation
314 coefficient (ICC_{2,1}) was also used on the collapsed data to
315 account for the between-individual variability.²⁷ Guidelines on
316 ICC models propose that values considered poor sit below 0.40,
317 whereas fair sit within 0.40-0.59, good between 0.60-0.74, and
318 excellent if or above 0.75.²⁸

319

320 *Correlation Analysis:* Pearson's correlation was used to
321 determine the correlation between the salivary and plasma
322 cortisol and testosterone concentrations, and the individual
323 absolute change in salivary cortisol and testosterone with HR_{max}.
324 As there was no change in the cortisol response to exercise, these
325 data have been collapsed. The correlation between plasma and
326 salivary testosterone has been examined at pre-, post- and 30 min
327 post-exercise. The level of significance was set as $P < 0.05$.

328

329

330 Results

331

332 *Acute Hormonal Responses*

333

334 All reproducibility data and average salivary
335 cortisol/testosterone concentrations are presented in Table 1.

336

337 *** Insert Table 1 near here ***

338

339 *Salivary cortisol.* A trial effect was observed ($P < 0.001$, $\eta^2 =$
340 0.463), with average responses being lower in T3 compared to
341 T2 ($P = 0.002$). A time effect ($P = 0.029$, $\eta^2 = 0.287$) was also
342 observed. Pairwise comparisons showed cortisol did not acutely
343 elevate in any exercise trial (ES = 0.10 in T1, ES = -0.11 in T2,
344 ES = 0.02 in T3, all $P > 0.05$), but a lower concentration was
345 observed at 30 min post-exercise when compared to post-
346 exercise in T1 ($P = 0.003$, ES = 0.32) and T2 ($P = 0.043$, ES =
347 0.11). Individual acute responses are presented in Figure 2.

348

349 *Salivary testosterone.* There was no effect of trial ($P = 0.789$, η^2
350 = 0.022), but a significant time effect was observed ($P < 0.001$,
351 $\eta^2 = 0.593$). Pairwise comparisons showed salivary testosterone
352 acutely elevated ($P < 0.001$) and remained elevated at 30 min
353 post-exercise ($P < 0.05$) in all exercise trials. Average acute
354 percentage-elevations were ~23% (ES = -0.94) in T1, ~40% (ES
355 = -1.10) in T2 and ~32% (ES = -0.87) in T3. Individual exercise-
356 induced changes are presented in Figure 2.

357

358 *Plasma cortisol and testosterone.* The plasma cortisol and
359 testosterone values can be examined in detail elsewhere.¹¹
360 Briefly, average raw data for plasma cortisol (nmol·L⁻¹) was
361 259.1 ± 105.3, 313.9 ± 125.8, and 292.7 ± 123.2 at pre-, post-,
362 and 30 min post-exercise, respectively. The average raw data for
363 plasma testosterone (nmol·L⁻¹) was 13.4 ± 2.6, 18.9 ± 3.7, and
364 15.0 ± 3.2 at pre-, post-, and 30 min post-exercise, respectively.

365

366 *Plasma and Salivary Hormone Correlation.* Plasma and salivary
367 cortisol were shown to positively correlate ($R = 0.581$, $P =$
368 0.037). However, no correlation was observed between plasma
369 and salivary testosterone concentration levels at pre-exercise (R
370 = 0.430, $P = 0.143$), post-exercise ($R = 0.250$, $P = 0.409$), and
371 30 min post-exercise ($R = 0.340$, $P = 0.256$), as presented in
372 Figure 3.

373

374 *** Insert Figure 2 near here ***

375

376 *** Insert Figure 3 near here ***

377

378 *Individual Absolute Change in Salivary Hormone and*
379 *Physiological Responses Correlation.* Salivary cortisol and
380 HR_{max} were shown to positively correlate ($R = 0.632$, $P < 0.001$).
381 A correlation between salivary testosterone and HR_{max} was not
382 observed ($R = 0.094$, $P = 0.671$), as presented in Figure 4.

383

384 ***** Insert Figure 4 near here *****

385

386 *Hormonal Responses During CTL*

387

388 Salivary cortisol concentrations were lower at post-CTL and 30
389 min post-CTL than Pre-CTL by $\sim 28\% \pm 17\%$ and $\sim 37\% \pm 19\%$,
390 respectively (both $P < 0.001$). Pre-CTL salivary testosterone was
391 not different from post-CTL ($P = 0.142$) but was $\sim 12\% \pm 5\%$
392 higher than 30 min post-CTL ($P = 0.003$) (Table 1).

393

394 ***** Insert Figure 5 near here *****

395

396 *Speed/HR Acute Responses and Urine Osmolality*

397 No differences between collapsed trials were found in speed or
398 HR ($P > 0.05$ for all) (see Figure 5). Reproducibility and average
399 data for speed and HR in response to the RPE_{TP} trials are
400 presented in Table 1. Urine osmolality did not differ between
401 trials ($P > 0.05$).

402

403

404 **Discussion**

405

406 This study's primary aim was to determine whether the same
407 RPE_{TP} cortisol and testosterone responses in plasma¹¹ could be
408 replicated in saliva, in healthy (i.e. non-overreached) adult, male
409 individuals. Indeed, the RPE_{TP} significantly and acutely elevated
410 salivary testosterone in T1 (515 to 630 pmol·L⁻¹, ES = -0.94), T2
411 (491 to 663 pmol·L⁻¹, ES = -1.10), and T3 (523 to 661 pmol·L⁻¹,
412 ES = -0.87). However, salivary cortisol did not significantly
413 elevate in any trial, as shown previously elsewhere¹¹, thus
414 accepting hypothesis (i). Furthermore, the CV_i in salivary
415 testosterone observed in this present study ($17 \pm 7\%$) is similar
416 to the exercise-induced variance observed for plasma
417 testosterone elsewhere ($12 \pm 9\%$)¹¹. This has not been observed
418 for salivary cortisol ($25 \pm 15\%$), whose variability in plasma to
419 the RPE_{TP} is moderately lower ($12 \pm 7\%$)¹¹, partially accepting
420 hypothesis ii). The secondary aim sought to assess the
421 relationship between saliva and blood cortisol and testosterone
422 responses to the RPE_{TP}, albeit in a subsection of previously
423 measured participants. A correlation between salivary and
424 plasma testosterone levels was not observed (see Figure 3).
425 Whilst salivary cortisol and plasma surrogates did correlate, their
426 lack of change across trials limits the utility in this cortisol

427 specific inference. Taken together (cortisol and testosterone
428 correlations) the data rejects hypothesis (iii).

429

430 Previous data demonstrates a correlation between plasma and
431 salivary testosterone^{29,30} – however – this is in resting samples
432 unlike the ‘exercise response’ data from the present study (see
433 Figures 2 and 3, and Table 1). Similarly, Hough et al. (2011)¹²
434 did not observe a correlation in exercise-induced responses of
435 plasma and salivary testosterone, although the authors suggest
436 caution is required when interpreting their data. Specifically,
437 they¹²: (i) missed some post-exercise blood samples; and (ii)
438 proposed that the correlation between plasma and salivary
439 testosterone might not have occurred as testosterone elevates to
440 exercise stress quicker in the blood than saliva. Indeed, it has
441 been observed elsewhere that despite parallel increases in blood
442 and saliva testosterone after oral testosterone undecanoate
443 administration in healthy men, the absorption curves showed a
444 high interindividual variability in the time at which maximum
445 concentrations were reached.³¹

446

447 Furthermore, Fiers et al. (2014)³² have observed that salivary
448 testosterone concentrations were not identical to comparable
449 serum free testosterone due to testosterone binding with salivary
450 proteins. In this present study, we speculate that the correlation
451 may not have been observed due to timing of the testosterone
452 entering the saliva. Supporting our speculation, we have
453 observed that post-exercise plasma testosterone correlates only
454 with the 30 min post-exercise salivary testosterone in T1 (data
455 not presented). However, **it should be noted that salivary
456 testosterone significantly increased in response to the RPE_{TP} in
457 all trials**, and that the intra-individual variability in the present
458 study was $17 \pm 7\%$. Importantly, this present exercise-induced
459 variability in salivary testosterone ($17 \pm 7\%$) is noticeably lower
460 than the 37% blunted elevation in cycling-induced salivary
461 testosterone responses after an 11-day period of intensified
462 training (compared with a mean $\sim 58\%$ elevation pre-training) in
463 active males suspected to be overreached.⁹ Suggesting the RPE_{TP}
464 may be useful when measuring testosterone responses in the
465 saliva of healthy male individuals.

466

467 A mean increase in salivary cortisol was not present, yet some
468 participants demonstrated an increase at the individual level (see
469 Figure 2 and Table 1; although responses are not replicated
470 across trials at the level of the individual), likely due to exercise
471 intensity variability between participants. Indeed, positive
472 correlations between HR_{max} observed during the RPE_{TP} and
473 absolute change values in salivary cortisol were observed (see
474 Figure 4). Cortisol has been reported to acutely elevate in
475 response to exercise.^{12,33} Although it has been **proposed** that
476 exercise must be at an intensity above $60\% \dot{V}O_{2max}$ for at least

477 20-30 min to induce an elevation in cortisol levels,³⁴ this is not
478 always observed.³⁵ As no acute elevation and a between-trial
479 difference was observed, the RPE_{TP} may not have provided a
480 sufficient physiological strain to activate an exercise-induced
481 salivary cortisol response in all participants; as also observed in
482 its plasma surrogate.¹¹ These data may suggest that the
483 variability observed in the salivary cortisol sensitivity to exercise
484 may be driven by exercise intensity. However, no correlation
485 was present between salivary testosterone and HR_{max} (see Figure
486 4) despite a consistent elevation in this hormone in response to
487 the RPE_{TP} and low inter- (see Figure 2, row D) and intra-
488 individual (Table 1) variability. The data reinforces the highly-
489 sensitive nature of salivary testosterone to exercise (certainly in
490 response to the RPE_{TP}, as observed elsewhere in its plasma
491 surrogate¹¹), highlighting its potential utility within
492 hypothalamic-pituitary-gonadal dysregulation associated with
493 NFOR/OTS.

494

495 **Practical Applications**

496

- 497 • Salivary testosterone is sensitive and a reproducible
498 biomarker to the RPE_{TP} indicating a triggered activation of
499 the hypothalamic-pituitary-gonadal complex after short-
500 duration, high-intensity running exercise.
- 501 • Salivary cortisol demonstrates a somewhat individualized
502 yet non-sensitive response to the RPE_{TP} and from the present
503 and related experimental designs, is currently an unproven
504 biomarker-related exercise-induced stress responsiveness.
- 505 • The RPE_{TP} elicits reproducible physiological and salivary
506 testosterone hormone responses with greater practical
507 application/integration than previous methods (e.g. saliva
508 and not plasma samples); with further proof of concept (i.e.
509 analysis of the effects of a period of intensified training on
510 the RPE_{TP}-induced responsiveness of salivary testosterone)
511 it may be a useful potential tool in NFOR/OTS paradigms.

512

513 **Conclusions**

514

515 Physiological responses (HR and speed at prescribed RPE) were
516 shown reproducible across all RPE_{TP} trials within the present
517 study; within a larger sample than used previously elsewhere.¹¹
518 Salivary cortisol was not sensitive to the RPE_{TP} (as shown
519 elsewhere albeit in plasma surrogates¹¹). Despite cortisol
520 salivary levels correlating with its plasma surrogate as
521 hypothesised, acute cortisol responses may be influenced by
522 diurnal variation and an individualised response to RPE_{TP} (likely
523 exercise intensity driven), rendering it an unreliable biomarker
524 to highlight exercise responsiveness, within the present design.
525 The present RPE_{TP} data suggests salivary testosterone to be a
526 more robust marker of a triggered endocrine mobilization during

527 exercise. Yet, and despite no strong correlation observed in the
528 exercise-induced salivary and plasma testosterone levels, the
529 variability between the acute responses of plasma and salivary
530 testosterone to the RPE_{TP} are relatively similar. Therefore, the
531 consistent and sensitive exercise responsiveness of salivary
532 testosterone to the RPE_{TP} suggest: (i) greater utility than cortisol;
533 (ii) a more practically compatible bio-sample than plasma; and
534 (iii) changes in salivary testosterone were no due to inconsistent
535 physiological strain. Nevertheless, future work is required to
536 detail proof of concept regarding the salivary testosterone
537 sensitivity to the RPE_{TP} in an active population following a
538 period of intensified training. Such data would demonstrate
539 quantitatively whether RPE_{TP}-induced salivary (and plasma)
540 testosterone responses and their blunting, are a suitable tool to
541 highlight the incidence of NFOR/OTS.

542

543

544 **Acknowledgements**

545 The authors would like to acknowledge all participants involved
546 in this study for their hard work and commitment, and the
547 technical staff at the Sports Sciences laboratories of the
548 University of Bedfordshire for their continuous support. The
549 authors have no conflict of interest to report. No external
550 financial support has been given.

551

552

553 **References**

554

- 555 1. Koutedakis Y, Metsios GS, A. S-K. Periodization of
556 Exercise Training in Sport. In: Whyte G, ed. *The*
557 *Physiology of Training: Advances in Sport and Exercise*
558 *Science Series*. Philadelphia: Churchill Livingstone,
559 Elsevier; 2006:1-21.
- 560 2. Halson SL, Bridge MW, Meeusen R, et al. Time course
561 of performance changes and fatigue markers during
562 intensified training in trained cyclists. *J Appl Physiol*.
563 2002;93(3):947-956.
564 doi:10.1152/jappphysiol.01164.2001
- 565 3. Meeusen R, Duclos M, Foster C, et al. Prevention,
566 diagnosis and treatment of the overtraining syndrome:
567 Joint consensus statement of the European College of
568 Sport Science (ECSS) and the American College of
569 Sports Medicine (ACSM). *Eur J Sport Sci*.
570 2013;13(1):1-24. doi:10.1080/17461391.2012.730061
- 571 4. Meeusen R, Duclos M, Foster C, et al. Prevention,
572 diagnosis and treatment of the overtraining syndrome:
573 Joint consensus statement of the European College of
574 Sport Science (ECSS) and the American College of
575 Sports Medicine (ACSM). *Eur J Sport Sci*.
576 2013;13(1):1-24. doi:10.1080/17461391.2012.730061

- 577 5. Kenttä G, Hassmén P, Raglin JS. Training practices and
578 overtraining syndrome in Swedish age-group athletes.
579 *Int J Sports Med.* 2001;22(6):460-465. doi:10.1055/s-
580 2001-16250
- 581 6. Morgan WP, O’Conner PJ, Sparling PB, Pate RR.
582 Psychological characterization of the elite female
583 distance runner. *Int J Sports Med.* 1987;8(2):124-131.
584 doi:10.1055/s-2008-1025717
- 585 7. Lucia A, Diaz B, Hoyos J, et al. Hormone levels of
586 world class cyclists during the tour of Spain stage race.
587 *Br J Sports Med.* 2001;35(6):424-430.
588 doi:10.1136/bjism.35.6.424
- 589 8. Grandys M, Majerczak J, Duda K, Zapart-Bukowska J,
590 Kulpa J, Zoladz JA. Endurance training of moderate
591 intensity increases testosterone concentration in young,
592 healthy men. *Int J Sports Med.* 2009;30(7):489-495.
593 doi:10.1055/s-0029-1202340
- 594 9. Hough J, Corney R, Kouris A, Gleeson M. Salivary
595 cortisol and testosterone responses to high-intensity
596 cycling before and after an 11-day intensified training
597 period. *J Sports Sci.* 2013;31(14):1614-1623.
598 doi:10.1080/02640414.2013.792952
- 599 10. Hough J, Robertson C, Gleeson M. Blunting of exercise-
600 induced salivary testosterone in elite-level triathletes
601 with a 10-day training camp. *Int J Sports Physiol
602 Perform.* 2015;10(7):935-938. doi:10.1123/ijsp.2014-
603 0360
- 604 11. Leal D V., Taylor L, Hough J. Reproducibility of Acute
605 Steroid Hormone Responses in Men to Short-Duration
606 Running. *Int J Sports Physiol Perform.*
607 2019;14(10):1430-1437. doi:10.1123/ijsp.2018-1004
- 608 12. Hough JP, Papacosta E, Wraith E, Gleeson M. Plasma
609 and salivary steroid hormone responses of men to high-
610 intensity cycling and resistance exercise. *J Strength
611 Cond Res.* 2011;25(1):23-31.
612 doi:10.1519/JSC.0b013e3181fef8e7
- 613 13. de Pauw K, Roelands B, Cheung SS, de Geus B,
614 Rietjens G, Meeusen R. Guidelines to Classify Subject
615 Groups in Sport- Science Research Guidelines. *Int J
616 Sports Physiol Perform.* 2013;8(2):111-122.
617 doi:10.1123/ijsp.8.2.111
- 618 14. Smyth JM, Ockenfels MC, Gorin AA, et al. Individual
619 differences in the diurnal cycle of cortisol.
620 *Psychoneuroendocrinology.* 1997;22(2):89-105.
621 doi:10.1016/S0306-4530(96)00039-X
- 622 15. Kraemer WJ, Loebel CC, Volek JS, et al. The effect of
623 heavy resistance exercise on the circadian rhythm of
624 salivary testosterone in men. *Eur J Appl Physiol.*
625 2001;84:13-18. doi:10.1007/s004210000322
- 626 16. Debono M, Ghobadi C, Rostami-Hodjegan A, et al.

- 627 Modified-release hydrocortisone to provide circadian
628 cortisol profiles. *J Clin Endocrinol Metab.* 2009.
629 doi:10.1210/jc.2008-2380
- 630 17. Kellmann M, Kallus KW. *Recovery-Stress*
631 *Questionnaire for Athletes: User Manual.* Champaign,
632 IL: Human Kinetics; 2001.
- 633 18. Sawka MN, Burke LM, Eichner ER, Maughan RJ,
634 Montain SJ, Stachenfeld NS. American College of
635 Sports Medicine position statement: Exercise and Fluid
636 Replacement. *Med Sci Sports Exerc.* 2007;39(2):377-
637 390. doi:10.1249/mss.0b013e31802ca597
- 638 19. Bird S, Davidson R. *Guidelines for the Physiological*
639 *Testing of Athletes.* (Bird S, Davidson R, eds.). Leeds:
640 British Association of Sport and Exercise Science; 1997.
- 641 20. Deighton K, Zahra JC, Stensel DJ. Appetite, energy
642 intake and resting metabolic responses to 60min
643 treadmill running performed in a fasted versus a
644 postprandial state. *Appetite.* 2012;58(3):946–954.
645 doi:10.1016/j.appet.2012.02.041
- 646 21. Borg G. Perceived exertion as an indicator of somatic
647 stress. *Scand J Rehabil Med.* 1970;2:92-98. doi:S/N
- 648 22. Navazesh M. Methods for Collecting Saliva. *Ann N Y*
649 *Acad Sci.* 1993;694(1):72-77. doi:10.1111/j.1749-
650 6632.1993.tb18343.x
- 651 23. Hopkins WG, Marshall SW, Batterham AM, Hanin J.
652 Progressive statistics for studies in sports medicine and
653 exercise science. *Med Sci Sports Exerc.* 2009;41(1):3.
654 doi:10.1249/MSS.0b013e31818cb278
- 655 24. Vincent W, Weir J. *Statistics in Kinesiology.* 4th ed.
656 Champaign, IL: Human Kinetics; 2012.
- 657 25. Cohen J. *Statistical Power Analysis for the Behavioral*
658 *Sciences.* 2nd ed. Lawrence Erlbaum Associates; 1988.
659 doi:10.1234/12345678
- 660 26. Sale D. Testing strength and power. In: MacDougall J.
661 D., Wenger HA, Green HJ, eds. *Physiological Testing of*
662 *the High-Performance Athlete.* Champaign, IL: Human
663 Kinetics; 1991:21-106.
- 664 27. Šerbetar I. Establishing Some Measures of Absolute and
665 Relative Reliability of a Motor Tests. *Croat J Educ.*
666 2015;17(1):37-48. doi:10.15516/cje.v17i0.1484
- 667 28. Fleiss JL. *Statistical Methods for Rates and Proportions.*
668 2nd ed. New York: Wiley; 1981.
- 669 29. Khan-Dawood FS, Choe JK, Dawood MY. Salivary and
670 plasma bound and “free” testosterone in men and
671 women. *Am J Obstet Gynecol.* 1984;148(4):442-445.
672 doi:10.1016/0002-9378(84)90723-3
- 673 30. Vittek J, L’Hommedieu DG, Gordon GG, Rappaport SC,
674 Louis Southren A. Direct Radioimmunoassay (RIA) of
675 salivary testosterone: correlation with free and total
676 serum testosterone. *Life Sci.* 1985;97:711-716.

- 677 doi:10.1016/0024-3205(85)90540-5
678 31. Schurmeyer T, Wickings EJ, Freischem CW, Nieschlag
679 E. Saliva and serum testosterone following oral
680 testosterone undecanoate administration in normal and
681 hypogonadal men. *Acta Endocrinol.* 1983;102(3):456-
682 462. doi:10.1530/acta.0.1020456
683 32. Fiers T, Delanghe J, T'Sjoen G, Van Caenegem E,
684 Wierckx K, Kaufman JM. A critical evaluation of
685 salivary testosterone as a method for the assessment of
686 serum testosterone. *Steroids.* 2014;86:5-9.
687 doi:10.1016/j.steroids.2014.04.013
688 33. Adebero T, McKinlay BJ, Theocharidis A, et al. Salivary
689 and Serum Concentrations of Cortisol and Testosterone
690 at Rest and in Response to Intense Exercise in Boys
691 Versus Men. *Pediatr Exerc Sci.* 2020;32(2):65-72.
692 34. Davies C, Few J. Effects of exercise on adrenocortical
693 function. *J Appl Physiol.* 1973;35(6):887-891.
694 35. Gass GC, McLellan TM, Gass EM. Effects of prolonged
695 exercise at a similar percentage of maximal oxygen
696 consumption in trained and untrained subjects. *Eur J*
697 *Appl Physiol Occup Physiol.* 1991;63(6):430-435.
698

699 **Figure Captions**

700

701 **Figure 1** Schematic presentation of (A) the experimental trial
702 day protocol and procedures, (B) the study design, and (C) the
703 RPE_{TP} design; Submax, submaximal treadmill-run test; $\dot{V}O_{2max}$,
704 maximal oxygen uptake test; CTL, control resting trial.

705

706 **Figure 2** Salivary hormone responses to the RPE_{TP} and CTL
707 protocols at Pre-Exercise (Pre-CTL), Post-Exercise (Post-CTL)
708 and 30 min Post-Exercise (30 min Post-CTL): (A) Salivary
709 cortisol; (B) Salivary testosterone; (C) Individual absolute
710 changes in salivary cortisol; (D) Individual absolute changes in
711 salivary testosterone.

712 #Trial difference (T3 different than T2). *Different than Pre-
713 exercise values ($P < 0.01$). **Different than Pre-exercise values
714 ($P < 0.05$). ‡Different than Post-exercise values ($P < 0.05$).
715 †Different than CTL ($P < 0.01$). δ - small effect size for trial; \clubsuit
716 - trivial effect size for trial.

717

718 **Figure 3** Collapsed salivary and plasma cortisol correlation (A),
719 and salivary and plasma testosterone correlation analysis at pre-
720 exercise (B), post-exercise (C), and 30 min post-exercise (D).

721

722 **Figure 4** Correlation analysis between HR_{max} observed during
723 RPE_{TP} and collapsed individual absolute changes in salivary
724 cortisol (top) and salivary testosterone (bottom).

725

726 **Figure 5** Heart rate and speed responses to the RPE_{TP} on each
727 separate experimental trial (all $P > 0.05$).

728

729

730

731

732

733

734

735

736

737

738

739

740

741

742

743

744

745

746

747

748

749 **Table Captions**

750

751 **Table 1** Average raw data for urine osmolality, the physiological
752 and hormone responses in the CTL, T1, T2 and T3 bouts and
753 reproducibility data (when applicable) data for T1, T2 and T3
754 only.

755