



UPLC–MS–QTOF analysis and antifungal activity of Cumaru (*Amburana cearensis*)

Maria Tatiana Alves Oliveira^{1,2} · Marcus Vinícius Oliveira Barros de Alencar² · Vicente de Paulo dos Anjos Landim² · Geovanna Maria Medeiros Moura¹ · Joelton Igor Oliveira da Cruz¹ · Elizeu Antunes dos Santos¹ · Henrique Douglas Melo Coutinho³ · Jacqueline Cosmo Andrade⁴ · Irwin Rose Alencar de Menezes⁵ · Paulo Riceli Vasconcelos Ribeiro⁶ · Edy Sousa de Brito⁶ · Erlânio Oliveira de Sousa⁷ · Adriana Ferreira Uchoa¹

Received: 22 April 2020 / Accepted: 9 November 2020
© King Abdulaziz City for Science and Technology 2020

Abstract

This study was aimed at investigating the phytochemical constituents, antifungal properties and antibiotic-modifying activity of the aqueous crude extract and fractions of *Amburana cearensis* seeds (CEFAC). The CEFAC were chemically characterized by LC–MS/MS–QTOF. In addition, the antifungal activity was assayed by the microdilution method against strains of *Candida albicans*. The phytochemical profile of CEFAC exhibited phenolic compounds, organic acids, and polyphenols. The results of the assessment of antifungal activity revealed an IC₅₀ ranging from 45.6 to 2048 µg/mL. Interestingly, when CEFAC was associated with Fluconazole, we evidenced a decreased IC₅₀ (1.81–11.9 µg/mL), suggesting a synergism with antibiotic. It was possible to identify in the crude extract and fractions several phenolic compounds, organic acids, and some polyphenols in positive ionization mode. These results suggest that CEFAC may present compounds with the ability to interact and act synergistically with antimicrobial drugs, highlighting its potential as an alternative source for the development of new antimicrobial agents.

Keywords *Amburana cearensis* · *Candida* spp. · Fungistatic · Fluconazole

✉ Henrique Douglas Melo Coutinho
hdmcoutinho@gmail.com

- ¹ Laboratory of Chemistry and Function of Bioactive Proteins, Federal University of Rio Grande do Norte, Natal, Rio Grande do Norte, Brazil
- ² Department of Biomedical Science, University Centre INTA-UNINTA, Sobral, Ceará, Brazil
- ³ Laboratory of Microbiology and Molecular Biology-LMBM, Regional University of Cariri-URCA, Rua Cel. Antonio Luis 1161, Pimenta, Crato, Ceará 63105-000, Brazil
- ⁴ Laboratory of Bioassays-LABIO, Educator Training Institute, Federal University of Cariri-UFCA, Campus Brejo Santo, Juazeiro do Norte, Ceará, Brazil
- ⁵ Department of Biological Chemistry, Regional University of Cariri, Crato, Ceará, Brazil
- ⁶ Multiuser Laboratory of Natural Products Chemistry-LMQPN, Embrapa Agroindustria Tropical, Fortaleza, Ceará, Brazil
- ⁷ Laboratory of Food Physicochemical Analysis-LAFIQ, Technology College of Cariri-FATEC, Juazeiro do Norte, Ceará, Brazil

Introduction

The fungi are part of the human microbiota (Siqueira and Sen 2004) and important etiological agents of human diseases, due to its ability to modulate the expression of virulence factors such as host cell adhesion, hyphae formation, phenotypic plasticity and production of hydrolytic enzymes. The genus *Candida* is the most widespread opportunistic fungal pathogen in the human body, causing mucosal and systemic infections, especially among immunosuppressed and hospitalized patients (Goulart et al. 2018). Due to the wide use of antifungal agents, drug resistance of *Candida albicans* is increasing, which poses a serious threat to antifungal therapy (Ksiezopolska and Gabaldón 2018). Therefore, exploring new effective antifungal agents is urgently needed.

In this sense, natural products have been reported as sources of various bioactive molecules that may pave the way for novel antifungal agents (Ksouri et al. 2017; Zida et al. 2017). In addition, studies are carried out to promote the combination of antifungal drugs and natural products against resistant *Candida*

strains (Calixto Jr. et al. 2015), which can be a strongly promise to face this challenge.

Amburana cearensis (Allemão) A. C. Sm., Fabaceae family (Papilionoideae, Leguminosae) is naturally widespread in Caatinga, northeastern Brazil (Bravo et al. 1999). The medicinal use of *A. cearensis* is common, especially in the treatment of headaches, muscle aches, constipation, and urinary tract infections (Leal et al. 2003; Agra et al. 2007; Lima et al. 2013). Several compounds with antioxidant and antimicrobial properties have been isolated from *A. cearensis* seeds, including coumarins, methyl esters, phytosterols (γ -sitosterol, stigmasterol, and campesterol) (Calixto Jr. et al. 2015; Pereira et al. 2017; Zida et al. 2017). Thus, the aqueous crude extract and fractions of *A. cearensis* seeds were subjected to chemical characterization and antifungal assessments to verify whether they present any potential to induce a synergistic effect when combined with Fluconazole against *C. albicans*.

Materials and methods

Plant material

Amburana cearensis seeds were identified and provided by Floresta Nacional de Nísia, a Conservation Unit managed by the Chico Mendes Institute for Biodiversity Conservation (ICMBio), located in the municipality of Nísia Floresta, state of the Rio Grande do Norte, Brazil, and the Caatinga Seed Network (UNIVASF).

Extraction and fractionation

The extraction was performed by homogenizing the seed powder of *A. cearensis* with 50 mM Tris-HCl solution, pH 7.5, ratio 1:10 (m/v), under constant agitation for 4 h at 4 °C and then centrifuged at 10,000×g for 30 min at the same temperature. The material was filtered, and the supernatant was identified as crude extract (CE). Then, CE was fractionated into three ammonium sulfate concentration ranges: 0–30% (FR-1), 30–60% (FR-2) and 60–90% (FR-3) of saturation. After each precipitation step, the sample was maintained at 4 °C for approximately 16 h and then centrifuged at 10,000×g for 30 min at 4 °C. The fractions were then resuspended and dialyzed for 20 h using distilled water. After dialysis, the fractions were lyophilized and stored at room temperature.

Identification of compounds by ultra-performance liquid chromatography-quadrupole/time-of-flight (UPLC-QTOF) system

The analysis was performed on ACQUITY UPLC (Waters), coupled to quadrupole/time-of-flight system (QTOF,

Waters) by Brazilian Agricultural Research Corporation (EMBRAPA). The chromatographs were performed on a column Waters ACQUITY UPLC BEH (150×2.1 mm, 1.7 μ m), fixed temperature of 40 °C, mobile water phases with 0.1% of formic acid (A) and acetonitrile with 0.1% formic acid (B), gradient ranging from 2 to 95% B (15 min), 0.4 mL/min flow and 5- μ L injection volume. The ESI- mode was purchased in the 110–1180 Da, fixed source temperature at 120 °C, desolvation temperature at 350 °C, desolvation gas flow at 500 L/h, extraction cone of 0.5 V and 2.6 kV of capillary voltage. The ESI+ mode was purchased in the 110–1180 Da, fixed source temperature at 120 °C, desolvation temperature at 350 °C, desolvation gas flow at 500 L/h and 3.2 kV of capillary voltage. Enkephalin leucine was used as lock mass. The acquisition mode was MS^E. The instrument was controlled by the software Masslynx 4.1 (Waters Corporation).

Antifungal assays

Strains and culture medium

The strains used in this study were standard type and isolated from Cultura Oswaldo Cruz (FIOCRUZ), Instituto Nacional de Controle de Qualidade em Saúde (INCQS), specifically CA INCQS 40006 and CA URM 4125, standard and isolated strain, respectively. The strains were inoculated on Sabouraud Dextrose Agar (SDA, KASVI) and incubated for 24 h at 37 °C. Subsequently, aliquots of the yeast were transported to test tubes, each containing 3 mL of sterile saline (0.9%). Inoculum concentration was standardized by comparison with the 0.5 McFarland scale (barium chloride) (NCCLS 2012). Sabouraud Dextrose (SD, HIMEDIA) in double concentration was used in the microdilution assay.

Chemicals and reagents

Dimethyl sulfoxide (DMSO, Merck, Germany) was used to fluidize the CE and fractions, while Fluconazole (Capsule—Prati Donaduzzi, Brazil) was diluted with distilled water and used as reference drug. The test solution was prepared by weighing 0.15 g of the CE and fractions and diluting with 1 mL of DMSO to obtain the desired concentration. Samples were diluted in sterile distilled water (16.384 μ g/mL) such that DMSO had no activity on the tested cells (Stoppa et al. 2009).

IC₅₀ and cell viability

To determine the concentration capable of inhibiting 50% of yeast growth (IC₅₀), the CE and fractions isolated or combined with Fluconazole were tested by microdilution method in Sabouraud Dextrose broth in 96-well microplates with

initial concentration of 16.384 $\mu\text{g/mL}$, and serially diluted to a final concentration of 8 $\mu\text{g/mL}$ (Javadpour et al. 1996; Morais-Braga et al. 2016). Controls were also prepared for the diluents using 0.9% sodium chloride solution instead of inoculum. All tests were performed in quadruplicates. The plates were incubated at 37 °C for 24 h and then read on a spectrophotometer ELISA (Thermoplate®) at 630 nm.

Determination of minimum fungicidal concentration (MFC)

To verify the MFC, the tip of a sterile rod was inserted into each well of the previously tested plate. After inserting into the middle of each well, the rod was taken to a Petri dish containing SD with the aid of a guide plate attached to the bottom of the plate for yeast subculture and cell viability evaluation. After 24 h of incubation, the plates were analyzed for colony formation (Ernst et al. 1999). The concentration in which there was no growth of fungal colony was considered the MFC.

Evaluation of the modifying effect on Fluconazole action

First, the intrinsic action of CE, fractions and Fluconazole on yeast growth were verified. Then, it was evaluated if the antifungal action of Fluconazole was modulated by the CE and fractions. Therefore, CE and fractions were used at sub-inhibitory concentrations (MFC/16) according to Coutinho et al. (2008) with minor modifications from Morais-Braga et al. (2016). If the CE or fractions potentiates the action of the Fluconazole, the verified effect was considered synergic type. If it interfered with the action of the Fluconazole, the verified effect was considered antagonistic type. The plates were filled with 100 μL of medium + inoculum + sample followed by microdilution with 100 μL of Fluconazole at concentration of 8192 $\mu\text{g/mL}$. The mixture was added to the first well of the plate to be subjected to serial dilutions (1:1), ranging from 8192 to 8 $\mu\text{g/mL}$. The last well was used as control. The plates were incubated at 37 °C for 24 h. The reading was performed on an ELISA spectrophotometer (Thermoplate®).

Statistical analysis

GraphPad Prism software v.5.0 was used for statistical analysis. The data obtained were verified for their normal distribution using a two-way ANOVA ($P < 0.05$; $*P < 0.1$; $****P < 0.0001$) and Bonferroni post hoc test. The IC_{50} values were obtained by nonlinear regression analysis with interpolation of the unknown standard curve obtained from fungal growth assays as a function of extract concentration and expressed in $\mu\text{g/mL}$.

Results

Characterization of CE and fractions from *A. cearensis* seeds

Figures 1, 2, 3 and 4 show the composition of the CE and fractions of *A. cearensis* seeds determined by LC–MS/MS–QTOF. The parameters analyzed were molecular weight, ionic mass, retention time, fragmentation pattern and compared with the available literature. In this study, we identified seven compounds in the CE (Table 1), six compounds in FR-1 (Table 2), eight compounds in FR-2 (Table 3) and five compounds in FR-3 (Table 4), varying from flavonoids, phenolic acids, flavan derivatives, organic acids and phenolic compounds. The chemical structures of the compounds are shown in Fig. 5.

In addition, it was possible to identify the presence of some organic acids such as cinnamic acid and some possible *trans*-isomers in FR-1 and FR-2 (Tables 2 and 3, respectively). We evidenced the presence of some polyphenols as 4'- or 5'-*O*-methyl-(*epi*)catechin I in CE, and a polyphenol monomer trigalloyl hexoside in FR-3. The presence of the flavonoid icariin glucuronide was observed in all samples, as well as dihydroxymethoxyisoflavone, 7,8,3'-trihydroxy-4'-methoxyisoflavone and trihydroxyflavanone.


Antifungal activity of CE and fractions of *A. cearensis* seeds

All samples used alone induced inhibition growth of *C. albicans* strains (Figs. 6, 7, 8, 9), and FR-3 of *A. cearensis* seeds showed the lowest IC_{50} (123.5 $\mu\text{g/mL}$) against INCQS 40.006 (Table 5). However, Fluconazole + FR-2 showed the lowest IC_{50} (2.10 and 1.81 $\mu\text{g/mL}$), indicating that the combination of antifungal drug and FR-2 has synergistic effect against *C. albicans* (Table 5). Similar effects have been evidenced for the CE and FR-1 combined with Fluconazole against URM 4127 strain. On the other hand, we evidenced antagonistic effect when CE, FR-1 and FR-3 were combined with Fluconazole (Table 5).

Fungicidal activity

The MFC for all samples tested alone was $\geq 16.384 \mu\text{g/mL}$ for INCQS 40.006. However, fluconazole + FR-2 showed the lowest MFC (128 $\mu\text{g/mL}$) against URM 4127 strain (Table 6), which indicated a synergic effect of antifungal drug and FR-2. Conversely, CE and FR-1 combined with antifungal drug showed antagonistic effect against URM 4127 strain (Table 6).

Fig. 1 Ultra-performance liquid chromatography with high-definition mass spectrometry (UPLC–MS) chromatography of CE from *A. cearensis* seeds



Discussion

Amburana cearensis characterization

Polyphenols are secondary metabolites ubiquitously distributed among plants. They are divided in phenolic compounds, phenolic acids (cinnamic acid derivatives) and glycosidic phenylpropanoid esters (Ferreira et al. 2010). Phenolic compounds were previously reported as having antimicrobial and anti-inflammatory activities (Klančnik et al. 2010).


In addition, they exhibit antioxidant properties (Moure et al. 2001; Giada 2013; Flores et al. 2013), due to hydroxyls attached to aromatic rings. These molecules include

flavonoids, phenolic acids, tannins, and tocopherols (Soares 2002; Angelo and Jorge 2007) that act as effective scavengers of reactive oxygen species, and chelating Fe^{3+} that catalyze lipid peroxidation (Andrade et al. 2007).

On the other hand, flavonoids represent a group of plant pigments abundantly distributed in nature. Its presence in plants may have a relationship with defensive ability, such as protection against UV radiation, antifungal and antibacterial actions, as well as attracting pollinators (Lavola 1998). These compounds have two aromatic rings connected by a bridge of three carbon atoms ($\text{C}_6\text{C}_3\text{C}_6$) (Simões et al. 2007).

Icariin glucuronide is categorized as a flavonol, one of the subclasses of flavonoid, which have ability to induce bone

Fig. 2 Ultra-performance liquid chromatography with high-definition mass spectrometry (UPLC–MS) chromatography of FR-1 from *A. cearensis* seeds




tissue repair (Wei et al. 2011), human stem cell proliferation (Fan et al. 2011), immunoregulation (Kim et al. 2001), increases of cGMP in smooth cavernous muscle cells (Ning et al. 2006), increases of nitric oxide (Bin and Huang 2007), and mimics testosterone's effects (Zhang and Yang 2006).

Cinnamic acid is a naturally occurring aromatic carboxylic acid whose molecular structure is usually presented in *trans*-form. It is part of the auxin group, plant hormones that regulate cell growth and differentiation, and some of its derivatives play an important role in plant defense against pest attack (Niero 2010). It should be noted that its derivatives exhibit antioxidant (Hussain et al. 2014), antibacterial (Chiriac et al. 2005), anticancer (Ekmekcioglu et al. 1998),

antifungal (Sadeghi et al. 2013), antitumor (Lee et al. 2003) and phytotoxic effects (Nishikawa et al. 2013).

Coumarins are derived from phenylalanine metabolism, being one of the early precursors of p-hydroxycinnamic acid (p-coumaric acid), which is hydroxylated at the C-2' (ortho-hydroxylation). The ortho-hydroxylated derivative undergoes a photocatalyzed double-bond (EZ) isomerization. The Z isomer spontaneously lactonizes, producing umbelliferone. Prenylation of the benzene ring at positions 6 or 8 of the 7-hydroxy coumarin derivative is the first step in furan- and pyranocoumarin biogenesis. Cyclization of 6- or 8-isoprenylcoumarin derivatives occurs by nucleophilic attack of the C-7 hydroxyl group to the epoxide formed by oxidation of

Fig. 3 Ultra-performance liquid chromatography with high-definition mass spectrometry (UPLC–MS) chromatography of FR-2 from *A. cearensis* seeds




the double bond of the isopentenyl residue. Based on the orientation of the nucleophilic attack, the product will be hydroxyisopropyl dihydrofuranocoumarin or hydroxy dimethyl dihydrofuranocoumarin (Bourgaud et al. 2006).

Most coumarins are biogenetically derived from the shikimic acid pathway, but a significant number of them appear to derive from a mixed pathway (shikimic acid and acetate) such as phenylcoumarin. 4-*n*-Propylcoumarin, e.g., are entirely derived from the acetate pathway (Strack 1997). Coumarin synthesis can be induced by a response to biotic and abiotic stress, nutritional deficiency, chemical messengers such as plant hormones and other external metabolites (Cabello-Hurtado et al. 1998; Dewick 2009; Bosqueiro

1996; Haida et al. 2007). Coumarins have a variety of pharmacological properties, especially anti-inflammatory, antioxidant, antibacterial, antiviral, antithrombotic, antimutagenic and antihypertensive activities (Hoult and Payá 1996).

Characterization studies performed on stem bark extracts of *A. cearensis*, evidenced coumarin (Carvalho 1994), responsible for its characteristic odor, isokaempferide (Maia 2004), kaempferol and afrormosin, amburoside A (Leal et al. 1997) and B (Leal et al. 2003), vanillic acid (Bravo et al. 1999) and protocatechuic acid (Canuto et al. 2004), and large amount of sucrose. In addition, other studies show that coumarin (Carvalho 1994), isokaempferide (Maia 2004) and amburoside A (Leal et al. 1997) have anti-inflammatory,

Fig. 4 Ultra-performance liquid chromatography with high-definition mass spectrometry (UPLC–MS) chromatography of FR-3 from *A. cearensis* seeds



antioxidant, and bronchodilator effects (Leal et al. 2008). Of which some coincide with precursors of those found in our study. However, it should be noted that some identified compounds in this study are not yet reported for its biological activity.

Antifungal, fungicidal, and modulating antifungal activity

It is noteworthy that the indiscriminate and increasing use of azole antifungal drugs, both to prevent and treat active infections, has increased the prevalence of Fluconazole-resistant strains of *Candida* (Day et al. 2013). In United

States, an annual increase of > 3400 cases of antifungal resistance were identified, as reported by the Center for Disease Control and Prevention, thus making fluconazole-resistant *Candida* a serious threat (Gajdacs 2019).

Fluconazole acts by preventing the biosynthesis pathway of the main constituent of the fungal cell membrane, ergosterol. It inhibits the enzyme lanosterol 14- α demethylase in the fungal cytochrome P-450 enzyme system, which is encoded by the *ERG11* gene. Thus, lanosterol cannot be converted to ergosterol and consequently the accumulation of precursors occurs causing fungal membrane instability (Carrillo-Munoz et al. 2006; Menozzi et al. 2017).

Table 1 Compounds identified by UPLC-ESI-TOFMS/MS in positive mode in the CE of *A. cearensis* seeds

Peak no.	Retention time (min)	[M-H] ⁻ observed	[M-H] ⁻ calculated	Product ions (MS/MS)	Empirical formula	Parts per million (error)	Putative name	References
1	1.18	689.2051	689.2082	689.2087, 367.1008, 352.8637	C ₃₃ H ₃₇ O ₁₆	4.5	Icariin glucuronide	Shunjun et al. (2017)
2	1.29	527.1564	527.1553	202.1848, 184.9314, 178.0209	C ₂₇ H ₂₇ O ₁₁	2.1	QQCA (quinic-quinic-caffeic acid ester)	Plazonić et al. (2009)
3	2.94	467.1732	467.1706	329.0866, 261.0790, 179.0338, 137.0256	C ₂₁ H ₂₃ O ₁₂	2.2	(Epi)gallo catechin hexose II	Ibrahim et al. 2014
4	3.11	305.0735	305.0720	179.0339, 167.0341, 137.0252, 125.0241	C ₁₅ H ₁₃ O ₇	4.9	Galocatechin	Ibrahim et al. (2014)
5	3.32	172.0376	172.0372	242.9876, 144.0423, 116.0501	C ₆ H ₂ N ₇	2.3	No identified	–
6	4.93	218.2143	218.2120	177.0513, 145.0170, 225.9959	C ₁₂ H ₂₈ NO ₂	10.5	No identified	–
7	6.09	246.2422	246.2433	258.8799, 184.0619, 156.0619	C ₁₄ H ₃₂ NO ₂	4.5	No identified	–
8	7.25	247.2733	247.2746	275.2783, 256.2661, 230.2536	C ₁₆ H ₃₆ NO ₂	4.7	No identified	–
9	7.96	283.1002	283.1029	295.0207, 240.0207, 133.0802	C ₁₆ H ₁₂ O ₅	9.5	Dihydroxy-methoxyisoflavone	Oliveira et al. (2017)
10	8.26	119.0829	119.0821	437.2041, 185.1608, 120.0852	C ₄ H ₁₁ N ₂ O ₂	6.7	No identified	–
11	8.46	303.0885	303.0874	137.0244	C ₁₆ H ₁₅ O ₆	3.6	4'- or 5'-O-Methyl-(epi)catechin I	Ibrahim et al. (2014)
12	9.27	393.3636	393.3639	334.3075, 249.1913, 133.0866	C ₁₃ H ₄₁ N ₁₄	0.8	No identified	–
13	9.69	332.3412	332.28	330.3311, 270.9351, 150.7104		1.0	Gallic-caffeic acid ester	Plazonić et al. (2009)
14	9.84	457.2762	457.2743	155.0720, 458.2881	C ₃₁ H ₃₇ O ₃	4.2	No identified	–
15	9.89	457.2829	457.2801	155.0675, 458.2868	C ₂₄ H ₄₁ O ₈	6.1	No identified	–

Alternatives for treating *Candida* infections are considered limited and many existing antifungal drugs have undesirable side effects such as high toxicity, especially in immunosuppressed individuals (Fica 2004). For these patients, antifungal therapies are a challenging situation, which represent an urgent need to research and develop novel antifungal drugs more effective and less toxic (Calabrese et al. 2013).

Bravo et al. (1999) already shown that the main component of *A. cearensis* was coumarin, mainly responsible for bronchodilator activity. It also highlights the presence of phenolic glycosides that presented antimalarial, anti-protozoal, antifungal, and antibacterial activity in vitro. In addition, Salas et al. (2011) using natural and enzymatically modified flavonoids from plants against *Penicillium*, *Aspergillus* and *Fusarium* species evidenced antifungal activity.

However, Steiner et al. (2008) showed that antifungal activity induced by isoflavone, especially genistein, demonstrated a better effect, but without a clear mechanism of action.

The combined use of drugs that can achieve synergistic results, showing increased of therapeutic effect, decreased minimal inhibitory concentration, regression of antibiotic resistance development, and a decrease in host toxicity has become increasingly studied as an alternative for the treatment of infections (Silva et al. 2015).

Some phenolic compounds such as gallic acid, catechin, luteolin and quercetin demonstrated in vitro antifungal activity against different *Candida* species, including *C. albicans* and *C. tropicalis* (Alves et al. 2014), corroborating with this study. In addition, Silva et al. (2014) evaluated the flavonoid combined with Fluconazole in in vitro microdilution against Fluconazole-resistant *C. tropicalis*, and observed a

Table 2 Compounds identified by UPLC–ESI–TOFMS/MS in positive mode in the FR-1 of *A. cearensis* seeds

Peak no.	Retention time (min)	[M–H] ⁺ observed	[M–H] ⁺ calculated	Product ions (MS/MS)	Empirical formula	Parts per million (error)	Putative name	References
1	1.18	689.2130	689.2117	689.2087, 367.1008, 352.8637	C ₃₃ H ₃₇ O ₁₆	4.5	Icarin glucuronide	Shunjun et al. (2017)
2	1.94	453.7779	453.7796	454.7751, 451.7924, 449.7784	C ₆ NO ₁₁ S ₆	3.7	No identified	–
3	5.35	147.0405	147.0446	147.0417, 62.1057	C ₉ H ₈ O ₂	–27.0	<i>Trans</i> -cinnamic acid	Ibrahim et al. (2014)
4	5.37	147.0408	147.0446	147.0417, 62.1057	C ₉ H ₈ O ₂	–25.0	Possible isomer <i>trans</i> -cinnamic acid	Ibrahim et al. (2014)
5	5.39	147.0407	147.0446	147.0417, 62.1057	C ₉ H ₈ O ₂	–	Possible isomer <i>trans</i> -cinnamic acid	Ibrahim et al. (2014)
6	5.40	147.0417	147.0446	147.0417, 62.1057	C ₉ H ₈ O ₂	–19.0	Possible isomer <i>trans</i> -cinnamic acid	Ibrahim et al. (2014)
7	5.62	147.0417	147.0446	147.0417, 62.1057	C ₉ H ₈ O ₂	–19.0	Possible isomer <i>trans</i> -cinnamic acid	Ibrahim et al. (2014)
8	6.65	299.0892	299.0919	284.0323, 256.0673, 184.0652	C ₁₇ H ₁₅ O ₃	9.0	7,8,3'-Trihydroxy-4'-methoxyisoflavone	Oliveira et al. (2017)
9	7.34	274.2764	274.2746	318.3075, 274.2844, 256.2657	C ₁₆ H ₃₆ NO ₂	6.6	No identified	–
10	7.70	313.1020	313.1017	269.0793, 252.0761, 184.0693	C ₁₈ H ₁₇ O ₃	1.0	7,3'-Dihydroxy-8,4'-dimethoxyisoflavone	Oliveira et al. (2017)
11	8.02	283.0938	283.0970	240.0811, 211.0820, 197.0556	C ₁₇ H ₁₅ O ₄	11.3	Dihydroxymethoxyisoflavone	Oliveira et al. (2017)
12	8.33	119.0855	119.0861	113.0861, 117.1045	C ₉ H ₁₁	–5.0	No identified	–
13	8.62	302.3014	302.3059	184.0782, 177.1111, 113.0846	C ₁₈ H ₄₀ NO ₂	–6.0	No identified	–
14	8.80	391.3424	391.3424	331.2966, 177.1157, 133.0889	C ₂₂ H ₄₇ O ₃	0.0	No identified	–
15	9.25	323.2748	323.2797	373.1831, 339.2472, 133.0887	C ₁₇ H ₃₉ O ₃	–15.2	No identified	–
16	9.44	393.3588	393.3588	393.3560, 334.3107, 177.1114	C ₂₂ H ₄₉ O ₃	2.0	No identified	–
17	9.96	330.3365	330.3372	258.0484, 312.3296, 331.3336	C ₂₀ H ₄₄ NO ₂	–2.1	No identified	–

Table 3 Compounds identified by UPLC–ESI–TOFMS/MS in positive mode in the FR-2 of *A. cearensis* seeds

Peak no	Retention time (min)	[M-H] ⁻ observed	[M-H] ⁻ calculated	Product ions (MS/MS)	Empirical formula	Parts per million (error)	Putative name	References
1	1.17	689.2051	689.2082	689.2087, 367.1008, 352.8637	C ₃₃ H ₃₇ O ₁₆	4.5	Icariin glucuronide	Shunjun et al. (2017)
2	4.03	591.3693	591.3686	401.1527, 172.8673, 133.0836	C ₃₇ H ₅₁ O ₆	1.2	No identified	–
3	5.37	147.0419	147.0446	147.0422, 62.2985	C ₉ H ₇ O ₂	– 18.4	<i>Trans</i> -cinnamic acid	Ibrahim et al. (2014)
4	5.40	147.0412	147.0446	147.0422, 62.2985	C ₉ H ₇ O ₂	– 23.1	Possible isomer <i>trans</i> -cinnamic acid	Ibrahim et al. (2014)
5	5.49	147.0408	147.0446	147.0422, 62.2985	C ₉ H ₇ O ₂	– 23.1	Possible isomer <i>trans</i> -cinnamic acid	Ibrahim et al. (2014)
6	6.12	246.2432	246.2433	247.4212, 184.0706, 156.0435	C ₁₄ H ₃₂ NO ₂	– 0.4	No identified	Ibrahim et al. (2014)
7	6.51	285.2693	285.2700	269.7, 163.8, 150.5	C ₁₅ H ₁₀ O ₆	– 2.5	Hydroxygenistein	Scheda-Hirschmann et al. (2019)
8	6.63	299.0910	299.0919	283.7, 230.7	C ₁₇ H ₁₅ O ₅	– 3.0	Methoxy trihydroxy flavanone	Scheda-Hirschmann et al. (2019)
9	7.26	274.2721	274.2719	275.2858, 256.2728, 184.0647	C ₁₂ H ₃₂ N ₇	0.7	No identified	–
10	7.28	274.2730	274.2719	275.2858, 256.2728, 184.0647	C ₁₂ H ₃₂ N ₇	4.0	No identified	–
11	7.66	313.1066	313.1076	269.0793, 252.0761, 184.0693	C ₁₈ H ₁₇ O ₅	– 3.2	7,3'-Dihydroxy-8,4'-dimethoxyisoflavone	Oliveira et al. (2017)
12	7.99	283.0953	283.0970	240.0811, 211.0820, 197.0556	C ₁₇ H ₁₅ O ₄	– 6.0	Dihydroxymethoxyisoflavone	Oliveira et al. (2017)
13	8.02	283.0921	283.0970	240.0811, 211.0820, 197.0556	C ₁₇ H ₁₅ O ₄	– 17.3	Dihydroxymethoxyisoflavone possible isomer	Oliveira et al. (2017)
14	8.29	119.0850	119.0861	113.0861, 117.1045	C ₉ H ₁₁	– 9.2	No identified	–
15	8.52	302.3076	302.3059	184.0782, 177.1111, 113.0846	C ₁₈ H ₄₀ NO ₂	5.6	No identified	–
16	8.67	391.3500	391.3509	184.0709, 177.1171	C ₁₇ H ₄₃ N ₈ O ₂	– 2.3	No identified	–
17	9.23	323.2783	323.2797	373.1831, 339.2472, 133.0887	C ₁₇ H ₃₉ O ₅	– 4.3	No identified	–
18	9.32	393.3557	393.3580	393.3560, 334.3107, 177.1114	C ₂₂ H ₄₉ O ₅	– 5.8	No identified	–
19	9.35	393.3589	393.3580	393.3560, 334.3107, 177.1114	C ₂₂ H ₄₉ O ₅	2.3	No identified	–
20	9.75	330.3365	330.3372	258.0484, 312.3296, 331.3336	C ₂₀ H ₄₄ NO ₂	– 2.1	No identified	–
21	9.90	457.2744	457.2743	155.0720, 458.2881	C ₃₁ H ₃₇ O ₃	0.2	No identified	–
22	9.95	457.2778	457.2743	155.0720, 458.2881	C ₃₁ H ₃₇ O ₃	7.7	No identified	–
23	10.11	395.3740	395.3750	335.3429, 184.0704, 133.0864	C ₂₃ H ₄₇ N ₄ O	– 2.5	No identified	–

Table 4 Compounds identified by UPLC–ESI–TOFMS/MS in positive mode in the FR-3 of *A. cearensis* seeds

Peak no	Retention time (min)	[M-H] ⁻ observed	[M-H] ⁻ calculated	Product ions (MS/MS)	Empirical formula	Parts per million (error)	Putative name	References
1	1.17	689.2147	689.2140	689.2087, 367.1008, 352.8637	C ₃₃ H ₃₇ O ₁₆	1.0	Icariin glucuronide	Shunjun et al. (2017)
2	1.25	136.0607	136.0610	152.0563, 136.0607	C ₄ H ₁₀ NO ₄	-2.2	No identified	-
3	1.72	136.0612	136.0610	152.0563, 136.0607	C ₄ H ₁₀ NO ₄	1.5	No identified	-
4	2.64	172.0363	172.0372	242.9876, 144.0423, 116.0501	C ₆ H ₂ N ₇	-5.2	No identified	-
5	3.11	433.1133	433.1135	300.9686	C ₂₁ H ₂₁ O ₁₀	-0.5	Ellagic acid pentoside 1	Scheda-Hirschmann et al. (2019)
6	3.54	287.0533	287.0556	287.0538, 241.0332, 175.0282	C ₁₅ H ₁₁ O ₆	8.0	Luteolin	Ibrahim et al. (2014)
7	3.71	133.0901	133.0865	207.0626, 151.0330, 133.0813	C ₆ H ₁₃ O ₃	2.7	No identified	-
8	3.81	133.0860	133.0865	207.0626, 151.0330, 133.0813	C ₆ H ₁₃ O ₃	-3.5	No identified	-
9	4.02	591.3591	591.3592	207.0608, 151.0291, 133.0836	C ₂₆ H ₅₅ O ₁₄	-0.2	No identified	-
10	4.13	635.3927	635.3927	465.0265	C ₃₉ H ₅₅ O ₇	-3.3	Trigalloyl hexoside	Scheda-Hirschmann et al. (2019)
11	4.35	207.0683	207.0657	369.1195, 207.0621, 175.0858	C ₁₁ H ₁₁ O ₄	12.6	No identified	-
12	4.37	207.0658	207.0657	369.1195, 207.0621, 175.0858	C ₁₁ H ₁₁ O ₄	0.5	No identified	-
13	4.91	207.0661	207.0657	240.0811, 211.0820, 197.0556	C ₁₁ H ₁₁ O ₄	1.9	No identified	-
14	6.11	246.2453	246.2433	258.8799, 184.0619, 156.0619	C ₁₄ H ₃₂ NO ₂	8.1	No identified	-
15	6.52	283.0364	283.1029	295.0207, 240.0207, 133.0802	C ₁₆ H ₁₂ O ₅	9.5	Dihydroxymethoxyisoflavone	Oliveira et al. (2017)
16	7.32	274.2742	274.2746	318.3075, 274.2844, 256.2657	C ₁₆ H ₃₆ NO ₂	-1.5	No identified	-
17	7.38	318.2995	318.2981	274.2706, 256.2648	C ₁₄ H ₃₆ N ₇ O	4.4	No identified	-
18	8.31	119.0870	119.0821	437.2041, 185.1608, 120.0852	C ₄ H ₁₁ N ₂ O ₂	6.7	No identified	-
19	8.56	302.3055	302.3059	184.0782, 177.1111, 113.0846	C ₁₈ H ₄₀ NO ₂	-1.3	No identified	-

Table 4 (continued)

Peak no	Retention time (min)	[M-H] ⁻ observed	[M-H] ⁻ calculated	Product ions (MS/MS)	Empirical formula	Parts per million (error)	Putative name	References
20	9.17	304.3001	304.3004	212.2469, 133.0871	C ₂₁ H ₃₈ N	-1.0	No identified	-
21	9.80	330.3346	330.3372	258.0484, 312.3296, 331.3336	C ₂₀ H ₄₄ NO ₂	-2.1	No identified	-
22	9.93	457.2736	457.2743	155.0720, 458.2881	C ₃₁ H ₃₇ O ₃	-1.5	No identified	-
23	9.95	457.2711	457.2743	155.0720, 458.2881	C ₃₁ H ₃₇ O ₃	-7.0	No identified	-


**Fig. 5** Main compounds identified in CE, FR-1, FR-2 and FR-3 of *A. cearensis* seeds. **a** aqueous crude extract; **b** fraction 1; **c** fraction 2; and **d** fraction 3

Fig. 6 Antifungal effects ($\mu\text{g}/\text{mL}$) of aqueous crude extract and fractions of *A. cearensis* seeds against *C. albicans* 40006. Fluconazole was used as a control against *C. albicans*




Fig. 7 Antifungal effects ($\mu\text{g}/\text{mL}$) of aqueous crude extract and fractions of *A. cearensis* seeds combined with fluconazole against *C. albicans* 40006. Fluconazole was also used as a control against *C. albicans*




Fig. 8 Antifungal effects ($\mu\text{g}/\text{mL}$) of aqueous crude extract and fractions of *A. cearensis* seeds against *C. albicans* 4127. Fluconazole was used as a control against *C. albicans*




Fig. 9 Antifungal effects ($\mu\text{g}/\text{mL}$) of aqueous crude extract and fractions of *A. cearensis* seeds combined with fluconazole against *C. albicans* 4127. Fluconazole was also used as a control against *C. albicans*




Table 5 IC₅₀ (µg/mL) of CE and fractions from *A. cearensis* seeds against *C. albicans* strains

Evaluated combination	CA INCQS 40006	CA URM 4127
FCZ	2.4 ± 1.2	7.68 ± 4.8
CE— <i>A. cearensis</i>	> 2048 ± 3.6	> 2048 ± 3.2
FR-1— <i>A. cearensis</i>	1968.6 ± 9.5	1138.4 ± 4.8
FR-2— <i>A. cearensis</i>	1173.3 ± 8.1	1273.3 ± 6.3
FR-3— <i>A. cearensis</i>	123.5 ± 5.5	45.6 ± 6.1
CE— <i>A. cearensis</i> + FCZ	9.54 ± 4.0	3.14 ± 2.0
FR-1— <i>A. cearensis</i> + FCZ	3.51 ± 2.5	2.41 ± 1.8
FR-2— <i>A. cearensis</i> + FCZ	2.1 ± 0.99	1.81 ± 0.8
FR-3— <i>A. cearensis</i> + FCZ	11.9 ± 8.7	18.8 ± 7.4

CA *Candida albicans*, FCZ fluconazole, CE aqueous crude extract, FR-1 fraction 1, FR-2 fraction 2, FR-3 fraction 3, MIC minimum inhibitory concentration, INCQS National Institute of Health Control, URM University Recife Mycology— $P \leq 0.05$ in relation to FCZ control

Table 6 Minimal fungicidal concentrations (MFC) in µg/mL to fluconazole (FCZ) alone and combined with CE and fractions from *A. cearensis* seeds

	CA URM 4127
Fluconazole	2048
<i>A. cearensis</i> CE + FCZ	2048
<i>A. cearensis</i> FR-1 + FCZ	8192
<i>A. cearensis</i> FR-2 + FCZ	128
<i>A. cearensis</i> FR-3 + FCZ	8192

considerable synergistic effect, reducing the MICs of the flavonoids from 64 to 0.25 µg/mL, similar to our results against *C. albicans*.

Conclusion

CE, FR-1, FR-2 and FR-3 of *A. cearensis* seeds presented phenolic compounds, organic acids, and polyphenols, that contributed to its antifungal capacity and synergistic effects. Our results show that *A. cearensis* seeds can be considered as promising source of antifungal agents and modulators of antifungal activity. However, further studies are needed to identify the active compounds and to prove their efficacy, safety, and mechanism of action.

Acknowledgements The authors are grateful to the Brazilian research agencies FUNCAP, CAPES, CNPq and FINEP for the support of this work.

Author contributions Methodology—antimicrobial assays (MTAO; MVOBA and GMMM); methodology—chemical analysis (PRVR; ESB and EOS); methodology—statistical analysis (IRAM; JCA and EAS); supervision of work—(HDMC and AFU); resources (VPAL and JIOC).

Compliance with ethical standards

Ethical statements This article is according to the international, national and institutional rules considering biodiversity rights.

References

- Agra MF, Baracho GS, Nurit K, Basílio IJLD, Coelho VPM (2007) Medicinal and poisonous diversity of the flora of “Cariri Paraíba”, Brazil. *J Ethnopharmacol* 111:383–395
- Alves CT, Ferreira ICFR, Barros L, Silva S, Azeredo J, Henriques M (2014) Antifungal activity of phenolic compounds identified in flowers from North Eastern Portugal against *Candida* species. *Future Microbiol* 9:139–146. <https://doi.org/10.2217/fmb.13.147>
- Andrade CA, Costa CK, Bora K, Miguel MD, Miguel OG, Kerber VA (2007) Determination of the phenolic content and evaluation of the antioxidant activity of *Acacia podalyriifolia* A. Cunn. ex G. Don, Leguminosae-mimosoideae. *Revista Brasileira de Farmacognosia* 17:231–235. <https://doi.org/10.1590/s0102-695x2007000200017>
- Angelo PM, Jorge N (2007) Phenolic compounds in foods—a brief review. *Rev Inst Adolfo Lutz* 66:1–9
- Bin XuH, Huang ZQ (2007) Icarin enhances endothelial nitric-oxide synthase expression on human endothelial cells in vitro. *Vascul Pharmacol* 47:18–24. <https://doi.org/10.1016/j.vph.2007.03.002>
- Bosqueiro ALD (1996) Plantas como fonte de fitofármaco. *Ciência Educação* 02:91–96
- Bourgaud F, Hehn A, Larbat R, Doerper S, Gontier E, Kellner S, Matern U (2006) Biosynthesis of coumarins in plants: a major pathway still to be unravelled for cytochrome P450 enzymes. *Phytochem Rev* 5:293–308. <https://doi.org/10.1007/s11101-006-9040-2>
- Bravo JAB, Sauvain M, Gimenez AT, Muñoz VO, Callapa J, Le Men-Olivier L, Massiot G, Lavaud C (1999) Bioactive phenolic glycosides from *Amburana cearensis*. *Phytochemistry* 50:71–74. [https://doi.org/10.1016/S0031-9422\(98\)00497-X](https://doi.org/10.1016/S0031-9422(98)00497-X)
- Cabello-Hurtado F, Durst F, Jorrín JV, Werck-Reichhart D (1998) Coumarins in *Helianthus tuberosus*: characterization, induced accumulation and biosynthesis. *Phytochemistry* 49:1029–1036. [https://doi.org/10.1016/S0031-9422\(97\)01036-4](https://doi.org/10.1016/S0031-9422(97)01036-4)
- Calabrese EC, Castellano S, Santoriello M, Sgheri C, Quartacci MF, Calucci L, Warrilow AGS, Lamb DC, Kelly SL, Milite C, Granata I, Sbardella G, Stefancich G, Maresca B, Porta A (2013) Antifungal activity of azole compounds CPA18 and CPA109 against azole-susceptible and -resistant strains of *Candida albicans*. *J Antimicrob Chemother* 68:1111–1119. <https://doi.org/10.1093/jac/dks506>
- Calixto Júnior JT, Morais SM, Martins CG, Vieira LG, Morais-Braga MFB, Carneiro JNP, Machado AJP, Menezes IRA, Tintino SR, Coutinho HDM (2015) Phytochemical analysis and modulation of antibiotic activity by *Luehea paniculata* Mart. & Zucc. (Malvaceae) in multiresistant clinical isolates of *Candida* Spp. *BioMed Research International* 2015:1–10
- Canuto KM, Leal LKAM, Viana GSB, Bezerra AME, Silveira E (2004) Aspectos químicos do estudo interdisciplinar (química, agronomia e farmacologia) de *A. cearensis*. In: XVIII Simpósio de Plantas Medicinais Do Brasil
- Carrillo-Muñoz AJ, Giusiano G, Ezkurra PA, Quindós G (2006) Antifungal agents: mode of action in yeast cells. *Rev Esp Quimioterap* 19:130–139
- Carvalho P (1994) Espécies Florestais Brasileiras : recomendações silviculturais, potencialidades e uso da madeira. EMBRAPA

- Chiriac CI, Tanasa F, Onciu M (2005) A novel approach in cinnamic acid synthesis: direct synthesis of cinnamic acids from aromatic aldehydes and aliphatic carboxylic acids in the presence of boron tribromide. *Molecules* 10:481–487. <https://doi.org/10.3390/10020481>
- Coutinho HDM, Costa JGM, Siqueira-Júnior JP, Lima EO (2008) In vitro anti-staphylococcal activity of *Hyptis martiusii* Benth against methicillin-resistant *Staphylococcus aureus*-MRSA strains. *Braz J Pharmacogn* 18:670–675
- Day JN, Chau TTH, Wolbers M, Mai PP, Dung NT, Mai NH, Phu NH, Nghia HD, Phong ND, Thai CQ, Thai LH, Chuong LV, Sinh DX, Duong VA, Hoang TN, Diep PT, Campbell JI, Sieu TPM, Baker SG, Chau NVV, Hien TT, Laloo DG, Farrar JJ (2013) Combination antifungal therapy for cryptococcal meningitis. *N Engl J Med* 368:1291–1302. <https://doi.org/10.1056/NEJMoa1110404>
- Dewick PM (2009) Medicinal natural products: a biosynthetic approach, 3rd edn. Wiley
- Ekmekcioglu C, Feyertag J, Marktl W (1998) Cinnamic acid inhibits proliferation and modulates brush border membrane enzyme activities in Caco-2 cells. *Cancer Lett* 128:137–144. [https://doi.org/10.1016/S0304-3835\(98\)00073-1](https://doi.org/10.1016/S0304-3835(98)00073-1)
- Ernst EJ, Klepser ME, Ernst ME, Messer SA, Pfaller MA (1999) In vitro pharmacodynamic properties of MK-0991 determined by time-kill methods. *Diagn Microbiol Infect Dis* 33:75–80
- Fan JJ, Cao LG, Wu T, Wang DX, Jin D, Jiang S, Zhang ZY, Bi L, Pei GX (2011) The dose-effect of icariin on the proliferation and osteogenic differentiation of human bone mesenchymal stem cells. *Molecules* 16:10123–10133. <https://doi.org/10.3390/molecules161210123>
- Ferreira JFS, Luthria DL, Sasaki T, Heyerick A (2010) Flavonoids from *Artemisia annua* L. As antioxidants and their potential synergism with artemisinin against malaria and cancer. *Molecules* 15:3135–3170. <https://doi.org/10.3390/molecules15053135>
- Fica CA (2004) Tratamiento de infecciones fúngicas sistémicas Primera parte: fluconazol, itraconazol y voriconazol. *Revista Chilena de Infectología* 21:26–38. <https://doi.org/10.4067/s0716-10182004000100004>
- Flores G, Dastmachi K, Wu SB, Whalen K (2013) Phenolic-rich extract from the Costa Rican guava (*Psidium friedrichsthalianum*) pulp with antioxidant and anti-inflammatory activity. Potential for COPD therapy. *Food Chem* 141:889–895. <https://doi.org/10.1016/j.foodchem.2017.03.040>
- Gajdác M (2019) The concept of an ideal antibiotic: implications for drug design. *Molecules*. <https://doi.org/10.3390/molecules24050892>
- Giada MLR (2013) Food phenolic compounds: main classes, sources and their antioxidant power. InTech, Croatia. <https://doi.org/10.5772/51687>
- Goulart LS, de Souza WWR, Vieira CA, de Lima JS, de Olinda RA, de Araújo C (2018) Oral colonization by *Candida* species in HIV-positive patients: association and antifungal susceptibility study. *Einstein*, Sao Paulo, p 16
- Haida K, Parzianello L, Werner S, Garcia D (2007) Avaliação in vitro da atividade antimicrobiana de oito espécies de plantas medicinais. *Arq Cienc Saude Unipar* 11:185–192
- Hoult JRS, Payá M (1996) Pharmacological and biochemical actions of simple coumarins: natural products with therapeutic potential. *Gen Pharmacol* 27:713–722. [https://doi.org/10.1016/0306-3623\(95\)02112-4](https://doi.org/10.1016/0306-3623(95)02112-4)
- Hussain SH, Latif A, Cox RJ, Simpson TJ, Ali M, Arfan M, Uddin G (2014) Phytochemicals from the aerial parts of *Ligularia thomsonii* and their radical scavenging activity. *Phytochem Lett* 7:6–10. <https://doi.org/10.1016/j.phytol.2013.09.002>
- Ibrahim M, Abu-Reidah A, Contreras M, Arraez-Roman D, Fernandez-Gutierrez A, Segura-Carretero A (2014) UHPLC-ESI-QTOF-MS-based metabolic profiling of *Vicia faba* L. (Fabaceae) seeds as a key strategy for characterization in foodomics. *Electrophoresis* 35:1571–1581. <https://doi.org/10.1002/elps.201300646>
- Javadpour MM, Juban MM, Lo WCJ, Bishop SM, Alberty JB, Cowell SM, Becker CL, McLaughlin ML (1996) De novo antimicrobial peptides with low mammalian cell toxicity. *J Med Chem* 39:3107–3113
- Kim JH, Mun YJ, Im SJ, Han JH, Lee HS, Woo WH (2001) Effects of the aqueous extract of Epimedium Herba on the antibody responses in mice. *Int Immunopharmacol* 1:935–944. [https://doi.org/10.1016/S1567-5769\(01\)00030-3](https://doi.org/10.1016/S1567-5769(01)00030-3)
- Klančnik A, Piskernik S, Jeršek B, Možina SS (2010) Evaluation of diffusion and dilution methods to determine the antibacterial activity of plant extracts. *J Microbiol Methods* 81:121–126. <https://doi.org/10.1016/j.mimet.2010.02.004>
- Ksiezopolska E, Gabaldón T (2018) Evolutionary emergence of drug resistance in *Candida* opportunistic pathogens. *Genes* 9:461–486
- Ksouri S, Djebir S, Bentorki AA, Gouri A, Hadeif Y, Benakhla A (2017) Antifungal activity of essential oils extract from *Origanum floribundum* Munby, *Rosmarinus officinalis* L. and *Thymus ciliatus* Desf. against *Candida albicans* isolated from bovine clinical mastitis. *J Mycol Med* 27:245–249
- Lavola A (1998) Accumulation of flavonoids and related compounds in birch induced by UV-B irradiance. *Tree Physiol* 18:53–58. <https://doi.org/10.1093/treephys/18.1.53>
- Leal LKAM, Matos ME, Matos FJA, Ribeiro RA, Ferreira FV, Viana GSB (1997) Antinociceptive and anti-inflammatory effects of the hydroalcoholic extract and coumarin from *Torresea cearensis* Fr. *All. Phytomedicine* 4:221–227. [https://doi.org/10.1016/S0944-7113\(97\)80071-2](https://doi.org/10.1016/S0944-7113(97)80071-2)
- Leal LKAM, Nechio M, Silveira ER, Canuto KM, Fontenele JB, Ribeiro RA, Viana GSB (2003) Anti-inflammatory and smooth muscle relaxant activities of the hydroalcoholic extract and chemical constituents from *Amburana cearensis* A. C. Smith. *Phytother Res* 17:335–340
- Leal LKAM, Fonseca FN, Pereira FA, Canuto KM, Felipe CFB, Fontenele JB, Pitombeira MV, Silveira ER, Viana GSB (2008) Protective effects of amburoside A, a phenol glucoside from *Amburana cearensis*, against CCl₄-induced hepatotoxicity in rats. *Planta Med* 74:497–502. <https://doi.org/10.1055/s-2008-1074501>
- Lee YJ, Kuo HC, Chu CY, Wang CJ, Lin WC, Tseng TH (2003) Involvement of tumor suppressor protein p53 and p38 MAPK in caffeic acid phenethyl ester-induced apoptosis of C6 glioma cells. *Biochem Pharmacol* 66:2281–2289. <https://doi.org/10.1016/j.bcp.2003.07.014>
- Lima LR, Cavalcante RRL, Martins MCC, Parente DM, Cavalcante AAMCC (2013) Avaliação da atividade anti-inflamatória, antimicrobiana e mutagênica das sementes de *Amburana cearensis* (A. C. Smith) (Imburana-de-cheiro). *Revista Brasileira de Plantas Medicinais* 15:415–422
- Maia GN (2004) Caatinga: Árvores e Arbustos e Suas Utilidades. Print-color gráfica e Editora, Fortaleza
- Menozi CAC, Castelo-Branco FS, França RRF, Domingos JLO, Boechat N (2017) Optimization of fluconazol synthesis: an important azole antifungal drug. *Revista Virtual de Química* 9:1216–1234. <https://doi.org/10.21577/1984-6835.20170071>
- Moras-Braga MFB, Carneiro JNP, Machado AJT, dos Santos ATL, Sales DL, Lima LF, Figueredo FG, Coutinho HDM (2016) *Psidium guajava* L., from ethnobiology to scientific evaluation: Elucidating bioactivity against pathogenic microorganisms. *J Ethnopharmacol* 194:1140–1152
- Moure A, Cruz JM, Franco D, Manuel Domínguez J, Sineiro J, Domínguez H, Núñez MJ, Carlos Parajó J (2001) Natural antioxidants from residual sources. *Food Chem* 72:145–171. [https://doi.org/10.1016/S0308-8146\(00\)00223-5](https://doi.org/10.1016/S0308-8146(00)00223-5)

- NCCLS (2012) Método de Referência para Testes de Diluição em Caldo para Determinação da Sensibilidade de Leveduras à Terapia Antifúngica : Norma Aprovada-Segunda Edição
- Niero ELO (2010) Efeitos de ácido cinâmico sobre melanócitos e células derivadas de melanoma humanos: avaliação do seu potencial antitumoral e de proteção contra danos celulares causados por radiação ultravioleta. Tese de doutorado. <https://doi.org/10.11606/T.42.2010.tde-28092010-111234>
- Ning H, Xin ZC, Lin G, Banie L, Lue TF, Lin CS (2006) Effects of icariin on phosphodiesterase-5 activity in vitro and cyclic guanosine monophosphate level in cavernous smooth muscle cells. *Urology* 68:1350–1354. <https://doi.org/10.1016/j.urology.2006.09.031>
- Nishikawa K, Fukuda H, Abe M, Nakanishi K, Taniguchi T, Nomura T, Yamaguchi C, Hiradate S, Fujii Y, Okuda K, Shindo M (2013) Substituent effects of *cis*-cinnamic acid analogues as plant growth inhibitors. *Phytochemistry* 96:132–147. <https://doi.org/10.1016/j.phytochem.2013.08.013>
- Oliveira GP, Silva TMG, Camara CA, Santana ALBD, Moreira MSA, Silva TMS (2017) Isolation and structure elucidation of flavonoids from *Amburana cearensis* resin and identification of human DNA topoisomerase II- α inhibitors. *Phytochem Lett* 22:61–70. <https://doi.org/10.1016/j.phytol.2017.09.006>
- Pereira EPL, Braga-De-Souza S, Santos CC, Santos LO, Cerqueira MD, Ribeiro PR, Fernandez LG, Silva VDA, Costa SL (2017) *Amburana cearensis* seed extracts protect PC-12 cells against toxicity induced by glutamate. *Braz J Pharmacogn* 27:199–205
- Plazonić A, Bucar F, Maleš Z, Mornar A, Nigović B, Kujundžić N (2009) Identification and quantification of flavonoids and phenolic acids in Burr Parsley (*Caucalis platycarpos* L.), using high-performance liquid chromatography with diode array detection and electrospray ionization mass spectrometry. *Molecules* 14:2466–2490
- Sadeghi M, Zolfaghari B, Senatore M, Lanzotti V (2013) Antifungal cinnamic acid derivatives from Persian leek (*Allium ampeloprasum* Subsp. *Persicum*). *Phytochem Lett* 6:360–363. <https://doi.org/10.1016/j.phytol.2013.04.007>
- Salas MP, Céliz G, Geronazzo H, Daz M, Resnik SL (2011) Antifungal activity of natural and enzymatically-modified flavonoids isolated from citrus species. *Food Chem* 124:1411–1415. <https://doi.org/10.1016/j.foodchem.2010.07.100>
- Schmeda-Hirschmann G, Burgos-Edwards A, Theoduloz C, Jiménez-Aspee F, Vargas-Arana G (2019) Male sexual enhancers from the Peruvian Amazon. *J Ethnopharmacol* 229:167–179. <https://doi.org/10.1016/j.jep.2018.10.007>
- Shunjun X, Yu J, Zhan J, Yang L, Guo L, Xu Y (2017) Pharmacokinetics, tissue distribution, and metabolism study of icariin in rat. *Biomed Res Int* 2017:1–17
- Silva CR, De Andrade NJB, De Sousa CR, Figueiredo NS, Sampaio LS, Magalhães HIF, Cavalcanti BC, Gaspar DM, De Andrade GM, Lima ISP, De Barros VGS, De Moraes MO, Lobo MDP, Grangeiro TB, Nobre HV (2014) Synergistic effect of the flavonoid catechin, quercetin, or epigallocatechin gallate with fluconazole induces apoptosis in *Candida tropicalis* resistant to fluconazole. *Antimicrob Agents Chemother* 58:1468–1478. <https://doi.org/10.1128/AAC.00651-13>
- Silva KMA, Chaves TP, Santos RL, Brandão DO, Fernandes FHA, de Ramos JFJL, Medeiros ACD, dos Santos VL, Felismino DC (2015) Modulation of the erythromycin resistance in *Staphylococcus aureus* by ethanolic extracts of *Ximenesia americana* L and *Schinopsis brasiliensis* Engl. *Bol Latinoam Caribe Plant Med Aromat* 14:92–98
- Simões C, Schenkel E, Gosman G (2007) Farmacognosia: da planta ao medicamento. Editora UFRGS 5
- Siqueira JF, Sen BH (2004) Fungi in endodontic infections. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 97:632–641
- Soares SE (2002) Phenolic acids as antioxidants. *Revista de Nutricao* 15:71–81. <https://doi.org/10.1590/s1415-52732002000100008>
- Steiner C, Arnould S, Scalbert A, Manach C (2008) Isoflavones and the prevention of breast and prostate cancer: new perspectives opened by nutrigenomics. *Br J Nutr*. <https://doi.org/10.1017/S0007114508965788>
- Stoppa MA, Casemiro LA, Vinholis AHC, Cunha WR, Andrade ML, Silva E, Martins CHG, Furtado NAJC (2009) Comparative study of the recommended methodologies by Clsi and Eucast for activity evaluation antifungal. *Quim Nova* 32:498–502
- Strack D (1997) Phenolic metabolism. In: Dey P, Harborne J (eds) *Plant biochemistry*. Elsevier, pp 387–416
- Wei H, Zili L, Yuanlu C, Biao Y, Cheng L, Xiaoxia W, Yang L, Xing W (2011) Effect of icariin on bone formation during distraction osteogenesis in the rabbit mandible. *Int J Oral Maxillofac Surg* 40:413–418. <https://doi.org/10.1016/j.ijom.2010.10.015>
- Zhang ZB, Yang QT (2006) The testosterone mimetic properties of icariin. *Asian J Androl* 8:601–605. <https://doi.org/10.1111/j.1745-7262.2006.00197.x>
- Zida A, Bamba S, Yacouba A, Ouedraogo-Traore R, Guiguemré RT (2017) Substances naturelles actives sur *Candida albicans*, sources de nouveaux médicaments antifongiques: revue de la littérature. *J Mycol Med* 27:1–19