SVII-P13





SALT TOLERANCE EFFECTS OF SOLANUM CHEESMANIAE HKT1;1 AND/OR HKT1;2 FUNCTION LOSS IN A RECIPROCAL GRAFTING TOMATO EXPERIMENT*

María Remedios Romero-Aranda¹, Jesús Espinosa², Paloma González-Fernández², Emilio Jaime-Fernández¹, José Ángel Traverso³, María José Asins⁴, <u>Andrés Belver</u>²

¹IMSH "La Mayora", UMA/CSIC, Málaga; ²Estación Experimental del Zaidín, CSIC, Granada; ³Faculty of Sciences, University of Granada;.⁴ IVIA, Moncada, Valencia Corresponding author: Andrés Belver (andres.belver@eez.csic.es)

HKT1-like Na⁺ transporters play an important role in Na⁺ and K⁺ homeostasis and salt tolerance by removing Na⁺ from the xylem controlling its accumulation in shoots (Asins et al. 2013). In previous studies, we used recombinant inbred lines (RILs) derived from Solanum lycopersicum cv. Cerasiform and S. cheesmaniae, a halotolerant wild species, combined with RNAi-induced loss-of-function in all HKT1;1 and HKT1;2 allelic variants. We showed that HKT1;2 lycopersicum and cheesmaniae alleles are functionally responsible for the major tomato QTL lkc7.1 related to Na⁺/K⁺ homeostasis in the aerial part of the plant under saline conditions (Asins et al. 2013; Jaime-Pérez et al. 2017). In the present study, we analyse the physiological impact of function loss in cheesmaniae alleles at the HKT1;1 and HKT1;2 loci in the roots and aerial parts of the tomato plant in order to determine the relative contributions of each locus in the different tissues to plant Na⁺/K⁺ homeostasis and subsequently to tomato salt tolerance. We generated different reciprocal rootstock/scion combinations with non-silenced, single RNAi-silenced lines for ScHKT1;1 and ScHKT1;2, as well as doubly silenced lines, at both loci from a near isogenic line (NIL14), which is homozygous for the S. cheesmaniae haplotype in a genomic region (31.1 Mb) containing both HKT1 loci (Jaime-Pérez et al. 2017). Grafted combinations were cultivated with a Hoagland nutrient solution supplemented with 0 and 70 mM of NaCl under natural greenhouse conditions during the fall-spring season following the usual cultural practices of commercial greenhouse tomato production (Romero-Aranda et al. 2020). Our results show that salt treatment reduced vegetative growth and altered the Na⁺/K⁺ ratio in leaves and flowers, negatively affecting fruit production, particularly in graft combinations containing ScHKT1;2-silenced and double-SIHKT1;2/ScHKT1;1 silenced lines, when used for both rootstock and scion. We concluded that the impact of the removal of Na⁺ from the xylem by ScHKT1;2 in the aerial part of the plant can be even greater than that on Na⁺ homeostasis at the root level in tomato under saline conditions. This suggests that ScHKT1;2 may also play a role in Na⁺ phloem loading and consequently in determining Na⁺ recirculation towards the root zone, as well as the salt content of sink tissues such as developing leaves, flowers and fruits.

Asins et al. (2013). Plant Cell and Environment 36: 1171-91. Jaime-Pérez et al. (2017) Plant Cell Environm 40, 658–671. Romero-Aranda et al. (2020) Plant Physiol Biochem 154, 341-352. *ERDF-cofinanced grants from AEI, AGL2017-82452-C2-1R (AB) and AGL2017-82452-C2-2R (MJA).





Book of Abstracts

26th & 27th November, 2020



ORGANIZING COMMITTEE



Dr. David Posé Padilla IHSM-UMA-CSIC Málaga

Prof. Miguel Ángel Botella Mesa IHSM-UMA-CSIC Málaga

Prof. Araceli Castillo Garriga IHSM-UMA-CSIC Málaga

Dra. Carmen Martín Pizarro IHSM-UMA-CSIC Málaga

Dra. Catharina Merchante Berg IHSM-UMA-CSIC Málaga **Dra. Sonia Osorio Algar** IHSM-UMA-CSIC Málaga

Prof. Eduardo Rodríguez Bejarano IHSM-UMA-CSIC Málaga

Dra. Noemí Ruíz López IHSM-UMA-CSIC Málaga

Dra. Victoria Sánchez Vera IHSM-UMA-CSIC Málaga



SCIENTIFIC COMMITTEE

Amparo Monfort – IRTA-CRAG José Luis Martínez Zapater – ICVV Javier Brumós – NCSU Jordi Moreno Romero – CRAG Crisanto Gutiérrez – CMBSO - CSIC -UAM Vicente Rubio – CNB Manuel Rodríguez-Concepción – IBMCP - UPV - CSIC Irene García Fernández – IBVF - US Javier Gallego-Bartolomé – IBMCP - UPV - CSIC José Luis Micol – UMH Mar Castellano – CBGP - UPM - INIA Emilio Gutiérrez Beltrán – IBVF - US Rosa Lozano Durán – PSC Antonio Di Pietro – UCO

