

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Publications from USDA-ARS / UNL Faculty

U.S. Department of Agriculture: Agricultural
Research Service, Lincoln, Nebraska

2018

Knowledge gaps that hamper prevention and control of Mycobacterium avium subspecies paratuberculosis infection

Herman W. Barkema

University of Calgary, barkema@ucalgary.ca

Karin Orsel

University of Calgary

S.S. Nielsen

University of Copenhagen

A.P. Koets

Utrecht University & Wageningen Bioveterinary Research

V.P.M.G. Rutten

Utrecht University & University of Pretoria

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/usdaarsfacpub>

 Part of the [Agriculture Commons](#)

Barkema, Herman W.; Orsel, Karin; Nielsen, S.S.; Koets, A.P.; Rutten, V.P.M.G.; Bannantine, J. P.; Keefe, G.P.; Kelton, D.F.; Wells, Scott J.; Whittington, Richard J.; Mackintosh, Colin G.; Manning, E.J.; Weber, M.F.; Heuer, C.; Forde, T.L.; Ritter, C.; Roche, S.; Corbett, C.S.; Wolf, R.; Griebel, P.J.; Kastelic, J.P.; and Buck, J. De, "Knowledge gaps that hamper prevention and control of Mycobacterium avium subspecies paratuberculosis infection" (2018). *Publications from USDA-ARS / UNL Faculty*. 2447. <https://digitalcommons.unl.edu/usdaarsfacpub/2447>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications from USDA-ARS / UNL Faculty by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Herman W. Barkema, Karin Orsel, S.S. Nielsen, A.P. Koets, V.P.M.G. Rutten, J. P. Bannantine, G.P. Keefe, D.F. Kelton, Scott J. Wells, Richard J. Whittington, Colin G. Mackintosh, E.J. Manning, M.F. Weber, C. Heuer, T.L. Forde, C. Ritter, S. Roche, C.S. Corbett, R. Wolf, P.J. Griebel, J.P. Kastelic, and J. De Buck

Knowledge gaps that hamper prevention and control of *Mycobacterium avium* subspecies *paratuberculosis* infection

H. W. Barkema¹  | K. Orsel¹  | S. S. Nielsen²  | A. P. Koets^{3,4}  |
 V. P. M. G. Rutten^{3,17}  | J. P. Bannantine⁵  | G. P. Keefe⁶  | D. F. Kelton⁷ |
 S. J. Wells⁸ | R. J. Whittington⁹ | C. G. Mackintosh¹⁰ | E. J. Manning¹¹ |
 M. F. Weber¹² | C. Heuer¹³  | T. L. Forde¹⁴  | C. Ritter¹ | S. Roche⁷ |
 C. S. Corbett¹ | R. Wolf¹⁵ | P. J. Griebel¹⁶  | J. P. Kastelic¹ | J. De Buck¹ 

¹Department of Production Animal Health, Faculty of Veterinary Medicine, University of Calgary, Calgary, AB, Canada

²University of Copenhagen, Copenhagen, Denmark

³Utrecht University, Utrecht, The Netherlands

⁴Wageningen Bioveterinary Research, Wageningen, The Netherlands

⁵National Animal Disease Center, USDA-ARS, Ames, IA, USA

⁶University of Prince Edward Island, Charlottetown, Canada

⁷University of Guelph, Guelph, Canada

⁸University of Minnesota, Minneapolis, MN, USA

⁹University of Sydney, Camperdown, NSW, Australia

¹⁰AgResearch, Hamilton, New Zealand

¹¹University of Wisconsin, Madison, WI, USA

¹²GD Animal Health, Deventer, The Netherlands

¹³Massey University, Palmerston North, New Zealand

¹⁴University of Glasgow, Glasgow, UK

¹⁵Amt der Steiermärkischen Landesregierung, Graz, Austria

¹⁶VIDO-Intervac, Saskatoon, SK, Canada

¹⁷Department of Veterinary Tropical Diseases, Faculty of Veterinary Science, University of Pretoria, Private Bag X04, Onderstepoort, 0110, South Africa

Correspondence

H. W. Barkema, Department of Production Animal Health, Faculty of Veterinary Medicine, University of Calgary, Calgary, AB, Canada.

Email: barkema@ucalgary.ca

Summary

In the last decades, many regional and country-wide control programmes for Johne's disease (JD) were developed due to associated economic losses, or because of a possible association with Crohn's disease. These control programmes were often not successful, partly because management protocols were not followed, including the introduction of infected replacement cattle, because tests to identify infected animals were unreliable, and uptake by farmers was not high enough because of a perceived low return on investment. In the absence of a cure or effective commercial vaccines, control of JD is currently primarily based on herd management strategies to avoid infection of cattle and restrict within-farm and farm-to-farm transmission. Although JD control programmes have been implemented in most developed countries, lessons learned from JD prevention and control programmes are underreported. Also, JD control programmes are typically evaluated in a limited number of herds and the duration of the study is less than 5 year, making it difficult to adequately assess the efficacy of control programmes. In this manuscript, we identify the most important gaps in knowledge hampering JD prevention and control programmes, including vaccination and diagnostics. Secondly, we discuss directions that research should take to address those knowledge gaps.

KEYWORDS

control, Johne's disease, *Mycobacterium avium* subspecies *paratuberculosis*, prevention

1 | INTRODUCTION

Johne's disease (JD) is an infectious chronic inflammatory disorder of the intestine in ruminants caused by *Mycobacterium avium* subspecies *paratuberculosis* (MAP). It is a major health problem, resulting in intermittent diarrhoea, loss of body condition and lower productivity (e.g., Tiwari, VanLeeuwen, McKenna, Keefe, & Barkema, 2006). In the terminal phase, which most animals will not reach due to premature culling, animals die in very poor body condition. Infected ruminants shed MAP in manure and milk in increasing quantities as the disease progresses (Whitlock & Buergelt, 1996). The disease is widespread in domestic and wild ruminant populations in almost all countries in the world and causes great economic losses, not only because of lower productivity, but also as a result of loss of future income due to premature culling (Garcia & Shaloo, 2015; McKenna, Keefe, Tiwari, VanLeeuwen, & Barkema, 2006). The herd-level prevalence of MAP infection is likely >50% in most countries with a substantial dairy industry (Barkema, Hesselink, McKenna, Benedictus, & Groenendaal, 2010). In the absence of control measures, JD typically spreads, as farms often purchase cattle, frequently from herds with unknown JD status. In a recent review, Garcia and Shaloo (2015) reported substantial losses due to MAP infection, which escalate as the within-herd MAP prevalence and incidence of clinical JD cases increase. In Canada, the economic damage caused by JD was estimated at \$50 CAN per cow per year in MAP-infected herds, resulting in an average loss per infected farm of nearly \$3,000 CAN annually (Chi, VanLeeuwen, Weersink, & Keefe, 2002; Tiwari, VanLeeuwen, Dohoo, Keefe, & Weersink, 2008). Raizman, Fetrow, and Wells (2009) estimated the income over feed cost losses at \$366 per MAP-shedding cow per lactation, whereas Bhattarai et al. (2014) estimated a loss of \$1,644 US per 100 cows in a herd with a true prevalence of 7%. The cost of the disease to the US cattle industry was estimated at \$250M US per year (Ott, Wells, & Wagner, 1999).

Meta-analyses have demonstrated that the association of MAP with Crohn's disease in humans is specific and cannot be denied (Abubakar, Myhill, Aliyu, & Hunter, 2008; Feller et al., 2007), although a causal role has not yet been demonstrated (Waddell, Rajic, Stärk, & McEwen, 2015, 2016). Furthermore, transmission from cattle to humans has never been proven. However, addressing JD worldwide should be considered a proactive step in ensuring consumer confidence if a link was to be established between JD and Crohn's disease.

The epidemiology of JD is very different among cattle, sheep and goats. These species have very distinct differences when it comes to the course of MAP infection; therefore, inferences from one species cannot be naively applied to other species. Production type (e.g., dairy versus beef) also has a role in the course of MAP infection. This is not only evident in cattle (dairy versus beef), but probably even more evident in sheep and goats (i.e., wool/meat producing sheep in Australia and milk producing sheep/goats in the Mediterranean).

In this manuscript, we identify the most pressing gaps in knowledge hampering JD prevention and control programmes (summarized in Table 1). Secondly, we discuss directions that research should take to solve these knowledge gaps.

2 | CONTROL PROGRAMMES

Due to economic losses, and its possible association with Crohn's disease, many control programmes for JD have been developed worldwide. Many of these programmes focused on MAP-infected herds and were based on testing and culling test-positive cows, plus management adaptations (Benedictus, 1984; Collins, 1994; Kennedy, 2001; Rossiter & Burhans, 1996). The focus of some other programmes was to identify MAP-negative herds with the aim of keeping these herds, and in the case of Australia, Norway and Sweden an entire region negative, and also having them as a source of MAP-negative replacement animals (e.g., Frössling et al., 2013; Kalis, Collins, Barkema, & Hesselink, 2004; Kennedy, 2011; Whist et al., 2014). Tests included delayed type hypersensitivity (skin) tests, serological tests, direct detection of MAP by microscopy and using culture or polymerase chain reaction (PCR) to detect MAP in faecal samples. Presently, faecal culture is considered the most sensitive and specific ante-mortem test to identify MAP infection (Kalis, Barkema, Hesselink, Van Maanen, & Collins, 2002; Whitlock, Wells, Sweeney, & Van Tiem, 2000). However, as individual faecal culture is expensive and time-consuming, most JD control programmes use ELISAs to detect potentially infected animals (e.g., Lavers, Barkema, Dohoo, McKenna, & Keefe, 2014). Another reason for using ELISAs is that the extent of shedding correlates well with the ELISA titres (Dargatz et al., 2001), and hence, culling ELISA positives may be an effective and relatively low-cost option for removing high shedders. Currently, faecal culture is often replaced with direct PCR on faeces (e.g., Laurin, Chaffer, McClure, McKenna, & Keefe, 2015; Plain et al., 2014).

JD control programmes have been implemented in most developed countries, with objectives based on the national economic situation of the cattle, sheep and goat industry and the herd-level prevalence of MAP infection (reviewed by Kennedy, 2011; Geraghty, Graham, Mullaney, & More, 2014). In general, objectives include the following: (i) prove and protect freedom of disease at the country, regional or farm-level, for example, in Norway (Whist et al., 2014), Sweden (Frössling et al., 2013) and northern and western Australia (Kennedy, 2011); (ii) protect export of milk or genetics, for example, Canada (McKenna, Vanleeuwen, et al., 2006); (iii) decrease prevalence of MAP infection and limit farm-level economic losses, for example, Denmark (Nielsen, Jepsen, & Aagaard, 2007), the UK (Pritchard, Coffey, Bond, Hutchings, & Wall, 2017), Ireland (McAloon et al., 2016) and the USA (Wells, Hartmann, & Anderson, 2008); (iv) eliminate or reduce MAP load in bulk milk, for example, the Netherlands (Weber & van Schaik, 2007); and (v) eliminate MAP infection, for example, Norway in goats (Nagel-Alne et al., 2014).

TABLE 1 Most important knowledge gaps that hamper prevention and control of *Mycobacterium avium* subspecies *paratuberculosis* (MAP) infection

Area	Knowledge gap
Control programmes	<ul style="list-style-type: none"> – (Long-term) efficacy of control programmes – Is eradication of MAP infection in a herd possible?
Prevalence	<ul style="list-style-type: none"> – Comparison of MAP prevalence over time in the same region – Comparison of prevalence in different regions using the same test (regime)
The pathogen	<ul style="list-style-type: none"> – Distribution of MAP genotypes – Differences in virulence, pathogenicity, immunogenicity, persistence, transmission, survival outside the host and host specificity between genotypes – Effect of mixed genotype infections and superinfections
Tests	<ul style="list-style-type: none"> – Characteristics of tests in a population that reflects the target population – Development of reliable early-stage diagnostics – Reliable on-farm tests – Value of detection of immunoglobulin isotypes other than IgG – Most accurate and cost-effective set of environmental samples and sampling interval (dairy) – Best DNA extraction protocols for PCR analyses – How to quantify MAP in tissues, blood, milk and faeces – Validation of biomarkers in naturally infected cattle
Transmission	<ul style="list-style-type: none"> – Role of MAP-shedding young stock – Relative importance of various transmission routes (including transmission through dust and drinking water) – Influences on survival of MAP in soil and on pasture) – Minimal infectious dose for animals of all ages – Consequence and importance of intrauterine transmission – Shedding pattern of MAP-infected animals, including role of supershedders – Effectiveness of commercial pasteurizers at reducing viable MAP bacteria in colostrum
Role of wildlife	<ul style="list-style-type: none"> – Validation of diagnostic tests for wildlife – Prevalence of MAP in wildlife – Impact of MAP infections on wildlife health – Magnitude of the role wildlife plays as MAP reservoir and in transmitting MAP to farmed animals – Economic benefits of reducing MAP transmission from wildlife to livestock
Susceptibility	<ul style="list-style-type: none"> – Influences on susceptibility and the progression of MAP infection (e.g., age, genetics) – Genetic heritability of MAP infection – Difference in susceptibility among breeds – Identification of genetic markers associated with MAP susceptibility, disease progression and response to vaccination – Benefit of the IFN-γ assay in genetic linkage studies – Interaction between host genotype and MAP genotype
Vaccination	<ul style="list-style-type: none"> – Development of a cattle vaccine that limits infection and/or shedding, and does not interfere with diagnostics for other mycobacterial infections – Better understanding of immune evasion and immune modulation strategies of MAP – Characterization of diversity of MAP strains to select the most appropriate vaccine strain – Definition of protective immune response by vaccination – Appropriate level of attenuation of vaccine strains to elicit immune response without causing disease – Differentiation between infected and vaccinated animals
Uptake	<ul style="list-style-type: none"> – Influences on farmers' motivation to enrol in (voluntary) disease prevention and control programmes and implement recommended management strategies – How to overcome the "wait-and-see" mindset – Veterinary practitioners' role in promoting control programme participation including communication – Potential of approaches that address Johne's disease in combination with other faecal-orally transmitted diseases

Some factors that hampered the success of such control programmes were lack of compliance with management protocols, use of tests with a sufficiently high sensitivity to identify infected cattle, persistence of MAP in the environment, inadequate test frequency, appearance of unexpected new infections and purchase of replacement animals causing new introductions (Benedictus, 1984; Collins, 2001). A complementary modelling study showed that consistent application of preventive measures was key to success, although only

"test and cull" was not crucial to control JD (Groenendaal, Nielen, & Hesselink, 2003). However, "test and cull" can support and hasten elimination of infection from a herd that also has good management practices (Kudahl, Nielsen, & Ostergaard, 2011). Because there is no cure and, except for sheep, there are no vaccines that effectively prevent MAP infection, control of JD is currently based primarily on herd management strategies that reduce the risk of infection in young calves and restrict within-farm and farm-to-farm transmission.

Additionally, control programmes which include testing of individual animals, in general advise culling of MAP-positive animals.

Although JD control programmes have been implemented in most developed countries, the lessons learned from the actual experience with JD prevention and control programmes are underreported worldwide. JD control programmes are typically evaluated in a small number of herds (e.g., Collins, Eggleston, & Manning, 2010; Pillars, Grooms, Gardiner, & Kaneene, 2011), and many cattle control programmes have not included beef cattle. Additionally, particularly in cattle, results of a JD control programme can only be judged after at least 5 y (Caldow & Gunn, 2001; Nielsen & Toft, 2011), although most of the current well-designed programmes have been implemented that long (Table 1). Except for goats in Norway (Nagel-Alne et al., 2014), no report could be found in a herd in which MAP infection has successfully been eradicated (Table 1). There are many explanations for this failure which are discussed in the following sections.

Finally, some programmes were implemented but abandoned due to the outbreak of another disease (e.g., bovine spongiform encephalopathy or foot-and-mouth disease) or loss of funding (e.g., USA and Alberta, Canada), making it difficult to regain trust and reinstitute the programme.

3 | PREVALENCE

It is essential that reliable estimates of disease prevalence (animal or herd level) are available as this will determine how to proceed and monitor the results of a control programme over time. The goals may vary from eradication in areas of low prevalence, control in areas with high prevalence or increased surveillance in an area with no prior history of disease. Prevalence estimates obtained by surveys are affected by test accuracy; therefore, comparisons among studies must be adjusted to better estimate the true prevalence.

Infection with MAP and cases of JD have been reported from all continents with ruminant populations. Prevalence in most regions is currently unknown, and prevalence studies have low design uniformity, making comparisons among regions unreliable due to different sampling strategies and case definitions. Additionally, in most studies, the true herd-level prevalence and animal-level prevalence have not been estimated. Thus, the results of these studies and the reported prevalence estimates of MAP infection can currently not be compared directly (Table 1). Additionally, in most regions, herd-level prevalence and animal-level prevalence are underestimated, as the sensitivity of ELISAs is overestimated, particularly for cattle in 1st lactation heifers (McKenna, Keefe, Barkema, & Sockett, 2005; McKenna, Sockett et al., 2005; Nielsen & Toft, 2008). Almost invariably, in regions where herd-level prevalence estimates were obtained using an ELISA and independently validated using faecal culture or environmental culture, true prevalence estimates (adjusted for test characteristics) are much lower based on testing with ELISA versus culture (e.g., Lavers et al., 2014; Wolf et al., 2014). Although MAP test comparisons have

been the subject of many studies, there is a lack of studies comparing test characteristics in populations that reflect the target population when estimating the prevalence of MAP infection (Nielsen & Toft, 2008). The sensitivity of MAP tests increases in dairy cattle with increasing days in milk and age but decreases with increased milk yield (Eisenberg, Veldman, Rutten, & Koets, 2015; Kirkeby, Græsboell, Halasa, Toft, & Nielsen, 2015), whereas sensitivity increases with age in sheep and goats (Lybeck, Storset, Djønnø, Valheim, & Olsen, 2011). Therefore, sensitivity estimates need to be adjusted for age in all ruminants, and days in milk and milk yield in dairy cattle when comparing between or within herds and also over time.

Comparing prevalence over time, among regions and countries is often unreliable due to the use of a variety of diagnostic tests, often with unknown or unreliable test characteristics. To compare herd- and animal-level prevalence estimates of MAP among countries, and to allow for the development of international trade standards, we recommend a supranational standardized study, comparable to a *Neospora caninum* seroprevalence study involving cattle from four countries (Bartels et al., 2006) (Table 1).

4 | THE PATHOGEN

Strain typing is useful in helping to clarify epidemiological and virulence questions that cannot otherwise be resolved; however, limited genetic variation among MAP isolates within host species has slowed progress. Several typing techniques have been used (Amonsin et al., 2004; Coffin et al., 1992; Collins & de Lisle, 1986; Motiwala et al., 2003; Thibault et al., 2007; Whittington, Hope, Marshall, Taragel, & Marsh, 2000; Whittington, Marsh, Choy, & Cousins, 1998). Whole-genome sequencing (WGS), because of its higher resolution, is the logical evolution of technology to assist epidemiological investigations and control programmes. The cost of WGS still has to decrease though for this technique to become the technique of choice in this kind of studies.

Variable number tandem repeat (VNTR) and mycobacterial interspersed repetitive unit (MIRU) typing have been moderately successful for characterizing MAP isolates. Diversity among isolates has been reported based on VNTR typing (Fritsch, Luyven, Köhler, Lutz, & Möbius, 2012; Sohal et al., 2014; Stevenson et al., 2009; Thibault et al., 2008), with most isolates being one of two dominant VNTR types. MIRU-VNTR genotyping does not, however, accurately reflect phylogenetic relationships, and repeat sequences are subject to homoplasy (Ahlstrom, Barkema, & De Buck, 2014; Bryant et al., 2016). Current VNTR typing includes loci that are too unstable and unreliable to be used for a molecular epidemiological analysis of MAP (Ahlstrom et al., 2015). Short sequence repeat (SSR) typing (Amonsin et al., 2004) further differentiates MAP isolates of the same VNTR type (Stratilo, Lewis, Bryden, Mulvey, & Bader, 2006; Thibault et al., 2008) but lacks sufficient discrimination and stability for epidemiological studies (Kasnitz, Köhler, Weigoldt, Gerlach, & Möbius, 2013). Nevertheless, in specific cases, VNTR (with or

without SSR typing) can still be useful for supporting source tracing investigations and JD control measures (Oakey, Gavey, Singh, Platell, & Waltisbuhl, 2014).

Whole-genome sequencing (WGS) provides unparalleled detail regarding genetic profiles of MAP. Genomic epidemiology of MAP is new (Ahlstrom et al., 2015, 2016) but has already provided important insights into transmission dynamics. Single nucleotide polymorphisms (SNPs) identified through WGS are evolutionarily stable and can be reliably used to identify true evolutionary relationships (Pearson, Okinaka, Foster, & Keim, 2009). In a MAP-endemic environment, this level of detail is invaluable in understanding the molecular epidemiology and transmission dynamics. For example, there are at least eight genetically distinct MAP subtypes in Canadian dairy cattle, with >80% of isolates belonging to a single dominant subtype (Ahlstrom et al., 2016). However, this information cannot be extrapolated to other countries because dominant strains vary between countries and continents (Kiehnbaum, Amonsin, Wells, & Kapur, 2005; Machackova et al., 2004; Motiwala et al., 2004; Whipple, Kapke, & Vary, 1990). Some strains may be more successful in spreading or persisting in different environments or management systems. It is difficult to know whether high MAP prevalence within a specific herd relates to a high MAP infection opportunity (e.g., poor biosecurity) or is due to MAP strains with increased transmission potential on these farms.

Strain typing can be used to track transmission of MAP in a variety of settings and host species. Studies of within-herd spread of MAP infection (Pradhan et al., 2011), between-herd transmission (Oakey et al., 2014) and the role of wildlife in spreading MAP (Fritsch et al., 2012) require molecular tools that can differentiate MAP types. WGS was recently used to identify multiple mixed genotype infections (Davidson, Ahlstrom, De Buck, Whitney, & Tahlan, 2016); these infections could affect disease dynamics and impact pathogen evolution and genetics and thereby impact the efficacy of therapeutic treatments or vaccines (Table 1).

An understanding of the full genetic diversity of MAP is needed to accurately assess subtype-specific phenotypes and virulence characteristics, and to develop vaccines and effective management practices (Table 1). Investigations into strain-specific differences in MAP virulence and pathogenicity have been mostly limited, however, to major strain types (Borrmann, Möbius, Diller, & Köhler, 2011; Gollnick et al., 2007; Janagama, Jeong, Kapur, Coussens, & Sreevatsan, 2006; Motiwala et al., 2006), with significant differences between Types I and II isolates reported for growth rates and intracellular survival (Table 1).

Investigations have determined phenotypic differences among MAP subtypes for a variety of traits, including growth rates and invasion efficiencies, immunogenicity, virulence as measured by macrophage invasion efficiencies and kinomic responses (Griebel et al., 2014; Whittington et al., 2011). Future vaccine development and molecular epidemiological studies should also consider the relative frequencies of MAP subtypes with a focus on dominant subtypes. There are differences among MAP strains in the immune response that they stimulate, as well as differences in host tropism, disease

phenotypes and ability to evade control by vaccines (Colavecchia et al., 2016; Sohal, Singh, Singh, & Singh, 2010) (Table 1).

5 | TESTS

5.1 | Purpose of testing

Test results used in control programmes should primarily assist decision-makers achieve a specific objective. Furthermore, the specific test strategies employed are likely to differ between regions, depending on the logistics of implementation and other practicalities, as well as herd size (Dorshorst, Collins, & Lombard, 2006). Identification of infected animals before they spread the infection while still maximizing the lifetime production of an animal is the goal of a cost-effective test strategy. Purposes of testing primarily include the following: (i) estimate infection prevalence to determine the best course of action; (ii) minimize financial losses as a result of impaired milk production or growth and increased culling rate; (iii) estimate the effects of proposed control measures or evaluate additional measures or management changes; (iv) eliminate infectious cattle to reduce spread of infection, (v) reduce the risk of MAP contamination of food products for human consumption; and (vi) eradicate MAP from a herd (or region). Tests for these purposes can focus on individual animals, herds or a subset of the herd. Control strategies that focus on the individual animal- and target-specific management conditions contributing to transmission include the following: (i) identification of infectious animals actively shedding MAP; and (ii) identification of infected animals at risk of shedding MAP (Table 1).

Ideal tests used for control should identify the infectious animal or predict its infectivity. However, as we do not know the infectious dose of MAP, shedding patterns of infected animals and factors contributing to disease progression, it is difficult to interpret test results (Table 1). Consequently, we struggle to identify MAP infection cases and non-cases, which is essential when evaluating the accuracy of diagnostic tests. This has resulted in decades of reported test evaluations that are of questionable quality (Nielsen & Toft, 2008).

Existing tests are performing reasonably well in detecting advanced stages of JD, although the specificity of test results may be challenged and the number of bacteria excreted in faeces or milk is rarely quantified. Detection of early-stage infection is only of interest for a control programme if these animals then become infectious (Marcé, Ezanno, Seegers, Pfeiffer, & Fourichon, 2011). Otherwise, these infections may disappear from a specific population. However, because the sensitivity of diagnostic tests in calves is generally low, young stock are rarely tested for MAP. As a result, no longitudinal studies have been published that followed infected calves or lambs and confirm how many subsequently test positive (Table 1).

5.2 | Early-stage diagnostics

Early-stage diagnostics primarily target pro-inflammatory immune responses. To be useful, however, they should differentiate infected

from recovered animals (Dennis, Reddacliff, & Whittington, 2011) and indicate if or when an animal will become infectious (de Silva et al., 2013). Whereas detection of faecal shedding by direct extraction followed by a MAP-specific PCR can detect if an animal is infectious (Fock-Chow-Tho, Topp, Ibeagha-Awemu, & Bissonnette, 2017), it cannot indicate whether or when it will become infectious. Such a test would likely be useful to achieve eradication in low prevalent populations (Table 1).

Identification and multiplexing of antigens that elicit an early humoral immune response could yield a serological test that overcomes the current delay in antibody-based detection of MAP infection. Analyses of longitudinal serum samples (from experimentally infected calves) were used to detect developing humoral immune responses against MAP proteins (Bannantine et al., 2008), with anti-MAP antibodies detected as early as 70 day after infection. Novel MAP antigens for detection of antibody responses in subclinical cattle were also reported (Facciuolo, Kelton, & Mutharia, 2013).

The interferon-gamma (IFN- γ) assay detects cell-mediated immune responses to early-stage MAP infection (Stabel, 2000), but needs improvements and standardization as differences in composition of the purified protein derivative (PPD) antigens can influence specificity (Capsel et al., 2016; Kalis, Collins, Hesselink, & Barkema, 2003). Single proteins to increase specificity of T-cell responses are being sought (Carlos et al., 2015; Huygen, 2014; Leite, Reinhardt, Bannantine, & Stabel, 2015; Rana, Rub, & Akhter, 2015). Blood samples must be tested within 8 hr for optimal results to ensure T-cell viability, although IL-12 and anti-IL-10 supplementation (Mikkelsen, Aagaard, Nielsen, & Jungersen, 2012) or IL-7 and IL-12 supplementation (Plain, Begg, de Silva, Purdie, & Whittington, 2012) can increase the interval to 48-hr interval before testing. Ultimately, an on-farm test would be most effective.

Leucocyte markers associated with MAP infection may also be used as a diagnostic test. For example, increased expression of CD25 and CD45RO on T cells may be an early indication of infection (Stabel & Robbe-Austerman, 2011), but these markers lack specificity. Furthermore, phenotype analysis of leucocytes is again dependent upon the isolation and analysis of viable cells. The lack of useable and rapid diagnostic tests for specific biomarkers is clearly a bottleneck in identifying animals with MAP infection.

To accommodate the discovery of biomarkers that can predict the onset of the infectious stage, experimental infection models presenting a natural disease outcome (e.g., Begg et al., 2010; Mortier et al., 2013) are essential for test development (Begg & Whittington, 2008; Hines et al., 2007). Ultimately, longitudinal sampling of herds with different prevalence profiles is necessary to validate the outcomes of experimental infections in an unbiased and representative set of samples from infected but not yet infectious animals.

5.3 | Late-stage diagnostics

Late-stage diagnostic tests detect anti-inflammatory immune responses, for example, ELISAs detecting IgG1. They are relatively well characterized and easy to automate, but sensitivity is in general

still low compared to viral infections, such as bovine leukosis (Nielsen & Toft, 2008). Ideally, these should be reliable point-of-care tests using blood or milk from individual cows (Lavers et al., 2014) (Table 1). Better characterization of disease progression in relation to the induction of IgG1 antibody responses is desired.

There might be value in the detection of immunoglobulin isotypes other than IgG (e.g., IgM and IgA) (Table 1). Although IgM is the first isotype to appear in response to an infection, this response has limited applicability as it is transient (Abbas & Riemann, 1988). Isotype switching results in local mucosal production of IgA and anti-MAP IgA responses have been detected in faecal samples from infected sheep (Begg et al., 2015). The challenge remains the identification of individual MAP proteins that detect antibodies specific to MAP and not environmental *Mycobacterium* spp.

Calves, kids and lambs were reported to be ELISA positive for MAP 3–4 month after infection (Kurade, Tripathi, & Rajukumar, 2004; Mortier, Barkema, Orsel, et al., 2015; Storset et al., 2001). However, in very young animals, such ELISA antibodies may be an indication of passive immunity (transfer of colostrum-derived maternal antibodies), as maternal antibodies reacting with MAP were detected in serum collected from calves up to 121 d of age (Schillinger et al., 2013).

The titre of an ELISA is predictive for the probability of MAP shedding (Weber & van Schaik, 2007). This association renders the utility of an ELISA as highly applicable to JD control in dairy herds when the aim is to reduce transmission by selective removal of shedders. In that instance, an ELISA may replace faecal culture or PCR, but it is then important to shorten the testing interval (Lu et al., 2008).

5.4 | Herd-level diagnostics

Environmental sampling is a quick sampling method to determine the MAP infection status of a herd (Wolf et al., 2017). However, this sampling method is only sufficiently sensitive in relatively intensive livestock operations such as housed dairy cattle. In extensive farming, such as sheep and cow-calf herds in many countries, environmental sampling currently does not have sufficient sensitivity (Whittington et al., 2003). For example, culture followed by PCR on six environmental samples detected 70% of MAP-infected dairy herds (Wolf, Barkema, De Buck, & Orsel, 2015). Samples collected from alleyways of lactating cow pens and manure lagoons were most sensitive (Wolf, Barkema, et al., 2015). On farms with only manure piles (or if an outdoor lagoon is inaccessible due to weather), additional samples should be collected from indoor manure pits or alleyways. In a recent German study in cow-calf herds, environmental samples collected in high cow traffic areas at the end of the winter had a sensitivity of 64% (Klawonn, Einax, Pützschel, Schmidt, & Donat, 2016), which is similar to housed dairy herds (Wolf et al., 2017). There is certainly room for improvement of environmental sampling; modifications in the number of collected samples per set and in sampling intervals may result in more accurate diagnostic protocols (Table 1). Further refinements of environmental sampling,

including determining impact of season, are needed (Wolf, Barkema, et al., 2015). Culture of pooled faecal samples, most often consisting of pools of 5 or 10 for cattle, or 10 to 50 for sheep faecal samples, has been extensively evaluated and proven to be relatively sensitive (Sergeant, Whittington, & More, 2002; Van Schaik, Pradenas, Mella, & Kruze, 2007; Verdugo, Jones, et al., 2014; Whittington, Fell, et al., 2000).

Processing pooled environmental samples with direct PCR instead of culture reduces processing time and costs (Bölske & Herthnek, 2010). An additional advantage of PCR compared to culture is that PCR is not dependent on viable MAP bacteria in the sample. That might result in a higher sensitivity with PCR, especially if samples are collected in the winter, or at locations where manure accumulates over extended times. Test characteristics for PCR (typically using IS900 or F57 as the target) relative to culture have been reported, with overall good results (e.g., Clark, Koziczowski, Radcliff, Carlson, & Ellingson, 2008; Cook & Britt, 2007; Fock-Chow-Tho et al., 2017) for both pooled and environmental samples (e.g., Kawaji, Taylor, Mori, & Whittington, 2007; Mita et al., 2016). Results are, however, highly depend on the extraction method and the primer(s) used (Mita et al., 2016) (Table 1).

Testing of bulk tank milk or a pool (whole herd set of Dairy Herd Improvement milk samples) using a commercial ELISA is an inexpensive method of herd screening for JD in dairy cattle (Kelton et al., 2014). To determine a herd's MAP infection status, bulk milk samples can either be analysed for the presence of MAP-specific antibodies using ELISA, or for the presence of MAP bacteria using direct PCR. Although repeated bulk tank test results are consistent over short intervals and correlated with herd prevalence (Nielsen & Toft, 2014), results can be influenced by herd size and within-herd prevalence of ELISA-positive individuals. This testing identifies herds with moderate (>5% within-herd milk ELISA positive) to high (>8% within-herd milk ELISA positive) prevalence (Serraino et al., 2014). This test is not sensitive enough to detect low prevalence herds (Jayarao et al., 2004; Khol et al., 2013; Van Weering et al., 2007) or to monitor subtle changes in within-herd prevalence over time.

As is the case with all JD testing strategies, given the low sensitivity of the tests, the proportion of infected herds is underestimated and negative test results can give dairy producers a false sense of security based on the belief that a negative test means their herd is uninfected. However, with increasing frequency statistical tools such as Bayesian statistics are used to correct for less than ideal test characteristics that are predominantly the result of the biology of MAP infection (e.g., McAloon et al., 2016; Verdugo, Pleydell, et al., 2014; Verdugo, Jones, et al., 2014).

5.5 | Recent developments and future directions

Novel findings and approaches in diagnostic test development and evaluations are rare, and so far no major breakthroughs have been provided. Mostly, incremental improvements to existing tests and test strategies have been identified, for example, repeated testing in a Danish JD control programme (Nielsen, 2008). Once prevalence is

reduced, cost-effective monitoring of the MAP status of a population can be difficult. Development of cost-effective strategies for surveillance and certification is still required while retaining a focus on the fit-for-purpose criterion (OIE, 2012).

Quantitative PCR (qPCR) for assessing the number of MAP bacteria in faeces has great promise in comparison with bacterial culture (Christopher-Hennings et al., 2003; Khare et al., 2004; Kralik, Beran, & Pavlik, 2012; Laurin et al., 2015) (Table 1). This technique has been validated on spiked faecal samples and proficiency panels, including faecal samples from naturally infected animals (e.g., Plain et al., 2014). This approach is strongly dependent on efficient DNA extraction, which has been studied (summarized in Bölske & Herthnek, 2010). However, quantification by qPCR is still only a relative method because a standard curve of spiked samples needs to be used to determine the approximate copy number of MAP organisms in a test sample. Digital (third-generation) PCR is expected to be more quantitative, more precise and predict absolute numbers of shed bacteria in faeces and therefore be useful to validate other tests and biomarkers as has been studied for *Mycobacterium tuberculosis* (Devonshire et al., 2016). microRNAs circulating in blood may provide another a promising method to detect MAP infection (Shaughnessy, Farrell, Riepema, Bakker, & Gordon, 2015), but this has not been confirmed.

The effect of MAP genotype on disease progression, shedding, immune responses or generation of other biomarkers is not well characterized (Table 1). A longitudinal follow-up of animals by ELISA, IFN- γ testing, faecal culture and eventually tissue culture is needed.

Using proteomic and transcriptomic analyses, several putative biomarkers for early infection with MAP have been proposed (Casey et al., 2011; David, Barkema, Luo Guan, & De Buck, 2014; David, Barkema, Mortier, et al., 2014; Seth et al., 2009; Shin et al., 2015; Skovgaard et al., 2006; Thirunavukkarasu et al., 2014; You et al., 2012; Zhong, Taylor, Begg, & Whittington, 2011). Metabolomic profiling detected MAP infection earlier than other diagnostic methods, with individual metabolites distinguishing infected from non-infected cattle (De Buck, Shaykhtudinov, Barkema, & Vogel, 2014). Furthermore, changes in faecal microbiota of MAP-infected cattle may have promise for identifying infected animals during subclinical stages of JD (Derakhshani et al., 2016). Analysis of volatile organic compounds emitted during culture may assist in identifying growth and also strain identification (Trefz et al., 2013). These and other emerging diagnostic technologies and approaches have recently been reviewed by Britton, Cassidy, O'Donovan, Gordon, and Markey (2016). Unfortunately, none of these biomarkers have been properly validated on naturally infected cattle of varying ages or in various stages of infection to reveal their true sensitivity and specificity (Britton et al., 2016) (Table 1).

6 | ROUTES OF TRANSMISSION

MAP can be transmitted by: (i) ingestion of faecal material, (ii) drinking contaminated colostrum or milk, (iii) intrauterine infection and,

potentially, (iv) aerogenic transmission. The best method to prevent transmission is expected to be limiting exposure of calves to adult faeces by systematic separation of adult cows and calves, in combination with good hygiene practices (Marcé et al., 2011).

There are limited data regarding the quantity of bacteria shed by animals in the different infection stages (Marcé et al., 2010; Whittington, Reddacliff, Marsh, McAllister, & Saunders, 2000), with substantial inter- and intra-animal variability (Crossley, Zagmutt-Vergara, Fyock, Whitlock, & Gardner, 2005; Whitlock et al., 2000). However, recent advances in PCR technologies (digital droplet PCR; Pinheiro et al., 2012) should enable a determination of the exact number of organisms associated with the different modes of transmission (faeces, milk, colostrum) for all different stages of infection (susceptible/uninfected, transiently infectious, latently infected, infectious, resistant stages) (Table 1). This knowledge would be especially meaningful if in parallel the corresponding infectious dose and the corresponding infection risk associated with the various transmission routes were determined for animals of all ages. However, it is important to note that single inoculations may be a poor model of natural infection as the frequency of exposure might be an important factor in transmission. Typically, several or consecutive-day inoculations are used (Begg et al., 2010; Hines et al., 2007). However, a trickle dose (limited numbers of organisms) was highly effective in causing intestinal infection (Eisenberg et al., 2011), suggesting that more complex dynamics play a role during natural infections and need to be taken into account for a better understanding of transmission and infection.

6.1 | Colostrum

Milk and colostrum can be contaminated with MAP, either through faecal contamination of teats or shedding from the udder (Stabel & Goff, 2004; Streeter, Hoffsis, Bech-Nielsen, Shulaw, & Rings, 1995; Sweeney, Whitlock, & Rosenberger, 1992). In a Danish study, calves that received colostrum from multiple cows had 1.24 times the odds of testing MAP ELISA positive as adults compared to cows that only received colostrum from their dam (Nielsen, Bjerre, & Toft, 2008). In contrast, the risk of infection from ingestion of colostrum was recently challenged in a cohort study concluding that MAP-contaminated colostrum did not increase the risk of MAP shedding in calves up to two years of age (Eisenberg, Rutten, & Koets, 2015). This reveals a knowledge gap regarding transmission and certainly the relative contribution of some transmission routes (Table 1).

A reduction in within-herd prevalence of MAP infection may be achieved by pasteurizing colostrum to reduce MAP transmission. Heat treatment reduces the number of viable MAP bacteria in milk (Eltholth, Marsh, Van Winden, & Guitian, 2009; Lund, Gould, & Rampling, 2002). However, most studies used huge numbers of bacteria, spiked milk (not naturally infected), milk instead of colostrum and a hot water bath, not a commercial pasteurizer. Therefore, the current knowledge is insufficient to answer the question whether on-farm pasteurization using commercial pasteurizers can effectively reduce the number of viable MAP bacteria in colostrum (Table 1).

6.2 | Calf-to-calf

Cattle farmers typically try to interrupt faecal-oral transmission by implementing best-hygiene management practices. In many studies, there were associations between specific management practices and the likelihood of animals being infected with MAP (e.g., Nielsen & Toft, 2011); however, many questions remain unanswered. In particular, the potential risk of calf-to-calf transmission is largely overlooked in JD prevention and control programmes; only one of eight MAP modelling studies included calf-to-calf transmission (reviewed by Marcé et al., 2010) (Table 1). Although many researchers associate a low risk with this potential transmission pathway, based on the assumption that calves will not shed MAP in their faeces (Groenendaal et al., 2002; Marcé et al., 2011), there is strong evidence of calves being infected by other calves (Benedictus et al., 2008; Van Roermund, Bakker, Willemsen, & de Jong, 2007; Wells & Wagner, 2000). One reason for these apparently discordant results is the difficulty in establishing an exposure-disease relationship, due to the delayed clinical onset of JD (Collins, 1996). Benedictus et al. (2008) reported that calf-to-calf transmission occurs, and that contact with infectious animals increases the likelihood of calves being infected with MAP, which supports previous data (Van Roermund et al., 2007). Unfortunately, this study (Benedictus et al., 2008) was carried out on a single farm and outcomes were measured long after exposure. Consistent with some studies (Bolton, Grooms, & Kaneene, 2005; Santema et al., 2012; Van Roermund et al., 2007; Weber, Kogut, Bree, & Van Schaik, 2006), in a recent challenge experiment on age- and dose-dependent susceptibility (Mortier et al., 2013), 47% of calves with a proven MAP infection shed the bacterium at least once from 2 to 6 mo after infection (Mortier, Barkema, Orsel, Wolf, & De Buck, 2014). In a recent study, co-mingling MAP-infected and non-infected calves for 3 mo resulted in infection of 50% of the naïve calves (Corbett, De Buck, Orsel, & Barkema, 2017). It was estimated that under the circumstances of the study, one MAP-infected calf on average infected three other calves. Additionally, on 17 MAP-infected dairy farms in Alberta, Canada, 3% of calves were shedding the bacterium, and on nine of these farms, MAP-positive environmental samples were collected from young stock pens (Wolf, Orsel, De Buck, & Barkema, 2015). Due to acidified milk feeding and automatic milk feeders (Barkema et al., 2015), many dairy calves are group-housed both before and after weaning. Consequently, there is a need for longitudinal studies quantifying the risk of MAP transmission among calves by measuring MAP shedding, environmental contamination and tissue levels in naturally infected calves (Table 1). If calf-to-calf transmission is deemed important, JD prevention and control programmes will require modifications.

6.3 | Intrauterine

Some calves from MAP-infected cows are infected at birth (reviewed by Whittington & Windsor, 2009). Intrauterine infection is more likely in cows with clinical JD (Whittington & Windsor, 2009) and may be lower in herds with a low MAP prevalence (Adaska &

Whitlock, 2012). Most studies that determined the proportion of intrauterine infection with MAP recovered foetuses of MAP-infected cows at various stages of pregnancy (Whittington & Windsor, 2009). As many of these foetuses were recovered prior to term, the proportion of calves born with a MAP infection is undoubtedly underestimated. Additionally, it is not known if or when intrauterine infected calves will start shedding, how infection progresses, and the nature of the immune response, compared to calves infected orally soon after birth (Table 1). The rate of 15% intrauterine infections may have been overestimated as a new analysis suggested it may only be 4% (Mitchell et al., 2015).

Intrauterine transmission from dam to foetus appears to be more common in red deer than in cattle and sheep, with 90% of clinically affected hinds having an infected foetus (Van Kooten, Mackintosh, & Koets, 2006). In another study, MAP was isolated from 78% of foetuses from 18 subclinically infected seropositive red deer hinds (Thompson, Clark, & Mackintosh, 2007). By contrast, MAP was isolated from only 39% of foetuses from clinically affected dairy cows and 9% of foetuses from subclinically infected cows (Whittington & Windsor, 2009). In sheep, intrauterine transmission is thought to occur in <10% of infected ewes. Intrauterine transmission of MAP has also been detected in free-ranging red deer and chamois (Deutz, Spergser, Wagner, Rosengarten, & Köfer, 2005). As in cattle, MAP-infected colostrum and milk may cause pseudo-vertical transmission in deer and other wildlife.

6.4 | Environment

If a cow is shedding MAP in the faeces, her manure is infectious and can remain so for at least 1 y (Whittington, Marshall, Nicholls, Marsh, & Reddacliff, 2004). The proportion of environmental manure samples that are culture-positive increases with an increasing prevalence of MAP-infected cows (Wolf, Barkema, et al., 2015). In sheep, shedding is dose- and age-at-infection dependent (McGregor, Dhand, Dhungyel, & Whittington, 2012). However, sheep of all ages and exposed at all doses are equally likely to be colonized by MAP, although the severity of histopathological lesions was strongly determined by age at exposure (McGregor et al., 2012). These findings stress the role of transmission at pasture, especially for young animals.

There is a need for more research to investigate the role of environmental transmission taking into account the survival characteristics of MAP and the contact structure between animals in a herd (Table 1). Survival of MAP in the environment has been suggested to depend on many factors, including soil pH, faecal content, concentrations of macro- and micronutrients (e.g., Fe, Mo and Cu), temperature and exposure to sunlight (reviewed by Elliott, Hough, Avery, Maltin, & Campbell, 2015). No viable (culturable) MAP was detected after 2 month of anaerobic digestion at a farm-scale biogas plant (Slana, Pribylova, Kralova, & Pavlik, 2011). MAP was present in settled dust samples on dairy farms under both experimental and field conditions (Eisenberg, Koets, et al., 2010; Eisenberg, Nielen, et al., 2010). Although current JD prevention programmes do not

consider dust, it could be a fomite and facilitate transmission (Cerner, Pfeiffer, & Abbott, 2004) (Table 1).

The results of many of the studies on survival in the environment are very difficult to generalize to different environmental circumstances (Elliott et al., 2015) and MAP strain types (Whittington et al., 2004) (Table 1). Secondly, most studies were carried out as an experimental model and not as a field study. Also, because it is not clear what dose of viable MAP will lead to infection in animals of different age groups, it is not clear whether ingestion of the surviving concentration of MAP bacteria would actually lead to a MAP infection. Additionally, it is not clear whether these factors only influence survival and/or virulence of MAP bacteria, or whether there might also be an influence on the host, through (in)direct effects of these factors on the immune system (Lugton, 2004). Finally, management practices relevant to the transmission of MAP may be associated with environmental factors such as soil type, exposure to sunlight and humidity (Table 1).

6.5 | Within-herd transmission

Understanding the routes of MAP transmission between cattle is very important for effective control of the disease. The best-established transmission route of MAP is oral uptake of bacteria by susceptible young stock, via colostrum, milk, water or food contaminated with faeces from MAP-shedding animals (Benedictus et al., 2008; Sweeney, 1996). In addition, intrauterine transmission has been described.

Although isolation of adult cattle from young calves has an important role in prevention of JD (Groenendaal et al., 2002), after 20 year of management and hygiene measures to prevent MAP transmission in a dairy herd, complete eradication was not achieved, implicating other non-identified and therefore non-controlled transmission routes (Table 1). In sheep, experimental intratracheal introduction of MAP caused infection (Kluge et al., 1968), whereas intestinal infection occurred after exposure to MAP-containing aerosols given intratracheally and intranasally (Eisenberg et al., 2011). As MAP has been detected in bioaerosols on dairy farms, transmission of MAP by bioaerosols should be further studied including estimations of the relative contributions of the various transmission routes identified. Additionally, there is evidence that infection of non-lactating heifers or adult cows can occur (e.g., Fecteau, Whitlock, Buergelt, & Sweeney, 2010), likely when infection pressure is high.

All existing recommendations (McKenna, Vanleeuwen, et al., 2006) for decreasing the risk of new infections of MAP in a dairy operation are meant to reduce infection rates in calves by decreasing contact with adult cows. Regardless, a better understanding of transmission and increased testing should improve disease control (Table 1).

6.6 | Between-herd transmission

Introduction of infected animals is the most important route of transmission of MAP between herds (Rangel et al., 2005). Frequent

cattle purchases from other herds without knowledge of their disease status increased the risk for MAP culture-positive environmental samples (Wolf, Barkema, De Buck, & Orsel, 2016). Although testing animals pre-purchase will prevent the introduction of positive animals, the long incubation period will result in many false-negative cattle. Thus, herd-level testing of the herd of origin in a certification-and-surveillance programme (Weber, Groenendaal, van Roermund, & Nielen, 2004; Weber, van Roermund, Vernooij, Kalis, & Stegeman, 2006) is likely to be more effective in reducing the risk associated with the trade of cattle between herds. Although MAP can survive for a long time in water (Elliott et al., 2015), the role of transmission through surface and drinking water is not known, and there are many unknowns about the role of transmission from other ruminants and wildlife to domestic ruminants (Table 1).

6.7 | Between-species transmission

Cross-grazing sheep and deer reduces the risk of clinical paratuberculosis in deer because the sheep strain of MAP is less pathogenic for deer than the cattle strain (Heuer et al., 2012; Verdugo, Pleydell, et al., 2014). Modelling exercises have shown that the similarity of strain types isolated from beef cattle and deer was 3-fold greater when direct contact between these species was considered compared to a scenario that ignored the contact structure. Transmission would be expected to go in both directions between co-grazed animals, thus grazing infected deer with sheep or cattle puts both species at risk. In the UK, the presence of farmed deer increased the risk of reporting clinical JD in dairy cattle kept on the same farm (ORs ranged from 15 to 209; Cetinkaya, Erdogan, & Morgan, 1997).

In many areas, there is co-grazing of wildlife and domestic livestock, which may allow cross-infection to occur under specific conditions (high animal density, neonates in population, etc.). In an alpine region in Italy, MAP-infected ibex are sympatric with seropositive cattle (Ferroglia, Nebbia, Robino, Rossi, & Rosati, 2000). In Spain, cattle, sheep and goats share pastures and waterholes with herds of fallow deer in which paratuberculosis has been diagnosed (Balseiro, García Marín, Solano, Garrido, & Prieto, 2008). In Germany, it appears that feral and farmed animals are reservoirs for specific MAP genotypes (Fritsch et al., 2012). Although other studies failed to demonstrate significant transmission between wildlife and livestock, false negatives can be common. It is important to keep in mind that many wildlife studies have limited observations and with that, limited power (Table 1). Also, there is no uniformity in MAP diagnostics which makes it challenging to compare studies from around the world (Table 1).

7 | ROLE OF WILDLIFE

A better understanding of the epidemiology of MAP in wildlife is essential for implementing effective disease prevention and infection control programmes for livestock. MAP has been recovered from a wide variety of wildlife species worldwide (with or without signs of

JD). Mostly, it has been found in wild ruminants (reviewed by Mackintosh & Griffin, 2010); however, other species such as lagomorphs may have an important role in MAP transmission (reviewed by Hutchings et al., 2010). Regardless, much remains to be learned about the distribution of MAP among free-ranging wildlife populations, impacts of infection on the health of these populations, potential for these populations to act as reservoirs for MAP and the extent of MAP transmission between wildlife and livestock in various environments (Table 1).

For most wildlife surveillance efforts, resources are scant. Current methods that reduce the cost of diagnostic testing, such as pooled faecal culture, have only been validated for cattle (Van Schaik et al., 2007). Commercially available ELISA tests require validation prior to their use for MAP surveillance in specific wildlife species (Pruvot et al., 2013). Furthermore, it is difficult to sample wildlife populations, particularly to obtain high-quality random samples, which negatively affects the quality of diagnostic results.

Although it has never been quantified, transmission from wildlife is considered a very low risk. Regardless, the livestock industry has opportunities to mitigate the likelihood of inter-species interactions; covering feed storage, keeping wildlife away from forage feeders and other attractants on the farm, and fencing ponds reduced the frequency with which wildlife visit livestock premises (Van Campen & Rhyan, 2010). The livestock industry could be motivated to proactively keep wildlife away from livestock if there were well-documented financial benefits. There is limited information available on disease introductions of endemic pathogens through wildlife (Table 1), and most published models focus on introductions of emerging and notifiable diseases with huge financial impact (e.g., tuberculosis and foot-and-mouth disease).

The main limitation in considering the role of wildlife species is understanding the circulation of MAP at the livestock-wildlife interface and identifying elements that allow certain wildlife populations to maintain MAP infection and potentially act as a reservoir for livestock. Certainly, it should not be assumed that sympatric wildlife and domestic cattle populations always exchange infection (e.g., Whittington, Marsh, & Whitlock, 2001). Increased knowledge on these aspects of the epidemiology of MAP would contribute to our understanding of infection dynamics and may improve JD control programmes (Table 1). Unfortunately, with current limited funding opportunities for wildlife-focused studies, as well as the challenges of designing appropriate multidisciplinary studies, these answers might not be easily generated.

Studies to date investigating the possibility of inter-species MAP transmission involving wildlife have focused simply on the presence or absence of shared strains. None have analysed the degree, directionality of transmission or incidence of infection caused by spillback or spillover. Therefore, a genotyping scheme with an appropriate level of discriminatory power, implemented within a well-designed sampling scheme, is essential for reliably investigating the role of wildlife in the epidemiology of MAP (Fritsch et al., 2012) (Table 1).

8 | SUSCEPTIBILITY

8.1 | Age

Dairy calves are exposed to MAP by the manure from infected adult cattle that shed the bacteria and contaminate water and feed (McKenna, Vanleeuwen, et al., 2006). Greater permeability of the neonatal intestine facilitated MAP entry (Sweeney, 1996), whereas there was increased resistance with age due to repeated exposure to the organism (Delgado et al., 2013) and the dilution effect of the growing rumen (Windsor & Whittington, 2010). However, in an experimental infection trial with 50 calves inoculated at 2 week, 3, 6, 9 or 12 month of age, calves were equally susceptible to infection with MAP up to 1 year of age, based on antibody production, faecal shedding, IFN- γ response, pathology and tissue culture (Mortier, Barkema, & De Buck, 2015; Mortier et al., 2013; Mortier, Barkema, Wilson, et al., 2014; Mortier, Barkema, Orsel, Wolf, et al., 2014; Mortier, Barkema, Orsel, Wolf, De Buck, et al., 2014). Additionally, infection of non-lactating heifers (1–2 year old) occurred when grazing on pasture contaminated with MAP (Fecteau et al., 2010). Therefore, although susceptibility of MAP infection clearly decreases with increasing age (Windsor & Whittington, 2010), control programmes should, depending on the purpose of the programme, consider including cattle of all ages (Table 1).

Although likely animals of all ages can be infected with MAP, age at infection seems to affect immune responses, faecal shedding and lesions at necropsy. Calves inoculated at 3 month of age shed more frequently and had a more robust humoral immune response and more severe lesions at necropsy than calves inoculated at an older age, or at the same age but with a low dose (Mortier, Barkema, Wilson, et al., 2014; Mortier, Barkema, Orsel, Wolf, et al., 2014). The cellular immune response was less marked in calves inoculated at 2 week of age than calves inoculated later in life (Mortier, Barkema, Orsel, Wolf, De Buck, et al., 2014), consistent with the need for a strong cellular immune response to confer better protection against infection (Stabel, 2006). In sheep, age of exposure also strongly affected the outcome of MAP infection, with lambs infected earlier in life starting to shed sooner, having more severe pathology and higher mortality rate (McGregor et al., 2012).

A strong initial cellular immune response, possibly in combination with humoral immunity, appeared to be key to controlling progression of JD (Stabel, 2006). Therefore, infection at a young age when the immune system is still immature and less effective (Chase, Hurley, & Reber, 2008) generally caused more severe lesions. Therefore, age at the time of infection affected the consequences of MAP infection. In a study analysing shedding patterns in naturally infected cows, it was clear that the majority of studied cows never developed high-shedding levels (Mitchell et al., 2015). Those that did, typically never reduced their shedding level to low or no shedding. Cows that eventually became high shedders had a pattern of continuous shedding. In contrast, cows with an intermittent shedding pattern had a low probability to ever become high shedders. In addition, cows that start shedding at a younger age (less than 3 years of age) have a

lower hazard of becoming high shedders compared to cows starting to shed at an older age. These data suggest the presence of three categories of immune control. Cows that are intermittent shedders have the infection process under control (no progressive infection). Cows that start shedding persistently at a young age partially control the infection, but eventually will be high shedders (slow progressive infection), whereas cows that start shedding persistently at an older age cannot effectively control the infection and become high shedders rapidly (Mitchell et al., 2015). However, little more is known about factors influencing the course of MAP infection.

8.2 | Heritability of susceptibility and resistance

Quantitative genetic studies predominantly performed in Holstein-Friesian cows have demonstrated that genetics play an important role in susceptibility to JD, and their heritability estimates indicate that the genetic basis of JD is likely a multigenic trait (Gonda, Chang, Shook, Collins, & Kirkpatrick, 2006; Koets et al., 2000). Channel Island cattle breeds are 1.4–8.3 times more likely to test positive for MAP compared to other dairy breeds (Cetinkaya et al., 1997; Norton, Heuer, & Jackson, 2009; Sorge et al., 2011). Outbreaks of JD occur in beef herds, but the prevalence of MAP infection is much lower than in dairy herds. *Bos indicus* purebreds and cross-breeds have odds ratios 17-fold and 3.5-fold greater than *Bos taurus* breeds for positive ELISA results (Roussel et al., 2005). Despite likely differences due to management (intensively farmed housed dairy cattle vs. extensively beef cattle kept on large pastures), the lower prevalence in beef cattle may also be the result of lower genetic susceptibility of some beef cattle breeds (Table 1). Genetics as an approach to disease control is an emerging discipline. Identifying the genetic basis of the lower susceptibility will provide possibilities for selection for resistance to MAP infection in dairy and beef cattle, as has been done with Red deer (Dobson, Liggett, O'Brien, & Griffin, 2013). While genetic improvement for disease resistance is slow, the results are permanent (van Hulzen et al., 2014; Kirkpatrick & Shook, 2011). There is good evidence in a range of ruminant species for genetic influence on susceptibility to mycobacterial infections (Kirkpatrick & Shook, 2011).

Heritability of test positivity for MAP infection in cattle ranges from 0.041 to 0.159 (Attalla, Seykora, Cole, & Heins, 2010; Gonda et al., 2006; Hinger, Brandt, Horner, & Erhardt, 2007; van Hulzen et al., 2011; Koets et al., 2000), indicating that part of the variability in response to exposure in the population is due to genetics (Table 1). Multiple recent studies also clearly suggest that susceptibility to MAP infection is multigenic or polygenic. The effect of genetic polymorphisms in candidate genes was recently reviewed (Kirkpatrick & Shook, 2011) as well as linkage analysis of genetic susceptibility using genomewide association studies (GWAS; Purdie, Plain, Begg, de Silva, & Whittington, 2011). Limited congruence between studies was attributed to: (i) definitions of case and control; (ii) phenotypic data recorded (tissue culture, faecal culture, blood ELISA, milk ELISA); and (iii) variability in diagnostic methods used between cattle at the same stage of infection.

The host response to MAP can be categorized as susceptible, resistant or tolerant. Attempts to locate loci associated with resistance to paratuberculosis have proven to be more challenging than finding loci associated with susceptibility. A major problem for investigating the influence of genetics on susceptibility against MAP infection is the difficulty in accurately classifying susceptible, resistant or tolerant animals. For example, it is yet unknown whether selection against ELISA positivity results in offspring more resistant to (progression of) the infection, or in offspring that is unable to mount an ELISA response given infection.

There is also an inherent uncertainty in phenotypes, due to the low sensitivity of diagnostic tests for MAP infection. Heritability estimates for resistance to JD are higher when faecal culture is used versus ELISA (Kupper, Brandt, Donat, & Erhardt, 2012). Possible genetic influences on host immunological responses could be a confounding factor (Table 1).

A comprehensive GWAS study (Settles et al., 2009) identified genetic loci associated with four phenotypes: presence of MAP in the tissues, presence in faeces, presence in both tissues and faeces, and presence in tissues but not in faeces. Based on the identification of loci associated with these groups, distinct loci may be important to specific stages of the disease. As different genes are associated with different steps in the infection and disease processes, it is expected that different diagnostics will identify different susceptibility genes corresponding to the observed phenotype. Misclassification can be avoided to some extent by parallel test interpretation.

So far, the IFN- γ assay has not been applied in genetic linkage studies. While it is debatable whether a positive test corresponds with active MAP infection or is the only evidence of exposure, it is likely that, in combination with other diagnostics, this additional information would help identify different phenotypes, particularly the resistant type (Table 1). Ideally, other biomarkers (e.g., transcriptomics) would be used to identify susceptible animals. For this purpose, biomarker outcomes need to be analysed with knowledge of the genetic make-up of exposed animals.

Consolidation of marker-assisted breeding approaches for protection of animal populations against paratuberculosis is the ultimate goal of genetic studies. Animal selection based on marker-assisted breeding might lead to cattle populations with enhanced disease resistance and favourable vaccine responses by stimulating protective immune responses (Fisher et al., 2011). However, before these markers can be used to guide selection, undesirable genetic linkages need to be identified. Such a counterproductive linkage occurred when high milk production was genetically associated with slightly increased susceptibility to MAP (Shook, Chaffer, Wu, & Ezra, 2012). In addition, genetic linkage studies can be expanded from identifying direct associations with susceptibility to infection to associations (e.g., likelihood of progression to clinical disease, specific responses to vaccination or tendency to progress to supershedding) (Table 1).

It is not surprising that studies of genetic influence on JD susceptibility of cattle have not had consistent outcomes (Table 1). Apparent causes include size of the study, population structures, markers used and case definitions. Because cattle populations often

have a high level of relatedness, the hidden presence of closely related animals in a sample would cause an *a priori* unequal distribution of allele frequencies between cases and controls, which could inflate the rate of false-positive associations between trait and marker (Minozzi et al., 2012). This calls for collaborative cross-border experimental designs, use of appropriate statistical models when analysing genetic data and awareness of the genetic structure of the population under study. An initial combined analysis of two distinct populations, in which different phenotypic definitions were used, resulted in discovery of novel putative genetic markers for susceptibility (Minozzi et al., 2012).

Future case-control studies should consider limitations of diagnostic tests and lack of knowledge on the family structure of the study objects. However, alternatives to case-control studies exist and involve cohort study design (classifying animals according to genotype), with disease outcome being determined after a period sufficient for signs/tests of disease to become apparent and be accurately determined in a longitudinal study (Purdie et al., 2011). Regardless, large numbers of animals are needed to detect small differences in susceptibility due to specific gene loci.

Now that multiple SNPs have been associated with susceptibility to MAP infection, a series of follow-up experiments are important next steps, the so-called post-GWAS functional characterization of susceptibility variants. Detailed molecular studies to determine the function of the specific nucleotide substitutions mechanistically should follow. Next, experimental infection trials with specific genotypes could be envisioned which would investigate actual effects on infection dose, susceptibility age, diagnostic outcomes, immune responses, immune cell profiles, etc. (Table 1).

Much effort has been spent identifying a connection between host genotype and susceptibility to MAP infection. However, a portion of the variation in disease manifestation and progression might also be correlated with different MAP genotypes (Table 1). Perhaps specific MAP genotypes associate with a specific host genotype. However, such multilevel genotype linkage analyses have not been carried out.

9 | VACCINATION

The Gudair vaccine has been widely applied in Australian sheep herds and has become the dominant JD control practice. Using this killed vaccine reduced the prevalence of MAP infection and faecal shedding, and mortality in Australian sheep herds considerably (Dhand, Eppleston, Whittington, & Windsor, 2016; Reddacliff, Eppleston, Windsor, Whittington, & Jones, 2006). This vaccine does not prevent MAP infection and can therefore not be used on its own to eradicate MAP infection. In contrast to sheep, in cattle no effective vaccine is available, and the lack of an efficacious vaccine that protects against infection with MAP is hampering control programmes. Existing JD cattle vaccines can reduce clinical impacts of infection, including sometimes reduced shedding, but they do not prevent infection. Additional obstacles are interference with tests to identify

animals infected with other mycobacterial species (e.g., *M. bovis*) and that they can cause severe reactions at the injection site (Kalis, Hesselink, Barkema, & Collins, 2001). Therefore, there is clearly a need for better vaccines in addition to improved diagnostic tests for JD (Table 1).

Ideally, a MAP vaccine would protect against infection, keeping negative herds MAP-free and safeguard young and susceptible animals in high-prevalence herds. Arguably, such an ideal vaccine could be live attenuated, can be administered orally, is sufficiently virulent to trigger protective cell-mediated immune responses, protects exposed animals against infection at the tissue level, is cleared relatively quickly from the vaccinated animal, can be differentiated from wild type strains, is not spread to other animals, protects against homologous and heterologous strains and generates immune responses that can be differentiated from *M. bovis* infections and ideally also from natural MAP infections (with appropriate diagnostic tests).

Current vaccines may partially reduce infectiousness or shedding load, prolong the latent period of infected animals, slow progression from low shedding to high shedding or decrease the cumulative incidence of clinical JD cases; however, they are not effective in preventing infection (Alonso-Hearn et al., 2012; Kalis et al., 2001; Kathaperumal et al., 2008; Kormendy, 1992; Romano & Huygen, 2009; Rosseels & Huygen, 2008; Santema, Hensen, Rutten, & Koets, 2009; Wentink, Bongers, Zeeuwen, & Jaartsveld, 1994). Bacterin vaccines, subunit vaccines (Faisal et al., 2013; Hoek, Rutten, van der Zee, Davies, & Koets, 2010; Koets et al., 2006; Thakur, Aagaard, Stockmarr, Andersen, & Jungersen, 2013) and vector vaccines (Bull et al., 2014; Roupie et al., 2012) have also been created, but have been less effective than expected. Their inability to prevent the establishment of infection leaves the potential for infected animals to break with disease if protective immunity wanes. Subunit vaccines have been reported to provide incomplete protection in murine models (Stabel, Barnhill, Bannantine, Chang, & Osman, 2012) or ruminant models (calves and goats) of infection (Kathaperumal et al., 2008, 2009; Koets et al., 2006). In addition, other subunit vaccines (Facciolo & Mutharia, 2014; Gurung, Begg, Purdie, & Whittington, 2013; Johnston et al., 2014) are being proposed as having potential to prevent infection. Most vaccines have been tested in a preventive pre-exposure setting. However, as calves are commonly born into herds with endemic MAP infection and a MAP-contaminated environment, post-exposure strategies should also be considered (Santema et al., 2013).

Globally, most current vaccines are based on MAP strain 316F formulated in mineral oil adjuvant, either as a live (Neoparasec) or dead strain (Mycopar, ID-Lelystad, Gudair, Silirum). Only Mycopar is licensed in the United States and used on only 5% of US dairy operations (Cho et al., 2012) and under strict control of local veterinary authorities, as vaccinated cattle are more likely to be false positive on a standard bovine tuberculosis test (Musken, van Zijderveld, Eger, & Bakker, 2002). The single intradermal tuberculin test is still the most widely used tuberculosis diagnostic test in cattle (Bastida & Juste, 2011). However, modification of the test, whereby two sites

are injected with either tuberculin from *M. bovis* or MAP and a difference in reactivity is recorded, has already been shown to solve the *M. bovis* interference problem in the vast majority of cases. This comparative intradermal tuberculin test has been available for many years and is an official test according to OIE and EU legislation (Bastida & Juste, 2011). So, false-positive *M. bovis* detection can be eliminated when next-generation vaccines are accompanied by compatible diagnostics (Table 1).

A critical aspect in the development of MAP vaccines is the protective immune responses they are supposed to elicit. The precise nature of a protective immune response is still to be determined (Table 1). An advantage of live-attenuated vaccines (LAV) is that they will stimulate both cell-mediated and humoral immune responses (Faisal et al., 2013; Park et al., 2011). Cell-mediated immune responses have been associated with protection (Settles, Kink, & Talaat, 2014; Stabel & Robbe-Austerman, 2011), and therefore, it has been postulated that if a LAV MAP vaccine could drive the immune response to a pro-inflammatory Th1 profile and prevent a shift to the humoral Th2 response, it might be more effective in delaying disease progression (Coussens, 2004; Stabel & Robbe-Austerman, 2011). In sheep, specific lymphocyte subsets play a role in protecting against MAP infection, including sheep vaccinated with killed vaccine (de Silva, Plain, Begg, Purdie, & Whittington, 2015). Although LAV induced strong protection in a mouse model (Ghosh, Shippy, & Talaat, 2015), in a goat model they showed no protective efficacy in some studies (Hines et al., 2014; Park, Allen, Barrington, & Davis, 2014) and strong reduction in shedding in a recent other study (Shippy et al., 2017); they have apparently not been tested in cattle.

The appropriate level of attenuation of LAV strains has not been established (Table 1). However, it is clear from past experiences that these models should not be restricted to *in vitro* experimentation or mouse models (Bannantine et al., 2014). Furthermore, single knock-out strains may not be optimal vaccine candidates because they may not be attenuated enough to stimulate the protective immune response without causing disease. Although a 1-gene knockout (KO) might yield a strain that cannot persist in the animal and/or environment and therefore not pose an infectious risk to spread and cause disease, a second attenuation should be introduced to eliminate the manipulative mechanisms that inhibit or counteract protected immune responses. So, a vaccine strain should not only trigger protective immune responses but also avoid immune modulation pathways to be activated, which will allow a wild type (WT) strain to establish an infection and ensure that the immune evasion strategies of MAP are rendered neutral.

Selection of the first type of essential gene can be carried out by screening transposon mutant libraries for an optimal LAV (Rathnaiah et al., 2014; Wang, Pritchard, Kreitmann, Montpetit, & Behr, 2014), ideally in a ruminant host. However, selection of the second type of KO requires further study of pathogenesis in *in vitro* and *in vivo* models. We need a better understanding of the immune evasion and immune modulation strategies that MAP clearly is capable of (Arsenault et al., 2014) (Table 1). Due to these immune evasion

properties, MAP can subvert both the induction of acquired immune responses and cell-mediated responses. Reasons for these are potentially 2-fold: First, because the LAV vaccine will have retained these immune evasion properties and therefore elicit inadequate immune responses; and secondly, because an infecting WT strain will be able to subvert immune responses generated by a vaccine. A future protective vaccine may need to prevent or counteract these evasion mechanisms.

9.1 | Diagnostic aspects of vaccination

Differentiation of infected and vaccinated animals (DIVA) is an important consideration. When using a subunit vaccine approach, DIVA can be obtained as has been shown in cattle vaccinated with a subunit vaccine against MAP (Santema et al., 2009). None of the three currently commercially available killed vaccines or LAV lend themselves well to DIVA, although engineered vaccine strains might be better. Complementary diagnostics will have to be developed to incorporate DIVA in an integrated platform of new diagnostic and vaccination strategies, particularly to differentiate *M. bovis* infections from JD vaccination (Table 1). Next-generation diagnostics for MAP infection will likely measure early cell-mediated immune responses. Immune responses against this positive marker of extraneous nature would clearly demonstrate that animals were vaccinated (Table 1). A negative marker, being an immunodominant antigen that is “deleted” from the vaccine strain, could be implemented in novel *M. bovis* diagnostics next to antigens specific for *M. bovis* (e.g., ESAT-6/CFP-10 and Rv3615c) (De Val et al., 2012). Thus, the problematic interference with *M. bovis* testing could be completely mitigated with a marked vaccine and compatible diagnostics.

Following complementary diagnostics for DIVA purposes, new biomarkers that act as correlates of protection and/or that indicate lack of protection are necessary for screening, development and testing of novel MAP vaccine candidates (de Silva et al., 2015). So far, good correlates of protection have remained elusive.

From exercises modelling impacts of imperfect MAP vaccines, it was concluded that vaccination should be integrated into a comprehensive control programme that includes test-and-cull intervention and improved calf rearing management (Bush, Windsor, Toribio, & Webster, 2008; Cho et al., 2012). Cost-benefit analysis of vaccination against MAP in dairy cattle was performed (Groenendaal, Zagmutt, Patton, & Wells, 2015). Vaccination was beneficial by reducing the frequency of heavy shedders and clinically affected animals. A meta-analysis also concluded that vaccination against MAP is a valuable tool in reducing MAP contamination risks and reducing or delaying production losses (Bastida & Juste, 2011).

Newly developed vaccines will need to be well characterized to fully understand their risks and benefits. For example, the risk of shedding in vaccinated cattle needs to be investigated, as was done in sheep vaccinated with the Gudair vaccine (Windsor, Eppleston, Dhand, & Whittington, 2014).

The success of a LAV vaccine will depend on which MAP strain is selected to generate the vaccine (~parent strain). Vaccine strains will likely have to be sufficiently homologous in antigen composition with the majority of field strains. In India, a vaccine based on a regional strain variant was more effective than a commercial JD vaccine (Singh et al., 2013). In our view, there should be a focused effort to characterize the diversity of MAP strains (Table 1). Genotypic and phenotypic variation needs to be investigated to derive quantitative and qualitative understanding of this diversity around the world.

10 | UPTAKE

Nearly all JD prevention and control programmes worldwide are voluntary, and their success depends on enrolment and retention, along with sufficient uptake of recommended best management practices. To establish high participation rates, JD control programmes need to account for farmers' motivators and barriers to enrol and implement recommended management changes (Table 1). In recent years, farmers' “mindset” and its influence on behaviour have become an important focus of research (e.g., Derks, van de Ven, van Werven, Kremer, & Hogeveen, 2012; Garforth, 2015; Jansen & Lam, 2012; Roche, Jones-Bitton, Meehan, Von Massow, & Kelton, 2015). Farmers' attitudes and beliefs towards the disease and the proposed approach for disease control have important roles in their motivation to adhere to suggested management strategies (Ritter et al., 2017). This relatively new avenue of research has important implications for our approaches to motivating individual farmers to adopt optimal JD management practices. Farmers' considerations such as improved herd health and concern over consumer health can be important motivators to participate in JD control or certification programmes (Kovich, Wells, & Friendshuh, 2006; Nielsen, 2011; Roche, 2014). External incentives can also be an important driver to control MAP; a decision analysis from the farmers perspective indicated that a milk-price differentiation of only 0.005/kg milk was sufficient to make enrolment of Dutch dairy farmers in a control programme attractive (Velthuis, Weber, Koeijer, & Van Roermund, 2006). However, management constraints (e.g., limited time or finances) and the perceived complexity of JD control programmes can be critical impediments for uptake of biosecurity measures (Rossiter & Burhans, 1996; Sorge et al., 2010; Wraight et al., 2000), and often farmers prefer to “wait-and-see” how the JD control programme works on other farms before they enrol (Ritter et al., 2015) (Table 1).

Several studies reported that farmers rely on various sources to obtain their farm management information and should be approached according to their specific needs to ensure successful knowledge uptake (Heffernan, Nielsen, Thomson, & Gunn, 2008; Jansen, van Schaik, Renes, & Lam, 2010; Russell & Bewley, 2011). In particular, herd health veterinarians have been regarded as trustworthy and reliable sources of advice on disease and disease risk management (Brennan & Christley, 2013; Ellis-Iversen et al., 2010). Because of this key role veterinarians play in farmers' management

decisions, it is important to employ them as mediators between industry and farmers. However, veterinary practitioners' attitudes towards JD control are often unknown and the extent to which they actively promote enrolment in JD control programmes and/or individual on-farm changes to reduce MAP transmission remains unclear (Table 1). Even farmers provided with veterinary advice on JD control often implemented less than half of the suggestions made (Sorge et al., 2010; Wraight et al., 2000). Miscommunication between dairy practitioners and farmers is likely an important cause of the lack of uptake. For example, veterinarians did not assess producers' expectations sufficiently but provided them with too many, potentially overwhelming, suggestions (Sorge et al., 2010). These apparent gaps in veterinary–farmer communication need to be addressed.

More recent applied research has investigated best practices for communicating with producers and motivating on-farm change, often employing group-based and peer learning approaches. Many of these efforts have emphasized the importance of the veterinarian for making tailored on-farm recommendations, but have also promoted the use of facilitators to better understand and respond to producer mindset. Recent examples from Australia (Kingham & Links, 2012), Denmark (Trier, Nielsen, & Krogh, 2012) and Canada (Roche et al., 2015) have yielded promising results for motivating on-farm change towards effective JD prevention and control.

In addition to a lack of knowledge regarding the discussion of JD control with producers, communication strategies used to inform decision-makers in government and policy authorities, farmer organizations and breeding associations are still unclear. Researchers' responsibility is to provide information that enables evidence-based decisions, for example, regarding JD certification and trade regulations. Clear communication between researchers, farmer organizations and authorities creating evidence-based policy will likely increase motivation to initiate and continue control programmes and improve policy acceptance and uptake by farmers.

Johne's disease is only one of many faecal-orally transmitted infectious diseases in cattle. Measures to prevent infection with MAP will likely also have positive effects on the incidence of infection with *Salmonella* spp., *Cryptosporidium parvum* and *Cryptosporidium bovis*, *Escherichia coli*, and rota- and corona virus (McKenna, Keefe, et al., 2006). Often, the latter bacteria cause more obvious clinical signs than MAP and convince the farmer of the presence and severity of illness. These "cues-to-action" might enhance farmers' openness to improve suboptimal management practices. Therefore, addressing calf health and biosecurity more holistically could help motivate producers that currently do not perceive JD control as high priority (Table 1).

11 | CONCLUSIONS












Nearly a century of JD control programmes has, particularly in the dairy industry, not resulted in sufficient progress. Except for goats in Norway, no reports can be found in a herd in which MAP infection has been eradicated, and in many countries, herd- and animal-level

prevalence has not decreased. As a result, JD continues to cause considerable losses to the livestock industry. The insufficient progress has been the result of gaps in our knowledge about this difficult disease. Research has focused on test development and evaluation, vaccine development, and design and evaluation of management strategies to prevent MAP infection. Many of the knowledge gaps identified are in these areas (Table 1). However, the authors are optimistic that if sufficient progress can be made addressing these knowledge gaps, progress in the control of this insidious disease in the next decades will be better. The introduction of a JD vaccine has made a huge impact in the Australian sheep industry. Development of a JD vaccine with accompanying diagnostic tests that prevents infection and shedding and does not impair tuberculosis diagnostics remains 1 of the most pressing gaps for the livestock industry. Reliably and proactively (pre-shedding) identifying infected animals that will very likely shed the pathogen, potentially involving biomarkers, is another research priority. Susceptibility for MAP infection differs among breeds. Identification of genetic markers that distinguish very susceptible from more resistant animals has the potential to advance JD control. Quantification of the role of calf-to-calf transmission will be necessary to improve cattle control programmes. Uptake of JD control programmes will improve if these knowledge gaps have been satisfactorily addressed. However, because of the voluntary nature of JD programmes, it will still be important to identify factors that motivate farmers to enrol in these programmes.

ACKNOWLEDGEMENTS

The origins of this manuscript are in the disease and product analysis conducted for Johne's disease in the DISCONTTOOLS project (<http://www.disconttools.eu>).

ORCID

- H. W. Barkema  <http://orcid.org/0000-0002-9678-8378>
- K. Orsel  <http://orcid.org/0000-0002-6499-5188>
- S. S. Nielsen  <http://orcid.org/0000-0003-2417-0787>
- A. P. Koets  <http://orcid.org/0000-0001-9809-3679>
- V. P. M. G. Rutten  <http://orcid.org/0000-0002-6171-1805>
- J. P. Bannantine  <http://orcid.org/0000-0002-5692-7898>
- G. P. Keefe  <http://orcid.org/0000-0003-2356-7344>
- C. Heuer  <http://orcid.org/0000-0003-0449-1540>
- T. L. Forde  <http://orcid.org/0000-0001-9058-7826>
- P. J. Griebel  <http://orcid.org/0000-0002-2450-9368>
- J. De Buck  <http://orcid.org/0000-0002-4824-9580>

REFERENCES

- Abbas, B., & Riemann, H. P. (1988). IgG, IgM and IgA in the serum of cattle naturally infected with *Mycobacterium paratuberculosis*. *Comparative Immunology, Microbiology and Infectious Diseases*, 11, 171–175.
- Abubakar, I., Myhill, D., Aliyu, S. H., & Hunter, P. R. (2008). Detection of *Mycobacterium avium* subspecies *paratuberculosis* from patients with

- Crohn's disease using nucleic acid-based techniques: a systematic review and meta-analysis. *Inflammatory Bowel Diseases*, 14, 401–410.
- Adaska, J. M., & Whitlock, R. H. (2012). Low rate of detectable in utero transmission of *Mycobacterium avium* subspecies *paratuberculosis* in a dairy herd with a low prevalence of Johne's disease. *Journal of Veterinary Diagnostic Investigation*, 24, 153–155.
- Ahlstrom, C., Barkema, H. W., & De Buck, J. (2014). Improved short-sequence-repeat genotyping of *Mycobacterium avium* subspecies *paratuberculosis* by using matrix-assisted laser desorption ionization-time of flight mass spectrometry. *Applied and Environmental Microbiology*, 80, 534–539.
- Ahlstrom, C., Barkema, H. W., Stevenson, K., Zadoks, R. N., Biek, R., Kao, R., ... De Buck, J. (2015). Limitations of variable number of tandem repeat typing identified through whole genome sequencing of *Mycobacterium avium* subsp. *paratuberculosis* on a national and herd level. *BMC Genomics*, 16, 161.
- Ahlstrom, C., Barkema, H. W., Stevenson, K., Zadoks, R. N., Biek, R., Kao, R., ... De Buck, J. (2016). Genome-wide diversity and phylogeography of *Mycobacterium avium* subsp. *paratuberculosis* in Canadian dairy cattle. *PLoS ONE*, 11, e0149017.
- Alonso-Hearn, M., Molina, E., Geijo, M., Vazquez, P., Sevilla, I. A., & Garrido, J. M. (2012). Immunization of adult dairy cattle with a new heat-killed vaccine is associated with longer productive life prior to cows being sent to slaughter with suspected paratuberculosis. *Journal of Dairy Science*, 95, 618–629.
- Amonsin, A., Li, L. L., Zhang, Q., Bannantine, J. P., Motiwala, A. S., Sreevatsan, S., & Kapur, V. (2004). Multilocus short sequence repeat sequencing approach for differentiating among *Mycobacterium avium* subsp. *paratuberculosis* strains. *Journal of Clinical Microbiology*, 42, 1694–1702.
- Arsenault, R. J., Maattanen, P., Daigle, J., Potter, A., Griebel, P., & Napper, S. (2014). From mouth to macrophage: mechanisms of innate immune subversion by *Mycobacterium avium* subsp. *paratuberculosis*. *Veterinary Research*, 45, 54.
- Attalla, S. A., Seykora, A. J., Cole, J. B., & Heins, B. J. (2010). Genetic parameters of milk ELISA scores for Johne's disease. *Journal of Dairy Science*, 93, 1729–1735.
- Balseiro, A., García Marín, J. F., Solano, P., Garrido, J. M., & Prieto, J. M. (2008). Histopathological classification of lesions observed in natural cases of paratuberculosis in free-ranging fallow deer (*Dama dama*). *Journal of Comparative Pathology*, 138, 180–188.
- Bannantine, J. P., Everman, J. L., Rose, S. J., Babrak, L., Katani, R., Li, L. L., ... Sreevatsan, S. (2014). Evaluation of eight live attenuated vaccine candidates for protection against challenge with virulent *Mycobacterium avium* subspecies *paratuberculosis* in mice. *Frontiers in Cellular and Infection Microbiology*, 4, 88.
- Bannantine, J. P., Waters, W. R., Stabel, J. R., Palmer, M. V., Li, L., Kapur, V., & Paustian, M. L. (2008). Development and use of a partial *Mycobacterium avium* subspecies *paratuberculosis* protein array. *Proteomics*, 8, 463–474.
- Barkema, H. W., Hesselink, J. W., McKenna, S. L. B., Benedictus, G., & Groenendaal, H. (2010). Global prevalence and economics of infection with *Mycobacterium avium* subsp. *paratuberculosis* in ruminants. In: M. A. Behr & D. M. Collins (Eds.), *Paratuberculosis: organism, disease, control* (pp. 10–21). Wallingford, Oxfordshire, UK: CABI ISBN 9781845936136.
- Barkema, H. W., von Keyserlingk, M. A. G., Kastelic, J. P., Lam, T. J. G. M., Luby, C., Roy, J.-P., ... Kelton, D. F. (2015). Invited review: Changes in the dairy industry affecting dairy health and welfare. *Journal of Dairy Science*, 98, 7426–7445.
- Bartels, C. J., Arnaiz-Seco, J. I., Ruiz-Santa-Quitera, A., Björkman, C., Frössling, J., von Blumröder, D., ... Ortega-Mora, L. M. (2006). Supranational comparison of *Neospora caninum* seroprevalences in cattle in Germany, The Netherlands, Spain and Sweden. *Veterinary Parasitology*, 137, 17–27.
- Bastida, F., & Juste, R. A. (2011). Paratuberculosis control: a review with a focus on vaccination. *Journal of Immune Based Therapies and Vaccines*, 9, 8.
- Begg, D. J., de Silva, K., Di Fiore, L., Taylor, D. L., Bower, K., Zhong, L., ... Whittington, R. J. (2010). Experimental infection model for Johne's disease using a lyophilised, pure culture, seedstock of *Mycobacterium avium* subspecies *paratuberculosis*. *Veterinary Microbiology*, 141, 301–311.
- Begg, D. J., de Silva, K., Plain, K. M., Purdie, A. C., Dhand, N., & Whittington, R. J. (2015). Specific faecal antibody responses in sheep infected with *Mycobacterium avium* subspecies *paratuberculosis*. *Veterinary Immunology and Immunopathology*, 166, 125–131.
- Begg, D. J., & Whittington, R. J. (2008). Experimental animal infection models for Johne's disease, an infectious enteropathy caused by *Mycobacterium avium* subsp. *paratuberculosis*. *The Veterinary Journal*, 176, 129–145.
- Benedictus, G. (1984). Evaluation of organized control of bovine paratuberculosis in Friesland province, The Netherlands. *Tijdschrift voor Diergeneeskunde*, 109, 905–916.
- Benedictus, A., Mitchell, R. M., Linde-Widmann, A., Sweeney, R., Fyock, T., Schukken, Y. H., & Whitlock, R. H. (2008). Transmission parameters of *Mycobacterium avium* subspecies *paratuberculosis* infections in a dairy herd going through a control program. *Preventive Veterinary Medicine*, 83, 215–227.
- Bhattarai, B., Fosgate, G. T., Osterstock, J. B., Fossler, C. P., Park, S. C., & Roussel, A. J. (2014). Comparison of calf weaning weight and associated economic variables between beef cows with and without serum antibodies against or isolation from feces of *Mycobacterium avium* subsp. *paratuberculosis*. *Journal of the American Veterinary Medical Association*, 243, 1609–1615.
- Bölske, G., & Herthnek, D. (2010). Diagnosis of paratuberculosis by PCR. In: M. A. Behr & D. M. Collins (Eds.), *Paratuberculosis: organism, disease, control* (pp. 267–283). Wallingford, Oxfordshire, UK: CABI. ISBN 9781845936136
- Bolton, M. W., Grooms, D. L., & Kaneene, J. B. (2005). *Fecal shedding of Mycobacterium avium ssp. paratuberculosis in calves: Implications for disease control and management*. In: Proc. 8th Int Colloq Paratuberculosis, Copenhagen, Denmark, p. 128.
- Borrmann, E., Möbius, P., Diller, R., & Köhler, H. (2011). Divergent cytokine responses of macrophages to *Mycobacterium avium* subsp. *paratuberculosis* strains of Types II and III in a standardized in vitro model. *Veterinary Microbiology*, 152, 101–111.
- Brennan, M. L., & Christley, R. M. (2013). Cattle producers' perceptions of biosecurity. *BMC Veterinary Research*, 9, 71.
- Britton, L. E., Cassidy, J. P., O'Donovan, J., Gordon, S. V., & Markey, B. (2016). Potential application of emerging diagnostic techniques to the diagnosis of bovine Johne's disease (paratuberculosis). *The Veterinary Journal*, 209, 32–39.
- Bryant, J. M., Thibault, V. C., Smith, D. G., McLuckie, J., Heron, I., Sevilla, I. A., ... Stevenson, K. (2016). Phylogenomic exploration of the relationships between strains of *Mycobacterium avium* subspecies *paratuberculosis*. *BMC Genomics*, 26, 79.
- Bull, T. J., Vrettou, C., Linedale, R., McGuinness, C., Strain, S., McNair, J., ... Hope, J. C. (2014). Immunity, safety and protection of an Adenovirus 5 prime-Modified Vaccinia virus Ankara boost subunit vaccine against *Mycobacterium avium* subspecies *paratuberculosis* infection in calves. *Veterinary Research*, 45, 112.
- Bush, R. D., Windsor, P. A., Toribio, J. A., & Webster, S. R. (2008). Financial modelling of the potential cost of ovine Johne's disease and the benefit of vaccinating sheep flocks in southern New South Wales. *Australian Veterinary Journal*, 86, 398–403.
- Caldow, G., & Gunn, G. J. (2001). *Assessment of surveillance and control of Johne's disease in farm animals in Great Britain*. Retrieved from www.defra.gov.uk/animalh/diseases/sac2.PDF (Accessed May 9, 2017).

- Capsel, R. T., Thoen, C. O., Reinhardt, T. A., Lippolis, J. D., Olsen, R., Stabel, J. R., & Bannantine, J. P. (2016). Composition and potency characterization of *Mycobacterium avium* subsp. *paratuberculosis* purified protein derivatives. *PLoS ONE*, *11*, e0154685.
- Carlos, P., Roupie, V., Holbert, S., Ascencio, F., Huygen, K., Gomez-Anduro, G., ... Angulo, C. (2015). In silico epitope analysis of unique and membrane associated proteins from *Mycobacterium avium* subsp. *paratuberculosis* for immunogenicity and vaccine evaluation. *Journal of Theoretical Biology*, *384*, 1–9.
- Casey, J. L., Sanalla, A. M., Tamvakis, D., Thalmann, C., Carroll, E. L., Parisi, K., ... Foley, M. (2011). Peptides specific for *Mycobacterium avium* subspecies *paratuberculosis* infection: diagnostic potential. *Protein Engineering, Design & Selection*, *24*, 589–596.
- Cetinkaya, B., Erdogan, H. M., & Morgan, K. L. (1997). Relationships between the presence of Johne's disease and farm and management factors in dairy cattle in England. *Preventive Veterinary Medicine*, *32*, 253–266.
- Chase, C. C., Hurley, D. J., & Reber, A. J. (2008). Neonatal immune development in the calf and its impact on vaccine response. *Veterinary Clinics of North America: Food Animal Practice*, *24*, 87–104.
- Chi, J., VanLeeuwen, J. A., Weersink, A., & Keefe, G. P. (2002). Direct production losses and treatment costs from bovine viral diarrhoea virus, bovine leukosis virus, *Mycobacterium avium* subspecies *paratuberculosis*, and *Neospora caninum*. *Preventive Veterinary Medicine*, *55*, 137–153.
- Cho, J., Tauer, L. W., Schukken, Y. H., Gomez, M. I., Smith, R. L., Lu, Z., & Gröhn, Y. T. (2012). Economic analysis of *Mycobacterium avium* subspecies *paratuberculosis* vaccines in dairy herds. *Journal of Dairy Science*, *95*, 1855–1872.
- Christopher-Hennings, J., Dammen, M. A., Weeks, S. R., Epperson, W. B., Singh, S. N., Steinlicht, G. L., ... Nelson, E. A. (2003). Comparison of two DNA extractions and nested PCR, real-time PCR, a new commercial PCR assay, and bacterial culture for detection of *Mycobacterium avium* subsp. *paratuberculosis* in bovine feces. *Journal of Veterinary Diagnostic Investigation*, *15*, 87–93.
- Clark, D. L. Jr, Koziczkowski, J. J., Radcliff, R. P., Carlson, R. A., & Ellingson, J. L. (2008). Detection of *Mycobacterium avium* subspecies *paratuberculosis*: comparing fecal culture versus serum enzyme-linked immunosorbent assay and direct fecal polymerase chain reaction. *Journal of Dairy Science*, *91*, 2620–2627.
- Coffin, J. W., Condon, C., Compston, C. A., Potter, K. N., Lamontagne, L. R., Shafiq, J., & Kunimoto, D. Y. (1992). Use of restriction fragment length polymorphisms resolved by pulsed-field gel electrophoresis for subspecies identification of mycobacteria in the *Mycobacterium avium* complex and for isolation of DNA probes. *Journal of Clinical Microbiology*, *30*, 1829–1836.
- Colavecchia, S. B., Fernández, B., Jolly, A., Minatel, L., Hajos, S. E., Paolicchi, F. A., & Mundo, S. L. (2016). Immunological findings associated with Argentinean strains of *Mycobacterium avium* subsp. *paratuberculosis* in bovine models. *Veterinary Immunology and Immunopathology*, *176*, 28–33.
- Collins, M. T. (1994). Clinical approach to control of bovine paratuberculosis. *Journal of the American Veterinary Medical Association*, *204*, 208–210.
- Collins, M. T. (1996). Diagnosis of paratuberculosis. *Veterinary Clinics of North America: Food Animal Practice*, *12*, 357–371.
- Collins, M. T. (2001). Prevention of paratuberculosis. *Bulletin of the International Dairy Federation*, *364*, 46–53.
- Collins, D. M., & de Lisle, G. W. (1986). Restriction endonuclease analysis of various strains of *Mycobacterium paratuberculosis* isolated from cattle. *American Journal of Veterinary Research*, *47*, 2226–2229.
- Collins, M. T., Eggleston, V., & Manning, E. J. (2010). Successful control of Johne's disease in nine dairy herds: results of a six-year field trial. *Journal of Dairy Science*, *93*, 1638–1643.
- Cook, K. L., & Britt, J. S. (2007). Optimization of methods for detecting *Mycobacterium avium* subsp. *paratuberculosis* in environmental samples using quantitative, real-time PCR. *Journal of Microbiol Methods*, *69*, 154–160.
- Corbett, C., De Buck, J., Orsel, K., & Barkema, H. W. (2017). Fecal shedding and tissue infections demonstrate transmission of *Mycobacterium avium* subsp. *paratuberculosis* in group-housed dairy calves. *Veterinary Research*, *48*, 27.
- Corner, L. A. L., Pfeiffer, D. U., & Abbott, K. A. (2004). The respiratory tract as a hypothetical route of infection of cattle with *Mycobacterium avium* subspecies *paratuberculosis*. *Australian Veterinary Journal*, *82*, 170–173.
- Coussens, P. M. (2004). Model for immune responses to *Mycobacterium avium* subspecies *paratuberculosis* in cattle. *Infection and Immunity*, *72*, 3089–3096.
- Crossley, B. M., Zagmutt-Vergara, F. J., Fyock, T. L., Whitlock, R. H., & Gardner, I. A. (2005). Fecal shedding of *Mycobacterium avium* subsp. *paratuberculosis* by dairy cows. *Veterinary Microbiology*, *107*, 257–263.
- Dargatz, D. D., Byrum, B. A., Barber, L. K., Sweeney, R. W., Whitlock, R. H., Shulaw, W. P., ... Stabel, J. R. (2001). Evaluation of a commercial ELISA for diagnosis of paratuberculosis in cattle. *Journal of the American Veterinary Medical Association*, *218*, 1163–1166.
- David, J., Barkema, H. W., Luo Guan, L., & De Buck, J. (2014). Gene-expression profiling of calves 6 and 9 months after inoculation with *Mycobacterium avium* subspecies *paratuberculosis*. *Veterinary Research*, *45*, 96.
- David, J., Barkema, H. W., Mortier, R., Ghosh, S., Luo Guan, L., & De Buck, J. (2014). Gene expression profiling and putative biomarkers of calves 3 months after infection with *Mycobacterium avium* subspecies *paratuberculosis*. *Veterinary Immunology and Immunopathology*, *160*, 107–117.
- Davidson, F. W., Ahlstrom, C., De Buck, J., Whitney, H. G., & Tahlan, K. (2016). Examination of *Mycobacterium avium* subspecies *paratuberculosis* mixed genotype infections in dairy animals using a whole genome sequencing approach. *PeerJ*, *4*, e2793.
- De Buck, J., Shaykhutdinov, R., Barkema, H. W., & Vogel, H. J. (2014). Metabolomic profiling in cattle experimentally infected with *Mycobacterium avium* subsp. *paratuberculosis*. *PLoS ONE*, *9*, e111872.
- De Val, B. P., Nofrarias, M., Lopez-Soria, S., Garrido, J. M., Vordermeier, H. M., Villarreal-Ramos, B., ... Domingo, M. (2012). Effects of vaccination against paratuberculosis on tuberculosis in goats: diagnostic interferences and cross-protection. *BMC Veterinary Research*, *8*, 191.
- Delgado, L., Marin, J. F., Muñoz, M., Benavides, J., Juste, R. A., García-Pariente, C., ... Pérez, V. (2013). Pathological findings in young and adult sheep following experimental infection with 2 different doses of *Mycobacterium avium* subspecies *paratuberculosis*. *Veterinary Pathology*, *50*, 857–866.
- Dennis, M. M., Reddacliff, L. A., & Whittington, R. J. (2011). Longitudinal study of clinicopathological features of Johne's disease in sheep naturally exposed to *Mycobacterium avium* subspecies *paratuberculosis*. *Veterinary Pathology*, *48*, 565–575.
- Derakhshani, H., De Buck, J., Mortier, R., Barkema, H. W., Krause, D. O., & Khafipour, E. (2016). The features of fecal and ileal mucosa-associated microbiota in dairy calves during early infection with *Mycobacterium avium* subspecies *paratuberculosis*. *Frontiers in Microbiology*, *7*, 426.
- Derks, M., van de Ven, L. M., van Werven, T., Kremer, W. D., & Hogeveen, H. (2012). The perception of veterinary herd health management by Dutch dairy farmers and its current status in the Netherlands: a survey. *Preventive Veterinary Medicine*, *104*, 207–215.
- Deutz, A., Sperser, J., Wagner, P., Rosengarten, R., & Köfer, J. (2005). *Mycobacterium avium* subsp. *paratuberculosis* in wild animal species and cattle in Styria/Austria. *Berliner und Munchener Tierärztliche Wochenschrift*, *118*, 314–320.

- Devonshire, A. S., O'Sullivan, D. M., Honeyborne, I., Jones, G., Karcmarczyk, M., Pavšič, J., ... Huggett, J. F. (2016). The use of digital PCR to improve the application of quantitative molecular diagnostic methods for tuberculosis. *BMC Infectious Diseases*, 16, 366.
- Dhand, N. K., Eppleston, J., Whittington, R. J., & Windsor, P. A. (2016). Changes in prevalence of ovine paratuberculosis following vaccination with Gudair: Results of a longitudinal study conducted over a decade. *Vaccine*, 34, 5107–5113.
- Dobson, B., Liggett, S., O'Brien, R., & Griffin, J. F. (2013). Innate immune markers that distinguish red deer (*Cervus elaphus*) selected for resistant or susceptible genotypes for Johne's disease. *Veterinary Research*, 44, 5.
- Dorshorst, N. C., Collins, M. T., & Lombard, J. E. (2006). Decision analysis model for paratuberculosis control in commercial dairy herds. *Preventive Veterinary Medicine*, 75, 92–122.
- Eisenberg, S. W., Koets, A. P., Hoeboer, J., Bouman, M., Heederik, D., & Nielen, M. (2010). Presence of *Mycobacterium avium* subsp. *paratuberculosis* in environmental samples collected on commercial Dutch dairy farms. *Applied and Environment Microbiology*, 76, 6310–6312.
- Eisenberg, S. W., Koets, A. P., Nielen, M., Heederik, D., Mortier, R., De Buck, J., & Orsel, K. (2011). Intestinal infection following aerosol challenge of calves with *Mycobacterium avium* subspecies *paratuberculosis*. *Veterinary Research*, 42, 117.
- Eisenberg, S. W., Nielen, M., Santema, W., Houwers, D. J., Heederik, D., & Koets, A. P. (2010). Detection of spatial and temporal spread of *Mycobacterium avium* subsp. *paratuberculosis* in the environment of a cattle farm through bio-aerosols. *Veterinary Microbiology*, 143, 284–292.
- Eisenberg, S. W., Rutten, V. P., & Koets, A. P. (2015). Dam *Mycobacterium avium* subspecies *paratuberculosis* (MAP) infection status does not predetermine calves for future shedding when raised in a contaminated environment: a cohort study. *Veterinary Research*, 46, 70.
- Eisenberg, S. W. F., Veldman, E., Rutten, V. P. M. G., & Koets, A. P. (2015). A longitudinal study of factors influencing the results of a *Mycobacterium avium* ssp. *paratuberculosis* antibody ELISA in milk of dairy cows. *Journal of Dairy Science*, 98, 2345–2355.
- Elliott, N. G., Hough, R. L., Avery, L. M., Maltin, C. A., & Campbell, C. D. (2015). Environmental risk factors in the incidence of Johne's disease. *Critical Reviews in Microbiology*, 41, 488–507.
- Ellis-Iversen, J., Cook, A. J., Watson, E., Nielen, M., Larkin, L., Wooldridge, M., & Hogeveen, H. (2010). Perceptions, circumstances and motivators that influence implementation of zoonotic control programs on cattle farms. *Preventive Veterinary Medicine*, 93, 276–285.
- Eltholth, M. M., Marsh, V. R., Van Winden, S., & Guitian, F. J. (2009). Contamination of food products with *Mycobacterium avium paratuberculosis*: a systematic review. *Journal of Applied Microbiology*, 107, 1061–1071.
- Facciuolo, A., Kelton, D. F., & Mutharia, L. M. (2013). Novel secreted antigens of *Mycobacterium paratuberculosis* as serodiagnostic biomarkers for Johne's disease in cattle. *Clinical and Vaccine Immunology*, 20, 1783–1791.
- Facciuolo, A., & Mutharia, L. M. (2014). Mycobacterial glycoproteins: a novel subset of vaccine candidates. *Frontiers in Cellular and Infection Microbiology*, 4, 133.
- Faisal, S. M., Chen, J. W., Yan, F., Chen, T. T., Useh, N. M., Yan, W., ... Chang, Y. F. (2013). Evaluation of a *Mycobacterium avium* subsp. *paratuberculosis* leuD mutant as a vaccine candidate against challenge in a caprine model. *Clinical and Vaccine Immunology*, 20, 572–581.
- Fecteau, M. E., Whitlock, R. H., Buergelt, C. D., & Sweeney, R. W. (2010). Exposure of young dairy cattle to *Mycobacterium avium* subsp. *paratuberculosis* (MAP) through intensive grazing of contaminated pastures in a herd positive for Johne's disease. *Canadian Veterinary Journal*, 51, 198–200.
- Feller, M., Huwiler, K., Stephan, R., Altpeter, E., Shang, A., Furrer, H., ... Egger, M. (2007). *Mycobacterium avium* subspecies *paratuberculosis* and Crohn's disease: a systematic review and meta-analysis. *The Lancet Infectious Diseases*, 7, 607–613.
- Ferroglio, E., Nebbia, P., Robino, P., Rossi, L., & Rosati, S. (2000). *Mycobacterium paratuberculosis* infection in two free-ranging Alpine ibex. *Revue Scientifique et Technique*, 19, 859–862.
- Fisher, C. A., Bhattarai, E. K., Osterstock, J. B., Dowd, S. E., Seabury, P. M., Vikram, M., ... Seabury, C. M. (2011). Evolution of the bovine TLR gene family and member associations with *Mycobacterium avium* subspecies *paratuberculosis* infection. *PLoS ONE*, 6, e27744.
- Fock-Chow-Tho, D., Topp, E., Ibeagha-Awemu, E. A., & Bissonnette, N. (2017). Comparison of commercial DNA extraction kits and quantitative PCR systems for better sensitivity in detecting the causative agent of paratuberculosis in dairy cow fecal samples. *Journal of Dairy Science*, 100, 572–581.
- Fritsch, I., Luyven, G., Köhler, H., Lutz, W., & Möbius, P. (2012). Suspicion of *Mycobacterium avium* subsp. *paratuberculosis* transmission between cattle and wild-living red deer (*Cervus elaphus*) by multitarget genotyping. *Applied and Environment Microbiology*, 78, 1132–1139.
- Frössling, J., Wahlström, H., Agren, E. C., Cameron, A., Lindberg, A., & Sternberg Lewerin, S. (2013). Surveillance system sensitivities and probability of freedom from *Mycobacterium avium* subsp. *paratuberculosis* infection in Swedish cattle. *Preventive Veterinary Medicine*, 108, 47–62.
- Garcia, A. B., & Shalloo, L. (2015). Invited review: The economic impact and control of paratuberculosis in cattle. *Journal of Dairy Science*, 98, 5019–5039.
- Garforth, C. (2015). Livestock keepers' reasons for doing and not doing things which governments, vets and scientists would like them to do. *Zoonoses Public Health*, 62(Suppl 1), 29–38.
- Geraghty, T., Graham, D. A., Mullowney, P., & More, S. J. (2014). A review of bovine Johne's disease control activities in 6 endemically infected countries. *Preventive Veterinary Medicine*, 116, 1–11.
- Ghosh, P., Shippy, D. C., & Talaat, A. M. (2015). Superior protection elicited by live-attenuated vaccines in the murine model of paratuberculosis. *Vaccine*, 33, 7262–7270.
- Gollnick, N. S., Mitchell, R. M., Baumgart, M., Janagama, H. K., Sreevatsan, S., & Schukken, Y. H. (2007). Survival of *Mycobacterium avium* subsp. *paratuberculosis* in bovine monocyte-derived macrophages is not affected by host infection status but depends on the infecting bacterial genotype. *Veterinary Immunology and Immunopathology*, 120, 93–105.
- Gonda, M. G., Chang, Y. M., Shook, G. E., Collins, M. T., & Kirkpatrick, B. W. (2006). Genetic variation of *Mycobacterium avium* ssp. *paratuberculosis* infection in US Holsteins. *Journal of Dairy Science*, 89, 1804–1812.
- Griebel, P., Arsenault, R., Facciuolo, A., Kusalik, A., Liang, G., Määttänen, P., ... Luo Guan, L. (2014). *Models and methods to dissect mucosal immune responses following MAP infection*. In: Proc. 12th Int. Colloq. Paratuberculosis 4–6.
- Groenendaal, H., Nielen, M., & Hesselink, J. W. (2003). Development of the Dutch Johne's disease control program supported by a simulation model. *Preventive Veterinary Medicine*, 60, 69–90.
- Groenendaal, H., Nielen, M., Jalvingh, A. W., Horst, S. H., Galligan, D. T., & Hesselink, J. W. (2002). A simulation of Johne's disease control. *Preventive Veterinary Medicine*, 54, 225–245.
- Groenendaal, H., Zagmutt, F. J., Patton, E. A., & Wells, S. J. (2015). Cost-benefit analysis of vaccination against *Mycobacterium avium* ssp. *paratuberculosis* in dairy cattle, given its cross-reactivity with tuberculosis tests. *Journal of Dairy Science*, 98, 6070–6084.
- Gurung, R. B., Begg, D. J., Purdie, A. C., & Whittington, R. J. (2013). Antigenicity in sheep of synthetic peptides derived from stress-regulated *Mycobacterium avium* subsp. *paratuberculosis* proteins and comparison with recombinant protein and complex native antigens. *Veterinary Immunology and Immunopathology*, 158, 46–52.
- Heffernan, C., Nielsen, L., Thomson, K., & Gunn, G. (2008). An exploration of the drivers to bio-security collective action among a sample

- of UK cattle and sheep farmers. *Preventive Veterinary Medicine*, 87, 358–372.
- Heuer, C., Mitchell, R. M., Schukken, Y. H., Lu, Z., Verdugo, C., & Wilson, P. R. (2012). Modelling transmission dynamics of paratuberculosis of red deer under pastoral farming conditions. *Preventive Veterinary Medicine*, 106, 63–74.
- Hines, M. E. 2nd, Stabel, J. R., Sweeney, R. W., Griffin, F., Talaat, A. M., Bakker, D., ... Whitlock, R. H. (2007). Experimental challenge models for Johne's disease: a review and proposed international guidelines. *Veterinary Microbiology*, 122, 197–222.
- Hines, M. E. 2nd, Turnquist, S. E., Ilha, M. R., Rajeev, S., Jones, A. L., Whittington, L., ... Kapur, V. (2014). Evaluation of novel oral vaccine candidates and validation of a caprine model of Johne's disease. *Frontiers in Cellular and Infection Microbiology*, 4, 26.
- Hinger, M., Brandt, H., Horner, S., & Erhardt, G. (2007). Short communication: Association analysis of microsatellites and *Mycobacterium avium* subspecies *paratuberculosis* antibody response in German Holsteins. *Journal of Dairy Science*, 90, 1957–1961.
- Hoek, A., Rutten, V. P., van der Zee, R., Davies, C. J., & Koets, A. P. (2010). Epitopes of *Mycobacterium avium* ssp. *paratuberculosis* 70 kDa heat-shock protein activate bovine helper T cells in outbred cattle. *Vaccine*, 28, 5910–5919.
- van Hulzen, K. J., Koets, A. P., Nielen, M., Heuven, H. C., van Arendonk, J. A., & Klinkenberg, D. (2014). The effect of genetic selection for Johne's disease resistance in dairy cattle: Results of a genetic-epidemiological model. *Journal of Dairy Science*, 97, 1762–1773.
- van Hulzen, K. J., Nielen, M., Koets, A. P., de Jong, G., van Arendonk, J. A., & Heuven, H. C. (2011). Effect of herd prevalence on heritability estimates of antibody response to *Mycobacterium avium* subspecies *paratuberculosis*. *Journal of Dairy Science*, 94, 992–997.
- Hutchings, M. R., Severson, K., Greig, A., Davidson, R. S., Marion, G., & Judge, J. (2010). Infection of non-ruminant wildlife by *Mycobacterium avium* subsp. *paratuberculosis*. In: M. A. Behr & D. M. Collins (Eds.), *Paratuberculosis: organism, disease, control* (pp. 188–200). Wallingford, Oxfordshire, UK: CABI. ISBN 9781845936136.
- Huygen, K. (2014). The immunodominant T-cell epitopes of the mycolyl-transferases of the antigen 85 complex of *M. tuberculosis*. *Frontiers in Immunology*, 5, 321.
- Janagama, H. K., Jeong, Ki., Kapur, V., Coussens, P., & Sreevatsan, S. (2006). Cytokine responses of bovine macrophages to diverse clinical *Mycobacterium avium* subspecies *paratuberculosis* strains. *BMC Microbiology*, 6, 10.
- Jansen, J., & Lam, T. J. G. M. (2012). The role of communication in improving udder health. *Veterinary Clinics of North America: Food Animal Practice*, 28, 363–379.
- Jansen, J., van Schaik, G., Renes, R. J., & Lam, T. J. (2010). The effect of a national mastitis control program on the attitudes, knowledge, and behavior of farmers in the Netherlands. *Journal of Dairy Science*, 93, 5737–5747.
- Jayarao, B. M., Pillai, S. R., Wolfgang, D. R., Griswolf, D. R., Rossiter, C. A., Tewari, D., ... Hutchinson, L. J. (2004). Evaluation of IS900 PCR assay for detection of *Mycobacterium avium* subspecies *paratuberculosis* infection in cattle using quarter milk and bulk milk samples. *Foodborne pathogens and disease*, 1, 17–26.
- Johnston, C. D., Bannantine, J. P., Govender, R., Endersen, L., Pletzer, D., Weingart, H., ... Sleator, R. D. (2014). Enhanced expression of codon optimized *Mycobacterium avium* subsp. *paratuberculosis* antigens in *Lactobacillus salivarius*. *Frontiers in Cellular and Infection Microbiology*, 4, 120.
- Kalis, C. H. J., Barkema, H. W., Hesselink, J. W., Van Maanen, C., & Collins, M. T. (2002). Evaluation of two absorbed ELISAs and a Complement Fixation Test as replacements for fecal culture in the detection of cows shedding *Mycobacterium avium* subspecies *paratuberculosis*. *Journal of Veterinary Diagnostic Investigation*, 14, 219–224.
- Kalis, C. H. J., Collins, M. T., Barkema, H. W., & Hesselink, J. W. (2004). Certification of herds as free of *Mycobacterium paratuberculosis* infection: actual pooled faecal results versus certification model predictions. *Preventive Veterinary Medicine*, 65, 189–204.
- Kalis, C. H. J., Collins, M. T., Hesselink, J. W., & Barkema, H. W. (2003). Specificity of two tests for the early diagnosis of bovine paratuberculosis based on cell-mediated immunity: the Johnin skin test and the gamma interferon assay. *Veterinary Microbiology*, 97, 73–86.
- Kalis, C. H. J., Hesselink, J. W., Barkema, H. W., & Collins, M. T. (2001). Use of long-term vaccination with a killed vaccine to prevent fecal shedding of *Mycobacterium avium* subsp. *paratuberculosis* in dairy herds. *American Journal of Veterinary Research*, 62, 270–274.
- Kasnitz, N., Köhler, H., Weigoldt, M., Gerlach, G. F., & Möbius, P. (2013). Stability of genotyping target sequences of *Mycobacterium avium* subsp. *paratuberculosis* upon cultivation on different media, in vitro- and in vivo passage, and natural infection. *Veterinary Microbiology*, 167, 573–583.
- Kathaperumal, K., Kumanan, V., McDonough, S., Chen, L. H., Park, S. U., Moreira, M. A., ... Chang, Y. F. (2009). Evaluation of immune responses and protective efficacy in a goat model following immunization with a cocktail of recombinant antigens and a polyprotein of *Mycobacterium avium* subsp. *paratuberculosis*. *Vaccine*, 27, 123–135.
- Kathaperumal, K., Park, S. U., McDonough, S., Stehman, S., Akey, B., Huntley, J., ... Chang, Y. F. (2008). Vaccination with recombinant *Mycobacterium avium* subsp. *paratuberculosis* proteins induces differential immune responses and protects calves against infection by oral challenge. *Vaccine*, 26, 1652–1663.
- Kawaji, S., Taylor, D. L., Mori, Y., & Whittington, R. J. (2007). Detection of *Mycobacterium avium* subsp. *paratuberculosis* in ovine faeces by direct quantitative PCR has similar or greater sensitivity compared to radiometric culture. *Veterinary Microbiology*, 125, 36–48.
- Kelton, D., Van De Water, D., Cantin, R., Hand, K., Innes, C., & Todd, C. (2014). A potential role for bulk milk or herd pool ELISA testing in a Johne's disease control program. Proc. 12 Intern. Colloq. Paratuberculosis, pp 71.
- Kennedy, D. J. (2001). Control of paratuberculosis. *Bulletin of the International Dairy Federation*, 364, 29–45.
- Kennedy, D. J. (2011). International efforts at paratuberculosis control. *Veterinary Clinics of North America: Food Animal Practice*, 27, 647–654.
- Khare, S., Ficht, T. A., Santos, R. L., Romano, J., Ficht, A. R., Zhang, S., ... Adams, L. G. (2004). Rapid and sensitive detection of *Mycobacterium avium* subsp. *paratuberculosis* in bovine milk and feces by a combination of immunomagnetic bead separation-conventional PCR and real-time PCR. *Journal of Clinical Microbiology*, 42, 1075–1081.
- Khol, J. L., Wassertheurer, M., Sodoma, E., Revilla-Fernandey, S., Damoser, J., Österreicher, E., ... Baumgartner, W. (2013). Long-term detection of *Mycobacterium avium* subspecies *paratuberculosis* in individual and bulk tank milk from a dairy herd with a low prevalence of Johne's disease. *Journal of Dairy Science*, 96, 3517–3524.
- Kiehnbaum, L. A., Amonsin, A., Wells, S. J., & Kapur, V. (2005). Amplified fragment length polymorphism to detect clonal diversity and distribution of *Mycobacterium avium* subspecies *paratuberculosis* in selected Minnesota dairy cattle. *Journal of Veterinary Diagnostic Investigation*, 17, 311–315.
- Kingham, L., & Links, I. (2012). *Development of a group approach to facilitate vaccination for ovine Johne's disease (OJD) in New South Wales*. Page 178 in 11th International Colloquium on Paratuberculosis Proc. Sydney, Australia
- Kirkeby, C., Græsbøll, K., Halasa, T., Toft, N., & Nielsen, S. S. (2015). Mean effective sensitivity for *Mycobacterium avium* subsp. *paratuberculosis* infection in cattle herds. *BMC Veterinary Research*, 11, 190.
- Kirkpatrick, B. W., & Shook, G. E. (2011). Genetic susceptibility to paratuberculosis. *Veterinary Clinics of North America: Food Animal Practice*, 27, 559–571.

- Klawonn, W., Einax, E., Pützschel, R., Schmidt, M., & Donat, K. (2016). Johne's disease: reliability of environmental sampling to characterize *Mycobacterium avium* subspecies *paratuberculosis* (MAP) infection in beef cow-calf herds. *Epidemiology and Infection*, *144*, 2392–2400.
- Kluge, J. P., Merkal, R. S., Monlux, W. S., Larsen, A. B., Kopecky, K. E., Ramsey, F. K., & Lehmann, R. P. (1968). Experimental paratuberculosis in sheep after oral, intratracheal, or intravenous inoculation lesions and demonstration of etiologic agent. *American Journal of Veterinary Research*, *29*, 953–962.
- Koets, A. P., Adugna, G., Janss, L. L., van Weering, H. J., Kalis, C. H., Wentink, G. H., ... Schukken, Y. H. (2000). Genetic variation of susceptibility to *Mycobacterium avium* subsp. *paratuberculosis* infection in dairy cattle. *Journal of Dairy Science*, *83*, 2702–2708.
- Koets, A., Hoek, A., Langelaar, M., Overdijk, M., Santema, W., Franken, P., ... Rutten, V. (2006). Mycobacterial 70 kD heat-shock protein is an effective subunit vaccine against bovine paratuberculosis. *Vaccine*, *24*, 2550–2559.
- Kormendy, B. (1992). Paratuberculosis vaccine in a large dairy herd. *Acta Veterinaria Hungarica*, *40*, 171–184.
- Kovich, D. A., Wells, S. J., & Friendshuh, K. (2006). Evaluation of the Voluntary Johne's Disease Herd Status Program as a source of replacement cattle. *Journal of Dairy Science*, *89*, 3466–3470.
- Kralik, P., Beran, V., & Pavlik, I. (2012). Enumeration of *Mycobacterium avium* subsp. *paratuberculosis* by quantitative real-time PCR, culture on solid media and optical densitometry. *BMC Research Notes*, *5*, 114.
- Kudahl, A. B., Nielsen, S. S., & Ostergaard, S. (2011). Strategies for time of culling in control of paratuberculosis in dairy herds. *Journal of Dairy Science*, *94*, 3824–3834.
- Kupper, J., Brandt, H., Donat, K., & Erhardt, G. (2012). Heritability estimates for *Mycobacterium avium* subspecies *paratuberculosis* status of German Holstein cows tested by fecal culture. *Journal of Dairy Science*, *95*, 2734–2739.
- Kurade, N. P., Tripathi, B. N., & Rajukumar, K. (2004). Sequential development of histologic lesions and their relationship with bacterial isolation, fecal shedding, and immune responses during progressive stages of experimental infection of lambs with *Mycobacterium avium* subsp. *paratuberculosis*. *Veterinary Pathology*, *41*, 378–387.
- Laurin, E. L., Chaffer, M., McClure, J. T., McKenna, S. L., & Keefe, G. P. (2015). The association of detection method, season, and lactation stage on identification of fecal shedding in *Mycobacterium avium* ssp. *paratuberculosis* infectious dairy cows. *Journal of Dairy Science*, *98*, 211–220.
- Lavers, C., Barkema, H. W., Dohoo, I. R., McKenna, S. L. B., & Keefe, G. P. (2014). Evaluation of milk ELISA for detection of *Mycobacterium avium* subspecies *paratuberculosis* in dairy herds and association with within-herd prevalence. *Journal of Dairy Science*, *97*, 299–309.
- Leite, F. L., Reinhardt, T. A., Bannantine, J. P., & Stabel, J. R. (2015). Envelope protein complexes of *Mycobacterium avium* subsp. *paratuberculosis* and their antigenicity. *Veterinary Microbiology*, *175*, 275–285.
- Lu, Z., Mitchell, R. M., Smith, R. L., Van Kessel, J. S., Chapagain, P. P., Schukken, Y. H., & Gröhn, Y. T. (2008). The importance of culling in Johne's disease control. *Journal of Theoretical Biology*, *254*, 135–146.
- Lugton, I. W. (2004). Review of possible links between the clinical expression of paratuberculosis and deficiency of macro and micronutrients. *Australian Veterinary Journal*, *82*, 490–496.
- Lund, B. M., Gould, G. W., & Rampling, A. M. (2002). Pasteurization of milk and the heat resistance of *Mycobacterium avium* subsp. *paratuberculosis*: a critical review of the data. *The International Journal of Food Microbiology*, *77*, 135–145.
- Lybeck, K. R., Storset, A. K., Dønne, B., Valheim, M., & Olsen, I. (2011). Faecal shedding detected earlier than immune responses in goats naturally infected with *Mycobacterium avium* subsp. *paratuberculosis*. *Research in Veterinary Science*, *91*, 32–39.
- Machackova, M., Svastova, P., Lamka, J., Parmova, I., Liska, V., Smolik, J., ... Pavlik, I. (2004). Paratuberculosis in farmed and free-living wild ruminants in the Czech Republic (1999–2001). *Veterinary Microbiology*, *101*, 225–234.
- Mackintosh, C. G., & Griffin, J. F. (2010). Paratuberculosis in deer, camelids and other ruminants. In: M. A. Behr & D. M. Collins, (Eds.), *Paratuberculosis: organism, disease, control* (pp. 179–187). Wallingford, Oxfordshire, UK: CABI. ISBN 9781845936136.
- Marcé, C., Ezanno, P., Seegers, H., Pfeiffer, D. U., & Fourichon, C. (2011). Within-herd contact structure and transmission of *Mycobacterium avium* subspecies *paratuberculosis* in a persistently infected dairy cattle herd. *Preventive Veterinary Medicine*, *100*, 116–125.
- Marcé, C., Ezanno, P., Weber, M. F., Seegers, H., Pfeiffer, D. U., & Fourichon, C. (2010). Invited review: modeling within-herd transmission of *Mycobacterium avium* subspecies *paratuberculosis* in dairy cattle: a review. *Journal of Dairy Science*, *93*, 4455–4470.
- McAloon, C. G., Doherty, M. L., Whyte, P., O'Grady, L., More, S. J., Messam, L. L., ... Green, M. J. (2016). Bayesian estimation of prevalence of paratuberculosis in dairy herds enrolled in a voluntary Johne's Disease Control Programme in Ireland. *Preventive Veterinary Medicine*, *128*, 95–100.
- McGregor, H., Dhand, N. K., Dhungyel, O. P., & Whittington, R. J. (2012). Transmission of *Mycobacterium avium* subsp. *paratuberculosis*: Dose-response and age-based susceptibility in a sheep model. *Preventive Veterinary Medicine*, *107*, 76–84.
- McKenna, S. L. B., Keefe, G. P., Barkema, H. W., & Sockett, D. C. (2005). Evaluation of three ELISAs for *Mycobacterium avium* subsp. *paratuberculosis* using tissue and fecal culture as comparison standards. *Veterinary Microbiology*, *110*, 105–111.
- McKenna, S. L., Keefe, G. P., Tiwari, A., VanLeeuwen, J., & Barkema, H. W. (2006). Johne's disease in Canada part II: disease impacts, risk factors, and control programs for dairy producers. *Canadian Veterinary Journal*, *47*, 1089–1099.
- McKenna, S. L. B., Sockett, D. C., Keefe, G. P., McClure, J., VanLeeuwen, J. A., & Barkema, H. W. (2005). Comparison of two enzyme-linked immunosorbent assays for diagnosis of *Mycobacterium avium* subsp. *paratuberculosis*. *Journal of Veterinary Diagnostic Investigation*, *17*, 463–466.
- McKenna, S. L. B., Vanleeuwen, J. A., Barkema, H. W., Jansen, J. T., Hauer, G., Hendrick, S. H., ... Empringham, R. E. (2006). Proposed Canadian voluntary national Johne's Disease prevention and control program. *Canadian Veterinary Journal*, *47*, 539–541.
- Mikkelsen, H., Aagaard, C., Nielsen, S. S., & Jungersen, G. (2012). Correlation of antigen-specific IFN- γ responses of fresh blood samples from *Mycobacterium avium* subsp. *paratuberculosis* infected heifers with responses of day-old samples co-cultured with IL-12 or anti-IL-10 antibodies. *Veterinary Immunology and Immunopathology*, *147*, 69–76.
- Minozzi, G., Williams, J. L., Stella, A., Strozzi, F., Luini, M., Settles, M. L., ... Neiberghs, H. L. (2012). Meta-analysis of two genome-wide association studies of bovine paratuberculosis. *PLoS ONE*, *7*, e32578.
- Mita, A., Mori, Y., Nakagawa, T., Tasaki, T., Utiyama, K., & Mori, H. (2016). Comparison of fecal pooling methods and DNA extraction kits for the detection of *Mycobacterium avium* subspecies *paratuberculosis*. *Microbiologyopen*, *5*, 134–142.
- Mitchell, R. M., Schukken, Y., Koets, A., Weber, M., Bakker, D., Stabel, J., ... Louzoun, Y. (2015). Differences in intermittent and continuous fecal shedding patterns between natural and experimental *Mycobacterium avium* subspecies *paratuberculosis* infections in cattle. *Veterinary Research*, *46*, 66.
- Mortier, R. A. R., Barkema, H. W., Bystrom, J. M., Illanes, O., Orsel, K., Wolf, R., ... De Buck, J. (2013). Evaluation of age-dependent susceptibility in calves infected with two doses of *Mycobacterium avium* subspecies *paratuberculosis* using pathology and tissue culture. *Veterinary Research*, *44*, 94.

- Mortier, R. A. R., Barkema, H. W., & De Buck, J. (2015). Susceptibility to and diagnosis of *Mycobacterium avium* subspecies *paratuberculosis* infection in dairy calves: A review. *Preventive Veterinary Medicine*, *121*, 189–198.
- Mortier, R. A. R., Barkema, H. W., Orsel, K., Muench, G. P., Bystrom, J., & De Buck, J. (2015). Longitudinal evaluation of diagnostics in experimentally infected young calves during subclinical and clinical paratuberculosis. *Canadian Veterinary Journal*, *56*, 1266–1270.
- Mortier, R. A. R., Barkema, H. W., Orsel, K., Wolf, R., & De Buck, J. (2014). Shedding patterns of dairy calves experimentally infected with *Mycobacterium avium* subspecies *paratuberculosis*. *Veterinary Research*, *45*, 71.
- Mortier, R. A. R., Barkema, H. W., Orsel, K., Wolf, R., Negron, M. E., Atkins, G. A., & De Buck, J. (2014). Antibody response early after experimental infection with *Mycobacterium avium* subspecies *paratuberculosis* in dairy calves. *Journal of Dairy Science*, *97*, 5558–5565.
- Mortier, R. A. R., Barkema, H. W., Wilson, T. A., Sajobi, T. T., Wolf, R., & De Buck, J. (2014). Dose-dependent interferon-gamma release in dairy calves experimentally infected with *Mycobacterium avium* subspecies *paratuberculosis*. *Veterinary Immunology and Immunopathology*, *161*, 205–210.
- Motiwala, A. S., Amonsin, A., Strother, M., Manning, E. J., Kapur, V., & Sreevatsan, S. (2004). Molecular epidemiology of *Mycobacterium avium* subsp. *paratuberculosis* isolates recovered from wild animal species. *Journal of Clinical Microbiology*, *42*, 1703–1712.
- Motiwala, A. S., Janagama, H. K., Paustian, M. L., Zhu, X., Bannantine, J. P., Kapur, V., & Sreevatsan, S. (2006). Comparative transcriptional analysis of human macrophages exposed to animal and human isolates of *Mycobacterium avium* subspecies *paratuberculosis* with diverse genotypes. *Infection and Immunity*, *74*, 6046–6056.
- Motiwala, A. S., Strother, M., Amonsin, A., Byrum, B., Naser, S. A., Stabel, J. R., ... Sreevatsan, S. (2003). Molecular epidemiology of *Mycobacterium avium* subsp. *paratuberculosis*: evidence for limited strain diversity, strain sharing, and identification of unique targets for diagnosis. *Journal of Clinical Microbiology*, *41*, 2015–2026.
- Muskens, J., van Zijderveld, F., Eger, A., & Bakker, D. (2002). Evaluation of the long-term immune response in cattle after vaccination against paratuberculosis in two Dutch dairy herds. *Veterinary Microbiology*, *86*, 269–278.
- Nagel-Alne, G. E., Asheim, L. J., Hardaker, J. B., Sølverød, L., Lindheim, D., & Valle, P. S. (2014). The Norwegian Healthier Goats programme—a financial cost-benefit analysis. *Preventive Veterinary Medicine*, *114*, 96–105.
- Nielsen, S. S. (2008). Transitions in diagnostic tests used for detection of *Mycobacterium avium* subsp. *paratuberculosis* infections in cattle. *Veterinary Microbiology*, *132*, 274–282.
- Nielsen, S. S. (2011). Dairy farmers' reasons for participation in the Danish control programme on bovine paratuberculosis. *Preventive Veterinary Medicine*, *98*, 279–283.
- Nielsen, S. S., Bjerre, H., & Toft, N. (2008). Colostrum and milk as risk factors for infection with *Mycobacterium avium* subspecies *paratuberculosis* in dairy cattle. *Journal of Dairy Science*, *91*, 4610–4615.
- Nielsen, S. S., Jepsen, O. R., & Aagaard, K. (2007). Control programme for paratuberculosis in Denmark. *Bull. Intern. Dairy Fed.* 410/207: Proc. 1st ParaTB Forum, pp. 23–29.
- Nielsen, S. S., & Toft, N. (2008). Ante mortem diagnosis of paratuberculosis: a review of accuracies of ELISA, interferon-gamma assay and faecal culture techniques. *Veterinary Microbiology*, *129*, 217–235.
- Nielsen, S. S., & Toft, N. (2011). Effect of management practices on paratuberculosis prevalence in Danish dairy herds. *Journal of Dairy Science*, *94*, 1849–1857.
- Nielsen, S. S., & Toft, N. (2014). Bulk tank milk ELISA for detection of antibodies to *Mycobacterium avium* subsp. *paratuberculosis*: correlation between repeated tests and within-herd antibody-prevalence. *Preventive Veterinary Medicine*, *113*, 96–102.
- Norton, S., Heuer, C., & Jackson, R. (2009). A questionnaire-based cross-sectional study of clinical Johne's disease on dairy farms in New Zealand. *The New Zealand Veterinary Journal*, *57*, 34–43.
- Oakey, J., Gavey, L., Singh, S. V., Platell, J., & Waltisbuhl, D. (2014). Variable-number tandem repeats genotyping used to aid and inform management strategies for a bovine Johne's disease incursion in tropical and subtropical Australia. *Journal of Veterinary Diagnostic Investigation*, *26*, 651–657.
- OIE (2012). *Validation and certification of diagnostic assays*. Paris, France: World Organisation for Animal Health. Retrieved from <http://www.oie.int/en/our-scientific-expertise/registration-of-diagnostic-kits/background-information/> (Accessed 9 May 2017).
- Ott, S. L., Wells, S. J., & Wagner, B. A. (1999). Herd-level economic losses associated with Johne's disease on US dairy operations. *Preventive Veterinary Medicine*, *40*, 179–192.
- Park, K. T., Allen, A. J., Bannantine, J. P., Seo, K. S., Hamilton, M. J., Abdellrazeq, G. S., ... Davis, W. C. (2011). Evaluation of two mutants of *Mycobacterium avium* subsp. *paratuberculosis* as candidates for a live attenuated vaccine for Johne's disease. *Vaccine*, *29*, 4709–4719.
- Park, K. T., Allen, A. J., Barrington, G. M., & Davis, W. C. (2014). Deletion of *relA* abrogates the capacity of *Mycobacterium avium paratuberculosis* to establish an infection in calves. *Frontiers in Cellular and Infection Microbiology*, *4*, 64.
- Pearson, T., Okinaka, R. T., Foster, J. T., & Keim, P. (2009). Phylogenetic understanding of clonal populations in an era of whole genome sequencing. *Infection, Genetics and Evolution*, *9*, 1010–1019.
- Pillars, R. D., Grooms, D. L., Gardiner, D. C., & Kaneene, J. B. (2011). Association between risk-assessment scores and individual-cow Johne's disease-test status over time on seven Michigan, USA dairy herds. *Preventive Veterinary Medicine*, *98*, 10–18.
- Pinheiro, L. B., Coleman, V. A., Hindson, C. M., Herrmann, J., Hindson, B. J., Bhat, S., & Emslie, K. R. (2012). Evaluation of a droplet digital polymerase chain reaction format for DNA copy number quantification. *Analytical Chemistry*, *84*, 1003–1011.
- Plain, K. M., Begg, D. J., de Silva, K., Purdie, A. C., & Whittington, R. J. (2012). Enhancement of the interferon gamma assay to detect paratuberculosis using interleukin-7 and interleukin-12 potentiation. *Veterinary Immunology and Immunopathology*, *149*, 28–37.
- Plain, K. M., Marsh, I. B., Waldron, A. M., Galea, F., Whittington, A. M., Saunders, V. F., ... Whittington, R. J. (2014). High-throughput direct fecal PCR assay for detection of *Mycobacterium avium* subsp. *paratuberculosis* in sheep and cattle. *Journal of Clinical Microbiology*, *52*, 745–757.
- Pradhan, A. K., Mitchell, R. M., Kramer, A. J., Zurakowski, M. J., Fyock, T. L., Whitlock, R. H., ... Schukken, Y. H. (2011). Molecular epidemiology of *Mycobacterium avium* subsp. *paratuberculosis* in a longitudinal study of three dairy herds. *Journal of Clinical Microbiology*, *49*, 893–901.
- Pritchard, T. C., Coffey, M. P., Bond, K. S., Hutchings, M. R., & Wall, E. (2017). Phenotypic effects of subclinical paratuberculosis (Johne's disease) in dairy cattle. *Journal of Dairy Science*, *100*, 679–690.
- Pruvot, M., Forde, T. L., Steele, J., Kutz, S. J., De Buck, J., van der Meer, F., & Orsel, K. (2013). The modification and evaluation of an ELISA test for the surveillance of *Mycobacterium avium* subsp. *paratuberculosis* infection in wild ruminants. *BMC Veterinary Research*, *9*, 5.
- Purdie, A. C., Plain, K. M., Begg, D. J., de Silva, K., & Whittington, R. J. (2011). Candidate gene and genome-wide association studies of *Mycobacterium avium* subsp. *paratuberculosis* infection in cattle and sheep: a review. *Comparative Immunology, Microbiology and Infectious Diseases*, *34*, 197–208.
- Raizman, E. A., Fetrow, J. P., & Wells, S. J. (2009). Loss of income from cows shedding *Mycobacterium avium* subspecies *paratuberculosis* prior to calving compared with cows not shedding the organism on two Minnesota dairy farms. *Journal of Dairy Science*, *92*, 4929–4936.
- Rana, A., Rub, A., & Akhter, Y. (2015). Proteome-wide B and T cell epitope repertoires in outer membrane proteins of *Mycobacterium avium*

- subsp. *paratuberculosis* have vaccine and diagnostic relevance: a holistic approach. *Journal of Molecular Recognition*, 28, 506–520.
- Rangel, S. J., Paré, J., Doré, E., Arango, J. C., Côté, G., Buczinski, S., ... Fecteau, G. (2005). A systematic review of risk factors associated with the introduction of *Mycobacterium avium* spp. *paratuberculosis* (MAP) into dairy herds. *Canadian Veterinary Journal*, 56, 169–177.
- Rathnaiah, G., Lamont, E. A., Harris, N. B., Fenton, R. J., Zinniel, D. K., Liu, X., ... Barletta, R. G. (2014). Generation and screening of a comprehensive *Mycobacterium avium* subsp. *paratuberculosis* transposon mutant bank. *Frontiers in Cellular and Infection Microbiology*, 4, 144.
- Reddacliff, L., Eppleston, J., Windsor, P., Whittington, R., & Jones, S. (2006). Efficacy of a killed vaccine for the control of paratuberculosis in Australian sheep flocks. *Veterinary Microbiology*, 115, 77–90.
- Ritter, C., Jansen, J., Roche, S., Kelton, D. F., Adams, C., Orsel, K., ... Barkema, H. W. (2017). Invited review: Determinants of farmers' adoption of management-based strategies for infectious disease prevention and control. *Journal of Dairy Science*, 100, 3329–3347.
- Ritter, C., Kwong, G. P., Wolf, R., Pickel, C., Slomp, M., Flaig, J., ... Barkema, H. W. (2015). Factors associated with participation of Alberta dairy farmers in a voluntary, management-based Johne's disease control program. *Journal of Dairy Science*, 98, 7831–7845.
- Roche, S. M. R. (2014). *Investigating the role of agricultural extension in influencing Ontario dairy producer behaviour for Johne's disease control*. PhD Thesis. Univ. Guelph, Guelph, Ontario, Canada.
- Roche, S. M., Jones-Bitton, A., Meehan, M., Von Massow, M., & Kelton, D. F. (2015). Evaluating the effect of Focus Farms on Ontario dairy producers' knowledge, attitudes, and behavior toward control of Johne's disease. *Journal of Dairy Science*, 98, 5222–5240.
- Romano, M., & Huygen, K. (2009). DNA vaccines against mycobacterial diseases. *Expert Review of Vaccines*, 8, 1237–1250.
- Rosseels, V., & Huygen, K. (2008). Vaccination against paratuberculosis. *Expert Review of Vaccines*, 7, 817–832.
- Rossiter, C. A., & Burhans, W. S. (1996). Farm-specific approach to paratuberculosis (Johne's disease) control. *Vet. Clin. North Am. Food Animal Practice*, 12, 383–415.
- Roupie, V., Viart, S., Leroy, B., Romano, M., Trincherio, N., Govaerts, M., ... Huygen, K. (2012). Immunogenicity of eight *Mycobacterium avium* subsp. *paratuberculosis* specific antigens in DNA vaccinated and Map infected mice. *Veterinary Immunology and Immunopathology*, 145, 74–85.
- Roussel, A. J., Libal, M. C., Whitlock, R. L., Hairgrove, T. B., Barling, K. S., & Thompson, J. A. (2005). Prevalence of and risk factors for paratuberculosis in purebred beef cattle. *Journal of the American Veterinary Medical Association*, 226, 773–778.
- Russell, R. A., & Bewley, J. M. (2011). Producer assessment of dairy extension programming in Kentucky. *Journal of Dairy Science*, 94, 2637–2647.
- Santema, W. J., Hensen, S., Rutten, V. P., & Koets, A. P. (2009). Heat shock protein 70 subunit vaccination against bovine paratuberculosis does not interfere with current immunodiagnostic assays for bovine tuberculosis. *Vaccine*, 27, 2312–2319.
- Santema, W. J., Poot, J., Segers, R. P., Van den Hoff, D. J., Rutten, V. P., & Koets, A. P. (2012). Early infection dynamics after experimental challenge with *Mycobacterium avium* subspecies *paratuberculosis* in calves reveal limited calf-to-calf transmission and no impact of Hsp70 vaccination. *Vaccine*, 30, 7032–7039.
- Santema, W., Rutten, V., Segers, R., Poot, J., Hensen, S., Heesterbeek, H., & Koets, A. (2013). Postexposure subunit vaccination against chronic enteric mycobacterial infection in a natural host. *Infection and Immunity*, 81, 1990–1995.
- Schillinger, S., Bridger, P. S., Bulun, H., Fischer, M., Akineden, O., Seeger, T., ... Menge, C. (2013). Flow cytometric detection of *Mycobacterium avium* subsp. *paratuberculosis*-specific antibodies in experimentally infected and naturally exposed calves. *Clinical and Vaccine Immunology*, 20, 1457–1465.
- Sergeant, E. S., Whittington, R. J., & More, S. J. (2002). Sensitivity and specificity of pooled faecal culture and serology as flock-screening tests for detection of ovine paratuberculosis in Australia. *Preventive Veterinary Medicine*, 52, 199–211.
- Serraino, A., Arrigoni, N., Ostanello, F., Ricchi, M., Marchetti, G., Bonilauri, P., ... Giacometti, F. (2014). A screening sampling plan to detect *Mycobacterium avium* subspecies *paratuberculosis*-positive dairy herds. *Journal of Dairy Science*, 97, 3344–3351.
- Seth, M., Lamont, E. A., Janagama, H. K., Widdel, A., Vulchanova, L., Stabel, J. R., ... Sreevatsan, S. (2009). Biomarker discovery in subclinical mycobacterial infections of cattle. *PLoS ONE*, 4, e5478.
- Settles, E. W., Kink, J. A., & Talaat, A. (2014). Attenuated strains of *Mycobacterium avium* subspecies *paratuberculosis* as vaccine candidates against Johne's disease. *Vaccine*, 32, 2062–2069.
- Settles, M., Zanella, R., McKay, S. D., Schnabel, R. D., Taylor, J. F., Whitlock, R., ... Neiberghs, H. (2009). A whole genome association analysis identifies loci associated with *Mycobacterium avium* subsp. *paratuberculosis* infection status in US Holstein cattle. *Animal Genetics*, 40, 655–662.
- Shaughnessy, R. G., Farrell, D., Riepema, K., Bakker, D., & Gordon, S. V. (2015). Analysis of biobanked serum from a *Mycobacterium avium* subsp. *paratuberculosis* bovine infection model confirms the remarkable stability of circulating miRNA profiles and defines a bovine serum miRNA repertoire. *PLoS ONE*, 10, e0145089.
- Shin, M. K., Park, H. T., Shin, S. W., Jung, M., Im, Y. B., Park, H. E., ... Yoo, H. S. (2015). Whole-blood gene-expression profiles of cows infected with *Mycobacterium avium* subsp. *paratuberculosis* reveal changes in immune response and lipid metabolism. *Journal of Microbiology and Biotechnology*, 25, 255–267.
- Shippy, D. C., Lemke, J. J., Berry, A., Nelson, K., Hines, M. E. 2nd, & Talaat, A. M. (2017). Superior protection from live-attenuated vaccines directed against Johne's disease. *Clinical and Vaccine Immunology*, 24, e00478–16.
- Shook, G. E., Chaffer, M., Wu, X. L., & Ezra, E. (2012). Genetic parameters for paratuberculosis infection and effect of infection on production traits in Israeli Holsteins. *Animal Genetics*, 43(Suppl 1), 56–64.
- de Silva, K., Begg, D. J., Plain, K. M., Purdie, A. C., Kawaji, S., Dhand, N. K., & Whittington, R. J. (2013). Can early host responses to mycobacterial infection predict eventual disease outcomes? *Preventive Veterinary Medicine*, 112, 203–212.
- de Silva, K., Plain, K. M., Begg, D. J., Purdie, A. C., & Whittington, R. J. (2015). CD4(+) T-cells, gammadelta T-cells and B-cells are associated with lack of vaccine protection in *Mycobacterium avium* subspecies *paratuberculosis* infection. *Vaccine*, 33, 149–155.
- Singh, K., Chandel, B. S., Chauhan, H. C., Dadawala, A., Singh, S. V., & Singh, P. K. (2013). Efficacy of 'indigenous vaccine' using native 'Indian bison type' genotype of *Mycobacterium avium* subspecies *paratuberculosis* for the control of clinical Johne's disease in an organized goat herd. *Veterinary Research Communications*, 37, 109–114.
- Skovgaard, K., Grell, S. N., Heegaard, P. M., Jungersen, G., Pudrith, C. B., & Coussens, P. M. (2006). Differential expression of genes encoding CD30L and P-selectin in cattle with Johne's disease: progress toward a diagnostic gene expression signature. *Veterinary Immunology and Immunopathology*, 112, 210–224.
- Slana, I., Pribylova, R., Kralova, A., & Pavlik, I. (2011). Persistence of *Mycobacterium avium* subsp. *paratuberculosis* at a farm-scale biogas plant supplied with manure from paratuberculosis-affected dairy cattle. *Applied and Environment Microbiology*, 77, 3115–3119.
- Sohal, J. S., Arsenault, J., Labrecque, O., Fairbrother, J.-H., Roy, J.-P., Fecteau, G., & L'Homme, Y. (2014). Genetic structure of *Mycobacterium avium* subsp. *paratuberculosis* population in cattle herds in Quebec as revealed by using a combination of multilocus genomic analyses. *Journal of Clinical Microbiology*, 52, 2764–2775.
- Sohal, J. S., Singh, S. V., Singh, A. V., & Singh, P. K. (2010). Strain diversity within *Mycobacterium avium* subspecies *paratuberculosis*-a review. *Indian Journal of Experimental Biology*, 48, 7–16.

- Sorge, U., Kelton, D., Lissemore, K., Godkin, A., Hendrick, S., & Wells, S. (2010). Attitudes of Canadian dairy farmers toward a voluntary Johne's disease control program. *Journal of Dairy Science*, *93*, 1491–1499.
- Sorge, U. S., Lissemore, K., Godkin, A., Hendrick, S., Wells, S., & Kelton, D. (2011). Associations between paratuberculosis milk ELISA result, milk production, and breed in Canadian dairy cows. *Journal of Dairy Science*, *94*, 754–761.
- Stabel, J. R. (2000). Transitions in immune responses to *Mycobacterium paratuberculosis*. *Veterinary Microbiology*, *77*, 465–473.
- Stabel, J. R. (2006). Host responses to *Mycobacterium avium* subsp. *paratuberculosis*: a complex arsenal. *Animal Health Research Reviews*, *7*, 61–70.
- Stabel, J. R., Barnhill, A., Bannantine, J. P., Chang, Y. F., & Osman, M. A. (2012). Evaluation of protection in a mouse model after vaccination with *Mycobacterium avium* subsp. *paratuberculosis* protein cocktails. *Vaccine*, *31*, 127–134.
- Stabel, J. R., & Goff, J. P. (2004). Efficacy of immunologic assays for the detection of Johne's disease in dairy cows fed additional energy during the periparturient period. *Journal of Veterinary Diagnostic Investigation*, *16*, 412–420.
- Stabel, J. R., & Robbe-Austerman, S. (2011). Early immune markers associated with *Mycobacterium avium* subsp. *paratuberculosis* infection in a neonatal calf model. *Clinical and Vaccine Immunology*, *18*, 393–405.
- Stevenson, K., Alvarez, J., Bakker, D., Biet, F., de Juan, L., Denham, S., ... Greig, A. (2009). Occurrence of *Mycobacterium avium* subspecies *paratuberculosis* across host species and European countries with evidence for transmission between wildlife and domestic ruminants. *BMC Microbiology*, *9*, 212.
- Storset, A. K., Hasvold, H. J., Valheim, M., Brun-Hansen, H., Berntsen, G., Whist, S. K., ... Larsen, H. J. (2001). Subclinical paratuberculosis in goats following experimental infection. An immunological and microbiological study. *Veterinary Immunology and Immunopathology*, *80*, 271–287.
- Stratilo, C. W., Lewis, C. T., Bryden, L., Mulvey, M. R., & Bader, D. (2006). Single-nucleotide repeat analysis for subtyping *Bacillus anthracis* isolates. *Journal of Clinical Microbiology*, *44*, 777–782.
- Streeter, R. N., Hoffsis, G. F., Bech-Nielsen, S., Shulaw, W. P., & Rings, D. M. (1995). Isolation of *Mycobacterium paratuberculosis* from colostrum and milk of subclinically infected cows. *American Journal of Veterinary Research*, *56*, 1322–1324.
- Sweeney, R. W. (1996). Transmission of paratuberculosis. *Veterinary Clinics of North America: Food Animal Practice*, *12*, 305–312.
- Sweeney, R. W., Whitlock, R. H., & Rosenberger, A. E. (1992). *Mycobacterium paratuberculosis* cultured from milk and supramammary lymph nodes of infected asymptomatic cows. *Journal of Clinical Microbiology*, *30*, 166–171.
- Thakur, A., Aagaard, C., Stockmarr, A., Andersen, P., & Jungersen, G. (2013). Cell-mediated and humoral immune responses after immunization of calves with a recombinant multiantigenic *Mycobacterium avium* subsp. *paratuberculosis* subunit vaccine at different ages. *Clinical and Vaccine Immunology*, *20*, 551–558.
- Thibault, V. C., Grayon, M., Boschiroli, M. L., Hubbans, C., Overduin, P., Stevenson, K., ... Biet, F. (2007). New variable-number tandem-repeat markers for typing *Mycobacterium avium* subsp. *paratuberculosis* and *M. avium* strains: comparison with IS900 and IS1245 restriction fragment length polymorphism typing. *Journal of Clinical Microbiology*, *45*, 2404–2410.
- Thibault, V. C., Grayon, M., Boschiroli, M. L., Willery, E., Allix-Béguec, C., Stevenson, K., ... Supply, P. (2008). Combined multilocus short-sequence-repeat and mycobacterial interspersed repetitive unit-variable-number tandem-repeat typing of *Mycobacterium avium* subsp. *paratuberculosis* isolates. *Journal of Clinical Microbiology*, *46*, 4091–4094.
- Thirunavukkarasu, S., Plain, K. M., de Silva, K., Begg, D., Whittington, R. J., & Purdie, A. C. (2014). Expression of genes associated with cholesterol and lipid metabolism identified as a novel pathway in the early pathogenesis of *Mycobacterium avium* subspecies *paratuberculosis*-infection in cattle. *Veterinary Immunology and Immunopathology*, *160*, 147–157.
- Thompson, B. R., Clark, R. G., & Mackintosh, C. G. (2007). Intra-uterine transmission of *Mycobacterium avium* subsp. *paratuberculosis* in subclinically affected red deer (*Cervus elaphus*). *The New Zealand Veterinary Journal*, *55*, 308–313.
- Tiwari, A., VanLeeuwen, J. A., Dohoo, I. R., Keefe, G. P., & Weersink, A. (2008). Estimate of the direct production losses in Canadian dairy herds with subclinical *Mycobacterium avium* subspecies *paratuberculosis* infection. *Canadian Veterinary Journal*, *49*, 569–576.
- Tiwari, A., VanLeeuwen, J. A., McKenna, S. L. B., Keefe, G. P., & Barkema, H. W. (2006). Johne's disease in Canada Part I: clinical symptoms, pathophysiology, diagnosis, and prevalence in dairy herds. *Canadian Veterinary Journal*, *47*, 874–882.
- Trefz, P., Koehler, H., Klepik, K., Moebius, P., Reinhold, P., Schubert, J. K., & Miekisch, W. (2013). Volatile emissions from *Mycobacterium avium* subsp. *paratuberculosis* mirror bacterial growth and enable distinction of different strains. *PLoS ONE*, *8*, e76868.
- Trier, L., Nielsen, S. S., & Krogh, K. (2012). Motivation of farmers to reduce between-herd transmission of paratuberculosis through repeated on-farm meetings with discussion on certification. Page 318 in 11th Proc. International Colloquium on Paratuberculosis Sydney, Australia.
- Van Campen, H., & Rhyan, J. (2010). The role of wildlife in diseases of cattle. *Veterinary Clinics of North America: Food Animal Practice*, *26*, 147–161.
- Van Kooten, H. C., Mackintosh, C. G., & Koets, A. P. (2006). Intra-uterine transmission of paratuberculosis (Johne's disease) in farmed red deer. *The New Zealand Veterinary Journal*, *54*, 16–20.
- Van Roermund, H. J. W., Bakker, D., Willemsen, P. T. J., & de Jong, M. C. M. (2007). Horizontal transmission of *Mycobacterium avium* subsp. *paratuberculosis* in cattle in an experimental setting: Calves can transmit the infection to other calves. *Veterinary Microbiology*, *122*, 270–279.
- Van Schaik, G., Pradenas, F. M., Mella, N. A., & Kruze, V. J. (2007). Diagnostic validity and costs of pooled fecal samples and individual blood or fecal samples to determine the cow- and herd-status for *Mycobacterium avium* subsp. *paratuberculosis*. *Preventive Veterinary Medicine*, *82*, 159–165.
- Van Weering, H., van Schaik, G., van der Meulen, A., Waal, M., Franken, P., & van Maanen, K. (2007). Diagnostic performance of the Pourquier ELISA for detection of antibodies against *Mycobacterium avium* subspecies *paratuberculosis* in individual milk and bulk milk samples of dairy herds. *Veterinary Microbiology*, *125*, 49–58.
- Velthuis, A. G. J., Weber, M. F., deKoeijer, A. A., & Van Roermund, H. J. W. (2006). A milk quality assurance program for Johne's disease: decision analysis from a farmer's perspective. Proc. 11th Symp. Intern. Soc. Vet. Epidemiol. Econ., Cairns, Australia, p. 313.
- Verdugo, C., Jones, G., Johnson, W., Wilson, P., Stringer, L., & Heuer, C. (2014). Estimation of flock/herd-level true *Mycobacterium avium* subspecies *paratuberculosis* prevalence on sheep, beef cattle and deer farms in New Zealand using a novel Bayesian model. *Preventive Veterinary Medicine*, *117*, 447–455.
- Verdugo, C., Pleydell, E., Price-Carter, M., Prattley, D., Collins, D., de Lisle, G., ... Heuer, C. (2014). Molecular epidemiology of *Mycobacterium avium* subsp. *paratuberculosis* isolated from sheep, cattle and deer on New Zealand pastoral farms. *Preventive Veterinary Medicine*, *117*, 436–446.
- Waddell, L. A., Rajic, A., Stärk, K. D. C., & McEwen, S. A. (2015). The zoonotic potential of *Mycobacterium avium* ssp. *paratuberculosis*: A systematic review and meta-analyses of the evidence. *Epidemiology and Infection*, *143*, 3135–3157.
- Waddell, L. A., Rajic, A., Stärk, K. D. C., & McEwen, S. A. (2016). The potential public health impact of *Mycobacterium avium* ssp. *paratuberculosis*: global opinion survey of topic specialists. *Zoonoses Public Health*, *63*, 212–222.

- Wang, J., Pritchard, J. R., Kreitmann, L., Montpetit, A., & Behr, M. A. (2014). Disruption of *Mycobacterium avium* subsp. *paratuberculosis*-specific genes impairs *in vivo* fitness. *BMC Genomics*, 15, 415.
- Weber, M. F., Groenendaal, H., van Roermund, H. J., & Nielen, M. (2004). Simulation of alternatives for the Dutch Johne's disease certification and-monitoring program. *Preventive Veterinary Medicine*, 62, 1–17.
- Weber, M. F., Kogut, J., deBree, J., & Van Schaik, G. (2006). *Survival analysis of age at onset of shedding of Mycobacterium avium ssp. paratuberculosis*. In: Proc. 11th Symp Int Soc Vet Epidemiol Econ, Cairns, Australia, p. 444.
- Weber, M. F., van Roermund, H. J., Vernooij, J. C., Kalis, C. H., & Stegeman, J. A. (2006). Cattle transfers between herds under paratuberculosis surveillance in The Netherlands are not random. *Preventive Veterinary Medicine*, 76, 222–236.
- Weber, M. F., & van Schaik, G. (2007). *Results of the Dutch milk quality assurance programme for paratuberculosis*. In: Proc. 8th Intern. Colloq. Paratuberculosis, pp. 325–327.
- Wells, S. J., Hartmann, W. L., & Anderson, P. L. (2008). Evaluation of progress made by dairy and beef herds enrolled in the Minnesota Johne's Disease Control Program. *Journal of the American Veterinary Medical Association*, 233, 1920–1926.
- Wells, S. J., & Wagner, B. A. (2000). Herd-level risk factors for infection with *Mycobacterium paratuberculosis* in US dairies and association between familiarity of the herd manager with the disease or prior diagnosis of the disease in that herd and use of preventive measures. *Journal of the American Veterinary Medical Association*, 216, 1450–1457.
- Wentink, G. H., Bongers, J. H., Zeeuwen, A. A., & Jaartsveld, F. H. (1994). Incidence of paratuberculosis after vaccination against *M. paratuberculosis* in two infected dairy herds. *Zentralbl Veterinarmed B*, 41, 517–522.
- Whipple, D., Kapke, P., & Vary, C. (1990). Identification of restriction fragment length polymorphisms in DNA from *Mycobacterium paratuberculosis*. *Journal of Clinical Microbiology*, 28, 2561–2564.
- Whist, A. C., Liland, K. H., Jonsson, M. E., Sæbø, S., Sviland, S., Østeras, O., ... Hopp, P. (2014). Designing a risk-based surveillance program for *Mycobacterium avium* ssp. *paratuberculosis* in Norwegian dairy herds using multivariate statistical process control analysis. *Journal of Dairy Science*, 97, 6835–6849.
- Whitlock, R. H., & Buergelt, C. (1996). Preclinical and clinical manifestations of paratuberculosis (including pathology). *Veterinary Clinics of North America: Food Animal Practice*, 12, 345–356.
- Whitlock, R. H., Wells, S. J., Sweeney, R. W., & Van Tiem, J. (2000). ELISA and fecal culture for paratuberculosis (Johne's disease): sensitivity and specificity of each method. *Veterinary Microbiology*, 77, 387–398.
- Whittington, R. J., Fell, S., Walker, D., McAllister, S., Marsh, I., Sergeant, E., ... Links, I. J. (2000). Use of pooled fecal culture for sensitive and economic detection of *Mycobacterium avium* subsp. *paratuberculosis* infection in flocks of sheep. *Journal of Clinical Microbiology*, 38, 2550–2556.
- Whittington, R. J., Hope, A. F., Marshall, D. J., Taragel, C. A., & Marsh, I. (2000). Molecular epidemiology of *Mycobacterium avium* subsp. *paratuberculosis*: IS900 restriction fragment length polymorphism and IS1311 polymorphism analyses of isolates from animals and a human in Australia. *Journal of Clinical Microbiology*, 38, 3240–3248.
- Whittington, R., Marsh, I., Choy, E., & Cousins, D. (1998). Polymorphisms in IS1311, an insertion sequence common to *Mycobacterium avium* and *M. avium* subsp. *paratuberculosis*, can be used to distinguish between and within these species. *Molecular and Cellular Probes*, 12, 349–358.
- Whittington, R. J., Marsh, I. B., Saunders, V., Grant, I. R., Juste, R., Sevilla, I. A., ... Whitlock, R. H. (2011). Culture phenotypes of genomically and geographically diverse *Mycobacterium avium* subsp. *paratuberculosis* isolates from different hosts. *Journal of Clinical Microbiology*, 49, 1822–1830.
- Whittington, R. J., Marsh, I. B., Taylor, P. J., Marshall, D. J., Taragel, C., & Reddacliff, L. A. (2003). Isolation of *Mycobacterium avium* subsp. *paratuberculosis* from environmental samples collected from farms before and after destocking sheep with paratuberculosis. *Australian Veterinary Journal*, 81, 559–563.
- Whittington, R. J., Marsh, I. B., & Whitlock, R. H. (2001). Typing of IS 1311 polymorphisms confirms that bison (*Bison bison*) with paratuberculosis in Montana are infected with a strain of *Mycobacterium avium* subsp. *paratuberculosis* distinct from that occurring in cattle and other domesticated livestock. *Molecular and Cellular Probes*, 15, 139–145.
- Whittington, R. J., Marshall, D. J., Nicholls, P. J., Marsh, I. B., & Reddacliff, L. A. (2004). Survival and dormancy of *Mycobacterium avium* subsp. *paratuberculosis* in the environment. *Applied and Environment Microbiology*, 70, 2989–3004.
- Whittington, R. J., Reddacliff, L. A., Marsh, I., McAllister, S., & Saunders, V. (2000). Temporal patterns and quantification of excretion of *Mycobacterium avium* subsp. *paratuberculosis* in sheep with Johne's disease. *Australian Veterinary Journal*, 78, 34–37.
- Whittington, R. J., & Windsor, P. A. (2009). *In utero* infection of cattle with *Mycobacterium avium* subsp. *paratuberculosis*: a critical review and meta-analysis. *The Veterinary Journal*, 179, 60–69.
- Windsor, P. A., Eppeleston, J., Dhand, N. K., & Whittington, R. J. (2014). Effectiveness of Gudair vaccine for the control of ovine Johne's disease in flocks vaccinating for at least 5 years. *Australian Veterinary Journal*, 92, 263–268.
- Windsor, P. A., & Whittington, R. J. (2010). Evidence for age susceptibility of cattle to Johne's disease. *The Veterinary Journal*, 184, 37–44.
- Wolf, R., Barkema, H. W., De Buck, J., & Orsel, K. (2015). Sampling location, herd size and season influence *Mycobacterium avium* ssp. *paratuberculosis* environmental culture results. *Journal of Dairy Science*, 98, 275–287.
- Wolf, R., Barkema, H. W., De Buck, J., & Orsel, K. (2016). Dairy farms testing positive for *Mycobacterium avium* ssp. *paratuberculosis* have poorer hygiene practices and are less cautious when purchasing cattle than test-negative herds. *Journal of Dairy Science*, 99, 4526–4536.
- Wolf, R., Barkema, H. W., De Buck, J., Slomp, M., Flaig, J., Hauptstein, D., ... Orsel, K. (2014). High herd-level prevalence of *Mycobacterium avium* subspecies *paratuberculosis* in Western Canadian dairy farms, based on environmental sampling. *Journal of Dairy Science*, 97, 6250–6259.
- Wolf, R., Donat, K., Khol, J. L., Barkema, H. W., Kastelic, J., & Wagner, P. (2017). Detection of *Mycobacterium avium* subspecies *paratuberculosis* infected cattle herds using environmental samples: a review. *Berliner und Munchener Tierarztliche Wochenschrift*, 130, 4–12.
- Wolf, R., Orsel, K., De Buck, J., & Barkema, H. W. (2015). Calves shedding *Mycobacterium avium* subspecies *paratuberculosis* are common on infected dairy farms. *Veterinary Research*, 46, 71.
- Wraight, M. D., McNeil, J., Beggs, D. S., Greenall, R. K., Humphris, T. B., Irwin, R. J., ... Mansell, P. D. (2000). Compliance of Victorian dairy farmers with current calf rearing recommendations for control of Johne's disease. *Veterinary Microbiology*, 77, 429–442.
- You, Q., Verschoor, C. P., Pant, S. D., Macri, J., Kirby, G. M., & Karrow, N. A. (2012). Proteomic analysis of plasma from Holstein cows testing positive for *Mycobacterium avium* subsp. *paratuberculosis* (MAP). *Veterinary Immunology and Immunopathology*, 148, 243–251.
- Zhong, L., Taylor, D., Begg, D. J., & Whittington, R. J. (2011). Biomarker discovery for ovine paratuberculosis (Johne's disease) by proteomic serum profiling. *Comparative Immunology, Microbiology and Infectious Diseases*, 34, 315–326.

How to cite this article: Barkema HW, Orsel K, Nielsen SS, et al. Knowledge gaps that hamper prevention and control of *Mycobacterium avium* subspecies *paratuberculosis* infection. *Transbound Emerg Dis*. 2018;65(Suppl. 1):125–148. <https://doi.org/10.1111/tbed.12723>