


SYSTEMATIC REVIEW PROTOCOL

Open Access



Management of UK priority invasive alien plants: a systematic review protocol

Philip A. Martin^{1,2*} , Gorm E. Shackelford^{1,2}, James M. Bullock³, Belinda Gallardo^{2,4,5}, David C. Aldridge^{2,4} and William J. Sutherland^{1,2}

Abstract

Background: Invasive alien plant species often have negative environmental and social impacts, such as loss of biodiversity and alteration of ecosystem services. As a result, managing the introduction, establishment, and abundance of invasive species is a major priority. To do this effectively, we need evidence on the effect of management interventions (such as using herbicide or cutting to control invasive plants). This evidence should not only include the effects of these management interventions on invasive alien species, but also on native species and other non-target outcomes such as ecosystem services. Such evidence would allow for comparison of the trade-offs between different management interventions. In the planned review we aim to assess how management interventions to control nine priority invasive alien plants species in England and Wales affect environmental outcomes. These species are: Japanese knotweed (*Fallopia japonica*), Nuttall's waterweed (*Elodea nuttallii*), Chilean rhubarb (*Gunnera tinctoria*), Giant hogweed (*Heracleum mantegazzianum*), Floating pennywort (*Hydrocotyle ranunculoides*), Himalayan balsam (*Impatiens glandulifera*), Curly waterweed (*Lagarosiphon major*), American skunk cabbage (*Lysichiton americanus*), Parrot's feather (*Myriophyllum aquaticum*).

Methods: Searches will be in English and use bibliographic databases (*Scopus*, *Web of Science Core Collection*, *Open Access Theses and Dissertations*, and *Conservation Evidence*) and internet searches (*Google Scholar*), as well as specialist databases. Our methodology will only use the names of each species (scientific names and common names, including synonyms) as our search string (we will not use qualifiers, such as "AND invasive"). This will give low specificity but will increase the likelihood of capturing all relevant information. We will use predefined criteria for study inclusion and data extraction. We will screen publications in two stages: (1) using titles and abstracts and (2) using full texts. Consistency of inclusion will be checked by two people screening a random sample of 10% of titles and abstracts. This dual-screening will be subject to kappa analysis and any disagreements resolved through discussion. We will use critical appraisal to assess study validity by identifying studies that are potentially prone to bias.

Keywords: Biosecurity, Evidence-based conservation, Invasive plant management, Meta-analysis, Subject-wide evidence synthesis, Species of union concern

Background

Invasive plant species can often have negative environmental and social impacts—reducing native biodiversity [1], altering ecosystem services [2, 3], causing losses

of crops [4], and potentially having negative effects on human health [5]. In the UK alone, invasive plant species cause over £400 million of losses per year [6]. These negative consequences mean that controlling the introduction, establishment, and abundance of invasive species is a major policy priority recognised by national [7], regional [8], and global policy instruments [9]. This requires significant expenditure. As such, to ensure that money allocated for control of invasive species is used

*Correspondence: phil.martin.research@gmail.com

¹ Conservation Science Group, Department of Zoology, University of Cambridge, Cambridge, UK

Full list of author information is available at the end of the article



© The Author(s) 2020. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

efficiently, it is vital that the most effective methods to manage invasive species are identified.

Numerous primary studies have tested the effect of management interventions on invasive plant species and the wider environment. To form generalisations based on these studies requires evidence synthesis, and a number of systematic reviews and meta-analyses have done this to address management of a range of species [10–13]. However, most syntheses of invasive plant management only concentrate on the effects of management on the abundance of invasive plants (target effects) [10–12], rather than on native biodiversity, ecosystem services, or other outcomes (non-target effects). Failure to consider these multiple metrics that may be affected by the management of invasive species can result in perverse outcomes. Indeed, one meta-analysis on this topic found that burning to control invasive plant species often also leads to a decline in the abundance of native plant species [13].

Recently the UK Department for Food and Rural Affairs (DEFRA) launched a consultation on the management of priority invasive alien species in England and Wales which focused on 14 priority species—8 plant species and six animal species. In response to this consultation, we aim to carry out a systematic review to investigate the effects of management on these priority alien plant species, and one additional plant species (Japanese knotweed), on a wide range of environmental outcomes.

Objective of the review

Our protocol is designed to be able to be applied to any management intervention, for any of the priority invasive plant species identified by Defra—Nuttall's waterweed (*Eloдея nuttallii*), Chilean rhubarb (*Gunnera tinctoria*), Giant hogweed (*Heracleum mantegazzianum*), Floating pennywort (*Hydrocotyle ranunculoides*), Himalayan balsam (*Impatiens glandulifera*), Curly waterweed (*Lagarosiphon major*), American skunk cabbage (*Lysichiton americanus*), Parrot's feather (*Myriophyllum aquaticum*). We will also investigate the management of an important invasive plant not identified by Defra: Japanese knotweed (*Fallopia japonica*). We will extract information on any relevant environmental outcome related to this. As such the overall objectives are phrased as 'open' questions. To improve the clarity of these questions, we have added examples of the type of specific questions we envisage being answered as part of our planned systematic review. All research questions follow a PICO (P = populations, I = interventions, C = comparators, O = outcomes) structure.

Primary questions:

1. What are the environmental impacts of interventions for managing invasive plant species?

The question consists of the following components (see study inclusion criteria for more detailed definitions):

Population: (i) The target invasive alien species; (ii) Habitats/environments that are invaded or might be invaded by the target invasive alien plant species.

Intervention: Any management intervention which aims to control or prevent the spread of the target invasive alien plant species.

Comparator: Any comparison between a location or time where a management intervention has been used and a location or time where it has not been used.

Outcome: Any environmental outcome.

Examples:

- What are the effects of herbicide on the abundance of Japanese knotweed (*Fallopia japonica*)?
- What are the effects of cutting to control Japanese knotweed (*Fallopia japonica*) on the abundance of native plants?
- What are the effects of biological control to control Japanese knotweed (*Fallopia japonica*) on soil carbon?

Secondary questions:

Secondary questions will investigate how effect modifiers alter the impacts of invasive alien plant management on environmental outcomes.

1. How does the intensity at which a management intervention is applied (e.g. application rate of herbicide, number of livestock per ha per year) alter its effects on environmental outcomes?

Example: How does application rate of herbicide alter its effects on the abundance of Japanese knotweed (*Fallopia japonica*)?

2. How do differences in study designs alter the observed effects of management interventions (e.g. greenhouse vs field or experimental vs observational)?

Example: Do the observed effects of herbicide application to control Japanese knotweed (*Fallopia japonica*) differ in field and greenhouse studies?

3. Does the time since invasion influence the management outcomes?
4. Does the time since a management intervention influence management outcomes?

5. How does the effectiveness of invasive plant control relate to impacts on other outcomes (e.g. native plant community)?

Example: How does the alteration of Japanese knotweed (*Fallopia japonica*) abundance affect the abundance of native plant species?

6. How do differences in climate affect the effectiveness of management interventions to control invasive plants?

Example: How do differences in precipitation alter the effects of herbicide application on the abundance of Japanese knotweed (*Fallopia japonica*)?

7. How does the scale which a management intervention was carried out alter its effects on invasive plant abundance?

Another aim of this review is to identify gaps in knowledge relating to the different interventions and outcomes for the invasive alien species that are the subject of the review.

Methods

Our methods describe a protocol that we will use for a systematic review of the management of nine priority invasive alien plant species in England and Wales. For the purposes of this protocol, an invasive alien plant is defined as a species that is considered to be outside of its native biogeographic range, that has spread as a result of human activity and causes negative ecological or economic impacts [14]. Because management interventions will be similar for different invasive species, this approach will allow us to answer broader scale questions about the management of invasive plant species. Our work will initially focus on global evidence about the eight invasive plant species identified as priorities by Defra in England and Wales, and one other species we have identified as a priority, but we envisage that similar protocols could be used to investigate invasive plant management of any species in any geographic region.

Our systematic review will make use of a new website for screening literature, extracting data, and presenting dynamic meta-analyses (<http://www.metadataset.com>) [15]. We aim to use this website to host multiple systematic reviews, using the current systematic review as a proof-of-concept. At present the website allows reviewers to screen papers, use automated kappa analysis, and enter study data. In addition, the website allows users to browse the included publications by management intervention, outcome, or country. In the future, users will be able to browse the data extracted from these publications and interact with meta-analyses, by selecting subsets of the data (e.g. data from selected countries or selected

management interventions). We welcome potential collaborators to please contact us.

The methods for this systematic review follow the Collaboration for Environmental Evidence guidelines for evidence synthesis [16], and a ROSES (RepOrting standards for Systematic Evidence Synthesis) checklist [17] has been completed (Additional file 1).

Stakeholder engagement

The protocol has been developed as part of the BioRISC (Biosecurity Research Initiative at St Catharine's) programme. This programme aims to build, integrate, and synthesise evidence, across the different domains of biosecurity, including invasive alien species. We have discussed this protocol with members of the BioRISC team and the Conservation Evidence group at the University of Cambridge [18]. The authors of this paper include experts in terrestrial and aquatic invasive species (BG, JMB, DA) and we consulted with one other academic expert. Although we did not engage directly with Defra, this protocol has been developed in response to their consultation on the management of priority invasive alien species in England and Wales. A previous systematic map protocol published in *Environmental Evidence* was used as the basis for this protocol [15].

Searching for articles

Our search strategy is designed to retrieve all the publications on each invasive plant species of interest. We will search for publications in four bibliographic databases that were identified as relevant by the review team. In addition to published, peer-reviewed literature, we will also search for unpublished research to minimize the risk of publication bias. Information on the studies screened will be held on the Metadataset website. We will also allow people to submit any relevant papers that they feel we have missed during our searches and we will subsequently screen these using the methodology outlined in this protocol.

Search strings

For each species, we will use its scientific name(s) and English common name(s) as the search string, based on a standard list of taxonomic synonyms [19], and details of all of these search strings are given in Table 1. These search strings should produce the most inclusive set of search results possible, because it will not use "AND" terms (e.g. "Impatiens glandulifera" AND control). Where Boolean search terms (e.g. "OR" or "AND") are not supported by databases (e.g. Google Scholar, which only allows basic Boolean searching) we will modify our search strategy to only use the most commonly used scientific name. This search strategy is likely to be very

Table 1 Search strings to be used for each species included in the systematic review

English name	Latin name	Search string
Chilean rhubarb	<i>Gunnera tinctoria</i>	"Chilean rhubarb" OR "giant rhubarb" OR (Gunnera AND (tinctoria OR scabra)) OR (Panke AND (acaulis OR caulescens OR tinctoria OR chilensis))
Giant hogweed	<i>Heracleum mantegazzianum</i>	"Giant hogweed" OR "cartwheel-flower" OR "giant cow parsley" OR "giant cow parsnip" OR "hogsbane" OR "wild parsnip" OR "hogsbane" OR "wild parsnip" OR "wild rhubarb" OR "Heracleum mantegazzianum"
Himalayan balsam	<i>Impatiens glandulifera</i>	"Himalayan balsam" OR "Policeman's Helmet" OR "Bobby Tops" OR "Copper Tops" OR "Gnome's Hatstand" OR "Kiss-me-on-the-mountain" OR "ornamental jewelweed" OR (Balsamina AND (glandulifera OR macrochila OR roylei)) OR (Impatiens AND (glandulifera OR candida OR cornigera OR glanduligera OR macrochila OR moschata OR royleana OR roylei OR taprobatica))
Curly waterweed	<i>Lagarosiphon major</i>	Lagarosiphon AND (major OR muscoides) OR "African elodea" OR "oxygen weed"
American skunk cabbage	<i>Lysichiton americanus</i>	"American skunk cabbage" OR "western skunk cabbage" OR "yellow skunk cabbage" OR "swamp lantern" OR "Lysichiton americanus"
Nuttall's waterweed	<i>Elodea nuttallii</i>	"Anacharis nuttallii" OR (Elodea AND (columbiana OR minor OR nuttallii)) OR (Philotria AND (minor OR nuttallii)) OR "Udora verticillata" OR "nutall* waterweed" OR "western waterweed"
Floating pennywort	<i>Hydrocotyle ranunculoides</i>	Hydrocotyle AND (ranunculoides OR adoensis OR batrachioides OR cymbalarifolia OR natans OR nutans) OR "floating pennywort" OR "floating marshpennywort"
Parrot's feather	<i>Myriophyllum aquaticum</i>	"Enydris aquatica" OR (Myriophyllum AND (brasiliense OR proserpinacoides OR aquaticum)) OR "parrots feather" OR "parrots-feather"
Japanese knotweed	<i>Fallopia japonica</i>	"Fallopia japonica" OR "fallopia compacta" OR "pleuropterus zuccarinii" OR (polygonum AND (cuspidatum OR cuspidatum OR hachidoense OR reynoutria OR sieboldii OR zuccarinii)) OR (reynoutria AND (japonica OR compacta OR hachidoensis OR hachijoensis OR hastata OR henryi OR uzenensis)) OR "tiniaria japonica" OR "japanese knotweed" OR "asian knotweed" OR "fleeceflower" OR "Himalayan fleece vine" OR "monkeyweed" OR "monkey fungus" OR "Hancock's curse" OR "elephant ears" OR "pea shooters" OR "donkey rhubarb" OR "sally rhubarb" OR "japanese bamboo" OR "american bamboo" OR "mexican bamboo"

Search strings were used for databases that allowed Boolean operators, but in cases where they were not allowed the Latin species name alone was used

comprehensive as it will find all available literature for the species of interest. It also has the advantage of making searches relatively easy to update, as well as allowing for changes in taxonomy to be dealt with easily in the future. Because our search string will have high 'sensitivity' (i.e. a high proportion of all the publications about each invasive alien species) we expect our search strategy to be very comprehensive and therefore the need for a benchmark list of articles to test comprehensiveness is minimised.

Because English common names may represent common words or phrases, we will subject all keyword searches to sensitivity tests. Where the inclusion of common names increases the number of papers returned by a search by ≥ 100 , we will restrict the search by adding "AND (control OR manag*)" for common names only.

Search languages

We will review publications in English only. We will tag all publications that are excluded because they were not in English. We acknowledge that excluding literature written in non-English languages is a shortcoming. However, we do not have the resources needed to work in other languages, especially since checking of consistency requires dual-screening of articles.

Publication databases to be searched

Publications will be collated from the following databases:

- Scopus
- Web of Science Core Collection
- Open Access Theses and Dissertations (<https://oatd.org/>)

Scopus and Web of Science searches will use the 'title, keyword, and abstract function' and will be performed using a University of Cambridge subscription. In addition to specialist bibliographic databases we will search one web-based search engine, *Google Scholar* (first 50 results sorted by relevance). Previous work has shown that this method is useful for finding both academic and grey literature [20].

Specialist searches

We will also search specialist websites for relevant studies. The first of these, <https://www.conservationevidence.com>, contains systematically collated evidence on the effects of management interventions taken both from the academic and grey literature [18, 21]. This website

has also already collated evidence on the control of three of the priority species that are the subject of this review (floating pennywort, American skunk cabbage, and Parrot's feather) and so represents a useful resource for the planned systematic review. In addition, we will examine accounts for each species on the CABI Invasive Species Compendium website (<https://www.cabi.org/isc/>), and on the IUCN 'EU regulation on invasive alien species' website (<https://bit.ly/2SjfgG>) to identify potential studies relating to management both from the academic and grey literature.

Additional search method

During screening of full texts, the reference lists of studies that meet inclusion criteria and any relevant secondary literature will be examined to identify other potentially useful studies, sometimes called 'snowballing' [22]. The same process will be carried out using any relevant studies found in this way until no relevant studies are found in reference lists. This method may help to find literature that was missed in key word searches as well as 'grey' literature [22].

Article screening and study eligibility criteria

Screening process

We will screen publications in two stages: [1] using titles and abstracts and [2] using full texts. At each stage we will decide whether to include or exclude a publication based on the eligibility criteria we set out below. At each stage we will record the number of publications included/excluded, and we will provide a list of the full texts that were excluded, together with reasons for their exclusion and a ROSES flow diagram [17]. To ensure a lack of bias, members of the review team will not screen any articles that they have authored—these papers will be screened by other members of the team.

Consistency checking

To check for consistency, a random sample of 10% of titles/abstracts will be screened, by two people, using our inclusion criteria. Any disagreements between the two people will be discussed, and eligibility criteria will be revised to show how disagreements were resolved. Kappa scores will be calculated to test the agreement between the two people [23]. If Kappa scores are below 0.6, another 10% of titles/abstracts will be screened by the same two people. Disagreements will be discussed and resolved again, Kappa scores recalculated, and the consistency checking process repeated until Kappa scores are greater than 0.6.

Following this, a sample of 10% of the full texts of publications that meet inclusion criteria based on their titles/

abstracts will be screened by two people. This process will mirror that detailed above for title/abstract screening.

Eligibility criteria

In this systematic review we will use a "PICO" approach to determine eligibility criteria. To be included in our systematic review studies must attain the criteria detailed below.

Populations We are interested in multiple populations in this systematic review. Firstly, we are interested in the following invasive species: Japanese knotweed (*Fallopia japonica*) Nuttall's waterweed (*Elodea nuttallii*), Chilean rhubarb (*Gunnera tinctoria*), Giant hogweed (*Heracleum mantegazzianum*), Floating pennywort (*Hydrocotyle ranunculoides*), Himalayan balsam (*Impatiens glandulifera*), Curly waterweed (*Lagarosiphon major*), American skunk cabbage (*Lysichiton americanus*), Parrot's feather (*Myriophyllum aquaticum*). In addition, we are also interested in the environments/ecosystems that have been impacted or may be impacted by these invasive alien species which may be affected by management interventions to control/prevent invasive alien species.

Interventions We are interested in all management interventions that attempt to exclude or control the invasive alien plant species detailed above. These interventions will be categorised using a provisional classification of practices that the review team have developed (see Additional file 2). We will adapt the categories in this classification scheme as we screen the literature and identify interventions that were not previously classified. The classification scheme is intended to be used as a means of classifying management interventions for invasive alien plant species that can be reused for any invasive plant species in the future. Previous general classifications for management interventions in conservation already exist [24] but these are defined at too low a resolution to be useful for the current systematic review.

Eligible comparators We will include any study that compares an area or time in which a management intervention has been carried out with one where no management intervention has been undertaken. To aid synthesis we will label studies based on different elements of their design. We will include both experimental and correlative studies. Experimental studies will be defined as those in which treatment and comparator groups were defined prior to a management intervention being carried out. Correlative studies will be defined as those in which comparators were defined after a management intervention was carried out. An example of a correlative study is one that compares areas grazed by livestock to control

an invasive plant with areas that were selected because they are similar to the grazed site but where no livestock grazing was used. This distinction is important because post hoc definition of comparators may result in biases in counterfactual comparisons.

Studies that compare an area where a management intervention has been used with an area where it has not been used will be labelled as controlled. This includes comparisons with areas where no management was used and with areas where alternative management interventions were used (e.g. a comparison of herbicide vs cutting). Studies where there is a comparison between a time when no management intervention was used and a time when the management intervention was used at a later time point will be labelled as a before-and-after study. These labels are not mutually exclusive—and so we will include studies that include both these design elements—commonly referred to as before-after-control-impact studies. Where experimental studies have used a blocked design, they will be labelled as such, and when correlative studies used paired sites, this will also be recorded. We will also include studies that have investigated different intensities of the same management intervention (e.g. comparison of different concentrations of herbicide).

Both replicated and unreplicated studies will be included in our systematic review, and we will label them as either replicated or not. We will also label studies as randomized or not. We will include studies carried out in natural habitats, under greenhouse conditions, or in laboratories. Further information on the methodological details of studies that we will record can be found in Table 2.

Outcomes We will include studies of all environmental outcomes, which we will categorise based on a provisional classification of outcomes that we have developed (see Additional file 3). This includes outcomes relating to invasive plant species and native plant species, as well as on other biodiversity, crop production, soil, water, and pollutants. Note that we will not include outcomes that relate to the adoption of management interventions by practitioners (e.g. the number of practitioners using herbicide to control invasive plants), which we see as being beyond the scope of this work. In addition, where biological control has been used to control invasive alien plants, we will not extract data about the biocontrol agents themselves. At the full-text screening stage studies that do not present tables and figures relating to outcomes will be excluded.

Similarly to our classification scheme of management interventions, this classification scheme will be used as a means of categorising different outcomes used in studies that have tested management interventions. This classification scheme will be updated if we find additional

relevant outcomes during screening and data extraction. This will allow us to develop a classification of outcomes that is reusable for other invasive plant species.

Study validity assessment

Critical appraisal strategy

We will critically appraise all studies that we consider to be suitable for quantitative synthesis to assess their validity. Doing this will allow us to identify studies that are likely to be more prone to biases that affect internal validity. We will use sensitivity analysis to further examine these biases.

Table 2 gives the criteria for study validity assessment. These criteria represent the variables that we consider to be critical in influencing the internal validity of study findings, which focus on the effects of selection bias and performance bias. We will not assess to what extent results from individual studies are generalisable (external validity) as this will vary depending on the context of the reader, given that generalisability is likely to vary geographically and taxonomically, as well as in other ways.

Data on criteria in Table 2 will be extracted from relevant studies. Any studies for which the answer is 'no' or 'unclear' to any of the questions will be assigned as having low validity; studies that are not assigned as having low validity will be assigned as having medium validity if any of the questions are answered as 'partially', the remainder of studies will be classified as having high validity. The study validity of a random sample of 10% of studies for each species will be determined by two reviewers. Any reasons for disagreement will be discussed.

Data on study validity will be presented alongside details extracted from the studies and descriptive statistics calculated to give an overview of study validity. Where possible, we will then use sensitivity analyses to examine how summary effect sizes are altered by exclusion of studies with particular characteristics. This will involve the exclusion of studies assessed as having low validity followed by exclusion of those with both low and medium validity.

Data coding and extraction strategy

Metadata extraction and coding

Metadata will be extracted from studies that meet our selection criteria. Metadata on the context and PICO elements that we will extract from each study are detailed in Table 3.

Data extraction strategy

We will extract the mean values of treatments (e.g. plots with management interventions) and controls (e.g. plots without management interventions or which used alternative management interventions). If available, we will

Table 2 Criteria for study validity assessment

Question/criterion	Response to question			Type of bias addressed
	Yes	Partially	No	
Did the study consist of both temporal and spatial comparisons?	Before-after-control-impact study	Before-and-after study or controlled study	N/A as study is not eligible for inclusion based on inclusion criteria	Lacking sufficient information to judge Selection bias
Did the study use randomization?	Study accounts for spatial heterogeneity by using appropriate randomisation of samples	N/A as study was either randomized with respect to the management intervention or not (e.g. random site selection but not random allocation of treatments/controls)	Study does not attempt to randomize sampling	Lacking sufficient information to judge Selection bias
Did the study avoid confounding factors?	Confounding factors were likely to be minimal as a result of blocking/pairing or stated attempts to match samples	Some confounding factors present, likely to have a moderate impact on outcome	Study was subject to confounding factors that could have a major impact on the outcome	Lacking sufficient information to judge Selection bias and performance bias
Can the study determine causality?	Experimental study in which comparator samples were selected prior to the management intervention being used	Correlative study in which comparators are selected after the management intervention has already been implemented, thereby limiting the ability of researchers to determine the similarity of comparators prior to management intervention use	N/A—studies with no comparator will be excluded	Lacking sufficient information to judge Selection bias and performance bias

Table 3 Data we plan to extract from studies and indication of which data will be used in study validity assessment

Variable	Description	Used in study validity assessment
Country	Country/countries where study was carried out	No
Latitude	Latitude of site(s) used in study	No
Longitude	Longitude of site(s) used in study	No
Start date	Date on which management intervention was first implemented (If only month is specified the date will only record the month and year)	No
End date	Date on which implementation of management intervention ceased/was carried out for the final time (If only month is specified the date will only record the month and year)	No
Invasive plant species	Accepted Latin binomial name for focal invasive species	No
Ecosystem type	Ecosystem type in which study was undertaken - following the IUCN habitat classification scheme (https://www.iucnredlist.org/resources/habitat-classification-scheme)	No
Management intervention	Name of management intervention - based on management intervention classification scheme (see Additional file 2 for more details of this)	No
Frequency of management intervention	Number of times year ⁻¹ that a management intervention was carried out	No
Intensity of management intervention	The intensity at which a management intervention was carried out (e.g. Grazed with 4 sheep ha ⁻¹ year ⁻¹)	No
Area over which intervention was applied	Area in ha over which a management intervention was applied	No
Controlled/not controlled	A study which compares outcomes in a location where a management intervention has been used with an area where no management intervention was used	Yes
Before-and-after	A study which compares outcomes between a time when no management intervention was used and a time when the management intervention was used at a later time point	Yes
Experimental/correlative study	Is the study experimental or correlative? (see text for definition of these terms).	Yes
Randomization	Randomized/not randomized	Yes
Pairing	Paired/unpaired (for correlative studies)	Yes
Blocking	Blocked/unblocked (for manipulative studies)	Yes
Size of sampling unit	Size in m ² of sampling unit (e.g. plot) used to collect data	No
Homogeneity of plot sizes used	Homogenous/not homogenous	Yes
Number of replicates	Number of replicate sample units in each comparative group	No
Distance between replicate plots	Distance in km between different plots used to investigate the same treatment	No
Distance between treatments	Distance in km between plots used to investigate different treatments	No
Greenhouse/field study	Was the study carried out in a greenhouse or in the field?	No
Duration of study	Number of years after management intervention for which results were monitored	No
Confounding factors	Were there any severely confounding factors? (yes, partially, no, unclear)	Yes
Outcomes	For a full list of outcomes that we will consider see our outcome classification scheme in Additional file 3	

In cases where there is not enough information in articles to extract information for a variable this will be left blank

also extract measures of variability around the mean (standard deviation, variance, standard error of the mean, or confidence intervals), number of replicates, and the *P* value of the comparison between treatments and controls (see Table 4 for a list of data that will be extracted). Comparisons will only be made within one figure/table but not between figures/tables. For example, if studies were done in two or more areas, but these results were presented in separate figures/tables, we will assume that a comparison cannot be made across tables unless specified otherwise in the article text. Where data are presented for multiple years, data from all years will be extracted. Where data is

presented for multiple sites, data for all sites will also be extracted. We will not contact primary authors to request additional data. All extracted data will be made available as additional files.

Approaches to missing data

Missing data can present a serious issue for meta-analyses in ecology and conservation as reporting standards in the literature are very variable. This often includes missing information about sampling variance or sample sizes [25]. Where possible, if information is missing for components required for effect size calculation and

Table 4 Data we will extract to allow calculation of effect sizes

Data to extract	Description of data
Treatment mean	Mean value for outcome in area/time period subject to management intervention
Control mean	Mean value for outcome in area/time period not subject to management intervention
Treatment standard deviation (SD)	Standard deviation for outcome in area/time period subject to management intervention
Control standard deviation (SD)	Standard deviation value for outcome in area/time period not subject to management intervention
Treatment N	Sample size for outcome in area/time period subject to management intervention
Control N	Sample size in area/time period not subject to management intervention
Treatment standard error (SE)	Standard error for outcome in area/time period subject to management intervention
Control standard error (SE)	Standard error value for outcome in area/time period not subject to management intervention
Unit	Units used to measure outcome (e.g. %, Kg ha ⁻¹ , or number of stems ha ⁻¹)
Least Significant Difference (LSD)	Metric allowing determination of statistical significance of a comparison between treatment mean and control mean
Correlation coefficient (r)	Measure of linear correlation between two variables (Pearson correlation coefficient)
Total N for correlation studies	Total sample size for studies that calculated the Pearson correlation coefficient
Approximate P value	Approximate P value where exact value of P is not given
Exact P value	Exact P value where it is given in the study
Exact Z value	Exact Z value where it is given in the study
Exact F value	Exact F value where it is given in the study. F values will only be taken from non-omnibus tests (i.e. not ANOVA tests) because these allow us to determine pairwise statistical significance

study weighting (i.e. mean values, sample sizes, measures of variability) we will calculate these values based on the available information. For example, P-values can be converted into t-test or F-statistics, which can subsequently be used to calculate effect sizes [26]. To investigate the influence of using variance to weight analyses we will use sensitivity analysis to compare results for both weighted and unweighted meta-analyses.

Consistency checking

Consistency of data extraction will be checked by randomly sampling 10% of those studies that meet inclusion criteria from which their data will be extracted by two people. Discrepancies in the data extracted will be discussed and extracted data modified accordingly. No Kappa analysis will be performed at this stage as our goal will be for complete agreement between reviewers.

Potential effect modifiers/reasons for heterogeneity

Potential reasons for heterogeneity will be explored using sub-group analysis, meta-regression, and sensitivity analysis (Table 5). Where relevant we will expand the list of potential modifiers during the synthesis process. This list was compiled following discussion within the review team.

Data synthesis and presentation

We will present quantitative data synthesis for intervention-outcome combinations that have ≥ 1 data points.

Interactive evidence synthesis will also be provided on Metadataset.

Narrative synthesis strategy

We will include a narrative synthesis to give an overview of the studies used in the systematic review. This will include presentation of tables containing relevant information about the studies that meet our inclusion criteria and further discussion of studies, where necessary.

Quantitative synthesis strategy

The minimum amount of data we will need to calculate an effect size will be the treatment mean and control mean, which will allow us to calculate the response ratio [27]. We will calculate the mean effect of each intervention on each outcome, using standard meta-analytic methods [28]. If we have only the minimum amount of data for the effect of an intervention on an outcome (i.e. treatment means and control means), we will only be able to do an unweighted meta-analysis. Otherwise, we will do a weighted meta-analysis (e.g. weighted by the inverse of variance, but sensitivity analyses using other weights will also be available on the website, as will unweighted meta-analysis). We will always use random-effects meta-analysis, as study-level effects are likely to vary because of true variation, rather than solely experimental error, as assumed by fixed-effects meta-analysis [29]. We will also explore heterogeneity using subgroup analysis and meta-regression, where appropriate. We will use standard

Table 5 Potential effect modifiers/reasons for heterogeneity

Variable	Potential effect	Method of testing
Study in greenhouse/field	Studies of management interventions such as herbicides in greenhouses may be more effective than those in the field because of the lack of exposure to wind/rain that may remove herbicide from target plants	Sub-group analysis
Rate of application of herbicide/cutting (management intensity)	Higher application rate of herbicide may lead to more effective control of invasive plant species	Sub-group analysis/meta-regression
Type/mechanism of herbicide	Different types of herbicide may differ in their effectiveness in controlling target invasive plant species	Sub-group analysis
Timing of herbicide application	Timing of herbicide application may alter their effectiveness in controlling target invasive plant species	Sub-group analysis/meta-regression
Frequency of management intervention (e.g. repeated spraying with herbicide)	More frequent application of herbicide or more regular cutting may lead to more effective control of invasive plant species	Sub-group analysis/meta-regression
Experimental vs observational	The results of experimental studies may be more consistent due to having fewer confounding factors	Sub-group analysis
Time since management intervention	Some management interventions may decline in success as time since they were last carried out increases	Meta-regression
Time since invasion	In sites where invasive plant species have been present for long periods of time they may be difficult to control because of a build-up of invasive propagules	Meta-regression

methods for estimation of publication bias (e.g. fail-safe numbers and funnel plots) [25]. Where relevant we will report the results of any sensitivity analyses.

Knowledge gap identification

We will identify knowledge gaps when the systematic review is completed. We will use our categorisation of both management interventions and outcomes to identify where there is a lack of knowledge for particular outcomes for each management intervention. If the systematic reviews are updated on the metadataset website, we will modify information on the website to identify where gaps remain.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s13750-020-0186-y>.

Additional file 1. Completed ROSES form.

Additional file 2. Classification of interventions.

Additional file 3. Classification of outcomes.

Acknowledgements

The authors thank Phil Hulme for useful comments on the protocol.

Authors' contributions

This protocol is based on a draft written by PM. GS and WS conceived the idea. All authors contributed to the improvement of the manuscript. All authors read and approved the final manuscript.

Funding

This work was funded by the David and Claudia Harding Foundation

Availability of data and materials

Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹ Conservation Science Group, Department of Zoology, University of Cambridge, Cambridge, UK. ² BioRISC (Biosecurity Research Initiative at St Catharine's), St Catharine's College, Cambridge CB2 1RL, UK. ³ Centre for Ecology and Hydrology, Maclean Building, Crowmarsh Gifford, Wallingford OX10 8BB, UK. ⁴ Aquatic Ecology Group, Department of Zoology, University of Cambridge, Downing St, Cambridge CB2 3EJ, UK. ⁵ Instituto Pirenaico de Ecología, Avenida Montañana, 1005, 50059 Saragossa, Spain.

Received: 17 June 2019 Accepted: 6 January 2020

Published online: 21 January 2020

References

- Vilà M, Espinar JL, Hejda M, Hulme PE, Jarošík V, Maron JL, et al. Ecological impacts of invasive alien plants: a meta-analysis of their effects on species, communities and ecosystems. *Ecol Lett*. 2011;14(7):702–8.
- Vilà M, Hulme PE, et al. Impact of biological invasions on ecosystem services, vol. 12. Berlin: Springer; 2017.
- Pejchar L, Mooney HA. Invasive species, ecosystem services and human well-being. *Trends Ecol Evol*. 2009;24(9):497–504.

4. Pimentel D, Zuniga R, Morrison D. Update on the environmental and economic costs associated with alien-invasive species in the United States. *Ecol Econ*. 2005;52(3):273–88.
5. Essl F, Biró K, Brandes D, Broennimann O, Bullock JM, Chapman DS, et al. Biological flora of the British Isles: *Ambrosia artemisiifolia*. *J Ecol*. 2015;103(4):1069–98.
6. Williams F, Eschen R, Harris A, Djeddour D, Pratt C, Shaw R, et al. The economic cost of invasive non-native species on Great Britain. *CABI Proj No VM10066*. 2010;1–99.
7. DEFRA. The Great British invasive non-native species strategy. 2015.
8. European Union. Combat invasive alien species—target 5 [Internet]; 2017. http://ec.europa.eu/environment/nature/biodiversity/strategy/target5/index_en.htm. Accessed 25 Apr 2019.
9. CBD. TARGET 9—technical rationale extended (provided in document COP/10/INF/12/Rev.1) [Internet]. 2014. <https://www.cbd.int/sp/targets/rationale/target-9/>. Accessed 25 Apr 2019.
10. Tyler C, Pullin A. Do commonly used interventions effectively control *Rhododendron ponticum*? CEE review 04-005 (SR6); 2005. <http://www.environmentalevidence.org/SR6.html>.
11. Kabat T, Stewart G, Pullin A. Are Japanese knotweed (*Fallopia japonica*) control and eradication interventions effective? CEE review 05-015 (SR21). *Collab Environ Evid*. <http://www.environmentalevidence.org/SR21.html>.
12. Roberts P, Pullin A. Effectiveness of the control of ragwort (*Senecio*) species: Are currently recommended herbicides effective for control of ragwort (*Senecio*) species? 2004. <http://www.environmentalevidence.org/SR5a.html>.
13. Kettenring KM, Adams CR. Lessons learned from invasive plant control experiments: a systematic review and meta-analysis. *J Appl Ecol*. 2011;48(4):970–9.
14. Essl F, Bacher S, Genovesi P, Hulme PE, Jeschke JM, Katsanevakis S, et al. Which taxa are alien? Criteria, applications, and uncertainties. *Bioscience*. 2018;68(7):496–509.
15. Shackelford GE, Haddaway NR, Usieta HO, Pypers P, Petrovan SO, Sutherland WJ. Cassava farming practices and their agricultural and environmental impacts: a systematic map protocol. *Environ Evid*. 2018;7(1):30.
16. Collaboration for Environmental Evidence. Guidelines and standards for evidence synthesis in environmental management. Version 5.0 [Internet]. 2018. <http://www.environmentalevidence.org/information-for-authors>. Accessed 23 Oct 2019.
17. Haddaway NR, Macura B, Whaley P, Pullin AS. ROSES Reporting standards for Systematic Evidence Syntheses: pro forma, flow-diagram and descriptive summary of the plan and conduct of environmental systematic reviews and systematic maps. *Environ Evid*. 2018;7(1):7.
18. Sutherland WJ, Taylor NG, MacFarlane D, Amano T, Christie AP, Dicks LV, et al. Building a tool to overcome barriers in research-implementation spaces: the conservation evidence database. *Biol Conserv*. 2019;1(238):108199.
19. The Plant List [Internet]. The plant list. <http://www.theplantlist.org/>. Accessed 3 Nov 2019.
20. Haddaway NR, Collins AM, Coughlin D, Kirk S. The role of Google Scholar in evidence reviews and its applicability to grey literature searching. *PLoS ONE*. 2015;10(9):e0138237.
21. Sutherland WJ, Wordley CF. A fresh approach to evidence synthesis. 2018.
22. Greenhalgh T, Peacock R. Effectiveness and efficiency of search methods in systematic reviews of complex evidence: audit of primary sources. *BMJ*. 2005;331(7524):1064–5.
23. Cohen J. A coefficient of agreement for nominal scales. *Educ Psychol Meas*. 1960;20(1):37–46.
24. Salafsky N, Salzer D, Stattersfield AJ, Hilton-Taylor C, Neugarten R, Butchart SHM, et al. A standard lexicon for biodiversity conservation: unified classifications of threats and actions. *Conserv Biol*. 2008;22(4):897–911.
25. Nakagawa S, Noble DW, Senior AM, Lagisz M. Meta-evaluation of meta-analysis: ten appraisal questions for biologists. *BMC Biol*. 2017;15(1):18.
26. Lajeunesse MJ, Koricheva J, Gurevitch J, Mengersen K. Recovering missing or partial data from studies: a survey of conversions and imputations for meta-analysis. In: Koricheva J, Gurevitch J, Mengersen K, editors. *Handbook of meta-analysis in ecology and evolution*. Princeton: Princeton University Press; 2013. p. 195–206.
27. Hedges LV, Gurevitch J, Curtis PS. The meta-analysis of response ratios in experimental ecology. *Ecology*. 1999;80(4):1150–6.
28. Borenstein M, Hedges LV, Higgins JP, Rothstein HR. *Introduction to meta-analysis*. Chichester: Wiley; 2011.
29. Overview. In: *Introduction to meta-analysis* [Internet]. Wiley; 2009. p. 59–62. <https://onlinelibrary.wiley.com/doi/abs/10.1002/9780470743386.ch10>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

