

# **A cross-sectional study of tuberculosis among workers in Tygerberg Academic Hospital, Western Cape province, South Africa**

by  
Julius Nkongho Ayuk

*Thesis presented in fulfilment of the requirements for the degree of  
Master of Medicine (Occupational Medicine) in the Faculty of Medicine  
and Health Sciences at Stellenbosch University*



Supervisor: Dr. Willem Albertus Jacobus Meintjes

December 2013

## Declaration

By submitting this thesis electronically, I declare that the entirety of the work contained therein is my own, original work, that I am the sole author thereof (save to the extent explicitly otherwise stated), that reproduction and publication thereof by Stellenbosch University will not infringe any third party rights and that I have not previously in its entirety or in part submitted it for obtaining any qualification.

December 2013

"

"

"

"

"

"

"

"

"

"

"

"

"

"

Eqr { tki j vÍ "4235"Uvngpdquej "Wpkxgtukv{

Cm'tki j w'tgugt xgf "

## Abstract

**Introduction:** The morbidity and mortality associated with tuberculosis (TB) disease is of grave consequences for the health and employment of afflicted individuals. Healthcare workers are identified amongst high risk groups in communities. The prevalence/incidence of TB is dependent on the presence of associated risk factors which varies in diversity and intensity in different communities and workplaces. Understanding the risk factors operating in any given environment is indispensable to any tuberculosis control programme.

**Objective:** The objective of this study was to describe the occurrence and trends of TB disease as well as to determine the risk factors associated with the disease among Tygerberg hospital employees.

**Method:** A cross-sectional descriptive study design with a nested case-control component was used to determine the occurrence (and trends) and risk factors of TB disease respectively.

**Occurrence and trends of tuberculosis:** The frequencies, distribution and trends of TB disease from 2008 to 2011 were obtained by calculating and comparing the annual incidence rates for each variable. Cases were identified from the occupational health clinic TB register, while the various denominator data were obtained from the Human Resource database.

**Determination of risk factors:** Cases were recruited from the occupational health clinic TB register and controls were randomly selected from unaffected workers during the study period. Self-administered risk factor questionnaires were completed by both cases and controls. Multivariate logistic regression analysis was used to determine the association between known and suspected risk factors and the occurrence of TB disease amongst employees.

**Results:** Sixty six cases of TB disease occurred in the workforce during the study period resulting in an annual average incidence rate of 397/100,000 population (95% CI: 307/100,000-505/100,000). Twenty three (34.8%) of the 66 cases occurred in Housekeeping staff, making them the most affected sub-group [1181/100,000 population (95% CI: 747/100,000-1768/100,000)]. The rate of TB disease in nurses was 1.7 times (95%CI: 1.4-2.0) that of doctors. Workers in the 40-49 years age-group experienced the highest incidence [490/100,000 population (95%CI: 329.6/100,000-706.8/100,000)] of TB disease compared to the other age-groups. There was no obvious difference in gender occurrences. Disease rates varied among different racial groups, with the highest rate in black employees [1473/100,000 population, (95%CI: 924/100,000-1981/100,000)]. Distribution of TB disease in the institution was widespread, with security department being the most affected [2500/100,000 population (95%CI: 311/100,000-9262/100,000)]. There was a downward but statistically insignificant (annual range 9-23; p=0.28) trend

in the rate of disease occurrence over the study period. No previous training on TB prevention (OR: 2.97, 95% CI: 1.15 - 7.71), HIV (OR: 67.08, 95% CI: 7.54 – 596.64) and working without knowledge of TB risk profile of the workplace (OR: 8.66, 95% CI: 1.10 – 67.96) were associated with TB disease occurrence.

**Conclusion:** Occurrence of TB disease among Tygerberg hospital employees was low compared to that of the general population of its drainage areas. Disease occurrence in the facility was wide and varied with respect to occupational groups, workplaces and time. Well-established risk factors for TB infection (and disease) were found to be determinants of disease occurrence in the facility.

## **Acknowledgements**

My appreciation goes to Dr WAJ Meintjes, division of Community Health, University of Stellenbosch through whose tutelage this work was guided and produced. I wish to also acknowledge the contributions to this work, the efforts of Mrs. Dawn Arendse and Jolene Samuels of the occupational health clinic, Tygerberg hospital, Cape Town, South Africa.

Finally, I thank God for his unlimited blessings

<b>Table of Contents</b>	<b>Page</b>
Title page.....	i
Declaration.....	ii
Acknowledgement.....	iii
Abbreviations.....	iv
Abstracts.....	v
Table of contents.....	vii
List of illustrations.....	ix
Introduction and Literature review.....	1
Study aims and objectives.....	3
Materials and Methods.....	3
Study designs.....	3
Ethical considerations.....	3
Setting.....	4
Study population.....	4
Methods.....	5
Determination of occurrence and trends of active tuberculosis.....	5
Determination of risk factors for active tuberculosis.....	6
Statistical analysis.....	6
Results.....	7
Frequency, distribution and trend of active tuberculosis.....	8
Frequency of occurrence.....	8
Trends of active TB disease occurrence.....	9
Distribution of active TB disease.....	10
Risk factors of active TB disease.....	17

Discussion and conclusion.....	21
Recommendations.....	26
References.....	28
Appendix.....	33

## List of Figures

<b>Number</b>	<b>Page</b>
Figure1. Annual incidence rates of TB disease in TBH employees.....	10
Figure2. Incidence rates of TB disease in different departments .....	15

## List of Tables

<b>Number</b>	<b>Page</b>
Table1. Incidence rates and trends of active TB disease in TBH employees.....	09
Table2. Frequency of occurrence of active TB disease based on the age of employees.....	12
Table3. Frequency of occurrence of active TB disease based on the race of employees.....	13
Table4. Frequency of occurrence of active TB disease based on the gender of employees...	14
Table5. Frequency of occurrence of active TB disease based on workplaces.....	16
Table6. Baseline characteristics of participants enrolled in risk factor determination.....	19
Table7. Multivariate logistic regression analysis of possible TB risk factors.....	20



## CHAPTER ONE: INTRODUCTION AND LITERATURE REVIEW

Tuberculosis (TB) is a systemic infectious disease caused by *Mycobacterium tuberculosis*<sup>[1]</sup>. Pulmonary disease is the most frequent clinical presentation of the contagious disease<sup>[1,2,3]</sup>. Extra-pulmonary diseases, which was only occasionally seen, is now increasing in frequency secondary to HIV<sup>[4, 5]</sup> and include tuberculosis of the skin, lymph nodes, gastrointestinal tract, meninges, bone & joints, pericardium, kidneys and pelvis<sup>[6-8]</sup>. The principal mode of transmission of *Mycobacterium tuberculosis* among humans is by inhalation of airborne bacteria<sup>[9, 10]</sup>.

Generally, there are two types of infections; latent TB which is a non-symptomatic infection and active TB, which is characterized by symptoms and/ or signs of active disease (and it is usually associated with the presence of active bacilli in appropriate samples)<sup>[11]</sup>. A recent classification introduces a third category which is an active but subclinical disease often seen in HIV positive individuals<sup>[12]</sup>. The commonest clinical symptoms of active TB disease are chronic cough, phlegm production, haemoptysis, fever, weight loss and night sweats<sup>[13]</sup>. There has been a rise in active TB disease globally since the beginning of the HIV/AIDS pandemic with sub-Saharan Africa, especially the Southern African countries, leading the epidemic<sup>[14,15,16]</sup>. The Global incidence rate of active TB disease in 2011 was estimated by WHO at 125/100,000 population and Africa with an incidence rate of 262/100,000 population accounted for 26% of all reported incident cases in the same year<sup>[17]</sup>. The incidence rate of active TB disease in South Africa in 2011 was estimated at 993/100,000 population and the Western Cape Province, with an incidence rate of 935/100,000 population is ranked as one of the most affected regions in the country<sup>[18]</sup>. In fact, of the 8.7 million incident cases of active TB disease reported globally in 2011, South Africa accounted for 0.4-0.5 million cases ranking it among the top 5 high burden countries in the World<sup>[17]</sup>.

There are several risk factors associated with the transmission of (and development of active) TB. These factors are classified into environmental and personal risk factors. The environmental risk factors include the concentration of airborne bacteria, ventilation & air circulation, contact with smear positive sputum individuals and overcrowding<sup>19</sup>. Personal risk factors include factors such as HIV status, cancer, malnutrition, smoking, age, substance abuse, diabetes, chronic renal failure with dialysis, body mass index of less than 18.5, genetic predisposition, silicosis, immunosuppressive drugs and other forms of immunosuppressive disease conditions<sup>[20, 21]</sup>. Certain identified groups within communities are more

vulnerable to TB and among these are healthcare workers and HIV infected patients<sup>[22-23]</sup>. Higher rates of both latent<sup>[24,25,26,27,28,29,30]</sup> and active TB disease<sup>[31]</sup> have been found in healthcare workers relative to their respective general populations both in developing countries<sup>[24-26,31]</sup> and developed countries<sup>[27-30]</sup>. Several of the above risk factors are present in workers within the healthcare work environment in South Africa. The risk of exposure by health care workers to TB has been shown to be related to the prevalence of TB in the population served and the degree of contact with TB patients<sup>[32]</sup>. The differential distribution of these risk factors in workplaces with regards to variables such as age, departments, number of patient contacts, profession, etc. are expected to influence the incidence of TB infections (and TB disease) relatively among workers. Studies have highlighted differences in the occurrence of TB amongst workers in different facilities based on some of the above variables<sup>[33, 34]</sup>.

Understanding the risk factors for TB operating in a given work environment is necessary for effective control of the disease among workers. The level of risks in healthcare facilities have been shown in some circumstances to vary depending on the setting, departments, occupation, patient loads and effectiveness of TB control measures. A review article on TB in healthcare workers by Menzies & co-workers concluded that, the risk of occupational TB varies considerably among and within institutions<sup>[35]</sup>. Engineering, administrative and personal protective devices have been used to reduce the risk of healthcare acquired infections but prompt diagnosis and rapid treatment of TB in addition to the above control measures is recommended by the WHO for controlling the spread of infection<sup>[14]</sup>. Preventing healthcare acquired infection is important not only because of the risk of transmission of infection to non-infected workers and patients, but because of the high level of transmission of multidrug resistant TB<sup>[36, 37]</sup>. In recent years the emergence of multidrug resistant (MDR-TB) and extensively drug resistant strains (XDR-TB) of TB has re-awakened the emphasis of protecting the healthcare workforce against TB infection. Based on the WHO global report 2010 (and MDR-TB update 2012), South Africa is listed among the high burden countries for MDR-TB transmission<sup>[17]</sup>. This has further heightened the need for healthcare worker protection against TB considering the severity of, and disability associated with active form of the disease. Furthermore, active TB disease is a compensable disease in healthcare workers in South Africa. Control of TB transmission within the workforce with consequent reduction in compensation claims in the current environment of scarce resources, should be encouraged.

Occupational TB is expected to be high among workers in Tygerberg hospital as it is located in the Western Cape Province which harbours one of the highest incidences of community based TB in South Africa. Understanding the distribution and predisposing risk factors of active TB disease in Tygerberg hospital is important in managing the disease in the establishment. Such information may lead to specific and targeted responses to disease control.

## **Study Aims and Objectives**

The aims of the study were to determine the following among workers in Tygerberg hospital:

1. The frequency, distribution and trends of active TB disease from 2008 to 2011.
2. The possible risks factors and their degree of association with active TB disease among employees.

## **CHAPTER TWO: MATERIALS AND METHODS**

### **Study design**

A cross-sectional descriptive study design with a nested case-control component (using a self-administered questionnaire) was used to answer the research questions.

### **Ethical considerations**

Ethical clearance was obtained from the Stellenbosch University Health Research Ethics Committee and the Western Cape Department of Health to conduct the research in Tygerberg Hospital (reference No.S12/02/032). Participants were educated on the purpose of the research and their rights with regards to the study. Written informed consent was obtained from every participant as a pre-condition for recruitment into the study. Confidentiality in respect to patients' information was ensured throughout the course of the study.

### **Study setting**

This study was performed at Tygerberg hospital, a tertiary referral hospital in the Western Cape Province of South Africa. The hospital is the second largest hospital in the country, with a staff complement of approximately four thousand workers. It has a well-developed Unit for Infection Prevention and Control working in collaboration with the occupational health unit in controlling infectious diseases among patients and staff. The hospital drainage area includes areas with one of the highest prevalence of active TB disease in the country. The hospital experienced an average of 17 cases of active TB annually from 2008 to 2011(unpublished data).

Employees with symptoms of active TB disease present at the occupational health clinic for evaluation and are given treatment when positively diagnosed. Employees who are diagnosed elsewhere (e.g. visiting their general practitioner), are referred to the clinic for the purpose of completion of their compensation application and other statutory documents. They are also evaluated for fitness to work in view of the risk they pose to hospital patients and other workers and the effects of their duties on their health. A register is kept of all persons treated for active TB disease at the occupational health clinic.

## **Study population**

The study population for the descriptive aspect (i.e. disease occurrence and trends of active TB disease) comprised of Tygerberg hospital employees from January 2008 to December 2011 and the analytic study population included employees of the institution from January 2008 to June 2012.

In the analytic study, the cases (Definition: employees of the hospital who were diagnosed with active TB disease from January 2008 to June 2012) were identified from the TB register of the occupational health clinic and controls (Definition: employees who reported no history of active TB disease during the study period) were identified by means of human resources data (a full list of the employees working at the hospital during the period of the study).

Selection of controls: The controls were selected from the list of employees who worked at the hospital during the same period when active TB disease was diagnosed in the cases. The controls were stratified in terms of their profession/occupation. For each active TB case, three controls were randomly selected (within the same professional stratum as the case) from the human resources database. The selection was performed by means of a list of random numbers generated by the PhStat add-in for Microsoft Excel, version 2.5.0. When a control was not available for participation (e.g. relocated to another province or didn't want to provide consent for participation), the control was replaced by the next employee on the list of randomly generated controls. When the replacement employee did not want to provide consent, the person was not replaced again.

## **Inclusion criteria:**

Participants met all of the following criteria of eligibility for inclusion in the study:

- Must have been a permanent employee of Tygerberg hospital during the period of review

- Must have been diagnosed with active TB disease between January 2008 and June 2012 (cases) OR randomly selected from the human resource database of the hospital (controls)
- Must have provided signed informed consent.

#### **Exclusion criteria:**

Potential participants who met any one of the following criteria were not eligible for participation in the study:

- Employees who were not willing to provide informed consent
- Employees who could not be contacted (e.g. relocated to other provinces)
- Contracted employees/employees provided by labour brokers

#### **Determination of frequency, distribution and trends of active TB**

The TB database in the occupational health clinic was reviewed for information relevant to computing the frequency, distribution and trends of active TB disease in the hospital. The information was entered into Microsoft Excel spread sheets for analysis. The population denominators for the entire Tygerberg workforce and different subset-populations within the workforce (used in the analysis) were obtained from the human resources department of the hospital for each year of analysis. The frequency of occurrence of active TB disease in both the entire and subsets of the workforce was obtained by calculating their respective annual incidence rates (dividing the total number of new cases of active TB disease for each category of the workforce in a particular year by the mid-year workers' population for that category in that year). The distribution of active TB disease was obtained by comparing the annual incidence rates with respect to person (occupation, age, gender, and race), place (departments) and time (year of occurrence). The trends of occurrence were obtained by serially comparing, the annual incidence rates of active TB disease in the Tygerberg workforce (and in subsets of the workforce) over four years (January 2008 to December 2011). The cases of active TB disease that occurred in 2012 were not included in this aspect of the study because of incomplete annual incidence data for the entire year (as data collection ended before the end of 2012).

### **Determination of possible risk factors for active TB among workers**

All workers diagnosed with active TB disease during the study period and their respective controls were approached to participate in the questionnaire component of the study. Both cases and controls completed the same structured self-administered TB risk factor questionnaire. The questionnaire consisted of work-related and community based risk factor variables, which were grouped into demographic, socio-economic and health-related parameters (see appendix). Assistance in completing the questionnaire was offered by trained assistants wherever help was needed. The responses were extracted from the questionnaires and captured on spread sheets for analysis.

### **Statistical analysis**

In the descriptive aspect, the difference in annual incidence rates as well as the nature of the trends of active TB disease occurrence during the study period were statistically analysed using Stata 12.0 (Statacorp, College Station, TX, USA). The relative occurrence was estimated using incidence rate ratio and population parameters were estimated using 95% confidence intervals, with alpha set at the 5% significance level.

Data in the analytic aspect of the study were also statistically analysed using Stata 12.0. The analytic study was designed to use a sample size capable of detecting an odds ratio of 2.5 with a power of 80% at 5% significance level (i.e. 2748 participants - 687cases & 2061 controls). Two hundred and forty nine (75 cases &174 controls) participants qualified for analysis at the time of the study. Of the 249, 216 (58 cases and 158 controls) were available and completed the questionnaires which were used for analysis (and estimation) to determine the risk factors.

Continuous baseline variables were summarised using descriptive statistics and compared using the Mann-Whitney U test. Categorical baseline data were summarised using frequency tables and compared using the Chi-squared test or the Fisher's exact test where applicable. All tests were 2-sided. The association of exposure variables and active TB disease were estimated using odds ratios. A logistic regression analysis was used to identify risk factors associated with active TB disease.

Simple logistic regression models were fitted for variables assumed to be associated with active TB disease. This resulted in unadjusted odds ratio for each variable with corresponding p-values and 95%CI. Variables in the simple logistic regression model which were significant ( $p < 0.05$ ), had a p-value  $< 0.1$  and those known to be associated with TB transmission/disease from literature but were not significant (irrespective of their p-values) were also included in the final multiple logistic regression model. A 95% confidence interval was used to estimate population parameters; alpha was set at 5% level.

## CHAPTER THREE: RESULTS

All 63 employees diagnosed with active TB disease (2008-2011) were included in the descriptive aspect of the study. Of the 63 employees affected, two had recurrent disease during the period; one case had an episode of recurrent active TB disease and the other had 2 recurrent episodes (i.e. 66 cases of active TB disease occurred during the period). The cases comprised of 83.3% (n=55) pulmonary TB and 16.7% (n=11) extra-pulmonary disease [TB of the skin (n=2), meninges (n=1), lymph nodes (n=1), abdomen (n=4), spine (n=1) and miliary (n=2)]. The ethnicity of the cases included 41.3% (n=26) blacks, 50.8% (n=32) coloureds, 6.3% (n=4) whites and 1.6% (n=1) Indian. The occupational representations of cases included 4 doctors (6.3%), 21 nurses (33.3%), 21 housekeeping staff members (33.3%), 7 clerks (11.1%), 4 kitchen staff members (6.3%), 2 security officers (3.2%) and 4 other staff members - a radiographer, porter, social worker and an interpreter (6.3%). Of the 61 workers with known marital status, the majority of cases were single (49.2%). The married, divorced and widowed represented 47.5%, 1.6% and 1.6% of the cases respectively. The mean age of the affected employees was 43.8 years (range 23-60 years; SD=10, 95%CI: 41.2-46.4). The gender distribution of cases were 71.4% (n=45) female and 28.6% (n=18) male. The majority of cases [71.2% (n=47)] were smear positive (50.0%) or culture positive (21.2%) with the remainder [28.8% (n=19)] diagnosed via histological (6.0%) and clinical (22.7%) methods. The cases were 30.2% (n=19) HIV-positive, 63.5% (n=40) HIV-negative and 6.3% (n=4) had an unknown HIV-status. Extra-pulmonary TB disease cases had a higher prevalence of HIV [36.4% (n=11)] compared to pulmonary disease cases [(29.4% (n=51)]. The proportion of cases with first experience of active TB disease was 79.4% (n=50) and recurrent disease (previous experience) accounted for 20.6% (n=13). Among the active TB disease cases (with known HIV status), those with recurrent disease had a higher prevalence of HIV infection (30.8%) compared to employees with first TB disease experience (19%). The proportion of multidrug resistant disease among the cases was 6.3% (n=4).

### **Determination of frequency, distribution and trends of active tuberculosis**

#### **Frequency of occurrence:**

The average annual incidence rate of active TB disease in Tygerberg hospital staff during the study period was 397/100,000 population (95%CI: 307-505 per 100,000). The frequencies of occurrence (in both the workforce and subsets populations) varied annually with maximum average incidence rate in 2008 (569/100,000 population; 95%CI: 361-853 per 100,000) and minimum rate in 2009 (222/100,000 population; 95%CI: 102-422 per 100,000).



Housekeeping staff belonged to the most affected sub-group of workers with an average annual incidence rate of up to 1181/100,000 population (95%CI: 747-1768 per 100,000). Their incidence rate was approximately 3 times (95%CI: 2.7-3.3) that of the entire workforce, 3.6 times (95%CI: 3.2-4.1) that of nurses and 6.1 times (95%CI: 5.2-7.1) the rate in doctors. The occurrence of active TB disease in nurses was more than that of doctors (324/100,000 population vs. 194/100,000 population) resulting in an incidence rate ratio of nurses to doctors of about 1.7 (95%CI: 1.4-2.0).

Table 1 summarizes the annual incidence rates of active TB disease with respect to both the entire as well sub-groups of the workforce.

**Table 1. Incidence rates and trends of active TB disease in Tygerberg hospital employees (2008-2011)**

Year	2008	2009	2010	2011	Average annual incidence rates (95%CI)
<b>Variables</b>					
<b>All Tygerberg Hospital Cases</b>					
<b>Cases</b>	23	9	20	14	
<b>Population</b>	4044	4049	4244	4297	
<b>Incidence rates</b>	569/100,000	222/100,000	471/100,000	326/100,000	397/100,000 (307/100,000-505/100,000)
<b>Nursing Personnel</b>					
<b>Cases</b>	8	2	6	6	
<b>Population</b>	1616	1592	1782	1787	
<b>Incidence rates</b>	495/100,000	126/100,000	337/100,000	336/100,000	324/100,000 (203/100,000-492/100,000)
<b>Medical Practitioners</b>					
<b>Cases</b>	0	0	3	1	
<b>Population</b>	527	512	510	532	
<b>Incidence rates</b>	0/100,000	0/100,000	588/100,000	188/100,000	194/100,000 (52/100,000-492/100,000)
<b>Housekeeping Staff</b>					
<b>Cases</b>	9	3	7	4	
<b>Population</b>	482	491	491	488	
<b>Incidence rates</b>	1867/100,000	612/100,000	1426/100,000	820/100,000	1181/100,000 (747/100,000-1768/100,000)
<b>Other Healthcare Workers</b>					
<b>Cases</b>	6	4	4	3	
<b>Population</b>	1419	1454	1461	1490	
<b>Incidence rates</b>	423/100,000	275/100,000	274/100,000	201/100,000	293/100,000 (170/100,000-467/100,000)

\* Other healthcare workers - defined as all other workers in the hospital excluding doctors, nurses and housekeeping staff

## Trends in active TB disease occurrence

There was a downward trend of active TB disease occurrence in both the entire workforce of the hospital and in the subgroups of employees analysed. The trend of occurrence of active TB disease depicted an irregular pattern with an initial decline in incidence rates from 2008 to 2009 in most cases, followed by a rise and fall in subsequent years. This undulating pattern was clearly exemplified in the entire workforce and housekeeping staff. The only exception was in doctors who experienced an occurrence of active TB disease in 2010 following a preceding 2 year period without any case of active TB disease in the group. However, the trend of active TB disease occurrence in this group was difficult to interpret because of the small numbers. The decline in the incidence of active TB disease in the workforce over time (i.e. during the study period) was not significant (annual range 9-23;  $p=0.28$ ). The study also showed an insignificant statistical difference in the occurrence of active TB disease with regards to occupation during the same period ( $p=0.46$ ).

A graphic representation of the trends of occurrence of active TB disease in different groups of employees is shown in figure 1.




Figure 1. Annual incidence rates of active TB in Tygerberg hospital workforce and different subsets of employees

**Distribution of active TB disease in employees:**

The occurrence of active TB disease based on the personal characteristics of age, race and gender are summarized in tables 2, 3 and 4 respectively.

There was no worker under 20 years of age amongst employees of Tygerberg hospital. Workers in the 40-49 years age-group were most affected (average annual incidence rate of 490/100,000 population; 95%CI: 329.6-706.8 per 100,000). The average incidence rate in this group was about 1.4 times (95%CI: 1.3-1.7) that of the 20-29 years age-group, 2.1 times (95%CI: 1.8-2.4) that of 30-39 years age-group and similar to that of  $\geq 50$  years age-group (incidence rate ratio of 1.0; 95%CI: 0.90-1.2). In fact, active TB disease occurrence varied relatively between the groups (during the 4 year study period) with the maximum incidence rates in 2008 and 2009 occurring in the  $\geq 50$  years age-group and that in 2010 and 2011 occurring in 40-49 years age-group. The rate of occurrence of active TB disease in employees aged  $\geq 40$  years (482/100,000; 95%CI: 354-641 per 100,000) was significantly higher than those  $< 40$  years (276/100,000 population; 95%CI: 166-431 per 100,000) resulting in an incidence rate difference of 206/100,000 population ( $p=0.038$ ). Male employees showed a slight but statistically insignificant difference in disease occurrence compared to female employees (438/100,000 vs. 383/100,000 population) resulting in an incidence rate ratio of 1.1 (95%CI: 1.0-1.3). Black employees had the highest incidence rate of active TB disease (1473/100,000 population; 95%CI: 925-1975 per 100,000). The incidence in blacks was higher than any other group with incidence rate ratios of 5 (95%CI: 4.4-5.7), 5 (95%CI: 4.4-5.7) and 12.4 (95%CI: 10.3-15.0) compared to coloureds, Indians and whites respectively.

**Table 2. Frequency of occurrence of active TB disease based on the age-groups of employees**

<b>Year</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>Average annual incidence rates (95%CI)</b>
<b>Age(years)</b>					
<b>Age 20 – 29</b>					
<b>Cases</b>	2	1	3	2	
<b>Population</b>	548	579	624	588	
<b>Incidence rate</b>	365/100,000	173/100,000	481/100,000	340/100,000	340/100,000 (148/100,000-674/100,000)
<b>Age 30 – 39</b>					
<b>Cases</b>	6	1	2	2	
<b>Population</b>	1186	1108	1120	1132	
<b>Incidence rate</b>	506/100,000	90/100,000	179/100,000	177/100,000	238/100,000 (121/100,000-433/100,000)
<b>Age 40 – 49</b>					
<b>Cases</b>	9	3	10	7	
<b>Population</b>	1455	1440	1494	1504	
<b>Incidence rate</b>	619/100,000	208/100,000	669/100,000	465/100,000	490/100,000 (330/100,000-707/100,000)
<b>Age ≥50</b>					
<b>Cases</b>	6	4	5	3	
<b>Population</b>	855	922	1006	1073	
<b>Incidence rate</b>	702/100,000	434/100,000	497/100,000	280/100,000	478/100,000 (277/100,000-738/100,000)
<b>Age ≥40</b>					
<b>Cases</b>	15	7	15	10	
<b>Population</b>	2310	2362	2500	2577	
<b>Incidence rate</b>	649/100,000	296/100,000	600/100,000	388/100,000	483/100,000 (354/100,000-641/100,000)
<b>Age &lt;40</b>					
<b>Cases</b>	8	2	5	4	
<b>Population</b>	1734	1687	1744	1720	
<b>Incidence rate</b>	461/100,000	119/100,000	287/100,000	233/100,000	275/100,000 (166/100,000-431/100,000)

**Table 3. Frequency of occurrence of active TB disease based on the race of employees**

<b>Year</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>Average annual incidence rates (95%CI)</b>
<b>Race</b>					
<b>White employees</b>					
<b>Cases</b>	2	0	2	0	
<b>Population</b>	873	850	807	826	
<b>Incidence rate</b>	229/100,000	0/100,000	248/100,000	0/100,000	119/100,000 (33/100,000-305/100,000)
<b>Coloured employees</b>					
<b>Cases</b>	10	6	8	8	
<b>Population</b>	2701	2660	2744	2739	
<b>Incidence rate</b>	370/100,000	226/100,000	292/100,000	292/100,000	295/100,000 (209/100,000-416/100,000)
<b>Black employees</b>					
<b>Cases</b>	11	3	9	6	
<b>Population</b>	389	454	608	651	
<b>Incidence rate</b>	2828/100,000	661/100,000	1480/100,000	922/100,000	1473/100,000 (925/100,000-1975/100,000)
<b>Indian employees</b>					
<b>Cases</b>	0	0	1	0	
<b>Population</b>	81	85	85	81	
<b>Incidence rate</b>	0/100,000	0/100,000	1176/100,000	0/100,000	294/100,000 (08/100,000-1665/100,000)

**Table 4. Frequency of occurrence of active TB based on the gender of employees**

<b>Year</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>Average annual incidence rates (95%CI)</b>
<b>Gender</b>					
<b>Male gender</b>					
<b>Cases</b>	7	4	4	3	
<b>Population</b>	1014	1017	1039	1052	
<b>Incidence rates</b>	690/100,000	393/100,000	385/100,000	285/100,000	438/100,000 (259/100,000-690/100,000)
<b>Female gender</b>					
<b>Cases</b>	16	5	16	11	
<b>Population</b>	3030	3032	3205	3245	
<b>Incidence rates</b>	528/100,000	165/100,000	499/100,000	339/100,000	383/100,000 (283/100,000-508/100,000)

The occurrence of active TB disease with regards to workplaces showed a wide variation in both distribution and incidence rates. Cases occurred in both clinical and non-clinical areas of the hospital. The lowest incidence rates in the hospital occurred in those working in administration (240/100,000 population; 95%CI: 117.6-697.2 per 100,000) and the highest in the security department (2500/100,000 population; 95%CI: 311-9262 per 100,000). Workplaces with the highest rates of occurrence were oncology (1,726/100,000 population; 95%CI: 209-6228 per 100,000), food services (1,603/100,000 population; 95%CI: 691-3153 per 100,000) and technical department (1110/100,000 population (95%CI: 470-2145 per 100,000). Relatively low rates were experienced mostly in the clinical departments of medicine (510/100,000 population 95%CI: 243.9-935.4 per 100,000), obstetrics and gynaecology (414/100,000 population; 95%CI: 107-1006 per 100,000) and surgery (409/100,000 population; 95%CI: 222-681 per 100,000). A detailed summary of active TB disease occurrence with regards to various departments is presented in table 5 and a diagrammatic representation is shown in figure 2.




Figure 2. Incidence rates of active TB disease in different departments in Tygerberg hospital.

**Table 5. Frequency of occurrence of active TB disease based on workplaces**

Year	2008	2009	2010	2011	Average annual incidence rates (95%CI)
<b>Departments</b>					
<b>Department of medicine &amp; medical wards</b>					
Cases	3	1	4	2	
Population	498	495	483	490	
Incidence rates	602/100,000	202/100,000	828/100,000	408/100,000	510/100,000 (244/100,000-935/100,000)
<b>Department of surgery &amp; surgical wards</b>					
Cases	6	0	5	3	
Population	834	849	849	919	
Incidence rates	719/100,000	0/100,000	589/100,000	326/100,000	409/100,000 (222/100,000-681/100,000)
<b>Department of obstetrics and gynaecology &amp; associated wards</b>					
Cases	3	0	1	0	
Population	235	250	265	268	
Incidence rates	1277/100,000	0/100,000	377/100,000	0/100,000	414/100,000 (107/100,000-1006/100,000)
<b>Department of paediatrics &amp; paediatric wards</b>					
Cases	2	0	2	3	
Population	239	298	293	304	
Incidence rates	837/100,000	0/100,000	683/100,000	987/100,000	627/100,000 (248/100,000-1272/100,000)
<b>Department of psychiatry &amp; psychiatric wards</b>					
Cases	0	0	0	1	
Population	30	31	32	32	
Incidence rates	0/100,000	0/100,000	0/100,000	3125/100,000	781/100,000 (20/100,000-4457/100,000)
<b>Radiology &amp; Nuclear medicine</b>					
Cases	0	2	0	0	
Population	152	151	152	152	
Incidence rates	0/100,000	1325/100,000	0/100,000	0/100,000	331/100,000 (40/100,000-1190/100,000)
<b>Oncology</b>					
Cases	1	0	1	0	
Population	30	29	28	29	
Incidence rates	3333/100,000	0/100,000	3571/100,000	0/100,000	1726/100,000 (209/100,000-6228/100,000)



**Table 5. Frequency of occurrence of active TB disease based on workplaces (cont.)**

<b>Food services department</b>					
<b>Cases</b>	4	0	3	1	
<b>Population</b>	128	125	120	127	
<b>Incidence rates</b>	3125/100,000	0/100,000	2500/100,000	787/100,000	1603/100,000 (691/100,000-3153/100,000)
<b>Administrative employees &amp; administrative department</b>					
<b>Cases</b>	0	2	2	2	
<b>Population</b>	589	608	629	636	
<b>Incidence rates</b>	0/100,000	329/100,000	318/100,000	314/100,000	240/100,000 (118/100,000-697/100,000)
<b>Security personnel</b>					
<b>Cases</b>	0	1	1	0	
<b>Population</b>	17	20	20	21	
<b>Incidence rates</b>	0/100,000	5000/100,000	5000/100,000	0/100,000	2500/100,000 (311/100,000-9262/100,000)
<b>Technical department and all technical &amp; maintenance personnel</b>					
<b>Cases</b>	4	2	1	1	
<b>Population</b>	177	177	185	196	
<b>Incidence rates</b>	2260/100,000	1130/100,000	541/100,000	510/100,000	1110/100,000 (470/100,000-2145/100,000)
<b>Assistance services (porters &amp; translators)</b>					
<b>Cases</b>	0	1	0	1	
<b>Population</b>	125	121	123	122	
<b>Incidence rates</b>	0/100,000	826/100,000	0/100,000	820/100,000	412/100,000 (49/100,000-1471/100,000)

### **Risk factors of active TB disease**

Seventy five workers were diagnosed with active TB disease from January 2008 to June 2012. Of the 75 affected employees, 58 completed the questionnaire for risk factor analysis accounting for a participation rate of 77.3%. Of those who did not enrol for the study, 5 declined participation, 1 had passed away and 11 could not be contacted (relocated to another province, working for another employer, etc.). Of the 174 original randomized controls chosen for the study, 112 of them completed the questionnaires and 46 were replaced (declined participation/unavailable) by the next employee on the list of random control resulting to a total of 158 controls (case to control ratio of 1:2.7). The remaining 16 questionnaires could not be completed as selected controls as well as their replacements refused participation. The baseline characteristics of participants are summarized on table 6. It indicated that of all the variables tested; only alcohol consumption differs statistically between cases and controls participants.

Apart from the variables indicated on the multivariate model, univariate logistic regression analysis of other tested possible risk factors (found in the study questionnaire) showed insignificant associations between active TB disease in participants and the number of patient contacts per participant per day [ $<10$  patients (OR: 2.6;  $p=0.9$ ),  $\geq 10$  patients (OR: 1.8;  $p=0.2$ )], co-morbidity of active TB disease with any other disease condition including HIV (OR: 1.5;  $p=0.18$ ), stress exposure during the study period (OR: 1.7;  $p=0.10$ ), regular protective device use (OR: 1.8;  $p=0.07$ ), race[(black OR: 2.5 ;  $p=0.10$ ), coloured(OR: 3;  $p=0.28$ ), Indian (OR: 1.4;  $p=0.49$ ), age (OR: 1.0;  $p=0.27$ ), prolonged working hours i.e.  $> 8$ hours (OR: 1;  $p=0.6$ ) duration of work in healthcare (OR:1;  $p=0.36$ ), gender (OR:1.2;  $p=0.60$ ), yearly flu attack (OR:1.2;  $p=0.65$ ), number of household contacts (OR:1.0;  $p=0.60$ ) and cooking methods(OR: 2.0;  $p=0.30$ ). The factors associated with increased risk of active TB disease based on univariate analysis were HIV (OR: 50.14, 95%CI: 6.41- 392.11), no previous training on TB prevention (OR: 1.94, 95%CI: 1.05 – 3.59) and unknown status of TB vaccination (OR: 2.8, 95% CI: 1.21-6.57).

However, in the multivariate logistic regression model, the risk factors identified for active TB disease after adjusting for all inter-variable interactions were; HIV (OR: 67.08, 95% CI:7.54 – 596.64), no previous training on TB prevention (OR: 2.97, 95% CI:1.15 - 7.71) and working with no knowledge of TB risk profile of the workplace (OR: 8.66, 95% CI: 1.10 – 67.96). A summary of the multivariate logistic regression analysis is displayed in table 7.

**Table 6. Baseline characteristics of participants enrolled in the study**

<b>Characteristics n (%)</b>	<b>Cases (n=58)</b>	<b>Controls (n=158)</b>	<b>p-value</b>
<b>Categorical variables</b>			
<b>Marital status</b>			
▪ married	31(53.4)	84(53.5)	0.99
▪ unmarried	27(46.6)	73(46.5)	
<b>Gender of participants</b>			
▪ male	17(29.3)	52(33.1)	0.60
▪ female	41(70.7)	105(66.9)	
<b>Highest level of education attained by participants</b>			
▪ higher education	18(31.6)	53(33.8)	0.57
▪ post primary education	32(56.1)	92(58.6)	
▪ primary education	7(12.3)	12(7.6)	
<b>Daily work-shift schedule</b>			
▪ routine day shift	38(65.5)	98(62.4)	0.34
▪ night shift	3(5.2)	3(1.9)	
▪ both day and night shifts	17(29.3)	56(35.7)	
<b>Yearly flu attack</b>			
▪ no	28(50)	80(52.6)	0.65
▪ yes	29(50)	72(47.4)	
<b>Alcohol consumption by participants</b>			
▪ no	44(77.2)	88(56.4)	0.00
▪ yes	13(22.8)	68(43.6)	
<b>Co-morbidity status of participants (TB with any other disease)</b>			
▪ no co-morbidity	21(37.5)	74(47.7)	0.19
▪ has co-morbidity	35(62.5)	81(52.3)	
<b>Nature of hobbies</b>			
▪ involves no contact with public	47(83.9)	114(77.0)	0.28
▪ involves contact with public	9(16.1)	34(23.0)	
<b>Use of personal protection (PPE) against TB</b>			
▪ yes	34(58.6)	85(54.8)	0.53
▪ no	24(41.4)	70(45.2)	
<b>Race of participants</b>			
▪ white	5(8.6)	23(14.6)	0.22
▪ coloured	29(50.0)	92(58.2)	
▪ Indian	2(3.4)	3(1.9)	
▪ black	22(37.9)	40(25.3)	
<b>Continuous variable [median(interquartile range)]</b>			
<b>Number of household contacts of participants</b>	4.0 (3)	4.0 (2)	0.91
<b>Number of room contacts of participants</b>	1.0 (2)	1.0 (1)	0.46
<b>Work duration in TBH (years)</b>	11.0 (19)	11.8 (21)	0.98
<b>Work duration in healthcare sector (years)</b>	20.5 (21)	13.0 (20.5)	0.25
<b>Time worked per day (hours)</b>	9.0 (3)	9.5 (4)	0.66
<b>No. of patients contacts per day</b>	21.0 (24)	12.0 (25)	0.05

**Table 7. Multivariate logistic regression analysis of possible risk factors for active TB disease in Tygerberg hospital**

Risk factors	Unadjusted OR	95%CI ( Unadjusted OR)	Adjusted OR	95% CI (adjusted OR)	p-value (adjusted OR)
Age(continuous data )	1.02	0.97 – 1.05	1.05	1.00 – 1.10	0.05
<b>smoking history</b>					
▪ never smoked	1		1		
▪ previous smoker	1.30	0.49 – 3.44	2.20	0.62– 7.76	0.22
▪ current smoker	1.03	0.50 – 2.10	1.30	0.47 – 3.46	0.63
<b>Training on TB prevention</b>					
▪ received training on TB prevention	1		1		
▪ never received training on TB prevention	1.94	1.05 – 3.59	2.57	1.10 – 5.99	0.03
<b>TB vaccination status of participants</b>					
▪ vaccinated participants	1		1		
▪ participants with unknown vaccination status	1.42	0.70 – 2.89	0.96	0.39 – 2.33	0.92
▪ participants with no history of vaccination	2.82	1.21 – 6.57	1.66	0.54 – 5.1	0.38
<b>Participants' assessment of work area TB risk profile</b>					
▪ low risk area	1		1		
▪ don't know	4.49	0.99 – 20.30	7.56	1.14 – 50.20	0.04
▪ high risk area	3.32	0.96 – 11.52	4.07	0.89 – 18.58	0.07
<b>Contact with known TB cases</b>					
▪ no contact	1		1		
▪ contact in TBH	1.71	0.88 – 3.36	1.35	0.54 – 3.37	0.51
<b>Alcohol consumption by participants</b>					
▪ none	1		1		
▪ yes	0.38	0.19 - 0.77	0.44	0.17-1.10	0.08
<b>Regular use of personal protective equipment</b>					
▪ none	1		1		
▪ yes	0.96	0.81 – 1.13	0.79	0.34 – 1.83	0.58
<b>HIV infection</b>					
▪ HIV negative participants	1		1		
▪ HIV positive participants	50.14	6.41 – 392.10	50.94	5.26 - 493.73	0.00
<b>Diabetes mellitus</b>					
▪ diabetics	1		1		
▪ non-diabetics	1.71	0.76 – 3.86	1.67	0.58 – 4.89	0.34
<b>Race</b>					
▪ white	1		1		
▪ Indian	1.45	0.51 – 4.16	1.12	0.33 – 3.74	0.86
▪ coloured	3.07	0.40-23.44	4.24	0.38 – 46.91	0.24
▪ black	2.53	0.84-7.59	2.44	0.55 – 10.81	0.24

## CHAPTER FOUR: DISCUSSION AND CONCLUSION

Although healthcare workers across the globe have been shown by many studies to have higher risks for TB than the populations they serve<sup>[26, 30]</sup>, this study did not reflect the assertion. The incidence rate of active TB disease in healthcare workers in Tygerberg hospital (397/100,000 population) was lower than the rate in the surrounding population (Cape Town(799/100,000 population)<sup>[18, 38]</sup> and Western Cape (935/100,000 population)<sup>[18]</sup>) and nationally<sup>[17]</sup>. The result of this study is consistent with the finding of an earlier study in Mpumalanga which showed a relatively lower rate of active TB disease in healthcare workers compared to the general population<sup>39</sup>. However, a study by Kranzer & co-workers in Cape Town rather found an increased risk of active TB disease among community healthcare workers<sup>40</sup>. The rate of disease occurrence in the institution also compares favourably with other study findings in similar settings in some parts of South Africa and the continent<sup>[25, 41-43]</sup>. This finding is interesting considering the background community TB risk profile of the population served by the institution and the institutional level in the healthcare structure. The institution is located in a region with one of the highest TB occurrences in the country and is a major reference centre in the province and as such, is expected to experience higher disease frequency amongst its workers. The lower rate of active TB disease occurrence may be ascribed to two possible reasons; the referral status (tertiary) of the institution in a province with the best primary and secondary healthcare delivery system in the country means that many cases of active TB disease may have been identified and treated at a lower healthcare level with consequently lower TB risk in patients referred for consultation in the institution. Secondly, the presence of an active (and competent) occupational health service and Unit for Infection Prevention and Control (UIPC) might have contributed in mitigating the risk within the institution. The occupational health department and the UIPC carry out education and training of workers on TB prevention, assesses the risk of TB associated with different work areas and procedures on an on-going basis and recommend risk mitigating measures including the provision of respirators where relevant. The department is also responsible for the diagnosis, treatment and removal of employees with active TB disease from the workplace as well as ensuring complete resolution of disease prior to job rehabilitation.

Contrary to expectation, the study showed that the annual incidence rates of active TB disease in doctors & nurses (groups with very close contact with patients) were comparatively (and statistically) lower with respect to the rate in non-clinical workers such as housekeeping staff. This finding may be interpreted bearing in mind, the multifactorial basis of TB disease causation.

The lower rates of disease in nurses & doctors may be due to a comparatively higher level of awareness in these groups of persons, the majority of whom have had information and training on TB prevention. It may be useful for subsequent studies to evaluate the rate of TB disease occurrence in this group of workers relatively to that of their respective social class in order to postulate the effect of the risk associated with the nature of their work. The high incidence of TB disease in housekeeping staff could be explained, considering that their workplace activities bring them in close contact with patients (as they clean) and the majority of them belong to the lower socio-economic group in the society (known for higher community based TB risk profile<sup>[44, 45]</sup>). This group should be particularly monitored as they form the major source of TB hazard in the workplace. Mitigating the risk in this group is likely to have a positive effect in limiting TB transmission within the facility (not only for other workers but for patients as well).

The decline in the rates of active TB disease in the institution during the study period, though not statistically significant in the short-term (i.e. after three years of implementation of control measures) may require subsequent re-evaluation to determine its long-term effect. This is necessary, considering that the relatively low rates of occurrence of active TB disease (associated with a small sample size) may require a longer duration to obtain a sample size large enough to detect any difference (due to the recently introduced control measures) in the rates of occurrence over time as statistically significant. Unfortunately, the study could not forecast the occurrence of active TB disease beyond the study period. Such a finding would have been important for evaluating the current intervention (which includes the identification of target areas, provision of respirators for use in high risk areas and procedures, and regular information and training of employees on TB prevention in targeted groups). It would have also provided information capable of impacting on management decisions with regards to active TB containment.

The distribution of active TB disease based on the personal characteristics of age, race and occupation showed useful information relevant for disease control in the institution. The occurrence of the disease was particularly high in the older age groups (40-49 and  $\geq 50$  years old), black employees and household workers in the institution. Employees with these characteristics should receive special attention during medical surveillance as well as during clinical evaluation of cases with relevant symptomatic presentations. The distribution of the disease in workers of the institution reflects the socio-demographic characteristics (age, race and socio-economic status) influencing TB occurrence in the general population<sup>[46]</sup>.

Other personal risk characteristics of TB occurrence relevant to the transmission, control (environmental and medical surveillance) and clinical management of the disease in the institution were the HIV background and TB infectivity of affected employees. Up to 30.2% of all TB affected employees in the study were HIV positive and up to half of all cases were sputum smear positive at presentation (highly infectious). Human immunodeficiency viral infection is a well-established risk factor for both transmission and development of active TB <sup>[47]</sup> and its presence in employees involved in healthcare work has implications for TB control. The effect of this background risk in TB affected workers of Tygerberg hospital is seen in the high prevalence of HIV in patients with recurrent active TB in this study. Several studies have also identified the HIV background of persons as a risk factor for extra-pulmonary disease <sup>[4, 5]</sup>. Therefore, the high incidence of extra-pulmonary TB and the high prevalence of HIV in extra-pulmonary TB affected employees in this study add to that evidence. Studies in South Africa have described this background risk (HIV) as high amongst certain categories of employees working in the healthcare sector<sup>[48, 49]</sup> and therefore requires considerations for disease control measures in Tygerberg hospital. The effects of active TB disease, HIV and other HIV-related complications on illness-related absenteeism in the workplace environment in South Africa cannot be over-emphasized. The emergence of multi-drug resistance TB, in addition to the aforementioned effects only increases the complexity in the clinical and workplace management of TB-affected employees. The presence of multidrug resistant TB (MDR-TB) strains in up to 6.3% of TB disease cases (and involving mostly healthcare workers) in the institution requires urgent attention. Infection with resistant strains of TB is associated with increase in disease severity, disability and illness-related absenteeism. A study carried out in a public TB referral hospital in Kwazulu Natal concluded that healthcare workers were substantially more likely to be hospitalized with either MDR-TB or XDR-TB compared to non-healthcare workers<sup>[50]</sup>. Current health policy in the facility requires MDR-TB affected employees to be removed from the workplace until at least two consecutive sputum cultures are negative, provided that the person is physically fit to return to work. If one considers that healthcare workers are highly skilled group of workers with acute shortage in the labour market, then the urgency in controlling the occurrence of TB disease in them warrants emphasis.

Workplace occurrence of active TB showed a widespread distribution in the institution, with higher rates occurring in non-clinical areas (security and kitchen). This is indicative of the wide-spread distribution of TB risks across different departments in the hospital. It further emphasizes the need for controls to be widely instituted within the facility rather than focussed only in areas where contact with patients occurs.

Generally, control measures and medical surveillance in most institutions have always targeted clinical areas with less emphasis on non-clinical areas but based on the findings of this study, a mind-shift is necessary to target both areas in Tygerberg hospital. The widespread distribution of active TB in the facility raises two fundamental issues: the sources of exposure and acquisition of TB among employees in the facility (workplace vs. community) and the criterion of patient contacts in the workplace as a major determinant for workers' benefits.

Understanding the risk factors driving TB occurrence in the facility is indispensable in controlling the disease. With regards to active TB disease among workers in the hospital; the lack of previous training on TB prevention, the HIV background of employees and the lack of awareness of the TB risk profile of their workplaces by some employees were identified as possible risk factors underlying disease occurrence. Unless further studies show contrary evidence, control measures against active TB in the institution require among others, the management of these risk factors. Studies performed in similar settings elsewhere have also identified some of the above risk factors as influencing transmission rates among workers<sup>[51]</sup>.

Many other known risk factors for TB transmission present in the facility did not show significant associations with TB transmission. Reasonably expected workplace risk factors such as the number of patient contacts per day, contact with patients and patients' specimens, direct contact with known active TB disease cases in the hospital and the length of daily working hours did not show significant association with active TB disease in the facility. However, caution needs to be exercised in the interpretation of these results. The matching of controls to cases based on occupation prior to randomization might have distributed the above variables in a more or less equal extent between the groups and hence their results. Other possible reasons may be due to the biases associated with subjectivity of the responses to the questions in the questionnaire and inadequate power to detect small differences between the groups due to the small sample size of participants. The effect of the small sample size of participants (cases & controls) is seen in the wide confidence intervals associated with the incidence rates of some variables in this study. The baseline socio-demographic characteristics (table 6) of gender, marital status, level of education, race, monthly income and number of household contacts were shown not to be statistically different between cases & controls and hence may not have affected the study result.

This study has some specific strength that warrants emphasis; it is the first in the hospital that quantifies the frequencies of occurrence and defines the distribution of active TB disease in the entire workforce.



By defining the occurrence of active TB disease with regards to place and person characteristics, the study has implications in medical surveillance as it identifies high risk groups and workplaces for targeted interventions. The findings from the study may be useful for administrative purposes. The study results may be used as an evaluation tool by management, in deciding the necessity for response with regards to managing TB disease in the facility and in so doing, meet the legislative obligation of ensuring a healthy work environment. By identifying the possible risk factors driving TB transmission (and disease occurrence) within the workforce, the study may contribute to risk mitigating strategies (and therefore, disease control) for the disease in the institution.

However, this study is not without weaknesses. Apart from the already mentioned weaknesses associated with the risk factor identification aspect of the study, the following weaknesses can affect either or both aspects of the study; Selection bias is a likely possibility in the risk factors aspect of the study. Only 77.3% of active TB cases took part in the risk factor identification aspect of the study and if their responses with respect to the tested risk factors are different from those who did not participate in the study, then some bias would have been introduced in the findings. The effect of such bias if present, might have introduced either type I or II errors depending on the responses of those who did not take part in the study. Measurement bias resulting from misclassification is also a possible source of bias in the risk factors determination aspect of the study. The answers to most questions in the study questionnaire were subjective and therefore prone to bias. Likely sources of misclassification may have been in responses to HIV status in controls, smoking history, alcohol consumption, regular use of personal protective devices, number of patient contacts per day, etc. The above responses which usually under-represent the magnitude of the variables is likely to biased the effect towards the null value. In the descriptive aspect of the study, measurement bias could have arisen from under-reporting of active TB disease in employees who sought medical care outside the facility (even though they are instructed to report TB disease in the occupational health clinic for statutory benefits). “Recall bias” is also a possibility in the risk factor identification aspect of the study. Recall was typically problematic in some individuals with regards to TB vaccination and average number of patient contacts per day in both cases and controls. Since difficulties with recall were not particular to either of the groups, the effect (if significant) is likely to bias the result in either direction (i.e. towards or away from the null value).

In conclusion, Occurrence of active TB disease among employees of Tygerberg hospital was low compared to that of the general population of its drainage areas. The occurrence was wide and varied with respect to occupational groups, workplaces and time. Well-established risk factors

for TB transmission (and disease) in other settings (and medical literature) were identified as variables associated with disease occurrence in the facility. However, the constant presence of active TB disease cases in the workplace requires continuous evaluation, monitoring and mitigation of risk factors associated with TB in the facility.

## RECOMMENDATIONS

The following recommendations may be invaluable in the management of TB in the hospital:

Regular TB risk assessment in the facility is necessary in view of the current disease burden. Such an assessment should be carried out throughout the hospital at least annually in the most affected areas. The frequency of assessment could be varied depending on the future incidence of disease and in any case as decided by management in conjunction with the health and safety committee of the hospital. The use of the Western Cape's Facility Risk Assessment Tool for TB (FRATT) is one of many tools which can be used for such an evaluation.

Information and training on TB prevention and symptom recognition should be carried out regularly especially in departments and occupations most affected. This will empower employees and encourage their participation in disease control. As shown in this study, the lack of training on TB prevention is identified as one of the possible risk factors associated with TB transmission in the facility. Early symptom recognition and consultation on an on-going basis will not only limit transmission within the workplace but will also reduce the level of disability associated with active TB disease amongst employees.

Medical surveillance should be undertaken especially in the most affected workplaces. A cheap, practical and useful program may take the form of screening questionnaire for symptoms of active TB disease, administered quarterly (or more frequently depending on magnitude of the disease) in respective departments. This will enable suspected cases to be referred for early diagnosis and removal from workplaces so as to prevent or lower transmissions in the workplace.

Accommodation of workers with high personal risk profiles in low risk areas should be considered with due regards to ethical and current legislative requirements especially regarding discrimination (see Employment Equity Act 1998, Basic Condition Employment Act 1997, Labour Relations Act 1995 and the bill of rights, South African constitution). Based on this study, workers with HIV and recurrent active TB disease should be considered for accommodation in work areas with low TB risk if they currently work in high risk areas. The workplace management of newly diagnosed HIV cases should include TB risk management (including workplace placements).

## REFERENCES

1. Finch RG, Moss P, Jeffries DJ, Anderson (2006). *Clinical Medicine* (6<sup>th</sup> ed.). Elsevier Saunders. pp. 86-87. ISBN 0702027634
2. WHO report 2012. Basic facts about Tuberculosis. [Cited 9 September 2013]; Available from: [http://apps.who.int/iris/bitstream/10665/75938/1/9789241564502\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/75938/1/9789241564502_eng.pdf).
3. Kobzik L, Schoen F (1994) *Robbins Pathologic Basis of Disease* (5<sup>th</sup> ed). W.B. Saunders. pp700. ISBN 0-7216-5032-5.
4. Huebner RE, Castro KG. The changing phase of tuberculosis. *Annu Rev Med* 1995;46: 463-471.
5. Yang Z, Kong Y, Wilson F, Foxman B, Fowler AH et al. Identification of Risk Factors for Extrapulmonary Tuberculosis. *Clinical infectious disease* 2003;38:199-205.
6. Leeds IL, Magee MJ, Kurbatova EV, et al. Site of extrapulmonary tuberculosis is associated with HIV infection. *Clin Infect Dis*. 55(1):75-81.
7. Noertjojo K, Tam CM, Chan SL and Chan-Yeung MM. Extra-pulmonary and pulmonary tuberculosis in Hong Kong. *Int J Tuberc Lung Dis*. 2002;6(10):879-886.
8. Sharma SK and Mohan A. Extrapulmonary tuberculosis. *Indian J Med Res*. 2004;120(4):316-353.
9. Escombe AR, Oeser C, Gilman RH, et al. The detection of airborne transmission of tuberculosis from HIV-infected patients, using an in vivo air sampling model. *Clin Infect Dis*. 2007;44(10):1349-1357.
10. Roy CJ and Milton DK. Airborne transmission of communicable infection--the elusive pathway. *N Engl J Med*. 2004;350(17):1710-1712.

11. WHO Media Centre. Factsheet No. 104. Tuberculosis. 2013 [cited 17 May 2013]; Available from: <http://www.who.int/mediacentre/factsheets/fs104/en/>
12. Achkar JM and Jenny-Avital ER. Incipient and subclinical tuberculosis: defining early disease states in the context of host immune response. *J Infect Dis.* 2011;204 Suppl 4:S1179-1186.
13. Centres for Disease Control. Basic TB facts. 2012 [cited 17 May 2013]; Available from: <http://www.cdc.gov/tb/topic/basics/default.htm>
14. Blanc L, Falzon D, Fitzpatrick C, et al. Global Tuberculosis Control: A short update to the 2009 report. Geneva: World Health Organization, 2009
15. Raviglione MC, Harries AD, Msiska R, Wilkinson D, Nunn P. Tuberculosis and HIV: current status in Africa. *AIDS* 1997; 11(suppl B): S115-S123.
16. Corbett EL, Watt CJ, Walker N et al. The growing burden of tuberculosis: global trends and interactions with HIV epidemic. *Arch Intern Med* 2003; 163:1009-1021.
17. Floyd K, Dias HM, Falzon D, et al. Global tuberculosis report: 2012. Geneva: World Health Organization.
18. Rossouw H. World TB Day, 24 March 2012. 2012 [cited 17 May 2013]; Available from: <http://www.westerncape.gov.za/news/world-tb-day-24-march-2012>.
19. Rieder HL (1999). *Epidemiologic Basis of Tuberculosis Control* (1<sup>st</sup> ed.). International Union Against Tuberculosis and Lung Disease. pp. 17-25. ISBN 2-9504238-8-4.
20. Gajalakshmi V, Peto R, Kanaka TS and Jha P. Smoking and mortality from tuberculosis and other diseases in India: retrospective study of 43000 adult male deaths and 35000 controls. *Lancet.* 2003;362(9383):507-515.

21. Restrepo BI. Convergence of the tuberculosis and diabetes epidemics: renewal of old acquaintances. *Clin Infect Dis*. 2007;45(4):436-438.
22. Joshi R, Reingold AL, Menzies D, Pai M. Tuberculosis among Health-Care Workers in Low- and Middle- Income Countries: A systematic Review. *Plos Med* 2006; 3(12): e494.
23. Lawn SD, Bekker LG, Middelkoop K, Myer L and Wood R. Impact of HIV infection on the epidemiology of tuberculosis in a peri-urban community in South Africa: the need for age-specific interventions. *Clin Infect Dis*. 2006;42(7):1040-1047.
24. Christopher DJ, Daley P, Armstrong L, et al. Tuberculosis infection among young nursing trainees in South India. *PLoS One*. 5(4):e10408.
25. Molina-Gamboa J, Fivera-Morales I and Ponce-de-Leon-Rosales S. Prevalence of tuberculin reactivity among healthcare workers from a Mexican hospital. *Infect Control Hosp Epidemiol*. 1994;15(5):319-320.
26. Tan LH, Kamarulzaman A, Liam CK and Lee TC. Tuberculin skin testing among healthcare workers in the University of Malaya Medical Centre, Kuala Lumpur, Malaysia. *Infect Control Hosp Epidemiol*. 2002;23(10):584-590.
27. Drobniewski F, Balabanova Y, Zakamova E, Nikolayevskyy V and Fedorin I. Rates of latent tuberculosis in health care staff in Russia. *PLoS Med*. 2007;4(2):e55.
28. Fraser VJ, Kilo CM, Bailey TC, Medoff G and Dunagan WC. Screening of physicians for tuberculosis. *Infect Control Hosp Epidemiol*. 1994;15(2):95-100.
29. Harada N, Nakajima Y, Higuchi K, et al. Screening for tuberculosis infection using whole-blood interferon-gamma and Mantoux testing among Japanese healthcare workers. *Infect Control Hosp Epidemiol*. 2006;27(5):442-448.

30. Schablon A, Harling M, Diel R and Nienhaus A. Risk of latent TB infection in individuals employed in the healthcare sector in Germany: a multicentre prevalence study. *BMC Infect Dis.* 2010;10:107.
31. Naidoo S and Jinabhai CC. TB in health care workers in KwaZulu-Natal, South Africa. *Int J Tuberc Lung Dis.* 2006;10(6):676-682.
32. CDC. Guidelines for Preventing the Transmission of *M. tuberculosis* in Health-Care Settings, 2005. *MMWR.* 2005;54(No. RR-17).
33. Kilinc O, Ucan ES, Cakan MD, et al. Risk of tuberculosis among healthcare workers: can tuberculosis be considered as an occupational disease? *Respir Med.* 2002;96(7):506-510.
34. Anoop M, Thambu D, Kurien T, et al. Risk factors for tuberculosis among health care workers in South India: a nested case-control study. *J Clin Epidemiol.* 2013;66(1):67-74.
35. Menzies D, Fanning A, Yuan L, Fitzgerald M. Tuberculosis Among Health Care Workers. *N Engl J Med.* 1995;332(2):92-98.
36. Blumberg HM, Watkins DL, Berschling JD, et al. Preventing the nosocomial transmission of tuberculosis. *Ann Intern Med.* 1995;122(9):658-663.
37. Drobniewski F, Balabanova Y, Nikolayevsky V, et al. Drug-resistant tuberculosis, clinical virulence, and the dominance of the Beijing strain family in Russia. *Jama.* 2005;293(22):2726-2731.
38. Claassens M, van Schalkwyk C, den Haan L, et al. High prevalence of tuberculosis and insufficient case detection in two communities in the Western Cape, South Africa. *PLoS One.* 2013;8(4):e58689.
39. Balt E, Durrheim DN, Weyer K. Nosocomial transmission of tuberculosis to healthcare workers in Mpumalanga. *S Afr Med J* 1998;88:1363-1366.

40. Kranzer K, Bekker LG, van Schaik N, Thebus L, Dawson M et al. Community health care workers in South Africa are at increased risk for tuberculosis. *SAMJ*. 2010;100(4):224-226.
41. Bjerregaard-Andersen M, da Silva ZJ, Ravn P, et al. Tuberculosis burden in an urban population: a cross sectional tuberculosis survey from Guinea Bissau. *BMC Infect Dis*. 10:96.
42. Eyob G, Gebeyhu M, Goshu S, et al. Increase in tuberculosis incidence among the staff working at the Tuberculosis Demonstration and Training Centre in Addis Ababa, Ethiopia: a retrospective cohort study (1989-1998). *Int J Tuberc Lung Dis*. 2002;6(1):85-88.
43. Kanyerere HS and Salaniponi FM. Tuberculosis in health care workers in a central hospital in Malawi. *Int J Tuberc Lung Dis*. 2003;7(5):489-492.
44. Cramm JM, Koolman X, Moller V and Nieboer AP. Socio-economic status and self-reported tuberculosis: a multilevel analysis in a low-income township in the Eastern Cape, South Africa. *J Public Health Africa*. 2011;2(e34):143-146.
45. Lienhardt C. From exposure to disease: the role of environmental factors in susceptibility to and development of tuberculosis. *Epidemiol Rev*. 2001;23(2):288-301.
46. Harling G, Ehrlich R and Myer L. The social epidemiology of tuberculosis in South Africa: a multilevel analysis. *Soc Sci Med*. 2008;66(2):492-505.
47. Wood R, Lawn SD, Caldwell J, et al. Burden of new and recurrent tuberculosis in a major South African city stratified by age and HIV-status. *PLoS One*. 2011;6(10):e25098.
48. Connelly D, Veriava Y, Roberts S, et al. Prevalence of HIV infection and median CD4 counts among health care workers in South Africa. *S Afr Med J*. 2007;97(2):115-120.
49. O'Donnell, Jarand J, Loveday M, Padayatchi N, Zelnick J et al. High Incidence Of Hospital Admissions with Multidrug and Extensively Drug Resistant Tuberculosis among South African Health Care Workers. *Ann Intern Med*. 2010;153(8):516-522.

50. Shisana O, Hall EJ, Maluleke R, Chauveau J and Schwabe C. HIV/AIDS prevalence among South African health workers. *S Afr Med J.* 2004;94(10):846-850.
  
51. Nava-Aguilera E, Andersson N, Harris E, et al. Risk factors associated with recent transmission of tuberculosis: systematic review and meta-analysis. *Int J Tuberc Lung Dis.* 2009;13(1):17-26.



## APPENDIX

Study questionnaire (see attachment)

## TUBERCULOSIS RISK QUESTIONNAIRE

[FOR SCREENING OF WORKERS IN TYGERBERG HOSPITAL (TBH)]

This study examines possible risk factors for TB in employees of Tygerberg Hospital. We want to use this information to make the workplace safer. Please, complete the questionnaire as accurately as you can. **This questionnaire is for study purposes ONLY and will not be shared with any other person.** Your name should not appear on this questionnaire.

DATE: \_\_\_\_\_

Study number \_\_\_\_\_

1. What is your gender? Male  Female

2. How old are you? \_\_\_\_\_ years Date of birth: day \_\_\_\_\_ month  year

3. What is your race? Black/ African  Coloured  White  Indian

Others (specify) \_\_\_\_\_

4. What is your Height  Weight

5. Did your weight change significantly in the past 4 years (if yes, describe)

	Yes/Increased	Yes/Decreased	No	Don't Know
2008	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
2009	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
2010	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
2011	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

6. What was your marital status in the following years? e.g. single, married, divorced

•2008

•2009

•2010

•2011

7. What is the highest level of education that you have COMPLETED?

None  Primary school  matric  diploma

Degree  postgraduate

Other (specify): -----

8. What was your occupation in the following years (e.g. Nurse, student, etc.):

•2008

•2009

•2010

•2011

9. Please list all your suburbs of residence in the past four years e.g. Khayelitsha, Boston, Kraaifontein, Parow, etc.

•2008

•2009

•2010

•2011

10. How long have you worked (your working career) in the health services?

- Months
- Years

11. How long have you been working as a TBH employee?

- Months
- Years

12. List all the departments that you have worked in at TBH in the last four years e.g.

- 2008
- 2009
- 2010
- 2011

13. On the average, how many hours did you work in TBH during a typical workday?

- 2008
- 2009
- 2010
- 2011

14. On average how many hours did you work in another health facility (other than TBH) every day (or every week) (e.g. for extra income/ “moonlighting”, etc.)?

	Per day	per week
•2008	<input type="text"/>	<input type="text"/>
•2009	<input type="text"/>	<input type="text"/>
•2010	<input type="text"/>	<input type="text"/>
•2011	<input type="text"/>	<input type="text"/>

15. Did your major daily work activities in TBH in the last 4 years involve;

- close contact with patients Yes/ No
- close contact with patients’ specimens Yes/ No
- close contact with both patients and patients’ specimens Yes/ No
- None of the above

16. How would you describe your daily work activities in the past four years?

2008	Worked mainly day shift <input type="text"/>	Night shift <input type="text"/>	Both day and night shift <input type="text"/>
2009	Worked mainly day shift <input type="text"/>	Night shift <input type="text"/>	Both day and night shift <input type="text"/>
2010	Worked mainly day shift <input type="text"/>	Night shift <input type="text"/>	Both day and night shift <input type="text"/>
2011	Worked mainly day shift <input type="text"/>	Night shift <input type="text"/>	Both day and night shift <input type="text"/>

17. On average, estimate how many patients (or patient specimens) you came into contact with during a typical day at work in the following years:

- 2008
- 2009
- 2010
- 2011

18. Have you been diagnosed with TB in the past four years?

•2008;	Yes	<input type="text"/>	No	<input type="text"/>
•2009;	Yes	<input type="text"/>	No	<input type="text"/>
•2010;	Yes	<input type="text"/>	No	<input type="text"/>
•2011;	Yes	<input type="text"/>	No	<input type="text"/>

19. Have you been involved in physical exercise for the purpose of improving your fitness in the following years? (if yes, how often were you involved in fitness exercises?)

	Yes/Daily	Yes/Weekly	Yes/Occasionally	No
•2008;	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
•2009;	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
•2010;	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
•2011;	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

20. How many persons lived with you (in the same house) in the following years?

(e.g. 0, 1, 2, 3 etc.)

- 2008
- 2009
- 2010
- 2011

-How many persons did you share your room with?

- 2008
- 2009
- 2010
- 2011

21. How many sexual partners did you have in the following years?

	0	1	2	3	4	5	>5
• 2008	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
• 2009	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
• 2010	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
• 2011	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

22. Do you suffer from flu almost every year? Yes  No

23. How much alcohol do you normally consume? (1 unit = 1 beer or 1 glass of wine or 1 tot of spirits)

- Nothing
- Less than 1 unit per month
- 1 unit per month
- 1 unit per week
- 1 unit per day
- >1 unit per day

24. Smoking History (complete all that applies to you and delete as appropriate)

- I never smoked
- I am a current smoker
- I used to smoke but stopped  days/weeks/months/years ago
- I have smoked for  days/weeks/months/years-(time)
- I currently smoke/ when I smoked before, I used to smoke
  - Less than 10 sticks per day
  - Between 10 and 20 sticks per day
  - More than 20 sticks per day

25. Recreational drug history e.g. tik, cocaine, mandrax, dagga, etc. (complete all that applies to you and delete as appropriate)

- I never used drugs
- I currently use drugs
- I used to use drugs but stopped  days/weeks/months/years ago
- I have used drug for  days/weeks/months/years
- I currently use/ I used to use drugs
  - Every day or almost every day
  - Every week or almost every week
  - Every month or almost every month
  - Less than once per month



26. Tick appropriately, if you have been diagnosed with any of the following diseases and indicate the year of diagnosis.

Diseases		Year of diagnosis		
HIV	Yes <input type="checkbox"/>	No <input type="checkbox"/>	<input type="checkbox"/>	<input type="text"/>
Diabetes	Yes <input type="checkbox"/>	No <input type="checkbox"/>	<input type="checkbox"/>	<input type="text"/>
Arthritis	Yes <input type="checkbox"/>	No <input type="checkbox"/>	<input type="checkbox"/>	<input type="text"/>
Renal disease	Yes <input type="checkbox"/>	No <input type="checkbox"/>	<input type="checkbox"/>	<input type="text"/>
Asthma	Yes <input type="checkbox"/>	No <input type="checkbox"/>	<input type="checkbox"/>	<input type="text"/>
COPD	Yes <input type="checkbox"/>	No <input type="checkbox"/>	<input type="checkbox"/>	<input type="text"/>
Cancer	Yes <input type="checkbox"/>	No <input type="checkbox"/>	<input type="checkbox"/>	<input type="text"/>
Allergies	Yes <input type="checkbox"/>	No <input type="checkbox"/>	<input type="checkbox"/>	<input type="text"/>
Hypertension	Yes <input type="checkbox"/>	No <input type="checkbox"/>	<input type="checkbox"/>	<input type="text"/>
Silicosis	Yes <input type="checkbox"/>	No <input type="checkbox"/>	<input type="checkbox"/>	<input type="text"/>
Others (specify)-----				<input type="text"/>

27. Have you had a significant stressful period in your life in the last 4 years?

Yes  No

• If yes, please describe the event in the specific year, including the duration thereof:

• 2008;-----

-----

• 2009;-----

-----

• 2010;-----

-----

• 2011;-----

-----

28. Have you lived with, worked closely with or attended to any person that you KNOW was diagnosed with TB in the past 4 years? (If yes, where did the contact occur?)

	Yes/at home	Yes/ at TBH	Yes/other places	No
• 2008;	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
• 2009;	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
• 2010;	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
• 2011;	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

Other (specify) -----

29. How do you normally get to work?

- Private car
- Taxi
- Major Bus
- Walking
- Bicycle
- Other (please specify)-----

When you come to work, how long does it normally take you? \_\_\_\_\_

30. What do you normally use to prepare your food at home?

- Electric stove
- Paraffin stove
- Fire / wood
- Coal
- Others (specify)-----

31. Please, list your main Hobbies

-----  
 -----  
 -----  
 -----

32. Please describe in your own words what tuberculosis (TB) is by completing the following:

TB is ----- and it is caused by -----

TB is transmitted through Air  Water  Sex

Others (specify)-----

33. Do you think you work in an area where there is a high risk for getting TB?

Yes  No  n't know

Name the area

34. Do you use any form of protection against TB daily when at work?

Yes

If yes; what do you use? Apron  Gloves  Face-cover

Others (specify)-----

35. Is there someone in the workplace in the past 4 years who reminded you frequently on the need to protect yourself against TB? (If yes, how often did this person remind you?)

	Yes/Daily	Yes/Weekly	Yes/Occasionally	No
2008	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2009	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2010	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2011	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

36. Have you received training on the following?

• How to prevent TB? Yes  No

• How to use personal protective equipment (PPE)? Yes  No

-How do you feel (what is your personal opinion) about using PPE?-----  
-----  
-----

- Do you use them regularly? Yes  No

37. Have you ever received (BCG) vaccination against TB before?

Yes  No  don't know

If yes, how long ago did you last receive the vaccination?

nths  ars

38. Did you receive the Flu vaccine during the following years?

• 2008; yes  No  Don't know

• 2009; yes  No  Don't know

• 2010; yes  No  Don't know

• 2011; yes  No  Don't know

Thank you for your participation!!!