

Manuscript ID ISP-20-0819

Title: The level of formal support received by people with severe mental illness living in supported accommodation and participation: a systematic review.

Authors

Akkara Lionel Jose

Ajose2@qmu.ac.uk

School of Health Sciences, Queen Margaret University, Queen Margaret University Drive, Edinburgh EH21 6UU, UK.

Dr Michele Harrison,

MHarrison@qmu.ac.uk

School of Health Sciences, Queen Margaret University, Edinburgh, UK.

ORCID ID: <https://orcid.org/0000-0001-6088-2998>

Dr Anusua Singh Roy

ASinghRoy@qmu.ac.uk

School of Health Sciences, Queen Margaret University, Edinburgh, UK.

ORCID ID:

Dr Linda Irvine Fitzpatrick,

Linda.irvinefitzpatrick@nhslothian.scot.nhs.uk

Strategic Programme Manager, Mental Health and Wellbeing, City of Edinburgh Health and Social Care Partnership, Edinburgh, UK

Professor Kirsty Forsyth

KForsyth@qmu.ac.uk

School of Health Sciences, Queen Margaret University, Edinburgh, UK

ORCID ID: <https://orcid.org/0000-0002-6732-1699>

Abstract

Aim

The review aimed to identify and explore the association of level of support received by people with severe mental illness in supported accommodation and participation.

Method

The authors conducted a systematic search in MEDLINE, PsychINFO, PsychARTICLES, CINAHL Plus and ASSIA. Searches were restricted to articles published in English and participants aged 18 years and over with severe mental illness. Articles were included based on level of support received in mental health supported accommodation, classified according to the Simple Taxonomy for Supported Accommodation, and three factors of participation: social participation, daily living functioning and personal empowerment. Studies of in-patient settings and nursing homes were excluded. The review protocol is registered on PROSPERO (registration number: CRD42019161808).

Results

Six articles were included in the review from USA, Australia, Sweden, and Taiwan. Factors of participation for people living in accommodation with moderate support and accommodation with high support were explored. Data indicated an association between level of support and participation showing that people living in accommodation with moderate support had increased participation compared to people living in accommodation with high support.

Conclusion

This review identified an association between level of formal support and participation. People with SMI living in accommodation with medium support participated in more community occupations, more activities and had a higher level of personal empowerment than people living in accommodation with high support.

Key words: Supported Accommodation, Social Participation, Empowerment, Daily Living Functioning, Severe mental illness

Abstract word count – 223 words

Article Manuscript Word count– 4147 words

Introduction

People with severe mental illness (SMI) have diagnoses such as schizophrenia, personality disorders, bipolar disorder and other psychosis-related disorders and have a range of complex needs which impact on different aspects of their everyday life. Supported accommodation (SA) provides residential, community-based support for individuals with SMI (McPherson et al. 2018a). SA provides individuals with SMI the opportunity to obtain a tenancy while receiving varying levels of staff support within the least restrictive settings in order to develop skills and abilities needed to participate in various daily living and social activities (Padmakar et al. 2020). SA can differ by type, staffing location, level of support provided and emphasis within the accommodation on moving on (McPherson et al. 2018b). Within SA, the support people with SMI receive is typically provided by formal carers inclusive of healthcare professionals, carers or other staff providing support.

Participation can vary over a person's lifetime dependant on life events affecting the person's confidence, abilities and motivation (Sánchez et al. 2016). For people with SMI, this can affect maintaining and creating relationships with friends and family (Cruce et al. 2012), how they engage with the support they receive and being in education or work and pursuing interests (Tjörnstrand et al. 2013). Participation therefore has several elements. These include engaging in daily living activities (self-care, meal planning and preparation, dressing, money management, medication management (Piškur et al. 2014); social participation, an individuals' involvement in society (Sanches et al. 2019) through roles they engage in within a group or in their community (Kaplan et al. 2012), including employment and vocational activities (van Eijk-Hustings et al. 2013), social functioning (Tobin et al. 2013) and building and maintaining relationships (Berkman, 2011); and personal empowerment, the feeling or sense of control an individual has over their own life alongside the level of responsibility and autonomy they possess to initiate and act on aspects of their participation (Bruschetta & Barone, 2016; Cavalieri & Almeida, 2018).

Literature focused on SA suggest when accommodation types or treatment environments are appropriate to people's needs there are improvements in activities of daily living and social participation (Siskind et al. 2012). It has also been shown that the therapeutic relationship between people living in SA and formal carers can improve social participation (Amati et al. 2017; Brunt & Rask, 2018; Krotofil et al. 2018). This results in increased personal and social responsibility for the individual and improved social functioning (Dixon et al. 2016; Green et al. 2009; Hitch et al. 2013). Previous systematic reviews have focused on the effect of the built and physical environment on mental health (Charlotte et al. 2007; Moore et al. 2018) and the impact of social climate, service delivery and quality of life for people with SMI living in SA. These factors have been shown to affect how care provided meets the person's needs (Macpherson et al. 2004), people with SMI's experience and satisfaction with SA (Harrison et al. 2020; Krotofil et al. 2018) and the impact on individual's feelings of stability and independence (Burgoyne, 2014). Reviews also focused on factors such as SA's links to psychosocial outcomes (McPherson et al. 2018a), quality or effectiveness of service delivery style (Rogers et al. 2010) and standardising service delivery models (Parker et al. 2019; Tabol et al. 2010). There is, however, no systematic review that considers formal support for people with SMI living in SA and its association with factors of participation.

The systematic review aimed to review formal care provided to people with SMI living in SA and its association with factors of participation specifically daily living functioning, social participation, and personal empowerment.

Methods

The following review protocol is registered on PROSPERO (registration number is CRD42019161808.) and follows the PRISMA guidance.

Eligibility Criteria

The review included articles published in academic journals with quantitative data relevant to the three participation factors: daily living functioning, social functioning and personal empowerment. Inclusion criteria were adults with SMI living in SA, receiving support from formal carers (nurses, paid carers and/or any health care professionals) and informal carers (family, friends or unpaid carers). Dissertations, book chapters, guidelines, policy and conference proceedings were excluded from the study. Studies reporting on people under 18 years old and those that were not published in English were excluded, however, no exclusions were made based on country of publication. Studies within in-patient settings, nursing homes and SA that was not being provided to people with SMI were excluded from the review.

Population

This review includes people with SMI aged 18 and above. The term SMI extends to the DSM-IV definition and includes the following conditions: schizophrenia, bipolar disorder, personality disorder or other psychosis-related disorders. Diagnoses were reviewed during the screening process to comply with inclusion criteria. Studies were excluded if they reported solely on the following diagnoses: substance misuse, eating disorders, learning/intellectual disability or dementia.

Supported Accommodation

SA was classified using the Simple Taxonomy for Supported Accommodation (STAX-SA) (McPherson et al. 2018b) which defines accommodation types by staffing location (on or off site), level of support (high/moderate/low/no), emphasis on move-on (limited or strong) and physical setting (congregate or individuals).

Formal Support

Formal support was defined using the STAX_SA level of support domain. The four levels of support (high/moderate/low/none) describe the frequency, nature and intensity of support (including staffing duration) required to meet service user need (e.g., for personal care, medication management). Studies identified with moderate support where available staff were identified as on or off site were combined for the purpose of the review.

Comparator

The search strategy reflects the authors' initial aim to explore formal care compared with informal care. Due to the limited information and lack of consistency of informal support provision detailed within studies, this comparator was not used. Studies were instead compared according to level of support and their association with participation for people with SMI living in SA.

Outcomes

Three participation factors, social participation, daily living functioning and personal empowerment were identified, with reported data in the included studies matched to these three factors (see Table 1). Detailed information about how factors were matched and measures used in included articles is available in Supplementary Material 1.

Search Strategy

An electronic database search was conducted between October 2019 and February 2020 using MEDLINE, Psychinfo, CINAHL Plus, ASSIA and PsychARTICLES. Alongside this, previous reviews were hand-searched to identify any relevant articles. The searches included a combination of MeSH terms and Boolean phrases that matched the population, interventions, comparators and outcomes. These included, but were not limited to, 'Mental health difficult*', 'Shared accommodation', 'social interaction/engage*', 'formal care provision', 'Formal Support', 'informal support', 'Factors of participation', 'improved skills and abilities' and 'engage*'. No time limit was placed on publication

date however articles were limited to population ages of 18 and above. The full search strategy used within the databases is detailed in Supplementary material 2.

Data Extraction

Data was extracted according to a form developed by ALJ, and included 1) Study title, year, location, study type; 2) Sample size, age, gender, condition/ inclusion criteria; 3) Accommodation type, support type (formal or informal); 4) Factors of participation, measures used, control/comparators; 5) Statistical analysis and findings/results. The extracted data was synthesised to include identified inclusion criteria and are detailed in Table 1. Statistical information pertaining to accommodation types with relevant level of support was included and is detailed in Table 2.

Insert Table 1: Data extraction table for included studies

Insert Table 2: Data extraction of results from selected studies

Quality Assessment

The quality of the 4 observational studies was assessed using the Research Triangle Institute (RTI) Item Bank (Viswanathan & Berkman et al. 2011; Viswanathan et al. 2013) due its ability to comprehensively assess bias (selection, performance, detection and confounding) across varying types of observational studies. 11 questions were selected as appropriate to assess the risk of bias for the included studies. Studies with 1 or more negative score were recorded as having a high risk of bias and those which scored 1 or more 'partially' or 'cannot determine' were recorded as having an unclear risk of bias (Viswanathan & Berkman, 2011). The ROBINS-I tool was used for the 2 quasi-experimental studies selected (Sterne et al. 2016). The tool is an update to the Cochrane collaboration risk of bias tool assessing 7 domains of bias at the pre-intervention, at-intervention and post-intervention stages of a study (Sterne et al. 2016). Risk of bias of individual studies was assessed independently by ST and ALJ. Overall scores are presented in Table 1 and detailed results are available in Supplementary Material 3.

Data Synthesis

Data collected could not be synthesised within a meta-analysis due to inconsistency of data reporting, unavailability of data required to calculate a common effect size (Cohen's d) and use of unstandardized measures of participation with no evidence of reliability or validity testing. Contacting authors for additional information or data was unsuccessful due to no response or the author no longer possessing the original data. Alternative methods to a meta-analysis recommended in the Cochrane guidelines (Deeks et al. 2019, Higgins et al. 2019) were used to include statistical data to support the systematic review, by calculating actual or estimated effect sizes where possible. Data synthesis was completed by ALJ. As seen in Table 3, data available was used to calculate Cohen's d (Dorer et al. 2009; Fossey et al. 2006) and a raw mean difference (Nelson et al. 1997) alongside the ANOVA value (Nelson et al. 1997) and a regression coefficient (Shu et al. 2001). These were used to estimate the magnitude, direction and statistical significance of association between level of support and level of participation as well as the association of the participation factors within specific levels of support. Cohen's d was calculated for one study (Fossey et al. 2006) using the 'dmetar' package (Harrer et al. 2019) in R (R Core Team, 2013), as t-tests were calculated for sub-group difference in this study. Reporting of data uses the SWIM guidelines (Campbell et al. 2020) which provides additional structure for reporting of the narrative synthesis of the systematic review while adhering to the PRISMA checklist.

Insert Table 3: Calculated effect sizes

Results

Study Selection

An initial search using the search strategy above presented 7892 articles from Medline, PsychINFO, PsychARTICLES, CINAHL using EBSCOHOST and 222 from ASSIA. After adding filters for age and language the results were 3948 (EBSCOHOST) and 221 (ASSIA). Duplicates were then removed from the initial search resulting in 1270 results. Following this titles and abstracts were screened and the results of this were reviewed for relevance by ST. ST reviewed 10% against the inclusion/exclusion criteria. Discrepancies (less than 5%) between ST's and ALJ's results were discussed and resolved. The screening process can be seen in Figure 1.

Insert Figure 1: PRISMA Flow diagram.

Following title, abstract and full-text screening the review identified 6 articles that addressed association between participation and at least two levels of support received by individuals with SMI. The review identified 4 observational studies using cross-sectional data and 2 quasi-experimental studies using longitudinal data. 1 of the observational studies was secondary analysis of cross-sectional data (ref). Studies were conducted in America (n=1), UK (n=1), Australia (n=1), Sweden (n=1), Taiwan (n=1) and Canada (n=1) (see Table 1).

Quality Appraisal

Heterogeneity was identified across the observational studies due to the difference in participants across the studies, with two studies reporting data from people with SMI only (Eklund & Tjörnstrand, 2019; Fossey et al. 2006); one study reporting data from staff only (Dorer et al. 2009) and one study reporting data from both people with SMI and staff (Kruzich & Berg, 1985). 2 studies were analyses of secondary data (Eklund & Tjörnstrand, 2019) (Kruzich & Berg, 1985). None of the included studies shared a common measurement tool. Length of stay in accommodation was only reported in 2 studies (Eklund and Tjörnstrand 2019; Nelson et al. 1997) so its potential effect on participation could not be considered.

Two studies (Eklund & Tjörnstrand, 2019; Fossey et al. 2006) were rated at low risk of bias; Kruzich and Berg's (1985) study was rated "Unclear" due to the recruitment strategy and attrition rate not being reported. Dorer et al.'s study (2009) was rated at high risk of bias as they did not use a validated measure. The studies were not excluded as Kruzich and Berg (1985) reported an internal consistency reliability coefficient and the measures used in Dorer et al.'s (2009) study were informed by existing standardized measures to improve content validity. Both quasi-experimental studies were found to have a moderate risk of bias. Nelson et al.'s (1997) study scored moderate on confounding bias due to measures for recording confounding variables being subjective, suggesting a higher risk of confounding bias. Shu et al.'s (2001) study scored a moderate in relation to reporting bias, as specific outcome data presented was not clearly labelled, so interpretation of values is assumed rather than stated.

Participation Factors

Social Participation

Social participation was the most frequently identified factor present within 5 out of 6 studies. 5 studies presented data comparing accommodation with high support (AHS) and accommodation with moderate support (AMS) and their association with social participation. Social participation described in these studies included people's involvement in roles related to education and employment, frequency of participation in social activities during the week and time spent in social activities. The social participation factors reported show overall that people with SMI living in AMS participated in more community activities than those living in AHS, identifying more social roles (Nelson et al. 1997) and social contacts (Fossey et al. 2006) including visiting family and friends more frequently (Dorer et

al. 2009). Actual and calculated effect sizes estimated demonstrated that people's social participation was statistically significant in relation to higher level of engagement in social activities (Shu et al. 2001). The combined effect size of time spent in community occupations in Dorer et al.'s (2009) study was statistically significant, showing a small effect size and people with SMI in AMS identified more social roles (Nelson et al. 1997) and had more social contacts (Fossey et al. 2006). Two elements were the exception to this inference; 'faith' in Dorer et al.'s (2009) study and 'day centre' in Eklund and Tjörnstrand's (2019) study with individuals in AHS spending more time participating in these social activities than those in AMS.

Only one study (Nelson et al. 1995) compared social participation between a combined value of AHS and AMS with accommodation with no support (ANS). A mixed two-way ANOVA produced a statistically significant difference ($p < 0.05$) that suggests those in AHS and AMS identify more social roles than those in ANS.

Daily Living Functioning

Daily living function was identified in four of the studies (Kruzich & Berg, 1985; Nelson et al. 1997; Fossey et al. 2006; Eklund & Tjörnstrand, 2019) and compared between AMS and AHS. The direction of inference across all but one study (Eklund & Tjörnstrand, 2019) suggests that people with SMI living in AMS participate in more daily living activities than those in AHS. While no statistically significant association was identified between support type and elements of daily living functioning, values calculated and reported support this direction of inference. Eklund and Tjörnstrand's (2019) study differs by reporting that 11% more individuals in AHS engage in household chores compared to those in AMS. Two studies reported data comparing a combined value of AHS and AMS with ANS. Kruzich and Berg (1985) calculated a raw mean difference of (0.38) for self-sufficiency, and Nelson et al. (1997) reported values from a two-way mixed ANOVA showing a statistically significant difference in daily living function, with people living in AHS and AMS being more independent in daily living functioning than those living in ANS.

Personal Empowerment

Three studies reported data on personal empowerment in AHS and AMS (Nelson et al. 1997; Shu et al. 2001; Fossey et al. 2006). Personal empowerment identified in these studies included mastery, autonomy and responsibility. Calculated and available data suggests the direction of inference shows that people with SMI living in AMS have higher levels of personal empowerment than those in AHS. Shu et al.'s study (2001) reported an unstandardized regression coefficient for autonomy favouring those in AMS. In Fossey et al.'s (2006) study, effect size d calculated for 'responsibility' showed a small effect size, with no statistical significance for subgroup difference. Nelson et al. (1997) compared mastery, using a combined value for AHS and AMS with ANS. A two-way mixed ANOVA identified a statistical significance which showed that those in AMS and AHS had a higher level of personal empowerment than those in ANS.

Discussion

The review identified an association between participation and level of support, particularly when comparing AHS and AMS. Social participation was the most frequently reported, followed by daily living functioning and personal empowerment. All factors demonstrated favourable results for people with SMI living in AMS who had higher levels of participation than those living in AHS, with few identified discrepancies in this direction of inference. Only two studies (Kruzich & Berg, 1985; Nelson et al. 1997) reported on ANS and compared this with a combined value for AHS and AMS, suggesting higher levels of social participation and daily living functioning in accommodation with support when compared to ANS.

The review identified that people with SMI living in AMS had higher levels of social participation. The studies included in this review suggest that those in AMS received more staff support to socially participate, particularly to attend community centres (Dorer et al. 2009; Eklund & Tjörnstrand, 2019) and enrol in vocational activities (Nelson et al. 1997) than those in AHS. Previous research suggests this may be due to range of factors including how services are structured, particularly facility size, whether staff are based on or off site, intensity of support provided and whether there is a focus on moving on to more independent living (Dalton-Locke et al. 2018; Hansson et al. 2002; Macpherson et al. 2004; Muir et al. 2010, Webber & Fendt-Newlin, 2017). It is reported that people with SMI living in AMS have higher levels of choice and freedom when compared to people living in AHS (Eklund & Tjörnstrand, 2019; Nelson et al. 1997). There is also discussion in the literature about whether higher levels of participation for people with SMI living in AMS are due to people having less complex needs including experiencing fewer symptoms and being on less medication (Segal et al. 1989; Shu et al. 2001, Killaspy et al. 2019), resulting in greater motivation to participate in activities (Nelson et al. 1997). However other studies have shown that level of disability in supported accommodation is comparable regardless of level of support received (Trauer, 2001; Trauer et al. 1997). Across the studies reviewed, the level of participation in employment or some form of education is low. This is reported in other studies of people with SMI living in supported accommodation (Bitter et al. 2016; Killaspy et al. 2016; Mirza et al. 2008). It is recognised that employment and education are important for social functioning for people with SMI (Modini et al. 2016) however the indication from this review is that this remains an area of social participation that is not available to many people with SMI when living in supported accommodation.

The review suggests that people with SMI living in AMS had higher levels of participation in daily living activities than those living in AHS. The minimal difference in mean scores and correlation statistics reviewed for daily living functioning factors between AMS and AHS is interesting as there is an increased focus in AMS on rehabilitation and increasing independence in daily living skills (Brunt & Hansson, 2002; Killaspy et al. 2016; Krotofil et al. 2018). All the studies described staff support in AHS as providing more guidance and support around daily living activities to people with SMI, with them receiving staff assistance with activities or high levels of prompting (Fossey et al. 2006; Kruzich & Berg, 1985; Nelson et al. 1997). Eklund and Tjörnstrand's (2009) study reported that people living in AHS participated in more household chores than those living in AMS. This may be due to people living in AHS spending more time in the accommodation as they had lower levels of social participation than people living in AMS, resulting in daily living activities being the main focus of daily time use. When results for daily living functioning in AHS and AMS were combined and compared to ANS, a positive association was demonstrated between accommodation with support and participation in daily living activities. It is generally assumed that people living in accommodation with no support are independently participating in all daily living activities (Trauer, 2001). However, research has shown that people with SMI receiving no or low support can experience difficulties organising daily living activities (Eklund et al. 2017).

Personal empowerment is the least explored factor among the selected studies. Results indicate a higher level of personal empowerment reported by people living in AMS compared to those living in AHS. Personal empowerment is an important aspect of recovery for people with SMI (Leamy et al. 2011). This review suggests people with SMI's experience of personal empowerment will vary depending on the level of support they receive in supported accommodation, with studies reporting increased personal empowerment as the level of support decreases. This may be indicative of peoples' perceptions of their own abilities and growth in confidence over time with people living in AMS in Nelson et al. (1997) and Shu et al.'s (2001) studies experiencing higher levels of personal empowerment as a result of increased independence and recovery. However, Nelson et al.'s (1997) study showed that personal empowerment related to skill mastery was lower in ANS compared to AMS and AHS. Research has shown that staff attitudes towards recovery influence people with SMI's level of participation, inhibiting their recovery (Bitter et al. 2017; Linhorst et al. 2005; Macpherson et

al. 2004; Pandiani et al. 1994), and likelihood of moving on to more independent living (Killaspy et al. 2013; Killaspy et al. 2019). Other personal and environmental factors can affect an individual's experience of personal empowerment including their illness experience, restrictions imposed by compulsory treatment orders and rules within SA which can restrict choices and involvement in decision making (Brolin et al. 2018; Fossey et al. 2006; Nelson et al. 2001; Sandhu et al. 2017; Valdes-Stauber and Kilian 2018). These environmental factors mean that staff have to uphold rules while also supporting individuals, limiting flexibility of approach which can hinder effective support for recovery (Bengtsson-Tops et al. 2014; Coffey et al. 2019; Nelson et al. 2007).

There is no indication in the included studies if participation in daily living, social participation and personal empowerment were assessed prior to people living in SA to inform decisions about which type of SA an individual moved to. Previous studies have shown that healthcare professionals can overestimate the level of support people with SMI require, which often differs from what the individual identifies as needing (Afilalo et al. 2012; Lasalvia et al. 2015; Piat et al. 2015). There is limited reporting on how an individual's level of participation is considered when selecting SA, resulting in people with differing participation needs residing in the same types of SA. This can create a disparity between individuals' needs, type of support provided and the extent to which people's participation is enabled (de Heer Wunderink et al. 2012; Sanches et al. 2019).

Limitations

The number of studies included in the review are small and highlight that there is limited published research available focusing on formal support on participation for people with SMI living in SA. Due to the lack of appropriate data, the authors were unable to conduct a meta-analysis by estimating overall effect sizes. Instead, unstandardized effect sizes such as raw mean difference were used to explore if there was an association between level of support and participation for people with SMI, affecting the robustness of the results. The original aim of the review was to compare the impact of informal and formal care on people with SMI living in SA. The role of informal carers is under explored in current literature, even though informal care networks such as family involvement (Allen et al. 2013; Dorer et al. 2009; Fossey et al. 2006) or supportive neighbourhoods (Kriegel et al. 2019) are indicated as beneficial to people with SMI's social participation and recovery. The review focuses on level of support and the authors acknowledge there are other factors that can influence participation for people with SMI living in SA including whether people are living in congregate settings or alone, and length of stay in accommodation which needs to be a consideration for future reviews.

Conclusion

This review identified an association between participation factors and level of formal support for people with SMI, between accommodation with moderate support and accommodation with high support. People living in accommodation with medium support participated in more community occupations, a higher number of daily living activities and experienced greater personal empowerment. The results suggest that further exploration of how formal and informal support enables participation for people with SMI in SA to support their recovery is needed.

Acknowledgements

The authors would like to thank Sumeyra Tayfur for assisting with screening and quality appraisal of studies.

Declaration of Conflict of Interest

The Authors declare that there is no conflict of interest.

Funding

This work was supported by a PhD scholarship from Queen Margaret University and National Health Service (NHS) Lothian.

References

- Afilalo, M., Soucy, N., Xue, X., Colacone, A., Jourdenais, E., & Boivin, J-F. (2015). Characteristics and needs of psychiatric patients with prolonged hospital stays. *Canadian Journal of Psychiatry, 60*, 181-188. <https://doi.org/10.1177/070674371506000405>
- Allen, J., Burbach, F., & Reibstein, J. (2013). 'A different world' individuals' experience of an integrated family intervention for psychosis and its contribution to recovery. *Psychology and psychotherapy, 86*(2), 212-28. <https://doi.org/10.1111/j.2044-8341.2011.02057.x>
- Amati, F., Banks, C., Greenfield, G., & Green, J. (2017). Predictors of outcomes for patients with common mental health disorders receiving psychological therapies in community settings: a systematic review. *Journal of Public Health, 40*(3), e375-e387. <https://doi.org/10.1093/pubmed/fox168>
- Bengtsson-Tops, A., Ericsson, U., & Ehliasson, K. (2014). Living in supportive housing for people with serious mental illness: a paradoxical life. *International Journal of Mental Health Nursing, 23*(5), 409-418.
- Berkman, L. (2001). Social Integration, Social Networks, and Health. *International Encyclopedia of the Social & Behavioral Sciences, 14327-14332*. 10.1016/B0-08-043076-7/03820-1
- Bitter, N., Roeg, D. P. K., van Nieuwenhuizen, C. & van Weeghel, J. (2016) Identifying profiles of service users in housing services and exploring their quality of life and care needs. *BMC Psychiatry, 16*, 419. <https://dx.doi.org/10.1186%2Fs12888-016-1122-0>
- Bitter, N., Roeg, D., van Assen, M., van Nieuwenhuizen, C., & van Weeghel, J. (2017). How effective is the comprehensive approach to rehabilitation (CARE) methodology? A cluster randomized controlled trial. *BMC Psychiatry, 17*(1), 396. 10.1016/B0-08-043076-7/03820-1
- Brolin, R., Syrén, S., Rask, M., Sandgren, A., & Brunt D. (2018) Residents' perceptions of the most positive and negative aspects of the housing situation for people with psychiatric disabilities *Scandinavian Journal of Caring Sciences, 32*, 603-611
- Brunt, D., & Hansson, L. (2002). Comparison of user assessed needs for care between psychiatric inpatients and supported community residents. *Scandinavian Journal of Caring Sciences, 16*(4), 406-413. <https://doi.org/10.1046/j.1471-6712.2002.00085.x>
- Brunt, D., & Rask, M. (2018). Resident and staff perceptions of the content of their relationship in supported housing facilities for people with psychiatric disabilities. *Journal of multidisciplinary healthcare, 11*, 673-681. <https://doi.org/10.2147/JMDH.S179322>
- Bruschetta, S., & Barone, R. (2016). Group-apartments for recovery of people with psychosis in Italy: Democratic therapeutic communities in post-modern social communities. *Therapeutic Communities, 37*(4), 213-226. <http://dx.doi.org/10.1108/TC-03-2016-0008>
- Burgoyne, J. (2014). Mental health and the settings of housing support – a systematic review and conceptual model. *Housing, Care and Support, 17*(1), 26-40. <https://doi.org/10.1108/HCS-10-2013-0018>

- Campbell, M., McKenzie, J., Sowden, A., Katikireddi, S. V., Brennan, S. E., Ellis, S., Hartmann-Boyce, J., Ryan, R., Shepperd, S., Thomas, J., Welch, V. & Thomson, H. (2020). Synthesis without meta-analysis (SWiM) in systematic reviews: reporting guideline. *BMJ (Clinical research ed.)*, 368, l6890. <https://doi.org/10.1136/bmj.l6890>
- Cavaliere, I., & Neves Almeida, H. (2018). Power, Empowerment and Social Participation- the Building of a Conceptual Model. *European Journal of Social Sciences Education and Research*, 12(1), 189. 10.2478/ejser-2018-0020
- Charlotte, C., Rowan, M., Stephen, S., & Bridget, C. (2007). A systematic review of the evidence on the effect of the built and physical environment on mental health. *Journal of Public Mental Health*, 6(2), 14-27. <https://doi.org/10.1108/17465729200700011>
- Coffey, M., Hannigan, B., Meudell, A., Jones, M., Hunt, J., & Fitzsimmons, D. (2019) Quality of life, recovery and decision-making: a mixed methods study of mental health recovery in social care. *Social Psychiatry and Psychiatric Epidemiology*, 54, 715-723 <https://doi.org/10.1007/s00127-018-1635-6>
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences*. Academic Press.
- Cruce, G., Öjehagen, A., & Nordström, M. (2012). Recovery-promoting Care as Experienced by Persons with Severe Mental Illness and Substance Misuse. *International Journal of Mental Health and Addiction*, 10(5), 660-669. <https://doi.org/10.1007/s11469-011-9363-0>
- Dalton-Locke, C., Attard, R., Killaspy, H., & White, S. (2018). Predictors of quality of care in mental health supported accommodation services in England: a multiple regression modelling study. *BMC Psychiatry*, 18(1), 344. <https://doi.org/10.1186/s12888-018-1912-7>
- De Heer-Wunderink, C., Visser, E., Caro-Nienhuis, A., Sytema, S., & Wiersma, D. (2012). Supported Housing and Supported Independent Living in the Netherlands, with a comparison with England. *Community Mental Health Journal*, 48, 321-327.
- Deeks, J., Higgins, J., & Altman, D. (2019). Chapter 10: Analysing data and undertaking meta-analyses. In D. Altman, J. Deeks, J. Higgins, J. Thomas, J. Chandler, M. Cumpston, T. Li, M. Page & V. Welch (Eds.), *Cochrane Handbook for Systematic Reviews of Interventions version 6.0 (updated July 2019)* (6 ed.). Cochrane.
- Dixon, L., Holoshitz, Y., & Nossel, I. (2016). Treatment engagement of individuals experiencing mental illness: review and update. *World psychiatry : official journal of the World Psychiatric Association (WPA)*, 15(1), 13-20. <https://dx.doi.org/10.1002%2Fwps.20306>
- Dorer, G., Harries, P., & Marston, L. (2009). Measuring Social Inclusion: A Staff Survey of Mental Health Service Users' Participation in Community Occupations. *British Journal of Occupational Therapy*, 72(12), 520-530. <https://doi.org/10.4276/030802209X12601857794691>
- Eklund, M., & Tjörnstrand, C. (2019). Associations between occupational and social interaction factors and well-being among people with psychiatric disabilities living in supported housing in Sweden. *Journal of Occupational Science*, 27(1), 54-68. <https://doi.org/10.1080/14427591.2019.1620121>
- Eklund, M., Argentzell, E., Bejerholm, U., Tjörnstrand, C., & Brunt, D. (2017) Wellbeing, activity and housing satisfaction – comparing people with psychiatric disabilities in supported housing and ordinary housing with support. *BMC Psychiatry* 2017, 17, 315. <https://doi.org/10.1186/s12888-017-1472-2>

- Fossey, E., Harvey, C., Plant, G., & Pantelis, C. (2006). Occupational Performance of People Diagnosed with Schizophrenia in Supported Housing and Outreach Programmes in Australia. *British Journal of Occupational Therapy*, 69(9), 409-419. <https://doi.org/10.1177/030802260606900904>
- Green, G., Hayes, C., Dickinson, D., Whittaker, A., & Gilheany, B. (2009). The role and impact of social relationships upon well-being reported by mental health service users: A qualitative study. *Journal of Mental Health*, 11(5), 565-579. <https://doi.org/10.1080/09638230020023912>
- Hansson, C., Middelboe, T., Sørgaard, K., Bengtsson-Tops, K., Bjarnason, O., Merinder, L., Nilsson, L., Sandlund, M., Korkeila, J. & Vinding, H. (2002). Living Situation, Subjective Quality of Life and Social Network Among Individuals With Schizophrenia Living in Community Settings. *Acta psychiatrica Scandinavica*, 106(5). <https://doi.org/10.1034/j.1600-0447.2002.02346.x>
- Harrer, M., Cuijpers, P., Furukawa, T., & Ebert, D. (2019). Doing meta-analysis in R: A hands-on guide. *PROTECT Lab Erlangen*.
- Harrison M, Singh Roy A, Hultqvist J, Pan, A-W, McCartney D, McGuire N, Irvine Fitzpatrick L, Forsyth K. (2020) Quality of life outcomes for people with serious mental illness living in supported accommodation: systematic review and meta-analysis. *Social Psychiatry and Psychiatric Epidemiology*, 55, 977-988. <https://doi.org/10.1007/s00127-020-01885-x>
- Higgins, J., Li, T., & Deeks, J. (2019). Chapter 6: Choosing effect measures and computing estimates of effect. In J. Higgins, T. Li, J. Deeks, & W. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ (Ed.), *Cochrane Handbook for Systematic Reviews of Interventions version 6.0 (updated July 2019)* (6 ed.). Cochrane.
- Hitch, D., Pepin, G., & Stagnitti, K. (2013). Engagement in Activities and Occupations by People Who Have Experienced Psychosis: A Metasynthesis of Lived Experience. *British Journal of Occupational Therapy*, 76(2), 77-86. <https://doi.org/10.4276/030802213X13603244419194>
- Kaplan, K., Salzer, M., & Brusilovskiy, E. (2012). Community participation as a predictor of recovery-oriented outcomes among emerging and mature adults with mental illnesses. *Psychiatric Rehabilitation Journal*, 35(3), 219-229. <https://doi.org/10.2975/35.3.2012.219.229>
- Killaspy, H., Marston, L., Omar, R. Z., Green, N., Harrison, I., Lean, M., Holloway, F., Craig, T., Leavey, G. & King M. (2013) Service quality and clinical outcomes: an example from mental health rehabilitation services in England. *British Journal of Psychiatry*, 202, 28-34. <https://doi.org/10.1192/bjp.bp.112.114421>
- Killaspy, H., Priebe, S., Bremner, S., McCrone, P., Dowling, S., Harrison, I., Krotofil, J., McPherson, P., Sandhu, S., Arbuthnott, M., Curtis, S., Leavey, G., Shepherd, G., Eldridge, S., & King, M. (2016) Quality of life, autonomy, satisfaction, and costs associated with mental health supported accommodation services in England: a national survey. *Lancet Psychiatry*, 3, 1129-1137. [https://doi.org/10.1016/s2215-0366\(16\)30327-3](https://doi.org/10.1016/s2215-0366(16)30327-3)
- Killaspy, H., Priebe, S., McPherson, P., Zenasni, Z., Greenberg, L., McCrone, P., Dowling, S., Harrison, I., Krotofil, J., Dalton-Locke, C., McGranahan, R., Arbuthnott, M., Curtis, S., Leavey, G., Shepherd, G., Eldridge, S. & King, M. (2019) Predictors of moving on from mental health supported accommodation in England: national cohort study. *British Journal of Psychiatry*, 3, 1-7 <https://doi.org/10.1192/bjp.2019.101>

- Kriegel, L., Townley, G., Brusilovskiy, E., & Salzer, M. (2019). Neighbors as distal support for individuals with serious mental illnesses. *American Journal of Orthopsychiatry*, *90* (1), 98-105. <https://doi.org/10.1037/ort0000403>
- Krotofil, J., McPherson, P., & Killaspy, H. (2018). Service user experiences of specialist mental health supported accommodation: A systematic review of qualitative studies and narrative synthesis. *Health & Social Care in the Community*, *26*(6), 787-800. <https://doi.org/10.1111/hsc.12570>
- Kruzich, J., & Berg, W. (1985). Predictors of self-sufficiency for the mentally ill in long-term care. *Community Mental Health Journal*, *21*(3), 198-207. <https://doi.org/10.1007/BF00754735>
- Lasalvia, A., Boggian, I., Bonetto, C., Saggiaro, V., Piccione, G., Zanoni, C., et al. (2012) Multiple perspectives on mental health outcome: needs for care and service satisfaction assessed by staff, patients and family members. *Soc Psychiatry Psychiatr Epidemiol*, *47*, 1035-1045 <https://doi.org/10.1007/s00127-011-0418-0>
- Leamy, M., Bird, V., Le Boutillier, C., Williams, J. & Slade, M. (2011). Conceptual framework for personal recovery in mental health: systematic review and narrative synthesis. *The British Journal of Psychiatry*, *199*(6), 445-452. <https://doi.org/10.1192/bjp.bp.110.083733>
- Linhorst, D., Eckert, A., & Hamilton, G. (2005). Promoting participation in organizational decision making by clients with severe mental illness. *Social work*, *50*(1), 21-30. <https://doi.org/10.1093/sw/50.1.21>
- Macpherson, R., Shepherd, G., & Edwards, T. (2004). Supported accommodation for people with severe mental illness: a review. *Advances in Psychiatric Treatment*, *10*(3), 180-188. <https://doi.org/10.1192/apt.10.3.180>
- Macpherson, R., Shepherd, G., & Thyarappa, P. (2012). Supported accommodation for people with severe mental illness: an update. *Advances in Psychiatric Treatment*, *18*(5), 381-391. <https://doi.org/10.1192/apt.bp.110.008714>
- McPherson, P., Krotofil, J., & Killaspy, H. (2018a). Mental health supported accommodation services: a systematic review of mental health and psychosocial outcomes. *BMC Psychiatry*, *18*(1), 128. <https://doi.org/10.1186/s12888-018-1725-8>
- McPherson, P., Krotofil, J., & Killaspy, H. (2018b). What Works? Toward a New Classification System for Mental Health Supported Accommodation Services: The Simple Taxonomy for Supported Accommodation (STAX-SA). *International journal of environmental research and public health*, *15*(2). <https://doi.org/10.3390/ijerph15020190>
- Middelboe, T., Mackeprang, T., Thalsgaard, A., & Christiansen, P. (1998). A housing support programme for the mentally ill: need profile and satisfaction among users. *Acta Psychiatrica Scandinavica*, *98*(4), 321-327. <https://doi.org/10.1111/j.1600-0447.1998.tb10091.x>
- Mirza M, Gossett A, Chan NK, Burford L, Hammel J (2008) Community reintegration for people with psychiatric disabilities: challenging systemic barriers to service provision and public policy through participatory action research. *Disabil Soc*. *23*, 323-336. <https://doi.org/10.1080/09687590802038829>
- Modini, M., Tan, L., Brinchmann, B., Wang, M.J., Killackey, E., Glozier, N, et al. (2016) Supported employment for people with severe mental illness: Systematic review and meta-analysis of the international evidence. *Br J Psychiatry*, *209*, 14–22. <https://doi.org/10.1192/bjp.bp.115.165092>

- Moore, T., Kesten, J., López-López, J., Ijaz, S., McAleenan, A., Richards, A., Gray, S., Savović, J. & Audrey, S. (2018). The effects of changes to the built environment on the mental health and well-being of adults: Systematic review. *Health & Place*, 53, 237-257. <https://doi.org/10.1016/j.healthplace.2018.07.012>
- Muir, K., Fisher, K., Abello, D., & Dadich, A. (2010). 'I didn't like just sittin' around all day': Facilitating Social and Community Participation Among People with Mental Illness and High Levels of Psychiatric Disability. *Journal of Social Policy*, 39(3), 375-391. <https://doi.org/10.1017/S0047279410000073>
- Nelson, G., Hall, G., & Walsh-Bowers, R. (1997). A comparative evaluation of supportive apartments, group homes, and board-and-care homes for psychiatric consumer/survivors. *Journal of Community Psychology*, 25(2), 167-188. [https://doi.org/10.1002/\(SICI\)1520-6629\(199703\)25:2<167::AID-JCOP6>3.0.CO;2-V](https://doi.org/10.1002/(SICI)1520-6629(199703)25:2<167::AID-JCOP6>3.0.CO;2-V)
- Nelson, G., Lord, J., & Ochocka, J. (2001). Empowerment and mental health in community: narratives of psychiatric consumer/survivors. *Journal of Community & Applied Social Psychology*, 11(2), 125-142. <https://doi.org/10.1002/casp.619>
- Nelson, G., Sylvestre, J., Aubry, T., George, L., & Trainor, J. (2007). Housing choice and control, and control over professional support as contributors to the subjective quality of life and community adaptation of people with severe mental illness. *Administration and Policy in Mental Health and Mental Health Services Research*, 34(2), 89-100.
- O'Malley, L., & Croucher, K. (2005). Supported Housing Services for People with Mental Health Problems: A Scoping Study. *Housing Studies*, 20(5), 831-845. <https://doi.org/10.1080/02673030500214126>
- Piat, M., Boyer, R., Fleury, M., Lesange, A., O'Connell, M., & Sabetti, J. (2015) Resident and proprietor perspectives of a recovery orientation in community-based housing. *Psychiatric Rehabilitation Journal*, 38, 88-95. <https://dx.doi.org/10.1037%2Fprj0000104>
- Padmakar, A., de Wit, E., Mary, S., Regeer, E., Bunders-Aelen, J., & Regeer, B. (2020). Supported Housing as a recovery option for long-stay patients with severe mental illness in a psychiatric hospital in South India: Learning from an innovative de-hospitalization process. (V. Gopichandran, Ed.) *PLOS ONE*, 15(4). <https://dx.doi.org/10.1371%2Fjournal.pone.0230074>
- Pandiani, J., Edgar, E., & Pierce, J. (1994). A longitudinal study of the impact of changing public policy on community mental health client residential patterns and staff attitudes. *The Journal of Mental Health Administration*, 21(1), 71-79. <https://doi.org/10.1007/BF02521347>
- Parker, S., Hopkins, G., Siskind, D., Harris, M., McKeon, G., Dark, F., & Whiteford, H. (2019). A systematic review of service models and evidence relating to the clinically operated community-based residential mental health rehabilitation for adults with severe and persisting mental illness in Australia. *BMC Psychiatry*, 19(1), 55. <https://doi.org/10.1186/s12888-019-2019-5>
- Piškur, B., Daniëls, R., Jongmans, M., Ketelaar, M., Smeets, R., Norton, M., & Beurskens, A. (2014). Participation and social participation: are they distinct concepts? *Clinical rehabilitation*, 28(3), 211-20. <https://doi.org/10.1177/0269215513499029>
- R Core Team (2013). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.

- Rogers E., S., Anthony, W., Kash, M., & Olschewski, A. (2010). Systematic Review of Supported Housing Literature 1993 – 2008. *OpenBU*. <http://www.bu.edu/drrk/research-syntheses/psychiatric-disabilities/supported-housing/>
- Sanches, S., Swildens, W., van Busschbach, J., & van Weeghel, J. (2019). Identifying social participation subgroups of individuals with severe mental illnesses: a latent class analysis. *Social Psychiatry and Psychiatric Epidemiology*, 54(9), 1067-1077. [https://doi.org/\(...\)7/s00127-019-01704-y](https://doi.org/(...)7/s00127-019-01704-y)
- Sánchez, J., Rosenthal, D., Tansey, T., Frain, M., & Bezyak, J. (2016). Predicting quality of life in adults with severe mental illness: Extending the International Classification of Functioning, Disability, and Health. *Rehabilitation Psychology*, 61(1), 19-31. <https://doi.org/10.1037/rep0000059>
- Sandhu, S., Priebe, S., Leavey, G., Harrison, I., Krotofil, J., McPherson, P., Dowling, S., Arbuthnott, M., Curtis, S., Michael, K., Shepherd, G. & Killaspy, H. (2017). Intentions and experiences of effective practice in mental health specific supported accommodation services: a qualitative interview study. *BMC Health Services Research*, 17(1), 471. <https://doi.org/10.1186/s12913-017-2411-0>
- Segal, S., Silverman, C., & Baumohl, J. (1989). Seeking Person-Environment Fit in Community Care Placement. *Journal of Social Issues*, 45(3), 49-64. <https://dx.doi.org/10.1111%2Fj.1540-4560.1989.tb01554.x>
- Shu, B., Lung, F., Lu, Y., Chase, G., & Pan, P. (2001). Care of Patients With Chronic Mental Illness: Comparison of Home and Half-Way House Care. *The International journal of social psychiatry*, 47(2). <https://doi.org/10.1177/002076400104700205>
- Siskind, D., Harris, M., Pirkis, J., & Whiteford, H. (2012). A Domains-Based Taxonomy of Supported Accommodation for People With Severe and Persistent Mental Illness. *Social psychiatry and psychiatric epidemiology*, 48(6). <https://doi.org/10.1007/s00127-012-0590-x>
- Sterne, J., Hernán, M., Reeves, B., Savović, J., Berkman, N., Viswanathan, M., David, H., Altman, D., Ansari, M., Boutron, I., Carpenter, J., Chan, A-W., Churchill, R., Deeks, J., Johnathan, J., Hróbjartsson, A., Kirkham, J., Jüni, P., Loke, Y. K., . . . Higgins, J. (2016). ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ*, 355. <https://doi.org/10.1136/bmj.i4919>
- Tabol, C., Drebing, C., & Rosenheck, R. (2010). Studies of “supported” and “supportive” housing: A comprehensive review of model descriptions and measurement. *Evaluation and Program Planning*, 33(4), 446-456. <https://doi.org/10.1016/j.evalprogplan.2009.12.002>
- Tjörnstrand, C., Bejerholm, U., & Eklund, M. (2013). Participation in Day Centres for People with Psychiatric Disabilities — A Focus on Occupational Engagement. *British Journal of Occupational Therapy*, 76(3), 144-150. <https://doi.org/10.4276/030802213X13627524435225>
- Tobin, M., Drager, K., & Richardson, L. (2013). A systematic review of social participation for adults with autism spectrum disorders: Support, social functioning, and quality of life. *Research in Autism Spectrum Disorders*, 8(3), 214-229. <https://doi.org/10.1016/j.rasd.2013.12.002>
- Trauer, T. (2001). Symptom Severity and Personal Functioning Among Patients with Schizophrenia Discharged from Long-Term Hospital Care into the Community. *Community Mental Health Journal*, 37(2), 145-155. <https://doi.org/10.1023/A:1002761732040>
- Trauer, T., Duckmanton, R., & Chiu, E. (1997). The Assessment of Clinically Significant Change Using the Life Skills Profile. *Australian & New Zealand Journal of Psychiatry*, 31(2), 257-263. <https://doi.org/10.3109/00048679709073829>

- Valdes-Stauber, J., & Killian, R. (2015). Is the level of institutionalisation found in psychiatric housing services associated with the severity of illness and the functional impairment of the patients? A patient record analysis. *BMC Psychiatry*, *15*, 215.
- van Eijk-Hustings, Y., Kroese, M., Tan, F., Boonen, A., Bessems-Beks, M., & Landewé, R. (2013). Challenges in demonstrating the effectiveness of multidisciplinary treatment on quality of life, participation and health care utilisation in patients with fibromyalgia: a randomised controlled trial. *Clinical Rheumatology*, *32*(2), 199-209. <https://doi.org/10.1007/s10067-012-2100-7>
- Viswanathan, M., & Berkman, N. (2011). Development of the RTI item bank on risk of bias and precision of observational studies. *Journal of Clinical Epidemiology*, *65*(2), 163-178. <https://doi.org/10.1016/j.jclinepi.2011.05.008>
- Viswanathan, M., Berkman, N., Dryden, D., & Hartling, L. (2013). Assessing Risk of Bias and Confounding in Observational Studies of Interventions or Exposures: Further Development of the RTI Item Bank.
- Webber, M., & Fendt-Newlin, M. (2017). A review of social participation interventions for people with mental health problems. *Social Psychiatry and Psychiatric Epidemiology*, *52*(4), 369. <https://dx.doi.org/10.1007%2Fs00127-017-1372-2>
- Wong, Y., Matejkowski, J., Lee, S., Wong, Y.-L., Matejkowski, J., & Lee, S. (2011, 1). Social integration of people with serious mental illness: network transactions and satisfaction. *Journal of Behavioral Health Services & Research*, *38*(1), 51-67. <https://dx.doi.org/10.1007%2Fs11414-009-9203-1>

Table 1: Data extraction table for included studies

Author, year, country	Study design	Level of support	Sample size	Mean Age (years)	Diagnoses	Gender	Factors of Participation			Study Quality: Measure/ Rating
							Social participation	Daily living functioning	Personal Empowerment	
Kruzich and Berg, 1985, America	Cohort study	High	42	Mean not reported. Over half were 35-65; 25% over 66 and a fourth under 35 *	66% schizophrenia, 7% affective disorder, 16% Chronic brain disorder, 10% with other mental health conditions	Male 51% Female 49%	-	Self sufficiency	-	RTI: Risk of bias unclear.
		Moderate	21							
		None	20							
Nelson, Hall and Walsh, 1997, Canada	Quasi-experimental Non-equivalent comparison group design.	High	30	32.5	Inclusion criteria: Psychiatric consumers/ survivors.	Male 60% Female 40%	Instrumental roles	Independent functioning	Mastery	ROBINS-I: Moderate risk of bias
		Moderate	52	34.1		Male 67% Female 33%				
		None	25	45		Male 44% Female 56%				
Fossey et al., 2006, Australia	Cross-sectional study	High	25	35.09	Schizophrenia	Male 56% Female 44%	Social contact	Self-care	Responsibility	RTI: Low risk of bias
		Moderate	18							
Dorer, Harries and Marston, 2009, UK	Cross-sectional study (staff-survey method)	High	91	42.7	Unspecified Long-term mental health problems	Male 66% Female 34%	Education, employment, day centre, local facilities, faith, family and friends	-	-	RTI: High risk of bias
		Moderate	62							
Shu et al 2001, Taiwan	Quasi-experimental (Longitudinal design)	High	23	37.8	Schizophrenia	Male 57% Female 43%	Social Activity	-	Autonomy	ROBINS-I: Moderate risk of bias
		Moderate	37	32.4						
Eklund and Tjornstrand, 2019, Sweden	Cross sectional (secondary data analysis)	High	155	48	62% psychosis, 13% anxiety/mood disorder and 25% other mental health disorder	Male 46% Female 64%	Employment status, education status, work training, day centre, leisure activities, cultural activities	Doing household work, managing self-care, gardening and repairs, Physical exercise.	-	RTI: Low risk of bias
		Moderate	111	46						

RTI= Research Triangle Institute item bank; ROBINS-I=Risk Of Bias In Non-randomised Studies of Interventions. *As reported by study

Table 2: Data extraction of results from selected studies

Study	Level of support	Sample size	Factors and Data Reported											
			Social Participation						Daily Living Functioning				Personal Empowerment	
Kruzich and Berg, 1985, America	High	42	---						Self-sufficiency (mean)				-	
	Moderate	21							1.23					
	None	24							2.19					
Nelson, Hall and Walsh, 1997, Canada	High	30	Instrumental Roles (mean)						Independent functioning (mean)				Mastery (mean)	
	Moderate	52	0.73						14.3				20.4	
	None	25	0.89						16.1				21.3	
Fossey et al, 2006, Australia	High	25	Social Contact (mean)						Self-care (mean)				Responsibility (mean)	
	Moderate	18	15.8						31.88				16.96	
			17						32				17.76	
Dorer, Harries and Marston, 2009, UK			Estimated time for participation in specific community occupations (Mean in Hours (SD))											
			<i>education</i>	<i>Employment</i>	<i>day centre</i>	<i>local facilities</i>	<i>faith</i>	<i>family and friends</i>					-	
	High	91	0.6(2.6)	0.4(1.5)	0.8(3.2)	6.2(7.7)	0.3(1.2)	8.1(16.9)					-	
Moderate	62	1.7(4.2)	1.0(4.1)	1.9(3.9)	8.0(9.1)	0.1(0.6)	13.9(24.3)							
Shu et al 2001, Taiwan			Social Activity Factor (GEE regression Coefficient)*											Autonomy Factor (GEE Regression Coefficient)**
	High	23	1.62, p=0.01											--
	Moderate	37												1.20, p=0.04
Eklund and Tjornstrand, 2019, Sweden			Percentage of participants involved in occupations						Percentage of participants involved in occupations				-	
	High	155	<i>employed or a student</i>	<i>training or enrolled in studies</i>	<i>attending a day centre</i>	<i>organised leisure/hobbies</i>	<i>work training</i>	<i>cultural occupations</i>	<i>doing household work</i>	<i>gardening or repairs</i>	<i>managing own personal hygiene</i>	<i>physical exercise</i>		
	Moderate	111	9%	10%	41%	23%	11%	79%	83%	10%	87%	88%		
			11%	14%	37%	34%	15%	79%	72%	37%	87%	88%		


GEE= Generalised Estimating Equations SD= Standard deviation *Social activity between group for level of support

Table 3: Calculated effect sizes

Factor	Level of support	No. of studies	Sample size (total)	Results
Social Participation	High vs Moderate	5	604	
Instrumental roles				RMD = -0.16
Social contact				d= - 0.3957 Sub-group difference (t)= -1.28 p=0.21 CI= -3.10 to 0.70
Estimated time for participation in specific community occupations				Education: d=- 0.321 Employment: d=-0.210 Day centre: d=-0.31 Local facilities: d=-0.21 Faith: d=0.199 Family and friends: d= -0.28
				Combined SMD* (d): -0.1926 p=0.004. Fixed effects model p<0.05, CI = -0.3250 to -0.0603
Social Activity factor				GEE regression coefficient between group: 1.20, p=0.04;
Percentage of participants involved in daily activities				employed or student(PD)= -2%,
				working or enrolled(PD) = -4%
				day centre(PD)= 4%
				leisure/hobbies(PD)= -11%,
				work training(PD) = -4%
				cultural occupation(PD)=0%
	High and Moderate vs No	1	25	
Instrumental roles				F(2,104)=3.0,p<0.05 **
Daily Living Function	High vs Moderate		454	
Self-sufficiency				RMD = -0.96
Independent functioning				RMD = -1.8
Self-care				d=-0.022 Sub-group difference (t)=-0.07 p=0.9, CI=-3.49 to 3.25
Percentage of participants involved in daily living activities				Household work (PD)= 11%
				Gardening(PD)=-27%
				Personal hygiene (PD)= 0%
				Physical exercise(PD)=0%
	High and Moderate vs No	2	194	
Self-sufficiency				RMD = 0.38
Independent Functioning				F(2,101)=6.4 p<0.001
Personal Empowerment	High vs Moderate	3	185	
Mastery				RMD= -0.9
Responsibility				d= -0.329 sub-group difference (t) = -1.04(p=0.30); CI: -2.37 to 0.77
Autonomy				GEE reg. coefficient between group = 1.62, p=0.01
	High and Moderate vs No	1	107	
Mastery				F(2,96)=3.1 p<0.05 **

d= Cohen's d; RMD= Raw mean difference; PD= percentage difference; *SMD=Standardised mean difference; **ANOVA for group

Figure 1: PRISMA Flow Diagram



Supplementary Material 1: Outcomes reported in included studies matched to participation factors

Participation factor	Outcome reported in study	Measure used in study
Social participation	Instrumental Roles (Nelson et al. 1997)	Total number of roles involved in
	Social contact (Fossey et al. 2006)	Life Skills Profile ¹
	Participation in community occupations (Dorer et al. 2009)	Survey developed by study authors
	Social Activity Factor (Shu et al. 2001)	Quality of Life Scale ²
	Percentage of participants involved in daily activities (employed or student, working or enrolled in studies, attending a day centre, organised leisure/hobbies at least once a week, work training, cultural occupations; Eklund and Tjörstrand 2019)	Satisfaction with Daily Living Occupations ³
Daily Living Functioning	Self-sufficiency (Kruzich and Berg 1985)	Self-sufficiency index developed by authors for study
	Independent Functioning (Nelson et al. 1987)	Independent Functioning Scale (adapted by authors from Rappaport et al. 1985)
	Self-care (Fossey et al. 2006)	Life Skill Profile ¹
	Percentage of participants involved in daily activities (doing household work, gardening or repairs, managing own personal hygiene on a daily basis, physical exercise; Eklund and Tjörstrand 2019)	Satisfaction with Daily Living Occupations ³
Personal Empowerment	Mastery (Nelson et al, 1985)	Mastery Scale ⁴
	Responsibility (Fossey et al. 2006)	Life Skills Profile ¹
	Autonomy (Shu et al. 2001)	Quality of Life Scale ²

1. Rosen A, Hadzi-Pavlovic D, Parker G (1989) The Life Skills Profile: a measure assessing function and disability in schizophrenia. *Schizophrenia Bulletin*, 15(2), 325-37.
2. Yu, W.Y. (1995). The development of quality of life for mental illness patient. *Public Health in Taiwan*, 22, 29-39.
3. Eklund, M., Bäckström, M., & Eakman, A. (2014). Psychometric properties and factor structure of the 13-item satisfaction with daily occupations scale when used with people with mental health problems. *Health and Quality of Life Outcomes*, 12(1), 1–9.
4. Pearlin, L. I., & Schooler, C. (1978). The structure of coping. *Journal of Health and Social Behavior*, 19, 2–21.

Supplementary material 2: Search strategy

<i>Database: Ebscohost - PsychINFO, PsychARTICLES, Medline, CINAHL Limited by Age over 18+, publications in English</i>	
S29	S11 AND S21 AND S22 AND S28
S28	S25 OR S27
S27	S23 OR S26
S26	occ* Participation OR process* skill* OR motor skill* OR cognitive ability OR roles OR occ* roles OR build* relation* OR organisation* skill* OR problem solving OR communicat* ability OR communicat* skill* OR Increase* responsibility OR improve* responsibility OR motivation to participate OR motivation to engage OR change in motivation
S25	S15 AND S24
S24	participation OR engag* OR involv*
S23	improved participation OR factors of participation OR participation OR level of participation OR empower* OR enabling relation* OR autonomy OR independ* OR promot* OR skill* OR abilit* OR occ* performance
S22	informal care* OR family OR relative* OR friend* OR support network
S21	S13 OR S14 OR S15 OR S17 OR S18 OR S19 OR S20
S20	facilitation OR encour* OR enabl*
S19	support OR care
S18	rapport OR rapport buil* OR buil* rapport
S17	S15 AND S16
S16	engage* OR interaction* OR environ*
S15	social
S14	S12 AND S13
S13	interaction OR relation* OR engage*
S12	staff OR care* OR professional*
S11	S9 AND S10
S10	S5 OR S7
S9	S1 AND S2
S8	S1 AND S2 AND S5 AND S7
S7	S4 AND S6
S6	shared
S5	S3 AND S4
S4	hous* OR accom* OR environment*
S3	support*
S2	client* OR resident* OR patient* OR service user*
S1	psyc* OR mental illness* OR mental health difficult* OR mental health

Database: ASSIA

((noft(hous* OR accom* OR environment*) AND noft(shared OR support*)) AND (noft(psyc* OR mental illness* OR mental health difficult* OR mental health) AND noft(client* OR resident* OR patient* OR service user*))) AND ((noft(interaction OR relation* OR engage*)) OR (noft(staff OR care* OR professional*) AND noft(interaction OR relation* OR engage*)) OR (noft(social)) OR (noft(social) AND noft(engage* OR interaction* OR environ*)) OR noft(rapport OR rapport buil* OR buil* rapport) OR noft(support OR care) OR noft(facilitation OR encour* OR enabl*)) AND (noft(informal care*) OR noft(family) OR noft(relative*) OR noft(friend*) OR noft(support network)) AND ((noft(social) AND noft(participation OR engag* OR involv*)) OR ((noft (improved participation OR factors of participation OR participation OR level of participation OR empower* OR enabling relation* OR autonomy OR independ* OR promot* OR skill* OR abilit* OR occ* performance)) OR (noft(occ* Participation OR process* skill* OR motor skill* OR cognitive ability OR roles OR occ* roles OR build* relation* OR organisation* skill* OR problem solving OR communicat* ability OR communicat* skill* OR Increase* responsibility OR improve* responsibility OR motivation to participate OR motivation to engage OR change in motivation)))))).

Supplementary material 3: Quality assessment of included studies

RTI item time bank													
Study	SD	1	2	3	5	6	7	8	9	11	12	13	Interpret
Fossey et al. 2006	CS	N(low)	N(low)	N(low)	N/A	Y(low)	-	-	N(low)	Y(low)	Y(low)	N(low)	Low
Kruzich and Berg, 1985	C	N(low)	CD (unclear)	N(low)	N/A	N(high)	N(low)	CD (unclear)	N(low)	Y(low)	Y(low)	N(low)	Unclear
Dorer, Harries and Marston, 2009	CS	N(low)	N(low)	N(low)	N/A	N(high)	-	-	N(low)	Y(low)	Y(low)	N(low)	High
Eklund and Tjornstrand, 2019	CS	N(low)	Y(high)	N(low)	N/A	Y(low)	-	-	N(low)	Y(low)	Y(low)	N(low)	Low
Question No.	Question												Type of bias identified
1	Do the inclusion/exclusion criteria vary across the comparison groups of the study?												Selection bias
2	Does the strategy for recruiting participants into the study differ across groups?												Selection bias, confounding
3	Is the selection of the comparison group inappropriate?												Selection bias, confounding
5	Was the assessor not blinded to the outcome, exposure, or intervention status of the participants?												Detection bias
6	Were valid and reliable measures implemented consistently across all study participants to assess inclusion/exclusion criteria, physical activity outcomes, and potential confounders?												Detection bias, confounding
7	Was the length of follow-up different across study groups?												Attrition bias
8	In cases of high loss to follow-up (or differential loss to follow-up), was the impact assessed (e.g., through sensitivity analysis or other adjustment method)?												Attrition bias, detection bias
9	Are any important primary outcomes missing from the results?												Reporting bias
11	Are results believable taking study limitations into consideration?												Overall assessment
12	Were there any attempts to balance the allocation between the groups or match groups?												Confounding
13	Were important confounding variables not taken into account in the design and/or analysis?												Confounding
SD= Study design C= Cohort study CS= Cross-section study Y= Yes N=No CD= Cannot determine													

ROBINS-I								
	D1	D2	D3	D4	D5	D6	D7	Overall Risk of bias
Nelson et al. 1985	Moderate	Low	Low	Low	Low	Low	Low	Moderate
Shu et al. 2001	Low	Low	Low	Low	Low	Low	Moderate	Moderate
Domains								
D1	Bias due to confounding							
D2	Bias due to selection of participants							
D3	Bias in classification of interventions							
D4	Bias due to deviations from intended interventions							
D5	Bias due to missing data							
D6	Bias due to measurement of outcomes							
D7	Bias in the selections of the reported result							
Low = low risk of bias; Moderate = moderate risk of bias								