



ISSN Print: 2394-7500  
 ISSN Online: 2394-5869  
 Impact Factor: 5.2  
 IJAR 2020; 6(6): 391-394  
[www.allresearchjournal.com](http://www.allresearchjournal.com)  
 Received: 24-04-2020  
 Accepted: 28-05-2020

**NG Shinde**  
 Department of Zoology,  
 K.J. Somaiya College,  
 Kopergaon, Ahmednagar,  
 Maharashtra, India

## Probable role of ablation of cerebral ganglia and injection of its extracts on O:N ratio of *Lamellidens corrianus* during summer season

**NG Shinde**

### Abstract

Amongst invertebrates, molluscs show great variability in their nervous system ranging from primitive arrangement in Chitons to the complex mass of fused ganglia forming the 'brain' of cephalopods. Most of the effector organs used for pharmacological or physiological experiments. The neurosecretory cells (NSCs) with their combination of neuronal and glandular capabilities are perfectly suited to translate a neuronal input into the hormonal output best suited to long-term process. In this capacity, the NSCs may produce hormones, which act directly upon the peripheral target or it may exert its effect indirectly by influencing the activity of other non-neural, endocrine organs. Neurosecretory cells have been detected in the cerebral, pedal and visceral ganglia of *Lamellidens corrianus*. The distribution and biology of freshwater bivalve like *Lamellidens corrianus* is influenced by local ecological factors (like temperature, pH, inorganic salts, type of soil etc.), water flow system and presence of micro-organisms, teleost fishes and seasonal variations in these parameters. Nitrogenous excretory changes which include ammonia in these molluscs is also influenced by these local ecological factors in which temperature is plays very crucial role. It is well known fact that the high stress conditions during drought or severe winter conditions have been conquered by several lineage of gastropods and sand bivalves and their ability to enter the resistant or dormant stages (low food) protein catabolism increased. Present investigation deals with the oxygen consumption and ammonia excretion ratio of these freshwater pelecypode *Lamellidens corrianus*. Study on *Lemellidens corrianus* directed in understanding the different behavioral and physiological aspects after cerebral ganglia removal and injection of their extracts revealed significant changes during summer (April-May).

**Keywords:** Cerebral ganglia ablation, Injection of cerebral ganglionic extract, *Lamellidens corrianus*, O:N ratio, summer season

### Introduction

Mollusca are the second largest phylum of Animal Kingdom after the phylum Arthropoda (Verma and Prakash, 2020) [25]. In tropical regions like India, in general the conditions are hot-dry from March to May, these weather conditions can have considerable impact on the life of bivalve molluscs, especially in the areas lacking heavy rain fall. The bivalve molluscs living especially in lentic environment have to face drastic environmental changes than those in lotic conditions. Temperature rise above the optimum requirement can affect the physiology of bivalves (Bayne, 1976) [3]. The studies with polluting substances also revealed that the freshwater bivalves from Godavari River at Paithan are very sensitive to pesticide toxicity in summer than in monsoon and winter (Akarte, 1985; Muley, 1985) [1, 18]. Pesticide toxicity is also recorded in case of fishes (Prakash and Verma, 2014; Verma and Prakash, 2018; Kaur and Mishra, 2019) [19, 25, 11]. An oxygen concentration in natural waters is influenced by temperature, (Cairns *et al.*, 1975) [6], thus in summer *Lamellidens corrianus* from Jayakwadi back water faces severe environmental conditions. The physiological aspects or the endocrine aspects should be directed towards the studies through different seasons because the aquatic animal itself show cyclic changes in the aspects (Kulkarni, 1987) [15]. Stickel (1973) [22] stated that, nutrient deposition or depletion is dependent upon the balance between anabolism and catabolism and one measure of metabolic activity often used is the rate of oxygen consumption which is affected by factors that fluctuate seasonally such as food availability and water temperature. Thus, in the present study this dual effect of changes in environmental parameters and starvation revealed that in summer the rate of oxygen consumption was decreased initially up to 12 days.

**Corresponding Author:**  
**NG Shinde**  
 Department of Zoology,  
 K.J. Somaiya College,  
 Kopergaon, Ahmednagar,  
 Maharashtra, India

Many authors have quoted that ammonia in general is a major nitrogenous excretory product of bivalves and there occurs a profound difference in loss of nitrogen between different sizes and seasons (Bishop *et. al.*, 1983) [4]. In mollusca, interactions with the environment are handled by the nervous system (including the sense organs) and the muscular system (the muscles that making the foot and those that attach the animal to its shell). Both systems, of course, have been of different forms in different group of animals. Basically, there are two extreme types of neural organization (with many intermediate forms).

The chord system, in which the neural tree tapers and branches in smooth lines; and the ganglia system, which is characterized by knots of nerve cells called ganglia, bound together by nervous tissue consisting of slender elongations of the cells.

The nervous and endocrine systems coordinate the activities of the various organs and the tissues in the body that the animal's function as individuals. Majority of the neurons with granular activity are known to be necessary for the transmission of transient impulses, with their highly localized production of chemicals such as neurohumors/neurohormones, which are rapidly destroyed. Neurons have similar properties throughout the animal kingdom, although their morphology and arrangement may vary.

The way in which nervous system operates may differ considerably between animal depending on the number of neurons involved and the individual shape, size and spatial arrangement of the component about fifty percent of the nervous system is composed of non-excitabile satellite glial cells which are packed between and around the neurons. They are supposed to transport neurosecretory substance (Lubet, 1955a; Umiji, 1969) [17, 23], provide physical support for the neurons and can modify action of nerve cells as well as a barrier or receiver for ions, metabolites and transmitters (Leak and Walker, 1980) [16].

The neurosecretory cells (NSCs) with their combination of neuronal and glandular capabilities are perfectly suited to translate a neuronal input into the hormonal output best suited to long-term process. In this capacity, the NSCs may produce hormones, which act directly upon the peripheral target or it may exert its effect indirectly by influencing the activity of other, non-neural, endocrine organs. In this later case, the NSCs organ may act via the production of blood-born hormone. Knowles and Bern (1966) [13] stated that the significance of NSCs as connecting link between nervous and endocrine systems and neurosecretory neurons participate either directly or indirectly in endocrine control and form all or part of endocrine organ. Yasmeeen (2019) [26] also worked a lot on gonads of *Lamellidens*.

Hormones are consequently well suited to exert their effects over extended period of time and the endocrine system controlling-term process within the body such as the coordinated growth of organs or the maintains of appropriate metabolite concentration in the blood and tissues. Neurosecretory cells of cerebral, pedal and visceral ganglia secrete its secretion which help in regulation of body activity.

Removal of both cerebral ganglia (bilateral cerebralctomy) troubled in neurocretion which might affect in metabolic activity and exert stress on animal which probably hamper the rate of respiration and ammonia excretion (O:N ratio).

## Materials and Methods

The freshwater bivalve molluscs, *Lamellidens corrianus* were collected from Jayakwadi backwaters (Nathsagar) at Paithan, 45 km. away from Aurangabad. After brought to the laboratory, the shells of the bivalves were brushed and washed with fresh reservoir water so as to remove the fouling algal biomass and mud. The animals of 80-85 mm shell length were selected for experiment and they were acclimatized for 24 h. at laboratory condition in fresh aerated reservoir water (with renewal of water at the interval of 12-13 h.) and stocking capacity was given during this period and no food was given to the bivalves during laboratory acclimatization and subsequent experimentation. After 24 h., reservoir water was once again renewed and aeration was given. After a lapse of 1 h. the animals extended their organs (foot, mantle, siphons) to maximum and soon surgical operations and injection of the ganglionic extract were done. For removal of both the cerebral ganglia (bilateral cerebralectomy) active animal were chosen from the aquarium and a wedge (4-5 mm thick) was kept between the valves of the shell. Both the cerebral ganglia were removed by performing minimum injury to the animals within 30 seconds, with the help of fine, pointed sterilized forceps. For injection of ganglionic extracts, cerebral ganglionic extract was prepared in ice cold distilled water (10 ganglia in 1ml cold distilled water was centrifuged and the supernatant (0.2 ml/animal i.e., equivalent to 2 ganglia/animal) were injected into the foot (muscular region) of normal control and gangliatomized (both cerebral ganglia ablated) bivalves. In sham operated control animals were injected by 0.2ml cold distilled water. The result for control and sham operated groups were similar and hence a comparison was made between gangliatomized and control group and extract injected to normal control as well as ablated and control group of animals only. Soon after the operation and injection of ganglionic extracts to normal control, extirpated 20 animals of cerebralectomy, 20 animals of extract injected, and 20 animals of extract injected to ablated bivalves were transferred to separate aquaria. Each aquarium contained 15 liter well aerated reservoir water, and experiment was run for 12 days. The water from each aquarium was changed at an approximate interval of 12-13 h. throughout experimental period. The behavior and mortality of the bivalves were recorded before each change of water from all the aquaria. The experiments were conducted for 12 days on freshly collected animals during summer season (April - May). The adult bivalves were placed individually in respiratory jars with 1 liter water. The rate of oxygen consumption of the animals in each group during summer seasons was determined according to Winkler's modified technique (Golterman *et.al.*, 1978) [10], in a specially prepared brown colored respiratory glass jar of one liter capacity. The rate of ammonia excretion was measured on 2<sup>nd</sup>, 7<sup>th</sup> and 12<sup>th</sup> day according to phenol-hypochlorite method of Solorzano (1969) [19]. Every time four individual animals of each group were used and mean of triplicate of water samples were estimated for each group. The statistical analysis was done to express final data. The atomic equivalent values of ammonia excretion obtained for the same individual. All the values were subjected to statistical analysis for confirmation using student 't' test (Dowdeswell, 1957) [8]. Statistical and percentage differences were also calculated in experimental animals.

## Results

**Table 1:** Effect of ablation of cerebral ganglia and injection of its extracts on O:N ratio of *Lamellidens carrianus* during summer seasons. (Bracket Values represents percentage difference)

Days	Summer season			
	Control	Ablation of cerebral ganglia	Injection of distilled water	Injection of cerebral ganglionic extract
On 2 <sup>nd</sup>	47.5575±19.4535	26.0221±09.6193(45.28)	24.8675±05.3141	17.8820±02.7288(62.39)
On 7 <sup>th</sup>	26.9100±05.3208(43.41)	15.6850±03.2732(41.37)	18.9100±03.0204	11.7844±01.9702(56.20)
On 12 <sup>th</sup>	10.0201±02.8612(78.93)	26.7002±11.8806(166.04)	07.2791±01.6874	15.6908±01.7687(56.58)

The changes in the O:N ratio from control, ablation of cerebral ganglia and injection of ganglionic extracts during different seasons, in freshwater bivalve mollusc *Lamellidens corrianus* were given in table-1. During summer season, in the control group, the O:N ratio was (47.5575 ± 19.4535) on 2<sup>nd</sup> day, (26.91 ± 5.3208) on 7<sup>th</sup> day, and (10.0201 ± 2.8612) on 12<sup>th</sup> day. The higher value of ratio was found on 2<sup>nd</sup> day. The ratio decreased (43.41%) on 7<sup>th</sup> and (78.93%) on 12<sup>th</sup> day, compared to 2<sup>nd</sup> day. Among the experimental groups O:N ratio in ganglia ablated animals was decreased (26.0221 ± 9.6193, 45.28%) on 2<sup>nd</sup>, (15.6850 ± 3.2732, 41.37%) on 7<sup>th</sup> and increased (26.7002 ± 11.8806, 116.4%) on 12<sup>th</sup> day. Whereas, the O:N ratio from ganglionic extract injected animals was decreased (17.882 ± 2.7288, 62.39%) on 2<sup>nd</sup> day, (11.7844 ± 1.9702, 56.20%) on 7<sup>th</sup> day, while it was increased (15.6908 ± 1.7687, 56.58%) on 12<sup>th</sup> day compared to respective controls.

## Discussion

The O:N ratio is an index of protein utilization in energy metabolism. A few investigators also demonstrated the probable role of ammonia in the settlement of larvae of different bivalves. According to Coon *et. al.*, (1990) [7] ammonia solution (pH 8.0) (2.5 mM concentration) induced stereotypical settlement behavior of larvae. The authors suggested that ammonia increased the intracellular pH. Fitt and coon (1992) [9] stated that the actual concentration of NH<sub>3</sub> was associated with the surface for the oysters *Crassostrea virginica* and *Crassostrea gigas*. Increased protein catabolism is indicated by high level of ammonia excretion and decline in oxygen: nitrogen ratio (Bayne, 1973) [2] and thus changes in the rates of nitrogen excretion are best understood in the context of physiological energetic and nitrogen balance, when related to overall metabolic rate by means of the oxygen: nitrogen (or O:N) ratio. According to Khalil (1994) [12] in *Tapes decussatus* ammonia excretion rate varied with body weight, temperature, and starvation. Ammonia excretion rates were steady during six days of starvation and higher excretion rate was depended on the temperature. The ammonia excretion rate was higher for starved clams than for fed clams of all sizes and at different temperatures, weight specific ammonia excretion rates were related to dry flesh weight of starved clams but were not related to fed ones in *T. decussatus* (Khalil, 1994) [12]. Increase in the rate of ammonia excretion might be due to starvation, because during starvation there is more protein catabolism hence ammonia excretion rate increases. Increased ammonia excretion indicated increased protein catabolism during starvation (Krishnakumar and Damodharan, 1986) [14]. The author also stated that the mussel is dependent on protein for energy production, which is less than healthy *Mytilus edulis* from Cochin backwater. In addition, the author further stated that the ratio might vary according to gametogenic cycle, nutritive reserves and

nature of food. It is well known fact that the high stress conditions during drought or severe winter conditions have been conquered by several lineage of gastropods and sand bivalves, and their ability to enter the resistant or dormant stages (low food) protein catabolism increased. Many workers have compared the excretion rates of bivalves related with their environment. Boucher and Boucher (1990) [5] have demonstrated the ammonia exchange processes of *Crassostrea gigas* and sediment. Authors compared the oyster biomass and lighting conditions (light and dark) with oxygen and ammonia exchange rates. The oxygen consumption increased with increasing biomass but the respiration rate was decreased, whereas, ammonia released never matched the rate predicted by adding sediment efflux to oyster excretion. The O:N ratio was influenced by oyster biomass. Further, the author suggested that both the organism and sediment were involved in regulation process (Boucher and Boucher, 1990) [5]. Food availability and the cells rapidly filled with material. With the close of monsoon, from October NM absorption phases extended till November NM. This was correlated with longer feeding period under short supply of food and further extended till February FM, showing tubules in either holding or absorption phases. The water temperature decreased with the onset of winter and no feeding occurred for a short period of November FM when the tubule showed re-organization phase (Saokar, 1994) [20]. The existing labor was undertaken to investigate the effect of endogenous parameters chiefly neuroendocrine on the aspects of O:N ratio. It reveals that cerebral ganglia ablation and cerebral ganglionic extract injection might exert some effect on O:N ratio along with the environmental parameters. As compare in cerebral ganglionic extract injection group, more profound effect of O:N ratio seen in the cerebral ganglia ablated group on 12<sup>th</sup> day of experimental period.

## References

1. Akarte SR. Effect of organophosphorus insecticides on bivalve molluscs. Ph.D. Thesis, Marathwada University 1985, 1-252.
2. Bayne BL. Physiological changes in *Mytilus edulis* (L.) induced by and nutritive stress. J. Mar. Biol. Ass. UK 1973;53:39-58.
3. Bayne BL. Marine mussels: their ecology and physiology Cambridge University Press, Cambridge London, New York, and Melbourne 1976a, 1-495.
4. Bishop SH, Ellis IL, Burchan JM. Ammino acid metabolism in molluscs in: The molluscs, (Ed. Wilbur, K.M.), Academic press, New York 1983;I:244-328.
5. Boucher-Rodoni R, Boucher G. In situ study of the effect of oyster biomass on benthic metabolic exchange rates. *Hydrobiologia* 1990;206:115-124.
6. Cairns J Jr, Health AG, Parker BC. Temperature influence of chemical toxicity to aquatic organisms. J WPCF 1975;47(2):267-280.

7. Coon SL, Walch M, Fitt WK, Weiner RM, Bonar DB. Ammonia induces settlement behavior in oyster larvae. *Biol. Bull* 1990;91:297-303.
8. Dowdeswell WH. *Practical Animal Ecology*, Methum and Co. Ltd., London 1957.
9. Fitt WK, Coon SL. Evidence for ammonia as a natural use for recruitment of oyster larvae to oyster beds in Georgia salt marsh. *Biol. Bull* 1992;182:401-408.
10. Golterman HL, Clymo RS, Ohnstad MAM. *Physical and chemical analysis of freshwaters*. IBP, Handbook No-8, Blackwell Scientific Publication, Oxford, London, Edinburg, Melbourne, 2<sup>nd</sup> Ed 1978, 172-178.
11. Kaur G, Mishra BKP. Histopathological changes in Liver of fish *Channa punctatus* exposed to sub lethal concentration of Hybrid Pesticide. *International Journal of Biological Innovations* 2019;1(2):83-86. <https://doi.org/10.46505/IJBI.2019.1209>
12. Khalil AM. Influence of starvation, body size and temperature on ammonia excretion in marine bivalve, *Tapes decussates* (L.). *Aquacult. Fish. Manage* 1994;25:839-847.
13. Knowles FGW, Bern HA. Function in neuroendocrine regulation. *Nature* 1966;210:271-272.
14. Krishnakumar PK, Damodaran R. Effect of the body size on the physiology of green mussel *Perna viridis* (Linnaeus), In: *Proceedings of the National Seminar Mussel Watch*. Cochin University of Science and Technology 1986; I:77-85.
15. Kulkarni DA. A study on the reproductive endocrinology of freshwater molluscs. Ph.D. Thesis, Marathwada University 1987, 1-192.
16. Leak LD, Walker RJ. *Invertebrate neuropharmacology* Thomson Litho Ltd., East Kilbride, Scotland, Blackie, Glasgow and London, Edward Arnold 1980, 1-357.
17. Lubet P. Cycle neurose' cre'torie chez *Chlamys varia* at *Mytilus edulis* L. (Mollusques: Lamellibranches). *C.R. Acad. Sci* 1955a;241:119-121.
18. Muley DV. Effect of pollutants on freshwater molluscs from Godavari river at Paithan. Ph.D. Thesis, Marathwada University (India) 1985, 1-313.
19. Prakash S, Verma AK. Effect of Organophosphorus Pesticide (Chlorpyrifos) on the Haematology of *Heteropnetues fossilis* (Bloch). *International Journal of Fauna and Biological Studies* 2014;1(5):95-98.
20. Saokar CD. Some aspects of the reproductive physiology of freshwater bivalve molluscs, *Indonaia caeruleus* from a pond in the Girna river near Chankapur dam, District Nashik, M.S., India., Ph.D. Thesis, Marathwada University Aurangabad 1994.
21. Solorazano L. Determination of ammonia in neutral waters by the phenol hypochlorite method *Limnology and Oceanography* 1969;14:799-801.
22. Stickle WB. The reproductive physiology of the intertidal proaobranch, *Thais lamellosa* (Gmelin) 1. Seasonal changes in the rate of oxygen consumption and body component indexes. *Biol. Bull* 1973;144:511-524.
23. Umiji S. Neurosecretas em *Mytilus perna*: (Molusco-Lamellibranquio). *Zool. Biol. Mar. (Sao Paulo) Vova Ser* 1969;26:131-254.
24. Verma AK, Prakash S. Haematotoxicity of Phorate, an Organophosphorous pesticide on a Freshwater Fish, *Channa punctatus* (Bloch). *International Journal on Agricultural Sciences* 2018;9(2):117-120.
25. Verma AK, Prakash S. Status of Animal Phyla in different Kingdom Systems of Biological Classification. *International Journal of Biological Innovations* 2020;2(2):149-154. <https://doi.org/10.46505/IJBI.2020.2211>
26. Yasmeen S. Cadmium induced histopathological alterations in female gonad of freshwater bivalve mollusks, *Lamellidens marginalis* during summer season. *International Journal of Biological Innovations* 2019;1(2):73-77. <https://doi.org/10.46505/IJBI.2019.1207>