



This is a repository copy of *Dilation and flow resistance of granular flows in a rotating drum*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/169397/>

Version: Published Version

---

**Proceedings Paper:**

Kaitna, R. [orcid.org/0000-0002-2289-723X](https://orcid.org/0000-0002-2289-723X), Shuai, L., Taylor-Noonan, A. et al. (4 more authors) (2020) Dilation and flow resistance of granular flows in a rotating drum. In: EGU General Assembly 2020 Abstracts. EGU General Assembly 2020 - Sharing Geoscience Online, 04-08 May 2020, Virtual conference. Copernicus GmbH .

<https://doi.org/10.5194/egusphere-egu2020-18418>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:  
<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

EGU2020-18418

<https://doi.org/10.5194/egusphere-egu2020-18418>

EGU General Assembly 2020

© Author(s) 2020. This work is distributed under the Creative Commons Attribution 4.0 License.



## Dilation and flow resistance of granular flows in a rotating drum

**Roland Kaitna**<sup>1</sup>, Li Shuai<sup>2</sup>, Alexander Taylor-Noonan<sup>3</sup>, William Andrew Take<sup>3</sup>, Brian McArdell<sup>4</sup>, James McElwaine<sup>5</sup>, and Elisabeth Bowmann<sup>6</sup>

<sup>1</sup>University of Natural Resources and Life Sciences, Vienna, Institute of Mountain Risk Engineering, Vienna, Austria (roland.kaitna@boku.ac.at)

<sup>2</sup>Institute of Mountain Hazards and Environment, Chinese Academy of Science, China

<sup>3</sup>Department of Civil Engineering, Queen's University, Canada

<sup>4</sup>Swiss Federal Institute for Forest Snow and Landscape Research WSL, Switzerland

<sup>5</sup>Department of Earth Sciences, Durham University, United Kingdom

<sup>6</sup>Department of Civil and Structural Engineering, University of Sheffield, United Kingdom

The flow resistance in granular mass flows can be due to frictional contacts or collisional interactions between particles. For a constant number of particles, the transition from a frictional to a collisional regime is expected to depend on flow velocity and is associated with an increase of volume and a decrease of bulk density, an effect termed dilation. The relation between velocity, dilation and flow resistance is not well understood. Here we present results of steady, non-uniform flows of ceramic beads ( $d = 4$  mm) in a rotating drum, a setup allowing observations and averaging of parameters measured over an extended period of time. We systematically varied flow mass between 12.3 and 49 kg and flow velocity between 0.2 and 1.2 m/s, while continuously measuring basal normal stress and flow depth. Flow resistance was assessed by calculating average bulk shear stress from torque measurements at the axis of the drum as well as from the deviation of the center of mass from the vertical. Additionally, the flows were captured by high-speed video recordings through the transparent side wall. We find bulk densities at the deepest section of the flow decreasing from 1430 kg/m<sup>3</sup> at low velocities to 1370 kg/m<sup>3</sup> at the highest velocity for the largest flow mass. At the same time flow resistance increased linearly. When the flow mass was reduced, also bulk density decreased, indicating the importance of overburden pressure for dilation. Video recordings revealed that shear is concentrated in depth zones of lower volume fraction. Our results shall contribute to a better understanding of the transition from a frictional to a collisional flow regime and may help to assess the importance of dilation for gravitational mass flows.