

# Galactosaminoglycans from normal myometrium and leiomyoma

A.G.A. Berto, S.M. Oba,  
Y.M. Michelacci  
and L.O. Sampaio

Departamento de Bioquímica, Escola Paulista de Medicina,  
Universidade Federal de São Paulo, São Paulo, SP, Brasil

## Abstract

### Correspondence

A.G.A. Berto  
Departamento de Bioquímica  
UNIFESP/EPM  
Rua Três de Maio, 100, 4º andar  
04044-020 São Paulo, SP  
Brasil  
Fax: +55-11-5573-6407

Presented at  
SIMEC 2000 - International  
Symposium on Extracellular  
Matrix, Angra dos Reis, RJ,  
Brazil, September 24-27, 2000.

Research supported by CNPq, CAPES,  
FAPESP and FINEP.

Received October 19, 2000  
Accepted February 5, 2001

In many tumors, the amount of chondroitin sulfate in the extracellular matrix has been shown to be elevated when compared to the corresponding normal tissue. Nevertheless, the degree of chondroitin sulfate increase varies widely. In order to investigate a possible correlation between the amount of chondroitin sulfate and tumor size, several individual specimens of human leiomyoma, a benign uterine tumor, were analyzed. The glycosaminoglycans from eight tumors were extracted and compared with those from the respective adjacent normal myometrium. The main glycosaminoglycan found in normal myometrium was dermatan sulfate, with small amounts of chondroitin sulfate and heparan sulfate. In leiomyoma, both dermatan sulfate and chondroitin sulfate were detected and the total amounts of the two galactosaminoglycans was increased in all tumors when compared to normal tissue. In contrast, the heparan sulfate concentration decreased in the tumor. To assess the disaccharide composition of galactosaminoglycans, these compounds were incubated with bacterial chondroitinases AC and ABC. The amounts of L-iduronic acid-containing disaccharides remained constant, whereas the concentration of D-glucuronic acid-containing disaccharides increased from 2 to 10 times in the tumor, indicating that D-glucuronic acid-containing disaccharides are responsible for the elevation in galactosaminoglycan concentration. This increase is positively correlated with tumor size.

### Key words

- Galactosaminoglycan
- Glycosaminoglycan
- Chondroitin sulfate
- Dermatan sulfate
- Leiomyoma
- Myometrium

Glycosaminoglycans (GAGs) are heteropolysaccharide chains made up of disaccharide-repeating units, in which one sugar is a hexosamine and the other is either a neutral sugar or a uronic acid. The GAG chains are sulfated to different extents and at different positions. The carboxyl groups of the uronic acids and the sulfate residues give the GAGs a polyanionic nature. With the exception of hyaluronic acid, which is not sulfated and occurs as free chains in tissues, every GAG is covalently linked to a protein, forming

proteoglycans.

Proteoglycans are important components of cell surface and extracellular matrices and individual proteoglycans interact specifically with other matrix components, such as collagen, laminin, and fibronectin, as well as with growth factors and cytokines (1-3). These interactions may affect cell growth, migration, adhesiveness, and differentiation, and many of these functions seem to be dependent on the GAG side chains (4). Sampaio and Dietrich (5) have first shown

that the type and quantity of GAGs in mature normal tissues differ from those found during embryonic development and in tumors (6). Chondroitin sulfate and dermatan sulfate, also referred to as galactosaminoglycans because they contain N-acetylgalactosamine as their hexosamine, are the most common GAGs in the extracellular matrix proteoglycans (7). The uronic acid in chondroitin sulfate is always D-glucuronic acid, whereas dermatan sulfate is a hybrid polymer, containing both L-iduronic acid and D-glucuronic acid residues.

Chondroitin sulfate was greatly increased in transformed cells in culture compared to normal cells (8). The amount of extracellular chondroitin sulfate in many tumors is also high when compared to the normal tissue of origin (6,9), but the magnitude of this increase varies widely, depending on the size and type of tumor (10).

The first study on the composition of sulfated GAGs in the uterus is from the 1960's (11), when it was shown that normal myometrium contains heparan sulfate, chondroitin sulfate, and dermatan sulfate. The proportions among these three GAGs did not vary significantly in pre- and postmenopause or in pre- and postgestation uterus. Although variations in the concentration of GAGs have been found, it was not possible to establish correlations between these variations and any of the analyzed parameters, such as patient age. The only correlations established were a significant decrease of uterine GAG during pregnancy (12,13) and an increase in chondroitin sulfate in leiomyoma, a benign tumor of the myometrium (6), and in leiomyosarcoma, a malignant tumor of the same tissue (14). Again, the magnitude of this increase varied from 2 to 20 times (6).

In order to determine if this variation in chondroitin sulfate concentration is a function of tumor class or size, or is due only to biological variability, several tumors of the same type were individually analyzed. The aim of the present investigation was to per-

form a more systematic study on the GAG composition of leiomyoma, in order to establish a possible correlation between GAGs and some parameters of neoplastic development in human myometrium.

Human leiomyomas were obtained shortly after surgical excision. Tumoral and adjacent normal myometrium were dissected and stored at -20°C until use. The frozen tissues were weighed and ground in 10 volumes of acetone. After standing overnight at room temperature, the tissues were collected by centrifugation and dried. About 0.45 g of the acetone powder thus obtained from normal myometrium and leiomyoma was incubated overnight with papain (2 mg/ml in 0.06 M phosphate-cysteine buffer, pH 6.5, containing 20 mM EDTA) at 60°C. Afterwards, debris was removed by centrifugation (5,000 g) for 15 min. Trichloroacetic acid and NaCl were added to the supernatant up to 10% and 1-M final concentrations, respectively. The mixture was left to stand for 10 min at 4°C and the precipitate formed was removed by centrifugation at 4,000 g for 10 min at 4°C. The GAGs were precipitated from the supernatant by the slow addition of 2 volumes of ethanol with shaking. After 18 h at -20°C the precipitate was collected by centrifugation, vacuum dried, resuspended in 0.5 ml of a solution containing deoxyribonuclease I (1 mg/ml) and 50 mM sodium acetate buffer, pH 6.0, and incubated at 30°C for 12 h. The GAGs were analyzed by a combination of agarose gel electrophoresis and degradation with specific bacterial mucopolysaccharidases.

Comparative results for leiomyoma and normal adjacent myometrium obtained from eight patients revealed important structural alterations of GAG content. Figure 1A shows the electrophoretic migration of GAGs extracted from one of these samples. The difference in GAG composition is evident. In normal myometrium, the main sulfated GAG is dermatan sulfate with small amounts of chondroitin sulfate and heparan sulfate,




Figure 1. Sulfated glycosaminoglycans (GAGs) from normal human myometrium (N) and leiomyoma (T). A, Agarose gel electrophoresis of GAGs extracted from leiomyoma and normal adjacent myometrium from patient 1: about 5 µl of the GAGs extracted as described in the text was submitted to agarose gel electrophoresis in 0.05 M 1,3-diaminopropane acetate buffer, pH 9.0, as previously described (6). Sd: mixture of standard GAGs; CS: chondroitin sulfate; DS: dermatan sulfate; HS: heparan sulfate. B, Amounts of CS plus DS (CS + DS) and HS extracted from eight tumors and from their respective adjacent normal myometrium. The GAGs were quantified by densitometry (525 nm) of the agarose gel slabs and the figure shows the individual results obtained. Data are reported as the mean ± SEM. \*P<0.01 compared to group N (Student t-test).

while in leiomyoma the main GAGs are chondroitin sulfate and dermatan sulfate. As chondroitin sulfate and dermatan sulfate migrate very close to each other, it was difficult to determine their individual concentrations. For this reason, the galactosaminoglycans (chondroitin sulfate and dermatan sulfate) were quantified together by densitometry of the agarose gel slabs. The concentration of galactosaminoglycans (expressed as mg per g dry tissue) increased in all samples of tumor tissue as compared to the respective adjacent normal myometrium (Figure 1B). In contrast, the heparan sulfate contents decreased in all cases (Figure 1B). The mean values and standard errors are also shown.

To investigate if this increase was due to both chondroitin sulfate and dermatan sulfate, the disaccharide units that compose these polymers were analyzed. The GAGs were incubated with chondroitinase AC (from *Flavobacterium heparinum*) (15) and chondroitinase ABC (from *Proteus vulgaris*) (16). The degradation products formed were analyzed by paper chromatography stained with alkaline silver nitrate (17) and quantified by densitometry. Both dermatan sulfate and chondroitin sulfate are hybrid polymers.

Chondroitin sulfate is composed of 4- and 6-sulfated disaccharide units, all of them containing D-glucuronic acid (18). Dermatan sulfate, on the other hand, contains both D-glucuronic and L-iduronic acid residues (19). The combined action of chondroitinases AC and ABC permits to assess the amounts of these disaccharide units in the polymers. Chondroitinase AC degrades chondroitin sulfate and the D-glucuronic acid-containing regions of dermatan sulfate. L-Iduronic acid-containing regions are not substrates for this enzyme but are degraded by chondroitinase ABC. Figure 2A shows that both 4-sulfated and 6-sulfated disaccharides were formed from normal and tumoral GAGs by chondroitinase AC (D-glucuronic acid) but more 4-sulfated disaccharides were produced by chondroitinase ABC, indicating the presence of L-iduronic acid. Nevertheless, the L-iduronic acid contents of tumors were unaltered when compared to the adjacent normal tissue, in contrast to the D-glucuronic acid contents that increased in all cases (Figure 2A) with magnitudes ranging from 1.7 to 11 times. Furthermore, Figure 2B shows that the amounts of D-glucuronic acid-containing disaccharides were positively correlated




Figure 2. Disaccharide composition of galactosaminoglycan from normal human myometrium (N) and leiomyoma (T). A, Concentration of unsaturated disaccharides produced by bacterial chondroitinases AC (D-glucuronic acid) and ABC (L-iduronic acid): incubation mixtures containing 30  $\mu$ g of glycosaminoglycans and  $20 \times 10^{-3}$  U of either *Flavobacterium heparinum* chondroitinase AC or *Proteus vulgaris* chondroitinase ABC were prepared in 0.1 M ethylenediamine-acetate buffer, pH 7, in 30  $\mu$ l final volumes. After overnight incubation at 37°C, the mixtures were applied to Whatman No. 1 filter paper and submitted to descending chromatography in isobutyric acid: 1.25 M  $\text{NH}_4\text{OH}$ , 5:3 (v/v). Reducing products were stained with alkaline silver nitrate (17). The disaccharides that contain D-glucuronic acid were estimated by densitometry of the disaccharides produced by the chondroitinase AC and the disaccharides that contain L-iduronic acid were estimated by the difference between the products formed by chondroitinase ABC and AC.  $\Delta$ Di-4S: Unsaturated 4-sulfated disaccharides;  $\Delta$ Di-6S: unsaturated 6-sulfated disaccharides. B, Correlation between tumor size and concentration of D-glucuronic acid-containing (circles and squares) and L-iduronic acid-containing (lozenges) disaccharide units in galactosaminoglycans. The values are reported as the leiomyoma/normal myometrium ratio.  $\Delta$ Di-6S (glc): unsaturated 6-sulfated disaccharides produced from D-glucuronic acid-containing disaccharide units;  $\Delta$ Di-4S (glc): unsaturated 4-sulfated disaccharides produced from D-glucuronic acid-containing disaccharide units;  $\Delta$ Di-4S (id): unsaturated 4-sulfated disaccharides produced from L-iduronic acid-containing disaccharide units.

with tumor size.

It is possible that the galactosaminoglycan chains, produced in progressively higher amounts as the tumor grows, were not properly processed, possibly due to a lower activity of uronic acid epimerase. This possibility was also raised by Sobue et al. (14), who analyzed benign and malignant tumors of the uterus, and observed that leiomyosarcomas contain considerably larger amounts of chon-

droitin sulfate than benign tumors, which contain dermatan sulfate with a small proportion of L-iduronic acid.

Elevated matrix chondroitin sulfate has also been correlated with the stage of prostate cancer, and the measurement of chondroitin sulfate concentrations at diagnosis has been proposed to allow stratification of patients with early-stage cancer for different therapies (20).

## References

1. Scott JE & Orford CR (1981). Dermatan sulphate-rich proteoglycan associates with rat tail-tendon collagen at the D band in the gap region. *Biochemical Journal*, 197: 213-216.
2. Couchman JR, Hook M, Rees DA & Timpl R (1983). Adhesion, growth and matrix production by fibroblast on laminin substrates. *Journal of Cell Biology*, 96: 177-183.
3. Iozzo RV (1998). Matrix proteoglycans: from molecular design to cellular function. *Annual Review of Biochemistry*, 67: 609-653.
4. Merle B, Durussel L, Delmas PD & Clezardin P (1999). Decorin inhibits cell migration through a process requiring its glycosaminoglycan side chain. *Journal of Cellular Biochemistry*, 75: 538-546.
5. Sampaio LO & Dietrich CP (1981). Changes of acidic mucopolysaccharides and mucopolysaccharidases during fetal development. *Journal of Biological Chemistry*, 256: 9205-9210.
6. Sampaio LO, Dietrich CP & Gianotti Filho O (1977). Changes in sulfated mucopolysaccharide composition of mammalian tissues during growth and cancerization. *Biochimica et Biophysica Acta*, 448: 123-131.
7. Vogel KG (1994). Glycosaminoglycans and proteoglycans. In: Yurchenco PD, Brik DE & Mechan RP (Editors), *Extracellular Matrix Assembly and Structure*. Academic Press, Inc., San Diego, CA.
8. Dietrich CP (1984). A model for cell-cell recognition and control of cell growth mediated by sulfated glycosaminoglycans. *Brazilian Journal of Medical and Biological Research*, 17: 5-15.
9. Iozzo RV (1988). Proteoglycans and neoplasia. *Cancer and Metastasis Reviews*, 7: 39-50.
10. Jerônimo SMB, Sales AO, Fernandes MZ, Melo FP, Sampaio LO, Dietrich CP & Nader HB (1994). Glycosaminoglycan structure and content differ according to the origins of human tumors. *Brazilian Journal of Medical and Biological Research*, 27: 2253-2258.
11. Loewi G & Consden R (1962). Acid mucopolysaccharides of the human uterus. *Nature*, 195: 148-150.
12. Cabrol D, Breton M, Berrou E, Visser A, Sureau C & Picard J (1980). Variations in the distribution of glycosaminoglycans in the uterine cervix of the pregnant woman. *European Journal of Obstetrics, Gynecology, and Reproductive Biology*, 10: 281-287.
13. Uldeberg N, Malmström A, Ekman G, Sheehan J, Ulmsten U & Wingerup L (1983). Isolation and characterization of dermatan sulphate proteoglycan from human uterine cervix. *Journal of Biochemistry*, 209: 497-503.
14. Sobue M, Takeuchi J, Yoshida K, Akao S, Fukatsu T, Nagasaka T & Nakashima N (1987). Isolation and characterization of proteoglycans from human nonepithelial tumors. *Cancer Research*, 47: 160-168.
15. Aguiar JAK & Michelacci YM (1999). Preparation and purification of *Flavobacterium heparinum* chondroitinases AC and B by hydrophobic interaction chromatography. *Brazilian Journal of Medical and Biological Research*, 32: 545-550.
16. Yamagata T, Saito H, Habuchi O & Suzuki S (1968). Purification and properties of bacterial chondroitinases and chondrosulfatases. *Journal of Biological Chemistry*, 243: 1523-1535.
17. Trevelyan WE, Procter DP & Harrison JS (1950). Detection of sugar on paper chromatograms. *Nature*, 166: 444-445.
18. Michelacci YM & Dietrich CP (1986). Structure of chondroitin sulphate from whale cartilage: distribution of 6- and 4-sulphated oligosaccharides in the polymer chains. *International Journal of Biological Macromolecules*, 8: 108-113.
19. Población CA & Michelacci YM (1986). Structural differences of dermatan sulfates from different origins. *Carbohydrate Research*, 147: 87-100.
20. Ricciardello C, Mayne K, Sykes PJ, Raymond WA, McCaul K, Marchall VR, Tilley WD, Skinner JM & Horsfall DJ (1997). Elevated stromal chondroitin sulfate glycosaminoglycan predicts progression in early-stage prostate cancer. *Clinical Cancer Research*, 3: 983-992.