

Original Article

Evaluation of lymphocyte levels in a random sample of 218 elderly individuals from São Paulo city

Daniela Teixeira¹
Ieda Maria Longo-Maugeri¹
Jair Lício Ferreira Santos²
Yeda Aparecida Oliveira Duarte³
Maria Lucia Lebrão³
Valquiria Bueno^{1,4}

¹Immunology Department,
Universidade Federal de São Paulo
– UNIFESP, São Paulo, SP, Brazil

²Faculdade de Medicina de
Ribeirão Preto, Universidade de
São Paulo – FMRP USP, Ribeirão
Preto, SP, Brazil

³Escola de Saúde Pública da
Universidade de São Paulo – USP
São Paulo, SP, Brazil

⁴Biosciences Department,
Universidade Federal de São
Paulo, Campus Baixada Santista –
UNIFESP, Santos, SP, Brazil

Conflict-of-interest disclosure:
The authors declare no competing
financial interest

Submitted: 4/18/2011
Accepted: 7/19/2011

Corresponding author:
Valquiria Bueno
Departamento de Biociências,
Universidade Federal de São Paulo
Rua Botucatu, 862 – 4º andar
04023-900 – São Paulo, SP, Brazil
Phone: 55 11 8962-2943
valquiria@nefro.epm.br

www.rbhh.org or www.scielo.br/rbhh

DOI: 10.5581/1516-8484.20110100

Background: Age-associated changes in the immune system cause decreased protection after vaccination and increased rates of infections and tumor development.

Methods: Lymphocyte percentages were compared by gender and age to establish differences between subtypes. Three mL blood samples were obtained from 218 randomly selected individuals (60-101 years old) who live in São Paulo city. Blood was lysed with Tris phosphate buffer and stained for 30 minutes with monoclonal antibodies (CD3PerCP, CD4FITC, CD8Pe, CD19Pe) for analysis by flow cytometry. Statistical analysis was by ANOVA.

Results: The percentage of CD4⁺ T cells (p -value = 0.005) and the CD4/CD8 ratio (p -value = 0.010) were lower in men, whereas the percentage of CD8⁺ T cells was lower (p -value = 0.002) in women; the percentage of B cells (CD19⁺) was similar between groups. Individuals grouped by gender and age range and compared showed a drop in CD4⁺ cells in 75 to 79-year-old men (female: 46.1% ± 8.1% and male: 38.8% ± 10.5%; p -value = 0.023). Also, the 80 to 84-year-old group of men had a higher percentage of CD8⁺ (female: 20.8% ± 8.2%, and male: 27.2% ± 8.2%; p -value = 0.032). Low percentages of B cells were detected in men in the 75 to 79-year-old (p -value = 0.003), 85 to 89-year-old (p -value = 0.020) and older than 90 year old (p -value = 0.002) age ranges.

Conclusion: Elderly men present with more changes in lymphocyte subsets compared to elderly women. These findings could demonstrate impairment in the immune response since the lower CD4⁺ in men would provide less help to B cells (also lower in men) in terms of antibody production. In addition, the increase in CD8⁺ cells in this group could represent chronic inflammation observed during the aging process.

Keywords: Aging; Sex distribution; Immune System; Lymphocytes; Flow Cytometry

Introduction

The rates of morbidity and mortality due to infectious diseases are high in elderly individuals. This population is more susceptible to severe infections, presents a slower recovery from infections, and the response to vaccination is not effective in all individuals. It is believed that changes in the immune system occurring in individuals after the age of 60 (immunosenescence) provide adequate conditions for susceptibility to infectious diseases, autoimmunity and the development of cancer. As an example, influenza vaccine is protective in 40-60% of over 65-year-old patients⁽¹⁾ while in younger individuals this percentage is higher.⁽²⁻⁵⁾

Immunosenescence affects all cells from the immune system to some extent. In addition, the contribution of this system to longevity and healthy aging remains unknown.⁽⁶⁻⁸⁾

Subclinical pathogenic infections, which are very common in the elderly, cause persistent inflammation and thus contribute to tumor development, heart attacks and strokes.⁽⁹⁾ It has already been demonstrated that in elderly individuals, persistent infections by herpes, cytomegalovirus (CMV) or parasites induce higher serum levels of pro-inflammatory factors (eg. IL-1, IL-6, tumor necrosis factor-alpha) leading to the aforementioned adverse clinical statuses.⁽¹⁰⁻¹³⁾ In addition, older CMV seropositive adults present up to 25% of the total CD8⁺ T cells pool specific for CMV immunodominant epitopes,⁽¹⁴⁾ the expansion of CMV-specific CD8⁺ T cells alters the capacity of the immune system to respond to other pathogens. These cells are able to secrete pro-inflammatory cytokines and contribute to an ongoing inflammatory process.⁽¹⁴⁾

Thymus involution, memory T cell accumulation, a decreased repertoire of naïve T cells and a diminished B cell population are changes that occur to the immune system

during aging; this can contribute to a deficient response in elderly individuals. Predicting responsiveness to vaccination, infectious diseases and tumor development using biological markers that distinguish between healthy and immunosenescent states is desirable as this might lead to more adequate therapies for the elderly population.

In order to determine possible changes in the subtypes of circulating lymphocytes in the elderly, these cells were evaluated in a random sample of 218 male and female individuals aged 60 to 101 years old from São Paulo city in Brazil. The percentages of CD4⁺ and CD8⁺ T cells, the CD4⁺/TCD8⁺ T cell ratio and the percentage of B cells (CD19⁺) were evaluated.

Methods

A random sample of 218 individuals from São Paulo city who agreed to participate in this project was investigated. This population (men and women) were aged from 60 to 101 years old. From 3 mL of blood collected in EDTA from each individual, 100 µL were used to determine each cell phenotype. Blood (100 µL) was lysed with Tris buffered solution for 10 minutes and centrifuged at 377 g for 5 minutes. Cells were washed in PBS twice and centrifuged at 377 g for 5 minutes. Cells were incubated with monoclonal antibodies (CD3PerCP, CD4FITC, CD8Pe - tritest, CD19Pe; BD Biosciences, San Jose, California) to determine the percentages of T and B lymphocytes using flow cytometry in a FACSCalibur cell counter (BD Biosciences).

Statistical analysis

ANOVA was used to compare CD4, CD8, CD4/CD8 and CD19 cells in respect to two variables (age and gender). Differences were evaluated between the following age ranges: 60-64, 65-69, 70-74, 75-79, 80-84, 85-89 and more than 90 years old. Bartlett's test was used for equal variances and significant differences between means were identified using Sidak's multiple comparisons.⁽¹⁵⁻¹⁷⁾ The level of statistical significance was defined for a p-value < 0.05.

Results

On comparing lymphocyte phenotypes in respect to age and gender, the percentages of CD4 and CD8 T cells and the CD4/CD8 T cell ratio were statistically similar for age but not for gender (Figure 1). On the other hand, significant differences were detected only for age when percentages of B cells (CD19⁺) were compared between age groups.

Possible differences between the percentages of cells of women and men in respect to age were then evaluated.

The percentage of CD3⁺CD4⁺ T cells in men was lower in all age groups compared to women (Table 1). However, a significant difference was only observed in the 75 to 79 age range (p-value = 0.023). An analysis of the percentage of




Figure 1 – Analysis of CD3⁺CD4⁺ T cells, CD3⁺CD8⁺ T cells, CD19⁺ B cells and the CD4⁺/CD8⁺ ratio in the blood of 218 over 60-year-old individuals. Women presented with a higher percentage of CD4⁺ T cells compared to men (p-value = 0.0005) whereas men presented with a higher percentage of CD8⁺ T cells compared to women. The percentage of CD19⁺ B cells was similar between women and men and the CD4⁺/CD8⁺ ratio was significantly higher in women.

Table 1. Percentage of TCD3⁺CD4⁺, TCD3⁺CD8⁺ and TCD4⁺/TCD8⁺ ratio determined in the peripheral blood of 60 to 101-year-old female and male individuals

Age range (years)	CD3 ⁺ CD4 ⁺		CD3 ⁺ CD8 ⁺		CD4 ⁺ /CD8 ⁺ ratio	
	Female %	Male %	Female %	Male %	Female	Male
60-64	45.6 ± 6.0 (n=9)	41.4 ± 15.3 (n=5)	24.6 ± 7.3 (n=9)	28.5 ± 11.4 (n=5)	2.1 ± 0.9 (n=9)	1.8 ± 1.1 (n=5)
65-69	43.9 ± 12.4 (n=19)	42.3 ± 11.0 (n=13)	18.8 ± 7.9 (n=19)	24.6 ± 9.8 (n=13)	2.8 ± 1.3 (n=19)	2.2 ± 1.4 (n=13)
70-74	44.7 ± 7.1 (n=16)	40.7 ± 7.8 (n=16)	22.3 ± 9.3 (n=16)	22.7 ± 11.4 (n=16)	2.5 ± 1.4 (n=16)	2.3 ± 1.2 (n=16)
75-79	46.1 ± 8.3 (n=21)	38.8 ± 10.5 (n=16)*	20.5 ± 5.2 (n=21)	23.6 ± 7.1 (n=16)	2.4 ± 0.8 (n=21)	1.9 ± 1.1 (n=16)
80-84	47.0 ± 12.5 (n=23)	40.5 ± 12.8 (n=13)	20.8 ± 8.2 (n=23)	27.2 ± 8.2 (n=13)#	2.6 ± 1.4 (n=23)	1.8 ± 1.3 (n=13)
85-90	41.6 ± 9.7 (n=18)	40.5 ± 10.5 (n=16)	19.6 ± 6.6 (n=18)	23.9 ± 12.2 (n=16)	2.4 ± 1.1 (n=18)	2.3 ± 1.4 (n=16)
90+	44.0 ± 9.4 (n=19)	38.7 ± 11.0 (n=14)	20.3 ± 8.1 (n=18)	24.7 ± 12.5 (n=14)	2.7 ± 1.6 (n=18)	1.9 ± 0.9 (n=14)

* In the 75 to 79-year-old age range there was a significant difference for CD3⁺CD4⁺ T cells between women and men (p-value = 0.023)

In the 80 to 84-year-old age range there was a significant difference for CD3⁺CD8⁺ T cells between women and men (p-value = 0.032)




Figure 2 – Percentages of CD3⁺CD4⁺ T cells, CD3⁺CD8⁺ T cells and CD19⁺ B cells of women (A, B and C) and men (D, E and F). No significant differences were seen for CD4 (A), CD8 (B) and CD19 (C) cell populations in women of different ages. Men presented with differences in CD19 cell populations (F) in the 70-74 age range compared with the 75-79 age range (p-value = 0.003), 85-89 age range (p-value 0.02) and the 90-94 age range (p-value 0.03). No significant differences were observed for the CD4 (D) and CD8 (E) cell populations in men

CD3⁺CD8⁺ T cell lymphocytes showed that this cell population was higher for men in all age groups with a significant difference for the 80-84 age range (p-value = 0.032). There was no significant difference in the CD4/CD8 T cell ratio on comparing women and men with similar ages.

A significant difference was found for the percentage of B cells (CD19) only for age. No statistical differences were found on comparing women of different age ranges. On the other hand, for men there was a statistical difference on comparing the 70 to 74-year-old age range with the 75 to 79-year-old (p-value = 0.003), 85 to 89-year-old (p-value = 0.020) and more than 90-year-old (p-value = 0.002) age ranges (Figure 2).

Discussion

Changes in the immune system occurring during aging have been associated with increases in morbidity and mortality.⁽¹⁸⁾ A Swedish study (Immune Longitudinal Study) established high levels of CD8 and low levels of CD4 and CD19 cells as predictive for mortality within 2 years.⁽¹⁹⁾ Using a random sample of elderly individuals from São Paulo city as the study group the same predictive parameters cited above were evaluated to see the effects of gender and age on the percentages of peripheral lymphocytes.

The percentage of CD3⁺CD4⁺ T cells was higher in women than in men when all individuals were compared without taking age into account. Moreover, the percentage of CD3⁺CD8⁺ T cells was higher in men than in women.

CD3⁺CD4⁺ T lymphocytes are essential to provide help for the proliferation (through the production of IL-2) and differentiation of other cells (i.e. IL-4 and IL-5 inducing B cell production of antibodies). Therefore, the low percentage of CD3⁺CD4⁺ T lymphocytes observed in men suggests that they are more likely to present impaired immune responses against pathogens and after vaccination.

Cytotoxic T lymphocytes increase in the elderly due to chronic infections (i.e. CMV) causing the accumulation of oligoclonal CD8⁺ T cells which interfere in immune responses either to vaccinations or infections.^(20,21) In this study men had a higher percentage of CD3⁺CD8⁺ T cells compared to women. On comparing ages this percentage was significantly higher in 80 to 84-year olds. De Fanis et al.⁽²²⁾ evaluated whether changes in the percentage of immune system cells could be related to frailty in 72 to 94-year old patients. Patients who were classified as frail had a significantly higher percentage of CD8⁺ T cells and a lower percentage of CD4⁺ T cells than the non-frail group. The results of this study suggest that elderly men present not only an impaired immune response but could also be frailer than women.

In this study there was a trend toward a higher CD4/CD8 ratio in women of all age groups. Hussein et al.⁽²³⁾ showed that young healthy individuals presented a mean CD4/CD8 ratio of 1.8 ± 0.8 which was slightly higher in over 50-year-old individuals with osteoarthritis (2.2 ± 0.4). Wikby et al.⁽²⁴⁾ observed similar CD4/CD8 ratios in healthy young men and women (1.55 for men and 1.77 for women). These authors showed that the CD4/CD8 ratio increases significantly during aging whereas in men this ratio remained relatively unchanged.

A Swedish study comparing age, matched 20 to 92-year-old male and female individuals and showed that women present a higher percentage of CD3⁺CD4⁺ cells than men during the entire adult lifespan. Moreover, the CD4/CD8 ratio was higher, albeit without statistical significance, in 20 to 59-year-old women whereas from 60 to 92 years old this ratio was significantly higher in women. The authors believe that there are specific gender differences in the processes that produce mature CD4⁺ and CD8⁺ cells in the thymus and that this could, at least in part, explain the higher longevity observed in women.⁽²⁴⁾

It has been shown that in over 60-year-old individuals, B lymphocytes (CD19⁺) are lower not only in numbers⁽²⁵⁾ but also in function. The capacity of these cells to produce antigen-specific antibodies after vaccination is less due to reduced expansion and differentiation. However the immune response generated before the immunosenescence process is maintained, as was showed by Yu et al.⁽²⁶⁾ during an evaluation of individuals who survived an influenza epidemic in 1918. The results of this study add new information to published data as the male population presented a trend toward a lower percentage of B cells than the female population. In the male group a more dramatic decrease in the B cell percentage was observed from the age of 75 years old. This result was different for women who presented a more constant percentage of B cells in all age ranges. This result suggests that men could be more susceptible to infections or produce lower antibody levels after vaccination.

In the elderly population the production of antibodies may also be impaired by changes in T helper cells which should send stimuli to B cells. In agreement this study, found that in men the percentage of CD3⁺CD4⁺ T cells was lower than in women. Therefore, the decreased percentage of B cells observed for men could be due to the decreased percentage (and maybe also function) of T cells found in male individuals.

Conclusion

This study showed that in a random sample of elderly from São Paulo, men suffered greater changes in lymphocyte subsets compared to women. These findings may represent impairment of the immune system as the lower percentage of CD4⁺ cells in men would provide less help to B cells (also

fewer in men) to produce antibodies. In addition, the increase in the percentage of CD8⁺ T cells in this group may represent chronic inflammation observed during the aging process. This profile observed in men reflects the parameters established in the Immune Longitudinal Study which are predictive of 2-year mortality rates. More studies are needed to confirm whether cell functions are also impaired differently according to gender and age in order to provide the best therapy for aging people.

Acknowledgments

The authors are thankful for financial support from FAPESP; Santos JLF and Bueno V are supported by fellowships from CNPq.

References

1. Vu T, Farish S, Jenkins M, Kelly H. A meta analysis of effectiveness of influenza vaccine in persons aged 65 years and over living in the community. *Vaccine*. 2002;20(13-14):1831-6.
2. Bernstein E, Kaye D, Abrutyn E, Gross P, Dorfman M, Murasko DM. Immune response to influenza vaccination in a large healthy elderly people population. *Vaccine*. 1999;17(1):82-94.
3. Prevention and control of influenza: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep*. 1999;48(RR-4):1-28.
4. Jefferson T, Rivetti D, Rivetti A, Rudin M, Di Pietrantonj C, Demicheli V. Efficacy and effectiveness of influenza vaccines in elderly people: a systematic review. *Lancet*. 2005;366(9492):1165-74. Comment in: *Lancet*. 2005;366(9503):2086. *Lancet*. 2005;366(9492):1139-40.
5. Goodwin K, Viboud C, Simonsen L. Antibody response to influenza vaccination in the elderly: a quantitative review. *Vaccine*. 2006;24(8):1159-69.
6. Miller RA. The aging immune system: primer and prospectus. *Science*. 1996;273(5271):70-4.
7. Hakim FT, Flomerfelt FA, Boyiadzis M, Gress RE. Aging, immunity and cancer. *Curr Opin Immunol*. 2004;16(2):151-6.
8. Pawelec G, Larbi A. Immunity and ageing in man: Annual Review 2006/2007. *Exp Gerontol*. 2008;43(1):34-8.
9. Falsey AR, Walsh EE, Francis CW, Looney RJ, Kolassa JE, Hall WJ, et al. Response of C-reactive protein and serum amyloid A to influenza A infection in older adults. *J Infect Dis*. 2001;183(7):995-9.
10. Luebke RW, Copeland CB, Andrews DL. Effects of aging on resistance to *Trichinella spiralis* infection exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Toxicology*. 1999;136(1):15-26.
11. Humphreys NE, Grenis RK. Effects of ageing on the immunoregulation of parasitic infection. *Infect Immun*. 2002;70(9):5148-57.
12. Saino N, Ferrari RP, Romano M, Rubolini D, Moller AP. Humoral immune response in relation to senescence and sexual ornamentation in the barn swallow (*Hirundo rustica*). *J Evol Biol*. 2003;16(6):1127-34.
13. Xainli J, Baisor M, Kastens W, Bockarie M, Adams JH, King CL. Age-dependent cellular immune responses to *Plasmodium vivax* Duffy binding protein in human. *J Immunol*. 2002;169(6):3200-7.
14. Pawelec G, Derhovanessian E, Larbi A, Strindhall J, Wikby A. Cytomegalovirus and human immunosenescence. *Rev Med Virol*. 2009;19(1):47-56.

15. Levene H. Robust tests for equality of variances. In: Olkin I, Ghurye SG, Hoeffding W, Madow WG, Mann HB, editors. *Contributions to probability and statistics: essays in honor of Harold Hotelling*. Menlo Park, CA: Stanford University Press; 1960. p. 278-92.
16. Sidak Z. Rectangular confidence regions for the means of multivariate normal distributions. *J Am Stat Assoc*. 1967;62:626-33.
17. Brown MB, Forsythe AB. Robust test for the equality of variances. *J Am Stat Assoc*. 1974;69:364-7.
18. Wayne SJ, Rhyne RL, Garry PJ, Goodwin JS. Cell mediated immunity as a predictor of morbidity and mortality in subjects over 60. *J Gerontol*. 1990;45(2):M45-8.
19. Ferguson F, Wikby A, Maxson P, Olsson J, Johansson B. Immune parameters in a longitudinal study of a very old population of Swedish people: a comparison between survivors and nonsurvivors. *J Gerontol A Biol Sci Med Sci*. 1995;50(6):B378-82.
20. Carson PJ, Nichol KL, O'Brien J, Hilo P, Janoff EN. Immune function and vaccine responses in healthy advanced elderly patients. *Arch Intern Med*. 2000;160(13):2017-24.
21. Goronzy JJ, Dulbright JW, Crowson CS, Poland GA, O'Fallon WM, Weyand CM. Value of immunological markers in predicting responsiveness to influenza vaccination in elderly individuals. *J Virol*. 2001;75(24):12182-7.
22. De Fanis U, Wang GC, Fedarko NS, Walston JD, Casalaro V, Leng SX. T-Lymphocytes expressing CC Chemokine receptor-5 are increased in frail older adults. *J Am Geriat Soc*. 2008;56(5):904-8.
23. Hussein MR, Fathi NA. Alterations of the CD4+, CD8+ T cells subsets, interleukins-1 β , IL-10, IL-17, tumor necrosis factor- α and soluble intercellular adhesion molecule-1 in rheumatoid arthritis and osteoarthritis: preliminary observations. *Pathol Oncol Res*. 2008;14(3):321-8.
24. Wikby A, Mansson IA, Johansson B, Strinbdhall J, Nilsson SE. The immune risk profile is associated with age and gender: findings from three Swedish population studies of individuals 20-100 years of age. *Biogerontology*. 2008;9(5):299-308.
25. Le Garff-Tavernier M, Béziat V, Decocq J, Siguret V, Gandjbakhch F, Pautas E, et al. Human NK cells display major phenotypic and functional changes over the life span. *Aging Cell*. 2010;9 (4):527-35.
26. Yu X, Tsibane PA, McGraw PA, House FS, Keefer CJ, Hicar MD, et al. Neutralizing antibodies derived from B cells of 1918 influenza pandemic survivors. *Nature*. 2008;455(7212):532-6.