

## Adhesin-Encoding Genes from Shiga Toxin-Producing *Escherichia coli* Are More Prevalent in Atypical than in Typical Enteropathogenic *E. coli*<sup>∇</sup>

Tânia A. T. Gomes,<sup>1\*</sup> Rodrigo T. Hernandez,<sup>1</sup> Alfredo G. Torres,<sup>2</sup> Fábila A. Salvador,<sup>1</sup>  
Beatriz E. C. Guth,<sup>1</sup> Tânia M. I. Vaz,<sup>3</sup> Kinue Irino,<sup>3</sup>  
Rosa M. Silva,<sup>1</sup> and Mônica A. M. Vieira<sup>1</sup>

Departamento de Microbiologia, Imunologia e Parasitologia, Universidade Federal de São Paulo, São Paulo, Brazil<sup>1</sup>; Department of Microbiology and Immunology and Department of Pathology, University of Texas Medical Branch, Galveston, Texas<sup>2</sup>; and Seção de Bacteriologia do Instituto Adolfo Lutz, São Paulo, Brazil<sup>3</sup>

Received 18 April 2011/Returned for modification 30 May 2011/Accepted 11 July 2011

**Four of six adhesin-encoding genes (*lpfA*, *paa*, *iha*, and *toxB*) from Shiga toxin-producing *Escherichia coli* strains were detected in typical and atypical enteropathogenic *E. coli* (EPEC) strains of various serotypes. Although the most prevalent gene was *lpfA* in both groups, *paa* was the only potential diarrhea-associated gene in atypical EPEC.**

The enteropathogenic *Escherichia coli* (EPEC) pathotype is subdivided into typical EPEC (tEPEC), which carries the large virulence EPEC adherence factor (EAF) plasmid (pEAF), and atypical EPEC (aEPEC), which lacks this plasmid (14). The pEAF encodes the bundle-forming pilus (BFP), which mediates a localized adherence (LA) pattern on HeLa/HEp-2 cells (13).

Both EPEC subgroups produce attaching and effacing (A/E) lesions on enterocytes, a phenotype associated with the chromosomal pathogenicity island termed the locus of enterocyte effacement (LEE). Intimin is an outer membrane protein encoded by the *eae* gene and is the fundamental LEE-encoded adhesin determining establishment of A/E lesions (reviewed in reference 13).

The LEE is also found in enterohemorrhagic *E. coli* (EHEC), a subgroup of the Shiga toxin-producing *E. coli* (STEC) pathotype (19). Various additional adhesins have been described in STEC, including Saa (STEC autoagglutinating adhesin) in LEE-negative STEC strains (18); Paa (porcine A/E-associated adhesin), involved in the initial adherence of porcine EPEC strains (4); Lpf (long polar fimbriae), an adhesin found in EHEC and other pathogenic *E. coli* (26, 27); ToxB (EHEC pO157 plasmid-encoded protein), which is involved in adherence of EHEC O157:H7 (24); Iha (IrgA homologue adhesin), an adhesin similar to *Vibrio cholerae* IrgA (23); and EspP (extracellular serine protease), which contributes to bovine intestinal colonization (9).

aEPEC strains are a highly heterogeneous group, with many strains presenting an assorted repertoire of virulence genes from various *E. coli* pathotypes (8, 11, 20, 25, 30). Interestingly, aEPEC strains carrying certain virulence genes or specific combinations are significantly associated with diarrheal disease (2, 11, 20, 21, 30). There is evidence indicating that some aEPEC

strains are in fact tEPEC or EHEC strains, which spontaneously lost either pEAF or the Shiga toxin-encoding genes (*stx* genes) (28). Furthermore, some comparative studies of the aEPEC genome have shown a closer relationship with EHEC than with tEPEC (16, 28), and others have shown that some clonal aEPEC groups are not related to any of these pathotypes (25).

In previous studies, we have searched for a number of *E. coli* adhesin-encoding genes in our EPEC collection, including the STEC *efa1/lifA* gene, but none of them were prevalent (11, 29,

TABLE 1. Adherence patterns and FAS test results from 100 aEPEC strains isolated from 73 patients and 27 controls

Phenotypic characteristic <sup>a</sup>	No. (%) of strains from:		
	Patients	Controls	Total
LAL	60 (82.2)	19 (70.4)	79 (79.0)
FAS <sup>+</sup>	55 (75.3) <sup>b</sup>	14 (51.8) <sup>b</sup>	69 (69.0)
FAS <sup>-</sup>	5 (6.8)	5 (18.5)	10 (10.0)
LAL/AA	3 (4.1)	0	3 (3.0)
FAS <sup>+</sup>	3 (4.1)	0	3 (3.0)
FAS <sup>-</sup>	0	0	0
AA	3 (4.1)	1 (3.7)	4 (4.0)
FAS <sup>+</sup>	2 (2.7)	1 (3.7)	3 (3.0)
FAS <sup>-</sup>	1 (1.3)	0	1 (1.0)
DA	2 (2.7)	4 (14.8)	6 (6.0)
FAS <sup>+</sup>	2 (2.7)	2 (7.4)	4 (4.0)
FAS <sup>-</sup>	0	2 (7.4)	2 (2.0)
NC	3 (4.1)	2 (7.4)	5 (5.0)
FAS <sup>+</sup>	2 (2.7)	2 (7.4)	4 (4.0)
FAS <sup>-</sup>	1 (1.3)	0	1 (1.0)
NA	1 (1.3)	1 (3.7)	2 (2.0)
D	1 (1.3)	0	1 (1.0)

\* Corresponding author. Mailing address: Departamento de Microbiologia, Imunologia e Parasitologia, Escola Paulista de Medicina, Universidade Federal de São Paulo, São Paulo, Brazil. Phone: 551150832980. Fax: 551155724711. E-mail: tatg.amaral@unifesp.br.

<sup>∇</sup> Published ahead of print on 27 July 2011.

<sup>a</sup> FAS, fluorescent actin staining; LAL, localized adherence-like; AA, aggregative adherence; DA, diffuse adherence; NC, noncharacteristic adherence; NA, nonadherent; D, cell detachment.

<sup>b</sup> Difference was statistically significant ( $P = 0.0302$ ).

TABLE 2. Prevalence of STEC adhesin-encoding genes in tEPEC and aEPEC strains from patients and controls

Adherence gene	No. (%) of strains:					
	aEPEC <sup>a</sup>			tEPEC <sup>b</sup>		
	Patients	Controls	Total	Patients	Controls	Total
<i>lpfA</i>	43 (58.9)	16 (59.3)	59 (59.0)	20 (50.0)	4 (66.7)	24 (52.2)
<i>paa</i>	35 (48.0) <sup>c</sup>	7 (25.9) <sup>c</sup>	42 (42.0) <sup>d</sup>	6 (15.0)	0	6 (13.0) <sup>d</sup>
<i>iha</i>	22 (30.1)	7 (25.9)	29 (29.0) <sup>e</sup>	3 (7.5)	1 (16.7)	4 (8.7) <sup>e</sup>
<i>toxB</i>	4 (5.5)	2 (7.4)	6 (6.0)	1 (2.5)	1 (16.7)	2 (4.35)
<i>espP</i>	3 (4.1)	0	3 (3.0)	0	0	0
<i>saa</i>	0	0	0	0	0	0
None	7 (9.6)	8 (29.6)	15 (15.0)	17 (42.5)	1 (16.6)	18 (39.1)

<sup>a</sup> The numbers of aEPEC strains studied were 73 from patients and 27 from controls.

<sup>b</sup> The numbers of tEPEC strains studied were 40 from patients and 6 from controls.

<sup>c</sup> *P* = 0.067.

<sup>d</sup> *P* = 0.0005.

<sup>e</sup> *P* = 0.0058.

30). Therefore, we aimed at investigating the prevalence of six other STEC adhesin-encoding genes and their potential association with diarrhea in aEPEC strains and compared them with those of tEPEC strains.

We examined a total of 146 *E. coli* strains, which were isolated from 113 diarrheic and 33 nondiarrheic children and adults in Brazil. The strains were characterized as EPEC based on the presence of the *eae* and absence of the *stx* genes (14).

TABLE 3. Distribution of STEC adhesin-encoding genes among aEPEC strains of distinct serotypes

Serotype	No. of strains	STEC adhesin-encoding gene(s)	<i>lpfA1</i> type	<i>lpfA2</i> type	Serotype	No. of strains	STEC adhesin-encoding gene(s)	<i>lpfA1</i> type	<i>lpfA2</i> type
O2ab:H45	1	<i>paa</i>	1		O177:H-	1			
O11:H2	1		2	1	NT:H-	3	<i>iha</i>		1
O11:H16	1					1			
O16:H-	1	<i>iha</i>				1	<i>toxB, iha</i>		
O19:H-	1	<i>iha</i>	1			1 <sup>c</sup>	<i>espP</i>		
O26:H-	1		2	1		1		2	1
	3	<i>paa</i>	2	1		1	<i>paa</i>	1	
	1	<i>iha</i>				1	<i>iha</i>	1	
	1 <sup>a</sup>	<i>iha, paa, espP</i>	2	1	NT:H2	1	<i>iha</i>		
O34:H-	1	<i>paa</i>	2	1	NT:H7	1 <sup>b</sup>	<i>toxB</i>		1
	1	<i>iha</i>	2	1	NT:H8	1			
O39:H-	1 <sup>b</sup>	<i>toxB, paa</i>				1 <sup>b</sup>			1
O49:H-	1	<i>iha</i>		1		1 <sup>b</sup>	<i>paa</i>	2	1
O49:H10	1 <sup>b</sup>	<i>toxB, paa</i>				1		2	1
O51:H-	1	<i>iha, paa</i>	2	1		1			1
O55:H7	6	<i>paa</i>	3	2	NT:H9	1	<i>iha</i>		
	1	<i>paa</i>	3		NT:H11	1			
O63:H6	2	<i>paa</i>				1	<i>iha</i>		1
O85:H-	1	<i>iha</i>		1	NT:H19	1		2	1
O93:H-	1	<i>iha, paa</i>	2	1	NT:H25	1		2	1
O98:H8	1	<i>iha, paa</i>		1	NT:H29,31	1			
O101:H33	1				NT:H33	1 <sup>b</sup>			
	1	<i>iha</i>			NT:H34	1		1	
O109:H9	1	<i>iha</i>		1		1	<i>iha</i>		
O111:H9	1					1	<i>paa</i>		
	1	<i>iha</i>		1	NT:H38	1			1
O119:H2	7 <sup>b</sup>	<i>paa</i>	2		NT:H40	1			
O123:H19	1		2	1	NT:H40,43	1			1
O124:H40	1		1			1			
O125:H6	2				NT:H46	1	<i>paa</i>		1
O128:H2	4	<i>paa</i>			NT:NT	1 <sup>b</sup>	<i>paa, toxB</i>		
O132:H8	2 <sup>b</sup>	<i>paa</i>	2	1		1 <sup>b</sup>	<i>paa</i>	1	1
O145:H-	1	<i>paa</i>			R:H-	1		1	1
O145:H34	1	<i>iha, paa</i>				1 <sup>c</sup>	<i>iha, paa, espP</i>	2	1
O154:H9	1	<i>iha</i>		1	R:H11	1			
O157:H-	2	<i>iha</i>			R:H28	1		1	
O157:H16	2				R:H33	1	<i>iha</i>		
O160:H19	1		2	1	R:H40	1 <sup>b</sup>			
O162:H-	1	<i>toxB, iha</i>							

<sup>a</sup> Strain carrying *ehxA* and *katP* genes.

<sup>b</sup> *bfpA*-positive strains lacking BFP production.

<sup>c</sup> Strain carrying *ehxA* gene.

Further classification as tEPEC or aEPEC was achieved by confirming BFP production by immunoblotting and by checking the *bfpA*-positive strains for their ability to produce LA on HeLa cells (after 3 h of incubation) (28). Strains lacking *bfpA* or carrying *bfpA* but lacking BFP production and showing an adherence pattern distinct from LA were classified as aEPEC (1, 12, 28). The aEPEC adherence patterns were defined after longer incubation periods (6 h) on HeLa cells (Table 1). The ability of all adherent strains to promote actin accumulation was evaluated by the fluorescent actin staining (FAS) test on HeLa cells (15). While all tEPEC strains were FAS positive (FAS<sup>+</sup>) (data not shown), this property was detected in only 83% of the aEPEC strains isolated from patients and control subjects (Table 1). Interestingly, none of the aEPEC adherence patterns were associated with diarrhea, but FAS<sup>+</sup> strains producing the LA-like (LAL) pattern were statistically associated with diarrhea.

The presence of five STEC adhesin-encoding genes (*espP*, *toxB*, *saa*, *iha*, and *paa*) was examined by colony hybridization under stringent conditions, using as probes fragments of these genes obtained by PCR (3, 4, 6, 17, 22) and labeled with [<sup>32</sup>P]dCTP. The presence of different *lpfA* alleles was investigated by PCR using primers and conditions previously described (26). Data were analyzed by two-tailed Fisher's exact test, and *P* values of  $\leq 0.05$  were considered statistically significant.

The *lpfA* genes were the most prevalent, followed by *paa*, *iha*, and *toxB* in both EPEC groups (Table 2). The *paa* and *iha* genes were significantly more frequent in aEPEC than in tEPEC strains, while *espP* was only found in three aEPEC strains, and *saa* was not detected (Table 2).

None of the adhesin-encoding genes were associated with diarrhea in both EPEC groups, but *paa* was more frequently found in aEPEC isolates from patients compared with isolates from controls, with this difference reflecting a trend to be statistically significant (Table 2). The association of *paa* with diarrhea has been previously observed in aEPEC isolates from Norwegian and Brazilian children (2, 21), but those studies did not examine tEPEC isolates.

Different combinations of the genes studied were found among diverse aEPEC and tEPEC serotypes (Tables 3 and 4). The distribution of *lpfA* in tEPEC agreed with our previous study (26); however, this is the first description of isolates *lpfA1-1* allele positive and *lpfA2* negative in serotype O142:H34 (Table 4). From the 100 aEPEC strains analyzed, 41 lack both *lpfA* alleles. Of the remaining 59 *lpfA*-positive strains, 20 contained the combination of *lpfA1-2* and *lpfA2-1* alleles. With the exception of one strain, all O55:H7 strains were *lpfA1-3* and *lpfA2-2*, thus confirming previous findings with EHEC O157:H7 and LEE-negative STEC strains (10, 26), that demonstrated that the presence of these two alleles is a reliable way to detect O157:H7 strains and the closely related O55:H- or O55:H7 serotypes. Although different combinations of these alleles were detected in aEPEC serotypes, we found that seven O119:H2 isolates carried the *lpfA1-2* allele and were negative for *lpfA2* (Table 3). Seven other isolates from different serotypes were also positive for the *lpfA1-1* allele and negative for *lpfA2*, and these alleles were found in rare tEPEC and aEPEC serotypes. Finally, a large number of EPEC strains were neg-

TABLE 4. Distribution of STEC adhesin-encoding genes among tEPEC<sup>a</sup> strains of distinct serotypes

Serotype	No. of strains	STEC adhesin-encoding gene	<i>lpfA1</i> type	<i>lpfA2</i> type
O55:H-	1	<i>paa</i>	3	2
	1		1	
O55:H6	4		1	
	1			
O86:H34	1			
	1		1	
O88:H25	5		2	1
	1	<i>iha</i>	2	1
O111:H-	6			
O111:H2	6			
O119:H6	3	<i>paa</i>		
	3			
O127:H6	1	<i>toxB</i>	1	
O127:H40	1			
	1	<i>toxB</i>		
O142:H6	2	<i>iha</i>	1	
	2 <sup>b</sup>	<i>paa</i>	1	
O142:H34	2		1	
	1	<i>iha</i>	1	
O145:H45	1		1,2	1
	2		1	

<sup>a</sup> *E. coli* strains carrying *eae* and *bfpA* and lacking *stx* genes, with all strains producing BFP.

<sup>b</sup> Only these two strains lacked the EAF probe sequence.

ative for both *lpfA1* and *lpfA2*, and no correlation with diarrhea was observed (Table 2).

We also found that all strains of the traditional aEPEC serotypes O55:H7, O119:H2, and O128:H2 carried *paa* and all O119:H2 strains carried only *lpfA1-2* and lacked *lpfA2*. In contrast, the O26:H- strains showed different allele combinations which may reflect distinct H types (serotypes); however, with one exception, they possessed the *lpfA1-2* and *lpfA2-1* alleles. In addition, tEPEC strains of serotypes O111:H- and O111:H2, which were the most prevalent isolates in São Paulo, Brazil, in the past (28), carried none of the genes investigated.

The three strains that harbored *espP* (O26:H-, NT:H-, and R:H-) carried this gene on large plasmids, which also contained *ehxA*, although only one possess *katP* (Table 3). These findings suggest that these strains could comprise EHEC strains that have lost the *stx* genes (EHEC-LST) (5, 7). In six strains (two tEPEC and four aEPEC strains) that carried both *bfpA* and *toxB*, Southern blot analysis indicated that *toxB* is either located in pEAF or in another plasmid of similar size (data not shown).

In conclusion, we showed that a subgroup of aEPEC strains, producing the LAL pattern and FAS positivity on HeLa cells, are statistically associated with diarrhea. In addition, the prevalence of the STEC adhesin-encoding genes studied is higher in aEPEC than in tEPEC, possibly reflecting an apparently closer relationship of some aEPEC strains to the STEC pathotype. Our case control analysis showed a trend of the *paa* gene to be associated with aEPEC diarrhea. However, more studies are necessary to confirm that these genes are expressed during human infections and to understand how they contribute to host colonization.

This work was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP; grant 08/53812-4) and Programa de Apoio a Núcleos de Excelência (PRONEX MCT/CNPq/FAPERJ). The work in A.G.T.'s laboratory was supported by NIH/NIAID grant no. 5R01AI079154-03.

The contents of this article are solely the responsibility of the authors and do not necessarily represent the official views of the NIAID or NIH.

#### REFERENCES

- Abe, C. M., et al. 2009. Virulence features of atypical enteropathogenic *Escherichia coli* identified by the *eae+* EAF-negative *stx*-genetic profile. *Diagn. Microbiol. Infect. Dis.* **64**:357–365.
- Afset, J. E., et al. 2006. Identification of virulence genes linked with diarrhea due to atypical enteropathogenic *Escherichia coli* by DNA microarray analysis and PCR. *J. Clin. Microbiol.* **44**:3703–3711.
- Badea, L., et al. 2003. Contribution of Efa1/LifA to the adherence of enteropathogenic *Escherichia coli* to epithelial cells. *Microb. Pathog.* **34**:205–215.
- Batisson, I., et al. 2003. Characterization of the novel factor Paa involved in the early steps of the adhesion mechanism of attaching and effacing *Escherichia coli*. *Infect. Immun.* **71**:4516–4525.
- Bielaszewska, M., et al. 2008. Shiga toxin-negative attaching and effacing *Escherichia coli*: distinct clinical associations with bacterial phylogeny and virulence traits and inferred in-host pathogen evolution. *Clin. Infect. Dis.* **47**:208–217.
- Brunder, W., H. Schmidt, M. Frosch, and H. Karch. 1999. The large plasmids of Shiga toxin-producing *Escherichia coli* (STEC) are highly variable genetic elements. *Microbiology* **145**:1005–1014.
- Bugarel, M., L. Beutin, F. Scheutz, E. Loukiadis, and P. Fach. 2011. Identification of genetic markers for differentiation of Shiga toxin-producing, enteropathogenic, and avirulent strains of *Escherichia coli* O26. *Appl. Environ. Microbiol.* **77**:2275–2281.
- Dulguer, M. V., et al. 2003. Atypical enteropathogenic *Escherichia coli* strains: phenotypic and genetic profiling reveals a strong association between enteroaggregative *E. coli* heat-stable enterotoxin and diarrhea. *J. Infect. Dis.* **188**:1685–1694.
- Dziva, F., A. Mahajan, P. Cameron, C. Currie, and I. J. McKen. 2007. EspP, a Type V-secreted serine protease of enterohaemorrhagic *Escherichia coli* O157:H7, influences intestinal colonization of calves and adherence to bovine primary intestinal epithelial cells. *FEMS Microbiol. Lett.* **271**:258–264.
- Galli, L., A. G. Torres, and M. Rivas. 2010. Identification of the long polar fimbriae gene variants in the locus of enterocyte effacement-negative Shiga toxin-producing *Escherichia coli* strains isolated from humans and cattle in Argentina. *FEMS Microbiol. Lett.* **308**:123–129.
- Gomes, T. A. T., et al. 2004. Emerging enteropathogenic *Escherichia coli* strains? *Emerg. Infect. Dis.* **10**:1851–1855.
- Hernandes, R. T., W. P. Elias, M. A. M. Vieira, and T. A. T. Gomes. 2009. An overview of atypical enteropathogenic *Escherichia coli*. *FEMS Microbiol. Lett.* **297**:137–149.
- Kaper, J. B., J. P. Nataro, and H. L. Mobley. 2004. Pathogenic *Escherichia coli*. *Nat. Rev. Microbiol.* **2**:123–140.
- Kaper, J. B. 1996. Defining EPEC. *Rev. Microbiol.* **27**:130–133.
- Knutton, S., T. Baldwin, P. H. Williams, and A. S. McNeish. 1989. Actin accumulation at sites of bacterial adhesion to tissue culture cells: basis of a new diagnostic test for enteropathogenic and enterohemorrhagic *Escherichia coli*. *Infect. Immun.* **57**:1290–1298.
- Moreira, F. C., et al. 2008. *Escherichia coli* strains of serotype O51:H40 comprise typical and atypical enteropathogenic *E. coli* strains and are potentially diarrheagenic. *J. Clin. Microbiol.* **46**:1462–1465.
- Paton, A. W., and J. C. Paton. 2002. Direct detection and characterization of Shiga toxin-producing *Escherichia coli* by multiplex PCR for *stx1*, *stx2*, *eae*, *ehxA*, and *saa*. *J. Clin. Microbiol.* **40**:271–274.
- Paton, A. W., P. Srimanote, M. C. Woodrow, and J. C. Paton. 2001. Characterization of Saa, a novel agglutinating adhesion produced by locus of enterocyte effacement-negative Shiga-toxigenic *Escherichia coli* strains that are virulent for humans. *Infect. Immun.* **69**:6999–7009.
- Paton, J. C., and A. W. Paton. 1998. Pathogenesis and diagnosis of Shiga toxin-producing *Escherichia coli* infections. *Clin. Microbiol. Rev.* **11**:450–479.
- Robins-Browne, R. M., et al. 2004. *Escherichia coli* and community-acquired gastroenteritis, Melbourne, Australia. *Emerg. Infect. Dis.* **10**:1797–1805.
- Scaletsky, I. C. A., K. R. S. Aranda, T. B. Souza, N. P. Silva, and M. B. Morais. 2009. Evidence of pathogenic subgroups among atypical enteropathogenic *Escherichia coli* strains. *J. Clin. Microbiol.* **47**:3756–3759.
- Szalo, I. M., et al. 2002. Presence in bovine enteropathogenic (EPEC) and enterohaemorrhagic (EHEC) *Escherichia coli* of genes encoding for putative adhesins of human EHEC strains. *Res. Microbiol.* **153**:653–658.
- Tarr, P. I., et al. 2000. Iha: a novel *Escherichia coli* O157:H7 adherence-conferring molecule encoded on a recently acquired chromosomal island of conserved structure. *Infect. Immun.* **68**:1400–1407.
- Tatsuno, I., et al. 2001. *toxB* gene on pO157 of enterohemorrhagic *Escherichia coli* O157:H7 is required for full epithelial cell adherence phenotype. *Infect. Immun.* **69**:6660–6669.
- Tennant, S. M., et al. 2009. Characterization of atypical enteropathogenic *E. coli* strains of clinical origin. *BMC Microbiol.* **9**:117.
- Torres, A. G., et al. 2009. The long polar fimbriae genes of pathogenic *Escherichia coli* strains as reliable markers to identify virulent isolates. *J. Clin. Microbiol.* **47**:2442–2451.
- Torres, A. G., et al. 2002. Identification and characterization of *lpfABCC'DE*, a fimbrial operon of enterohemorrhagic *Escherichia coli* O157:H7. *Infect. Immun.* **70**:5416–5427.
- Trabulsi, L. R., R. Keller, and T. A. T. Gomes. 2002. Typical and atypical enteropathogenic *Escherichia coli*. *Emerg. Infect. Dis.* **8**:508–513.
- Vieira, M. A. M., et al. 2010. Prevalence and characteristics of the O122 pathogenicity island in typical and atypical enteropathogenic *Escherichia coli* strains. *J. Clin. Microbiol.* **48**:1452–1455.
- Vieira, M. A. M., et al. 2001. Phenotypic and genotypic characteristics of *Escherichia coli* strains of non-enteropathogenic *E. coli* (EPEC) serogroups that carry *eae* and lack the EPEC adherence factor and Shiga toxin DNA probe sequences. *J. Infect. Dis.* **183**:762–772.