

Letter to the Editor

First description of a Shiga toxin-producing *Escherichia coli* O103:H2 strain isolated from sheep in Brazil

Fernando H. Martins¹, Beatriz E.C. Guth², Roxane M.F. Piazza³, Jorge Blanco⁴, Jacinta S. Pelayo¹

¹ Departamento de Microbiologia, Universidade Estadual de Londrina, Londrina, Brazil

² Departamento de Microbiologia, Parasitologia e Imunologia, Universidade Federal de São Paulo, São Paulo, Brazil

³ Laboratório de Bacteriologia, Instituto Butantan, São Paulo, Brazil

⁴ Laboratorio de Referencia de *E. coli*, Departamento de Microbiología y Parasitología, Facultad de Veterinaria, Universidad de Santiago de Compostela, Lugo, Spain

Key words: STEC O103:H2; sheep; virulence features

J Infect Dev Ctries 2014; 8(1):126-128. doi:10.3855/jidc.3615

(Received 01 April 2013 – Accepted 12 June 2013)

Copyright © 2014 Martins *et al.* This is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Shiga toxin-producing *Escherichia coli* (STEC) comprise an important group of zoonotic pathogens causing a broad spectrum of disorders in humans, including mild to severe diarrhea, hemorrhagic colitis (HC) and the life-threatening condition hemolytic uremic syndrome (HUS). Although most outbreaks and sporadic cases of STEC have been attributed to strains belonging to serotype O157:H7, the frequency of non-O157 STEC infections is increasing in several regions [1]. More than 400 non-O157 STEC serotypes have been described so far and a considerable proportion of them have been linked to human illness [1,2]. In Brazil, human infections due to STEC are mainly associated with non-O157 serotypes, of which O26:H11, O103:H2 and O111:H8/H- (non-motile) accounted for the majority of cases [3-5].

STEC O103:H2 was first described as a causative agent of HUS in 1992 [6]. Since then, serotype O103:H2 has been implicated either in outbreaks or in sporadic cases of gastroenteritis and HUS in Europe [7,8], Japan [9] and the United States [10,11]. In Brazil, STEC O103:H2 was isolated for the first time in 1986, and re-emerged years later as causative agent of infantile diarrhea and hemolytic anemia [4,5]. Despite the fact that multiple STEC strains have already been isolated from farm animals, such as cattle [12] and pigs [13] in Brazil, none of them belonged to the known pathogenic serotype O103:H2 and did not display the virulence repertoire commonly associated

with severe human disease (i.e., *stx* plus *eae* and/or *ehxA* genes). This is the first report of STEC O103:H2 isolated and characterized from animals in Brazil.

The Study

Ten sheep flocks located on southern Brazil were tested between April and September 2010. Fecal samples were collected from 130 healthy animals, streaked onto MacConkey agar and lactose-fermenting colonies were biochemically characterized as *E. coli* [14]. STEC strains were identified by detection of *stx1*, *stx2*, *eae* and *ehxA* virulence genes [15] and serotyped as previously described [16]. Twenty-three different STEC serotypes were detected (data not shown), including O103:H2, carrying *stx1*, *eae* and *ehxA* genes which was isolated from a 4-week-old lamb. Intimin type ϵ was identified in this strain [16]. The *stx1 eae- ϵ ehxA* virulence gene profile is commonly observed among STEC O103:H2 strains [4,5,7,9]. *Stx1* expression was confirmed by cytotoxicity and neutralization assays on Vero cells [17], and enterohemolysin production was evidenced by the appearance of lysis zone on washed sheep blood agar plates [17]. The genotypic and phenotypic characteristics observed were similar to those of O103:H2 strains previously isolated from patients with diarrhea and hemolytic anemia in Brazil (Table 1).

Table 1. Characteristics of ovine STEC O103:H2 compared to human O103:H2 strains isolated in Brazil.

Strain	Serotype	Origin ⁽¹⁾	Genotypic and phenotypic characteristics ⁽²⁾				Reference
			Virulence profile	Intimin type	Stx	Ehly	
JN2b	O103:H2	S	<i>stx1 eae ehxA</i>	ε	+	+	This study
437/01	O103:H2	HA	<i>stx1 eae ehxA</i>	ε	+	+	[4]
651-1	O103:H2	HD	<i>stx1 eae ehxA</i>	ε	+	+	[4]
495-12	O103:H2	HD	<i>stx1 eae ehxA</i>	ε	+	+	[4,5]
91	O103:H2	HD	<i>stx1 stx2 eae ehxA</i>	ε	+	+	[4]

⁽¹⁾S: sheep; HA: hemolytic anemia; HD: human diarrhea

⁽²⁾Stx: cytotoxic activity; Ehly: production of enterohemolysin

A low prevalence of STEC O103:H2 in sheep has been also documented in other studies [18], supporting our results and suggesting that the occurrence of this serotype in ovine seems to be uncommon. However, STEC O103:H2 outbreaks and sporadic cases have been traced to contact with animals [7] as well as to consumption of contaminated meat [11], indicating that transmission of strains belonging to this serotype can occur between animals and humans.

Conclusion

The potential role of animals in the epidemiology of STEC O103:H2 human infection is still poorly understood in Brazil. In the present study, we described for the first time the isolation and characterization of STEC O103:H2 of animal origin in this country. This strain exhibited virulence features similar to those of human clinical strains, suggesting that sheep may be carriers, albeit at low frequency, of potentially human-pathogenic STEC O103:H2. However, more studies are needed to establish sheep and other animal species as source of STEC O103:H2 infection in our settings.

Acknowledgements

This work was supported by grants from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES-Brazil) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq-Brazil).

References

- Johnson KE, Thorpe CM, Sears CL (2006) The emerging clinical importance of non-O157 Shiga-toxin producing *Escherichia coli*. Clin Infect Dis 43: 1587-1595.
- WHO (World Health Organization) (1998) Zoonotic non-O157 Shiga toxin-producing *Escherichia coli* (STEC), 23-26 June 1998. Berlin, Germany, Geneva, Switzerland: World Health Organization. W.H.O./CSR/APH/98.8-30
- Vaz TM, Irino K, Kato MA, Dias AM, Gomes TA, Medeiros MI, Rocha MM, Guth BE (2004) Virulence properties and characteristics of Shiga toxin-producing *Escherichia coli* in São Paulo, Brazil, from 1976 through 1999. J Clin Microbiol 42: 903-905.
- Guth BE, Vaz TM, Gomes TA, Chinarelli SH, Rocha MM, de Castro AF, Irino K (2005) Re-emergence of O103:H2 Shiga toxin-producing *Escherichia coli* infections in São Paulo, Brazil. J Med Microbiol 54: 805-806.
- Irino K, Vaz TM, Medeiros MI, Kato MA, Gomes TA, Vieira MA, Guth BE (2007) Serotype diversity as a drawback in the surveillance of Shiga toxin-producing *Escherichia coli* infections in Brazil. J Med Microbiol 56: 565-567.
- Mariani-Kurkdjian P, Denamur E, Milon A, Picard B, Cave H, Lambert-Zechovsky N, Loirat C, Goullet P, Sansonetti PJ, Elion J (1993) Identification of a clone of *Escherichia coli* O103:H2 as a potential agent of hemolytic-uremic syndrome in France. J Clin Microbiol 31:296-301.
- Prager R, Liesegang A, Voigt W, Rabsch W, Fruth A, Tshape H (2002) Clonal diversity of Shiga toxin-producing *Escherichia coli* O103:H2/H- in Germany. Infect Genet Evol 1: 265-275.
- Heinikainen S, Pohjanvirta T, Eklund M, Siitonen A, Pelkonen S (2007) Tracing Shigatoxigenic *Escherichia coli* O103, O145 and O174 infections from farm residents to cattle. J Clin Microbiol 45: 3817-3820.
- Igushi A, Iyoda S, Ohnishi M, EHEC Study Group (2012) Molecular characterization reveals three distinct clonal groups among clinical shiga toxin-producing *Escherichia coli* strains of serogroup O103. J Clin Microbiol 50: 2894-2900.
- Brooks JT, Sowers EG, Wells JG, Greene KD, Griffin PM, Hoekstra RM, Strockbyne MA (2005) Non-O157 Shiga toxin-producing *Escherichia coli* infections in the United States, 1983-2002. J Infect Dis 192: 1422-1429.
- Rounds JM, Rigdon CE, Muhl LJ, Forstner M, Danzeisen GT, Koziol BS, Taylor C, Shaw BT, Smith KE (2012) Non-O157 Shiga toxin-producing *Escherichia coli* associated with venison. Emerg Infect Dis 18: 279-282.
- Aidar-Ugrinovich L, Blanco J, Blanco M, Blanco JE, Leomil L, Dahbi G, Mora A, Onuma DL, Silveira WD, Pestana de Castro AF (2007) Serotypes, virulence genes, and intimin types of Shiga toxin-producing *Escherichia coli* (STEC) and enteropathogenic *E. coli* (EPEC) isolated from calves in São Paulo, Brazil. Int J Food Microbiol 115:297-306.
- Borges CA, Beraldo LG, Maluta RP, Cardozo MV, Guth BE, Rigobelo EC, de Ávila FA (2012) Shiga toxigenic and atypical enteropathogenic *Escherichia coli* in the feces and carcasses of slaughtered pigs. Foodborne Pathog Dis 9: 1119-1125.
- Irino K, Kato MA, Vaz TM, Ramos II, Souza MA, Cruz AS, Gomes TA, Vieira MA, Guth BE (2005) Serotypes and virulence markers of Shiga toxin-producing *Escherichia coli* (STEC) from dairy cattle in São Paulo State, Brazil. Vet Microbiol 105: 29-36.
- Paton AW and Paton JC (1998) Detection and characterization of Shiga toxigenic *Escherichia coli* by using

- multiplex PCR assays for *stx1*, *stx2*, *eaeA*, enterohemorrhagic *E. coli hlyA*, *rfbO111* and *rfbO157*. *J Clin Microbiol* 36: 598-602.
16. Blanco M, Blanco JE, Mora A, Rey J, Alonso JM, Hermoso M, Hermoso J, Alonso MP, Dahbi G, González EA, Bernárdez MI, Blanco J (2003) Serotypes, virulence genes, and intimin types of Shiga toxin (verotoxin)-producing *Escherichia coli* isolates from healthy sheep in Spain. *J Clin Microbiol* 41: 1351-1356.
 17. Mendes-Ledesma MR, Rocha LB, Bueris V, Krause G, Beutin L, Franzolin MR, Trabulsi LR, Elias WP, Piazza RM (2008) Production and characterization of rabbit polyclonal sera against Shiga toxins Stx1 and Stx2 for detection of Shiga toxin-producing *Escherichia coli*. *Microbiol Immunol* 52: 484-491.
 18. Sekse C, Sunde M, Hopp P, Bruheim T, Cudjoe KS, Kvitle B, Urdahl AM (2013) Occurrence of potentially human-pathogenic *Escherichia coli* O103 in Norwegian sheep. *Appl Environ Microbiol* 79: 7502-7509.

Corresponding author

Jacinta Sanchez Pelayo. PhD.

Departamento de Microbiologia, Universidade Estadual de

Londrina, Rodovia Celso Garcia Cid Pr 445 Km 380, Brazil

Tel: (55-043) 3371-4494(Work)

Fax: (55-043) 3371-4291

Email: jspelayo@gmail.com

Conflict of interests: No conflict of interests is declared.