



Role of adenosine in the antiepileptic effects of deep brain stimulation

Maisa F. Miranda¹, Clement Hamani^{2,3,4*}, Antônio-Carlos G. de Almeida¹, Beatriz O. Amorim², Carlos E. Macedo⁵, Maria José S. Fernandes⁶, José N. Nobrega³, Mayra C. Aarão¹, Ana Paula Madureira¹, Antônio M. Rodrigues¹, Monica L. Andersen⁵, Sergio Tufik⁵, Luiz E. Mello² and Luciene Covolan²

¹ Laboratório de Neurociência Experimental e Computacional, Universidade Federal de São João del-Rei, São João del-Rei, Brazil

² Disciplina de Neurofisiologia, Universidade Federal de São Paulo, São Paulo, Brazil

³ Behavioural Neurobiology Laboratory, Centre for Addiction and Mental Health, Toronto, Canada

⁴ Division of Neurosurgery, Toronto Western Hospital, University of Toronto, Toronto, Canada

⁵ Departamento de Psicobiologia, Universidade Federal de São Paulo, São Paulo, Brazil

⁶ Disciplina de Neurologia Experimental, Universidade Federal de São Paulo, São Paulo, Brazil

Edited by:

Tycho M. Hoogland, Netherlands
Institute for Neuroscience,
Netherlands

Reviewed by:

Scott Krahl, VA Greater Los Angeles
Healthcare System, USA
Hongyu Sun, University of
Pennsylvania, USA

*Correspondence:

Clement Hamani, Division of
Neurosurgery, Toronto Western
Hospital, University of Toronto, 399
Bathurst Street, Toronto,
ON M5T2S8, Canada
e-mail: Clement.Hamani@uhn.on.ca;
c.hamani@sympatico.ca

Despite the effectiveness of anterior thalamic nucleus (AN) deep brain stimulation (DBS) for the treatment of epilepsy, mechanisms responsible for the antiepileptic effects of this therapy remain elusive. As adenosine modulates neuronal excitability and seizure activity in animal models, we hypothesized that this nucleoside could be one of the substrates involved in the effects of AN DBS. We applied 5 days of stimulation to rats rendered chronically epileptic by pilocarpine injections and recorded epileptiform activity in hippocampal slices. We found that slices from animals given DBS had reduced hippocampal excitability and were less susceptible to develop ictal activity. In live animals, AN DBS significantly increased adenosine levels in the hippocampus as measured by microdialysis. The reduced excitability of DBS *in vitro* was completely abolished in animals pre-treated with A1 receptor antagonists and was strongly potentiated by A1 receptor agonists. We conclude that some of the antiepileptic effects of DBS may be mediated by adenosine.

Keywords: deep brain stimulation, epilepsy, thalamus, anterior nucleus, seizures, adenosine

INTRODUCTION

Deep brain stimulation (DBS) involves the delivery of current to the brain parenchyma through implanted electrodes. Over the last decade, preclinical and clinical studies have shown that DBS applied to the anterior thalamic nucleus (AN) reduces seizure rate and increases the latency for the development of seizures and status epilepticus (SE; Mirski et al., 1997; Hodaie et al., 2002; Hamani et al., 2004, 2008; Kerrigan et al., 2004; Andrade et al., 2006; Takebayashi et al., 2007a,b; Fisher et al., 2010). Despite the efficacy of AN DBS, mechanisms involved in the antiepileptic effects of this therapy remain elusive.

We have recently shown that AN DBS reduced the frequency of spontaneous recurrent seizures in chronic epileptic animals (Covolan et al., 2014). In addition, animals receiving stimulation had a significant decrease in hippocampal excitability. This proved to be a more reliable measure of the effectiveness of DBS than frequency of seizures, due to the variability in seizure rate that occurs not only across animals but also in the same animals over time (Covolan et al., 2014).

Adenosine and other purines are known modulators of neuronal excitability and seizure activity in experimental animals and humans (Dunwiddie, 1980; Turski et al., 1985; O'Brien, 1988; Arvin et al., 1989; During and Spencer, 1992; Boison

et al., 1999; Huber et al., 2001; Zuchora et al., 2001; Ansel et al., 2004; Boison, 2005, 2006, 2008, 2009; Cunha, 2005; Vianna et al., 2005; Pagonopoulou et al., 2006; Li et al., 2007, 2011; Boison and Stewart, 2009; Van Dycke et al., 2010, 2011; Hargus et al., 2012). In rats, adenosine injections directly onto the epileptic focus (Ansel et al., 2004) or the hippocampus improve seizure control (Van Dycke et al., 2010). Synthetic polymers capable of releasing adenosine into the ventricle have also been shown to reduce seizure activity (Boison et al., 1999). In preclinical models, SE up-regulates adenosine kinase (ADK), a key metabolic enzyme for the regulation of extracellular adenosine (Aronica et al., 2011). Fibroblasts engineered to release adenosine by inactivating ADK and adenosine deaminase grafted into the ventricles of kindled rats protected the animals from behavioral seizures and afterdischarges (Huber et al., 2001).

In thalamic slices from ferrets, stimulation-induced glutamate and adenosine release abolished spontaneous spindle oscillations (Tawfik et al., 2010). In slices from mice, thalamic stimulation increased ATP release and induced an accumulation in extracellular adenosine (Bekar et al., 2008). In mice, A1 receptor activation depressed excitatory transmission *in vitro* and reduced both tremor- and DBS-induced side effects *in vivo* (Bekar et al., 2008).

Bearing in mind the fact that adenosine modulates seizures in animal models and is involved in mechanisms of thalamic DBS for tremor (Bekar et al., 2008), we hypothesized that this nucleoside could be one of the substrates involved in the antiepileptic effects of AN DBS.

RESULTS

AN-DBS INDUCES ADENOSINE RELEASE IN HIPPOCAMPUS

As shown in **Figure 1**, AN DBS increased hippocampal levels of adenosine in both epileptic and non-epileptic animals (DBS main effect $F_{(1,12)} = 527.59$, $p < 0.0001$). This increase was significantly higher in Pilo + DBS rats compared to Pilo + Sham animals (**Figure 1A**). The same pattern was observed after 5 days of stimulation (DBS main effect $F_{(1,12)} = 297.90$, $p < 0.0001$; **Figure 1B**), with the additional observation that mean adenosine levels in the DBS + Pilo group were almost twice as high as their own values on stimulation day 1 ($p < 0.001$). In both days 1 and 5, adenosine remained increased for at least 1 h following DBS discontinuation. Levels of this nucleoside were significantly higher in chronic epileptic animals as compared to non-epileptic controls (Pilo main effect $F_{(1,12)} = 114.88$, $p < 0.001$).

ANTIEPILEPTIC EFFECTS OF AN DBS *IN VITRO*


We have previously found that animals receiving DBS for 5 days had a significant decrease in hippocampal excitability measured *in vitro* with electrophysiology (Covolan et al., 2014). In preclinical models, the antiepileptic effects of adenosine occur largely through A1 receptors (Simonato et al., 1994; Ekonomou et al., 1998; Gouder et al., 2003; Rebola et al., 2003; Avsar and Empson, 2004; Mohammad-Zadeh et al., 2005, 2009; Fedele et al., 2006;


Rosim et al., 2011; Silva et al., 2011). Based on these premises, we hypothesized that the increased adenosine levels recorded after DBS might be interacting with A1 receptors to reduce hippocampal excitability. To test this hypothesis, we recorded activity from slices of rats previously given 5 days of stimulation with or without co-administration of the A1 antagonist 8-Cyclopentyl-1,3-dipropylxanthine (DPCPX) or the A1 agonist R-isomer of N6-phenylisopropyladenosine (R-Pia). In these experiments, we did not record *in vivo* seizure rate due to the marked variability observed across animals in our previous report (Covolan et al., 2014).

Confirming our previous findings (Covolan et al., 2014), epileptiform activity in slices of animals previously given AN DBS had longer latencies ($p = 0.04$), with shorter ($p < 0.01$) and smaller direct current (DC) shifts ($p < 0.01$), and a smaller spike amplitude ($p < 0.01$), as compared to non-stimulated epileptic controls (**Figure 2**). These effects were completely abolished in animals given DPCPX (**Figure 2**). In contrast, the protective effects of DBS were somewhat potentiated in animals given DBS plus R-Pia. Slices from this group had lower spike amplitudes ($p < 0.01$) and a longer latency for the development of epileptiform events ($p < 0.01$) as compared to those from rats receiving DBS alone (**Figure 2**). No differences in activity have been recorded in slices from animals given saline, DPCPX or R-Pia alone.

DISCUSSION

Reduced hippocampal excitability and susceptibility for the development of ictal activity after AN DBS in slices from chronic





epileptic animals were completely blocked by treatment with A1 antagonists and potentiated by the co-administration of the A1 agonist R-Pia. This suggests that hippocampal adenosine released after AN stimulation may activate A1 receptors and subsequently lead to structural and/or functional changes capable of rendering the hippocampus less susceptible to ictal events.

The first interesting aspect of our study was the long-lasting stimulation-induced release of adenosine. This nucleoside has been shown to inhibit seizure activity both *in vitro* and in animal models of epilepsy (Dunwiddie, 1980; Van Dycke et al., 2010). We found that in chronic epileptic animals, the increase in adenosine release after DBS not only outlasted the administration of current but was significantly more pronounced at long-term. Bearing in mind the antiepileptic effects of adenosine, one may speculate that the frequency of seizures and epileptiform activity after AN stimulation could become less pronounced with time. Though this has not been demonstrated in animals, it has been reported in clinical trials (Fisher et al.,

2010). In a recent randomized controlled multicenter study, the effects of AN DBS have been shown to build up with time, being far more pronounced at long-term (Fisher et al., 2010). Our results obviously do not allow us to conclude that adenosine is involved in the increased efficacy of DBS over time but they do suggest a path to be explored in future studies.

Due to the high variability in seizure rate observed in the pilocarpine model, we did not record the frequency of behavioral or electrographic seizures in our dialysis experiments. That said, animals selected for that part of the study were not those with high number of seizures at baseline, as we were afraid of them damaging the dialysis set up while having generalized ictal events (this in fact did happen with one of our rats). A few animals in the epileptic groups did have seizures that lasted a few seconds during the experiments. As our samples were collected over 30 min, associated changes in levels of adenosine could not be captured. Another interesting finding in our study was the fact that epileptic animals had higher baseline levels of adenosine in

the hippocampus compared to non-epileptic controls. Whether this is a compensatory mechanism to hinder seizure activity remains to be investigated. Finally, one may argue that the simple implantation of electrodes in the brain might have contributed to our results. We find this unlikely for two main reasons. First, electrodes were implanted a few millimeters away from the collection site (AN vs. hippocampus). Second, no differences in baseline adenosine levels have been recorded between DBS treated animals and their respective controls (i.e., prior to stimulation onset).

We have focused on A1 receptors as these are highly expressed in the hippocampus (Swanson et al., 1995) and have been extensively demonstrated to be involved in the inhibition of ictal activity (Fedele et al., 2006). Overall, our electrophysiological experiments have shown that (1) the decrease in hippocampal excitability observed after AN stimulation was almost completely reversed in slices from animals previously given A1 antagonists; and (2) previous administration of A1 agonists to animals receiving DBS potentiated the *in vitro* antiepileptic effects of stimulation. Our results suggest that adenosine released after DBS would activate A1 receptors and induce functional and/or structural changes that might have decreased hippocampal excitability and the propensity for seizures. In our study, electrophysiology results in slices from animals receiving R-Pia alone were not identical to those recorded in the DBS group. This may be due to the fact that the effects of DBS are fairly complex, involving synaptic and non-synaptic mechanisms as well as multiple neurotransmitter systems. A more suggestive finding supporting the role of A1 receptors was the reduced hippocampal excitability recorded in animals previously given DBS was completely abolished by the co-administration of DPCPX.

As a final remark we would like to point out two caveats of our study. First, we have only recorded electrophysiological data 5 days after DBS onset, an interval that was based on our previous report (Covolan et al., 2014). In contrast, our dialysis experiments were conducted at two time points. As in our previous studies DBS was shown to induce an increase in neurotransmitter release right after stimulation onset (Hamani et al., 2010), we have decided to record adenosine levels within the first hours of DBS administration. To understand whether changes in adenosine levels after stimulation were sustained and still noticeable within the timeframe of our electrophysiological recordings, we have also conducted dialysis experiments after 5 days of DBS. At present, it is still unclear whether the administration of A1 antagonists blocks the effects of 1-day DBS treatment. As a second remark, we did not record behavioral seizure rate, dialysis samples and *in vitro* electrophysiology in the same animals for the reasons described above. As a result, we could not correlate these events and are unable to provide a clear causal relationship between hippocampal excitability, adenosine levels and the frequency of behavioral seizures.

In summary, we found that the purinergic system may be involved in the antiepileptic effects of AN DBS. In the clinical scenario, 40–60% of patients with treatment refractory epilepsy are responsive to AN DBS (Fisher et al., 2010). Therapeutic strategies for individuals considered as being non-responders remain unclear. Though no pure A1 agonist is available for clinical use,

our findings may pave the way for future investigation on the use of medications that influence A1 receptors to potentiate the effects of stimulation.

MATERIALS AND METHODS

Protocols were approved by the Animal Care committee of the Universidade Federal de São Paulo (CEUA 1482/11).

SURGERY AND AN STIMULATION

Four months after pilocarpine injections (Pilo; 320 mg/Kg i.p.), adult male Wistar rats (250–300 g) were divided in groups to receive DBS or sham-surgery (holes drilled in the skull). Animals with implanted electrodes that did not receive stimulation have not been included as such treatment did not lead to significant differences as compared to non-implanted controls in our previous work (Covolan et al., 2014). Under ketamine/xylazine anesthesia (100/7.5 mg/kg i.p.), animals in the DBS group had insulated stainless steel electrodes (cathodes; 250 μ m diameter; 0.5 mm exposed length) bilaterally implanted into the AN (anteroposterior -1.5 mm, lateral ± 1.5 mm, depth 5.2 mm) (Paxinos and Watson, 1998). A screw implanted over the right somatosensory cortex was used as the anode. On the second postoperative week, DBS was administered for 5 days (8 h/day) using a portable stimulator (St Jude Medical, Plano, TX) at 130 Hz, 90 μ s, and 100 μ A. These settings were chosen as they were effective against pilocarpine-induced chronic epileptic seizures in our previous study (Covolan et al., 2014).

ELECTROPHYSIOLOGY

Prior to the experiments, chronic epileptic rats were given 5 days of DBS with or without the concomitant administration of the A1 receptor antagonist DPCPX (50 μ g/kg/day i.p.) or agonist R-Pia (25 μ g/kg/day i.p.) (Vianna et al., 2005). Following CO₂ narcosis, animals were decapitated, brains were removed and 400 μ m hippocampal slices were cut on a vibratome. Slices were then individually transferred to an interface-type chamber, placed on a membrane (0.4 μ m Millicell culture plate inserts; Millipore, Badford, MA) and continuously bathed with artificial cerebrospinal fluid (aCSF; 127 mM NaCl, 2 mM KCl, 1.5 mM MgSO₄, 1.1 mM KH₂PO₄, 26 mM NaHCO₃, 2 mM CaCl₂, and 10 mM glucose) at 33°C under a stream of moisturized 95% O₂—5% CO₂. Slices were left undisturbed for 40 min before being perfused with a zero calcium and 8 mM potassium solution for approximately 1 h and 30 min (de Almeida et al., 2008).

Extracellular field potentials were recorded from the hippocampal dentate gyrus (DG), as previously described (de Almeida et al., 2008). For each group, slices containing the dorsal hippocampus of six different animals were analyzed (14 extracellular potentials per animal). The following parameters were studied: latency for epileptiform discharges after perfusion with a zero calcium high potassium solution; DC shift; amplitude of population spikes; and event duration. A digital Fourier transform was used to quantify DC shifts. Once in the frequency domain, the event signal was recalculated taking into account only components below 10 Hz. This process allowed the analysis of DC shifts without interference from population spikes. Event duration was calculated by subtracting the final time from the initial time of an event (de Almeida et al., 2008).

MICRODIALYSIS

Along with the implantation of DBS electrodes, a microdialysis cannula was placed into the right dorsal hippocampus (antero-posterior -3.7 mm, lateral 2.4 mm, depth 5.1 mm) (Paxinos and Watson, 1998). Five days later, a microdialysis probe (CMA/12-2 mm, CMA Microdialysis, Sweden) was inserted into the target and perfused with a Ringer's solution at $1 \mu\text{L}/\text{min}$ (Macedo et al., 2012). Dialysate samples were collected every 30 min in vials containing 0.25 mM ascorbic acid, Na_2EDTA 0.27 mM, 0.1 M acetic acid on the first and last days of treatment at baseline (2 h), during DBS (3 h), and in the 3 h following stimulation offset ($n = 4$ animals per group).

Adenosine levels were detected with capillary electrophoresis (Beckman-Coulter PACE/MDQ, Beckman Coulter, USA) equipped with a laser-induced fluorescence detector (ZETALIF Picometrics, France; He-Cd laser wavelength 410 nm). Dialysate samples ($20 \mu\text{l}$) were derivatized with $20 \mu\text{l}$ of a chloroacetaldehyde solution (0.15 M) (Sigma, USA). This mixture was allowed to react at 100°C for 20 min, subsequently refrigerated at 4°C and then injected into the capillary system. Adenosine analysis was carried out using a 20 mM sodium-phosphate buffer (containing 1% SDS, pH 8). The applied voltage was 20 kV. Samples were hydrodynamically injected (0.4 nl) in a fused-silica capillary (Beckman-Coulter, $25 \mu\text{m}$ i.d. and $375 \mu\text{m}$ o.d., 60 cm in length, and an effective length of 40 cm). The capillary was sequentially flushed with sodium hydroxide (0.1 M), ultra-pure water, and running buffer between each analysis. Electropherograms were acquired at 15 Hz using a PACE MDQ software (Beckman-Coulter).

STATISTICAL ANALYSES AND HISTOLOGY

Dialysis results for baseline, DBS and post-DBS periods were each analyzed with a mixed model ANOVA, with pilocarpine and DBS as between-subject factors and time as a within-subjects factor. As electrophysiological data was non-parametric, results were compared using the Kruskal Wallis and Mann Whitney tests. Dialysis data is displayed as mean \pm standard error. Electrophysiology results are shown as median \pm quartiles. Statistical significance was considered when $p \leq 0.05$.

ACKNOWLEDGMENTS

Support to conduct the work was provided by FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo) 2011150680-2; 50950-2, Fapemig (Fundação de Amparo à Pesquisa do Estado de Minas Gerais), CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico), CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior), and AFIP (Associação Fundo de Incentivo à Pesquisa).

REFERENCES

- Andrade, D. M., Zumsteg, D., Hamani, C., Hodaie, M., Sarkissian, S., Lozano, A. M., et al. (2006). Long-term follow-up of patients with thalamic deep brain stimulation for epilepsy. *Neurology* 66, 1571–1573. doi: 10.1212/01.wnl.0000206364.19772.39
- Anschel, D. J., Ortega, E. L., Kraus, A. C., and Fisher, R. S. (2004). Focally injected adenosine prevents seizures in the rat. *Exp. Neurol.* 190, 544–547. doi: 10.1016/j.expneurol.2004.07.017
- Aronica, E., Zurolo, E., Iyer, A., de Groot, M., Anink, J., Carbonell, C., et al. (2011). Upregulation of adenosine kinase in astrocytes in experimental and human temporal lobe epilepsy. *Epilepsia* 52, 1645–1655. doi: 10.1111/j.1528-1167.2011.03115.x
- Arvin, B., Neville, L. F., Pan, J., and Roberts, P. J. (1989). 2-chloroadenosine attenuates kainic acid-induced toxicity within the rat striatum: relationship to release of glutamate and Ca^{2+} influx. *Br. J. Pharmacol.* 98, 225–235. doi: 10.1111/j.1476-5381.1989.tb16886.x
- Avsar, E., and Empson, R. M. (2004). Adenosine acting via A1 receptors, controls the transition to status epilepticus-like behaviour in an in vitro model of epilepsy. *Neuropharmacology* 47, 427–437. doi: 10.1016/j.neuropharm.2004.04.015
- Bekar, L., Libionka, W., Tian, G. F., Xu, Q., Torres, A., Wang, X., et al. (2008). Adenosine is crucial for deep brain stimulation-mediated attenuation of tremor. *Nat. Med.* 14, 75–80. doi: 10.1038/nm1693
- Boison, D. (2005). Adenosine and epilepsy: from therapeutic rationale to new therapeutic strategies. *Neuroscientist* 11, 25–36. doi: 10.1177/1073858404269112
- Boison, D. (2006). Adenosine kinase, epilepsy and stroke: mechanisms and therapies. *Trends Pharmacol. Sci.* 27, 652–658. doi: 10.1016/j.tips.2006.10.008
- Boison, D. (2008). The adenosine kinase hypothesis of epileptogenesis. *Prog. Neurobiol.* 84, 249–262. doi: 10.1016/j.pneurobio.2007.12.002
- Boison, D. (2009). Engineered adenosine-releasing cells for epilepsy therapy: human mesenchymal stem cells and human embryonic stem cells. *Neurotherapeutics* 6, 278–283. doi: 10.1016/j.nurt.2008.12.001
- Boison, D., Scheurer, L., Tseng, J. L., Aebischer, P., and Mohler, H. (1999). Seizure suppression in kindled rats by intraventricular grafting of an adenosine releasing synthetic polymer. *Exp. Neurol.* 160, 164–174. doi: 10.1006/exnr.1999.7209
- Boison, D., and Stewart, K. A. (2009). Therapeutic epilepsy research: from pharmacological rationale to focal adenosine augmentation. *Biochem. Pharmacol.* 78, 1428–1437. doi: 10.1016/j.bcp.2009.08.005
- Covolán, L., de Almeida, A. C., Amorim, B., Cavarsan, C., Miranda, M. F., Aarão, M. C., et al. (2014). Effects of anterior thalamic nucleus deep brain stimulation in chronic epileptic rats. *PLoS One* 9:e97618. doi: 10.1371/journal.pone.0097618
- Cunha, R. A. (2005). Neuroprotection by adenosine in the brain: from A(1) receptor activation to A(2A) receptor blockade. *Purinergic Signal.* 1, 111–134. doi: 10.1007/s11302-005-0649-1
- de Almeida, A. C., Rodrigues, A. M., Scorza, F. A., Cavalheiro, E. A., Teixeira, H. Z., Duarte, M. A., et al. (2008). Mechanistic hypotheses for nonsynaptic epileptiform activity induction and its transition from the interictal to ictal state—computational simulation. *Epilepsia* 49, 1908–1924. doi: 10.1111/j.1528-1167.2008.01686.x
- Dunwiddie, T. V. (1980). Endogenously released adenosine regulates excitability in the in vitro hippocampus. *Epilepsia* 21, 541–548. doi: 10.1111/j.1528-1157.1980.tb04305.x
- During, M. J., and Spencer, D. D. (1992). Adenosine: a potential mediator of seizure arrest and postictal refractoriness. *Ann. Neurol.* 32, 618–624. doi: 10.1002/ana.410320504
- Ekonomou, A., Angelatou, F., Vergnes, M., and Kostopoulos, G. (1998). Lower density of A1 adenosine receptors in nucleus reticularis thalami in rats with genetic absence epilepsy. *Neuroreport* 9, 2135–2140. doi: 10.1097/00001756-199806220-00042
- Fedele, D. E., Li, T., Lan, J. Q., Fredholm, B. B., and Boison, D. (2006). Adenosine A1 receptors are crucial in keeping an epileptic focus localized. *Exp. Neurol.* 200, 184–190. doi: 10.1016/j.expneurol.2006.02.133
- Fisher, R., Salanova, V., Witt, T., Worth, R., Henry, T., Gross, R., et al. (2010). Electrical stimulation of the anterior nucleus of thalamus for treatment of refractory epilepsy. *Epilepsia* 51, 899–908. doi: 10.1111/j.1528-1167.2010.02536.x
- Gouder, N., Fritschy, J. M., and Boison, D. (2003). Seizure suppression by adenosine A1 receptor activation in a mouse model of pharmacoresistant epilepsy. *Epilepsia* 44, 877–885. doi: 10.1046/j.1528-1157.2003.03603.x
- Hamani, C., Diwan, M., Macedo, C. E., Brandão, M. L., Shumake, J., Gonzalez-Lima, F., et al. (2010). Antidepressant-like effects of medial prefrontal cortex deep brain stimulation in rats. *Biol. Psychiatry* 67, 117–124. doi: 10.1016/j.biopsych.2009.08.025
- Hamani, C., Ewerton, F. I., Bonilha, S. M., Ballester, G., Mello, L. E., and Lozano, A. M. (2004). Bilateral anterior thalamic nucleus lesions and high-frequency stimulation are protective against pilocarpine-induced seizures and status epilepticus. *Neurosurgery* 54, 191–195; discussion 195–197. doi: 10.1227/01.neu.0000097552.31763.ae

- Hamani, C., Hodaie, M., Chiang, J., del Campo, M., Andrade, D. M., Sherman, D., et al. (2008). Deep brain stimulation of the anterior nucleus of the thalamus: effects of electrical stimulation on pilocarpine-induced seizures and status epilepticus. *Epilepsy Res.* 78, 117–123. doi: 10.1016/j.eplepsyres.2007.09.010
- Hargus, N. J., Jennings, C., Perez-Reyes, E., Bertram, E. H., and Patel, M. K. (2012). Enhanced actions of adenosine in medial entorhinal cortex layer II stellate neurons in temporal lobe epilepsy are mediated via A(1)-receptor activation. *Epilepsia* 53, 168–176. doi: 10.1111/j.1528-1167.2011.03337.x
- Hodaie, M., Wennberg, R. A., Dostrovsky, J. O., and Lozano, A. M. (2002). Chronic anterior thalamus stimulation for intractable epilepsy. *Epilepsia* 43, 603–608. doi: 10.1046/j.1528-1157.2002.26001.x
- Huber, A., Padrun, V., Déglon, N., Aebischer, P., Möhler, H., and Boison, D. (2001). Grafts of adenosine-releasing cells suppress seizures in kindling epilepsy. *Proc. Natl. Acad. Sci. U S A* 98, 7611–7616. doi: 10.1073/pnas.131102898
- Kerrigan, J. F., Litt, B., Fisher, R. S., Cranstoun, S., French, J. A., Blum, D. E., et al. (2004). Electrical stimulation of the anterior nucleus of the thalamus for the treatment of intractable epilepsy. *Epilepsia* 45, 346–354. doi: 10.1111/j.0013-9580.2004.01304.x
- Li, Y., Fan, S., Yan, J., Li, B., Chen, F., Xia, J., et al. (2011). Adenosine modulates the excitability of layer II stellate neurons in entorhinal cortex through A1 receptors. *Hippocampus* 21, 265–280. doi: 10.1002/hipo.20745
- Li, T., Steinbeck, J. A., Lusardi, T., Koch, P., Lan, J. Q., Wilz, A., et al. (2007). Suppression of kindling epileptogenesis by adenosine releasing stem cell-derived brain implants. *Brain* 130, 1276–1288. doi: 10.1093/brain/awm057
- Macedo, C. E., Angst, M. J., Gobaille, S., Schlee, C., Guignard, B., Guiberteau, T., et al. (2012). Prefrontal dopamine release and sensory-specific satiety altered in rats with neonatal ventral hippocampal lesions. *Behav. Brain Res.* 231, 97–104. doi: 10.1016/j.bbr.2012.02.041
- Mirski, M. A., Rossell, L. A., Terry, J. B., and Fisher, R. S. (1997). Anticonvulsant effect of anterior thalamic high frequency electrical stimulation in the rat. *Epilepsy Res.* 28, 89–100. doi: 10.1016/s0920-1211(97)00034-x
- Mohammad-Zadeh, M., Amini, A., Mirnajafi-Zadeh, J., and Fathollahi, Y. (2005). The role of adenosine A(1) receptors in the interaction between amygdala and entorhinal cortex of kindled rats. *Epilepsy Res.* 65, 1–9. doi: 10.1016/j.eplepsyres.2005.03.012
- Mohammad-Zadeh, M., Mirnajafi-Zadeh, J., Fathollahi, Y., Javan, M., Jahanshahi, A., Noorbakhsh, S. M., et al. (2009). The role of adenosine A(1) receptors in mediating the inhibitory effects of low frequency stimulation of perforant path on kindling acquisition in rats. *Neuroscience* 158, 1632–1643. doi: 10.1016/j.neuroscience.2008.11.008
- O'Brien, D. R. (1988). The adenosine hypothesis of epilepsy. *Med. Hypotheses* 27, 281–284. doi: 10.1016/0306-9877(88)90007-2
- Pagonopoulou, O., Efthimiadou, A., Asimakopoulos, B., and Nikolettos, N. K. (2006). Modulatory role of adenosine and its receptors in epilepsy: possible therapeutic approaches. *Neurosci. Res.* 56, 14–20. doi: 10.1016/j.neures.2006.05.010
- Paxinos, G., and Watson, C. (1998). *The Rat Brain in Stereotaxic Coordinates*. San Diego, London: Academic Press.
- Rebola, N., Coelho, J. E., Costenla, A. R., Lopes, L. V., Parada, A., Oliveira, C. R., et al. (2003). Decrease of adenosine A1 receptor density and of adenosine neuromodulation in the hippocampus of kindled rats. *Eur. J. Neurosci.* 18, 820–828. doi: 10.1046/j.1460-9568.2003.02815.x
- Rosim, F. E., Persike, D. S., Nehlig, A., Amorim, R. P., de Oliveira, D. M., and Fernandes, M. J. (2011). Differential neuroprotection by A(1) receptor activation and A(2A) receptor inhibition following pilocarpine-induced status epilepticus. *Epilepsy Behav.* 22, 207–213. doi: 10.1016/j.yebeh.2011.07.004
- Silva, I. R., Nehlig, A., Rosim, F. E., Vignoli, T., Persike, D. S., Ferrandon, A., et al. (2011). The A1 receptor agonist R-Pia reduces the imbalance between cerebral glucose metabolism and blood flow during status epilepticus: could this mechanism be involved with neuroprotection? *Neurobiol. Dis.* 41, 169–176. doi: 10.1016/j.nbd.2010.09.004
- Simonato, M., Varani, K., Muzzolini, A., Bianchi, C., Beani, L., and Borea, P. A. (1994). Adenosine A1 receptors in the rat brain in the kindling model of epilepsy. *Eur. J. Pharmacol.* 265, 121–124. doi: 10.1016/0014-2999(94)90421-9
- Swanson, T. H., Drazba, J. A., and Rivkees, S. A. (1995). Adenosine A1 receptors are located predominantly on axons in the rat hippocampal formation. *J. Comp. Neurol.* 363, 517–531. doi: 10.1002/cne.903630402
- Takebayashi, S., Hashizume, K., Tanaka, T., and Hodozuka, A. (2007a). Anti-convulsant effect of electrical stimulation and lesioning of the anterior thalamic nucleus on kainic acid-induced focal limbic seizure in rats. *Epilepsy Res.* 74, 163–170. doi: 10.1016/j.eplepsyres.2007.03.007
- Takebayashi, S., Hashizume, K., Tanaka, T., and Hodozuka, A. (2007b). The effect of electrical stimulation and lesioning of the anterior thalamic nucleus on kainic acid-induced focal cortical seizure status in rats. *Epilepsia* 48, 348–358. doi: 10.1111/j.1528-1167.2006.00948.x
- Tawfik, V. L., Chang, S. Y., Hitti, F. L., Roberts, D. W., Leiter, J. C., Jovanovic, S., et al. (2010). Deep brain stimulation results in local glutamate and adenosine release: investigation into the role of astrocytes. *Neurosurgery* 67, 367–375. doi: 10.1227/01.neu.0000371988.73620.4c
- Turski, W. A., Cavalheiro, E. A., Ikonomidou, C., Mello, L. E., Bortolotto, Z. A., and Turski, L. (1985). Effects of aminophylline and 2-chloroadenosine on seizures produced by pilocarpine in rats: morphological and electroencephalographic correlates. *Brain Res.* 361, 309–323. doi: 10.1016/0006-8993(85)91302-2
- Van Dycke, A., Raedt, R., Dauwe, I., Sante, T., Wyckhuys, T., Meurs, A., et al. (2010). Continuous local intrahippocampal delivery of adenosine reduces seizure frequency in rats with spontaneous seizures. *Epilepsia* 51, 1721–1728. doi: 10.1111/j.1528-1167.2010.02700.x
- Van Dycke, A., Raedt, R., Vonck, K., and Boon, P. (2011). Local delivery strategies in epilepsy: a focus on adenosine. *Seizure* 20, 376–382. doi: 10.1016/j.seizure.2011.03.003
- Vianna, E. P., Ferreira, A. T., Doná, F., Cavalheiro, E. A., and Da Silva Fernandes, M. J. (2005). Modulation of seizures and synaptic plasticity by adenosinergic receptors in an experimental model of temporal lobe epilepsy induced by pilocarpine in rats. *Epilepsia* 46(Suppl. 5), 166–173. doi: 10.1111/j.1528-1167.2005.01027.x
- Zuchora, B., Turski, W. A., Wielosz, M., and Urbańska, E. M. (2001). Protective effect of adenosine receptor agonists in a new model of epilepsy—seizures evoked by mitochondrial toxin, 3-nitropropionic acid, in mice. *Neurosci. Lett.* 305, 91–94. doi: 10.1016/s0304-3940(01)01816-x

Conflict of Interest Statement: Clement Hamani is a consultant for St Jude Medical. The other authors do not have a conflict of interest related to this article.

Received: 19 July 2014; accepted: 17 September 2014; published online: 02 October 2014.

Citation: Miranda MF, Hamani C, de Almeida A-CG, Amorim BO, Macedo CE, Fernandes MJS, Nobrega JN, Aarão MC, Madureira AP, Rodrigues AM, Andersen ML, Tufik S, Mello LE and Covolan L (2014) Role of adenosine in the antiepileptic effects of deep brain stimulation. *Front. Cell. Neurosci.* 8:312. doi: 10.3389/fncel.2014.00312

This article was submitted to the journal *Frontiers in Cellular Neuroscience*. Copyright © 2014 Miranda, Hamani, de Almeida, Amorim, Macedo, Fernandes, Nobrega, Aarão, Madureira, Rodrigues, Andersen, Tufik, Mello and Covolan. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.