



Title	Generation and chromosome mapping of expressed sequence tags (ESTs) from a human infant thymus
Author(s)	Hwang, MY; Kang, YJ; Kim, YH; Scherer, SW; Tsui, LC; Sohn, U
Citation	Genome, 1999, v. 42 n. 3, p. 457-464
Issued Date	1999
URL	http://hdl.handle.net/10722/42308
Rights	Creative Commons: Attribution 3.0 Hong Kong License

Generation and chromosome mapping of expressed sequence tags (ESTs) from a human infant thymus

Mee-Yul Hwang, Yoon-Joong Kang, Young-Ho Kim, Steve W. Scherer, Lap-Chee Tsui, and Uik Sohn

Abstract: In an effort to identify novel genes that are expressed differentially in an infant thymus, we constructed an oligo-d(T) primed cDNA library from a human infant thymus followed by single-run partial sequencing to generate expressed sequence tags (ESTs). Characterization of more than 1400 sequences enabled us to convert human thymus transcripts into 1223 useful ESTs. These ESTs consisted of 613 (50.1%) showing homology to known human genes, 51 (4.2%) matching to genes from other species, 289 (23.6%) matching ESTs of unknown functions, and 182 (14.9%) being novel transcripts. The expression profile of an infant thymus features a high number of genes related to cell division-DNA synthesis and gene-protein expression, indicating the active growth stage of an infant thymus. To identify the chromosomal localization of 43 thymus ESTs, PCR-based mapping was performed using a human-rodent somatic cell hybrid or radiation hybrid mapping panel. The results indicated that several novel genes were determined to be located in the vicinity of previously mapped disease loci; histidinemia loci, plasminogen Tochigi disease loci, Ehlers-Danlos syndrome, hypertriglyceridemia, thyroid resistance locus, ocular albinism, galactosemia, porphyria variegata, Charcot-Marie-tooth disease, FEOM (fibrosis of extraocular muscles), Prader-Willi syndrome.

Key words: cDNA, expression profile, radiation hybrid mapping, disease locus.

Résumé : Afin d'identifier de nouveaux gènes qui sont exprimés de manière différentielle dans le thymus d'un nouveau-né, une banque d'ADNc provenant du thymus d'un nouveau-né et dont la synthèse a été initiée avec une amorce oligo-d(T) a été produite. Ces ADNc ont ensuite fait l'objet d'une seule réaction de séquençage de façon à générer une collection de séquences exprimées (EST). Une caractérisation de plus de 1400 séquences a permis de convertir des transcrits humains en 1223 EST utiles. Ces EST comprenaient 613 (50,1%) séquences montrant une homologie à des gènes humains connus, 51 (4,2%) étaient homologues à des gènes connus chez d'autres espèces, 289 (23,6%) étaient apparentés à des EST de fonction inconnue et 182 (14,9%) correspondaient à de nouvelles séquences. Le profil d'expression d'un thymus de nouveau-né montre un nombre élevé d'expression de gènes impliqués dans la division cellulaire-synthèse de l'ADN et dans l'expression des gènes-protéines, ce qui est le reflet du stade de croissance active du thymus chez le nouveau-né. Afin de déterminer la position chromosomique de 43 EST du thymus, une cartographie a été réalisée à l'aide d'une technique PCR employée sur des hybrides somatiques ou une collection d'hybrides résultant d'irradiation. Les résultats indiquent que plusieurs nouveaux gènes sont localisés à proximité de loci de maladies : des loci d'histidinémie, de la maladie du plasminogène Tochigi, du syndrome d'Ehlers-Danlos, de l'hypertriglycéridémie, de la résistance thyroïdienne, de l'albinisme oculaire, de la galactosémie, du porphyra variegata, de la maladie dentaire Charcot-Marie, de la FEOM (fibrose des muscles extraoculaires) et du syndrome de Prader-Willi.

Mots clés : ADNc, profil d'expression, cartographie à l'aide d'hybrides résultant d'irradiation, locus de maladie.

[Traduit par la Rédaction]

Corresponding Editor: R.S. Singh.

Received July 8, 1998. Accepted November 11, 1998.

M.-Y. Hwang, Y.-J. Kang, and U. Sohn.¹ Department of Genetic Engineering, College of Natural Sciences, Kyungpook National University, Taegu, 702-701, Korea.

Y.-H. Kim. Department of Microbiology, College of Natural Sciences, Kyungpook National University, Taegu, 702-701, Korea.

S.W. Scherer and L.-C. Tsui. Department of Genetics, The Hospital For Sick Children, Toronto, Ontario, M5G 1X8, Canada.

¹Author to whom all correspondence should be addressed (e-mail: usohn@bh.kyungpook.ac.kr).

Introduction

Since ESTs have been used as a starting point for the functional and structural analysis of the entire human genome, many efforts to identify all the human genes have focused on studies of expressed sequence tags (ESTs) generated by single-run partial cDNA sequencing (Kirkness 1996). This large-scale sequencing has not only provided comprehensive information on the expression pattern of a variety of human tissues and cells (Adams et al. 1993 *a*, 1993*b*; Affara et al. 1994; Choi et al. 1995; Frigerio et al. 1995; Hwang et al. 1995; Kawamoto et al. 1996; Liew et al. 1994; Lee et al. 1995; Okubo et al. 1995; Sudo et al. 1994; Tanaka et al. 1996) but has also generated gene-based sequence-tagged sites for rapid large scale PCR mapping (Berry et al. 1995). The utility of EST mapping has been shown by the uncovering of new insights into genome organization, evolution, and expression, as well as by the facilitating of the positional cloning of human genes linked to diseases (Boguski and Schuler 1995; Collins et al. 1995).

The thymus is a central lymphoid organ in the upper chest which is responsible for the development of immunocompetent cells from immature T cells. In 1995, the first report appeared on the generation and chromosomal localization of human adult thymus ESTs from a normalized cDNA library (Lamerdin et al. 1995). In addition, the TIGR collection also reported over 1100 ESTs from primary or substracted cDNA libraries of human adult normal and tumor thymus tissues (Adams et al. 1995). However, there has not been any report on a gene expression profile reflecting the specialized functions and physiological characteristics of the human infant thymus. Accordingly, we describe here the generation and analysis of 1223 ESTs from a human infant thymus, and the chromosome mapping of 43 ESTs using PCR-based mapping strategy.

Methods

Construction of cDNA library and sequencing

Total RNA was isolated from a 14 mo old human thymus according to the guanidine thiocyanate method (Sambrook et al. 1989). Poly (A)+ RNA was obtained from total RNA by an oligo d(T)-cellulose column. cDNAs were synthesized using a cDNA synthesis kit (Amersham, U.K.). The cDNAs were anchored with a *NotI*-*EcoRI* adaptor and digested by the *EcoRI* restriction enzyme, and subsequently inserted into an *EcoRI*-digested λ ZAPII vector and packaged utilizing a Gigapack gold II packaging system (Stratagen, La Jolla, Cal.). The cDNA library was in vivo excised with a R408 helper phage and transfected into an *E. Coli* strain, XL1-Blue. For sequencing, DNA templates were prepared by the alkaline lysis method (Sambrook et al. 1989). Sequencing reactions were performed with a Sequenase v. 2.0 kit (Amersham, UK).

Database analysis

Sequences obtained by the single-run partial sequencing were subject to a BLAST search (Altschul et al. 1990) against non-redundant nucleotide and protein databases, as well as dbEST. When the *P*-value of BLAST reports was less than 1.0×10^{-5} , the sequences were considered as database-matched clones.

Chromosome mapping

PCR primers were designed by the PRIMER program available on the Web site (<http://www-genome.wi.mit.edu>). PCR reactions were

Table 1. Classification of human infant thymus ESTs.

Classification	Percent	No. of clones
Human matched	50.1	613
Non-human matched	4.2	51
EST	23.6	289
rRNA	4.7	58
<i>Alu</i> repeat	2	24
MtDNA	0.5	6
Unknown	14.9	182
Total	100	1223

performed with specific primers against a NIGMS (Coriell Institute for Medical Research, Camden, N.J.) human-rodent somatic cell hybrid mapping panel 2 (Drwina et al. 1993) and against a GENEBRIDGE 4 RH panel (Research Genetics of Huntsville, Alta.). All PCR reactions were performed in a 50 μ L volume containing 25 ng of template DNA and 2.5 U of AmpliTaq Gold DNA polymerase (Perkin-Elmer Cetus, Conn.). Cycling parameters were 30 s at 94°C, 30 s at 55–60°C, and 1 min at 72°C for 35 cycles, followed by an additional 7 min at 72°C in a DNA thermal cycler (Perkin-Elmer Cetus). The radiation hybrid scoring data resulting from the PCR screenings were statistically analyzed to localize unknown genes on the framework map using the RHMAPPER program at the Whitehead/MIT center for Genome Research (<http://www-genome.wi.mit.edu/cgi-bin/contig/rhmapper.pl>).

Results and discussion

Random sequencing and database analysis

A bidirectional cDNA library was constructed and a total of 1223 ESTs were generated by single-run partial sequencing of over 1400 randomly selected cDNA clones. The average insert size was 0.7 kb and the average length of sequence was 220 bp, which is considered to be useful ESTs. Hillier et al. (1996) reported that the highest quality portion of the EST sequence was between 100 and 300 bases. We classified the results of the database comparisons according to materials and methods. A summary of the database searches for the 1223 sequences is shown in Table 1. Of the 1223 clones, 613 (50.1%) showed significant similarities to human genes and 51 (4.2%) to non-human genes. These 664 clones matching known genes represented 416 unique genes. The other database-matched clones contained 289 (23.6%) ESTs of unknown functions, 24 (2.0%) *Alu*-repeats, 58 (4.7%) rRNA, and 6 (0.5%) mitochondrial DNA. The frequencies of *Alu*-like repeats and mitochondrial DNA are significantly lower than those from previous reports (Sudo et al. 1994; Choi et al. 1995). Of the 1223 cDNA clones, 182 (14.9%) were found to be novel transcripts not showing any significant homology to known genes.

The 664 ESTs showing significant homology to known genes were classified into 7 groups according to their functions (Adams et al. 1995); cell signaling-communication, gene-protein expression, cell division-DNA synthesis, cell structure-motility, cell-organism defense and homeostasis, metabolism, and unclassified (Appendix 1). Among these 7 groups, the gene-protein expression group contained the greatest percentage (34.6%) of total putative identifications from the infant thymus ESTs. In order to investigate the changes of the gene expression in relation to the aging of the

thymus, we compared the expression profile of infant thymus ESTs with that of adult thymus ESTs (Adams et al. 1995). Compared to the adult thymus ESTs, the infant thymus ESTs contained a much higher percentage of ESTs that are involved in gene–protein expression and cell division–DNA synthesis (Fig.1). These differences in the EST profiles between infant and adult thymus may reflect the active growth state of the infant thymus and involution of thymic tissue with aging. Despite the fact that it is an immune-related organ, the infant thymus appeared to express a proportion of ESTs for cell–organism defense similar to other organs (Adams et al. 1995) and twice as small as that of the adult thymus. This is likely to indicate that the relatively high levels of ESTs of gene–protein expression and cell division–DNA synthesis in the infant thymus are more closely related to rapid growth than immune function of thymus.

Regarding the EST frequency, eEF-1 a was the most frequently represented in the infant thymus followed by thymosin beta 4. eEF-1 a was abundantly expressed in cancer, infant tissue, and fetal tissue (Frigerio et al. 1995; Adams et al. 1993b; Hwang et al. 1995; Sudo et al. 1994). Ribosomal proteins also showed high redundancies. Abundant expression of ribosomal proteins was observed in several tissues and differentially expressed in different tissues (Kawamoto et al. 1996; Choi et al. 1995; Tanaka et al. 1996; Frigerio et al. 1995). Consequently, these data coincided with the expression profile of infant thymus ESTs.


We found that some EST sequences corresponding to putative ribosomal proteins (RPs), RTH108 (RP L7a), KTH086 (RP S3), JTH138 (RP S27), and CJC060 (RP S4), showed striking similarities or identities with PLA-X, Fte-1 (v-fos transformation factor), metallopanstimulin, and the SCAR protein, respectively. Interestingly, the sequence of RTH100 (RP S11) was identical to the antisense sequence of isocitrate dehydrogenase. In 1996, Chan et al. reported that rat ribosomal protein L10 showed a 99% and 100% amino acid identity with a human QM and mouse QM, respectively, and suggested that their data was a presumptive example of the extraribosomal function of the ribosomal protein (Wool 1996).

When we assembled ESTs of unknown functions into a redundant EST group with a TIGR Assembler (Sutton et al. 1995), the group (ThyGR-15) had the highest redundancy ($n = 7$), which was equal to that of ribosomal L30. Thus, we can assume that the ThyGR-15 group can be considered as a moderately expressed transcript in a human infant thymus.

Chromosome mapping of unknown ESTs


Forty-three clones were successfully localized onto a human chromosome as the first stage to identify gene-linked genetic disorders (Fig. 2). Of 43 clones, 35 (81.8%) were mapped onto a single chromosome, 7 (15.9%) onto 2 chromosomes, and 1 (2.2%) onto 4 chromosomes. Chromosome 6 was most predominantly hit by the infant thymus ESTs. To obtain a more refined map of novel ESTs, 17 clones were screened against a GENEBRIDGE 4 radiation hybrid panel using a PCR-based mapping method and then mapped through the statistical analysis of radiation hybrid mapping data. To survey the ESTs linked to disease loci, the RH maps of thymus ESTs were combined with the cytogenetic map available at the National Center for Biotechnology Information (NCBI).

Fig. 1. Comparison of expression profile of infant thymus ESTs with that of adult thymus ESTs. The 664 ESTs showing significant homology to known genes were classified into 6 groups based on putative functional categories (Appendix 1). After the adult thymus ESTs were extracted from the Genome Directory, the expression profile was compared by calculating the number of the genes of each group of both infant and adult thymus ESTs.



JTH107 placed 6.51 cR from D6S2350, which corresponds to 3p24 on the cytogenetic map, was linked to the thyroid hormone resistance locus. PTH261 placed 4.81 cR from WI-10875 (12q23), which is related to the histidinemia; ATH341 2.94 cR from D6S442 (6q26) related to the region of plasminogen deficiency I & II, plasminogen Tochigi disease, and dysplasmingogenic thrombophilla. TH572 was mapped 2.74 cR from D6S455 (6q14.1) on chromosome 6 (ocular albinism); TH320, 8.12 cR from WI-6400 (2q31.2–32.1) on chromosome 2 (Ehlers-Danlos syndrome, fibromuscular dysplasia of arteries, familiar aneurysm); HTH093, 6.4 cR from CHLCC GATAA 28C02 (9p13.2–13.3) on chromosome 9 (galactosemia); HTH210, 1.71 cR from WI-6355 (11q23.3) on chromosome 11 (hypertriglyceridemia, amyloidosis); HTH167, 4.6 cR from D14S267 (14q32) on chromosome 14 (porphyria variegata); HTH314, 4.19 cR from WI5273 (1p35) on chromosome 1 (Charcot-Marie-tooth disease, galactose epimerase deficiency); PTH056, 6.4 cR from WI-5254 on chromosome 16; ATH209, 40.8 cR from WI-4263 on chromosome 12 (FEOM), HTH191, 16.61cR from D15S144 on chromosome 15 (Prader-Willi syndrome), and HTH311, 12.56 cR from WI-6622 on chromosome 20 (neuraminidase deficiency, diabetes mellitus). The combination of EST sequences and map locations are currently in the process of providing numerous new candidate genes for those human pathologies that can be mapped genetically (Boguski and Schuler 1995). Several novel genes from the infant thymus were mapped on to disease loci, including the histidinemia loci, the plasminogen Tochigi disease loci, the Ehlers-Danlos syndrome, the hypertriglyceridemia, etc. Consequently, the radiation hybrid mapping of unknown ESTs from an infant thymus could facilitate the discovery of novel genes involved in genetic diseases or immune disorders. In summary, our cDNA approach enabled us to convert transcripts of a human infant thymus into 1223 expressed sequence tags which featured a high level of genes related to

Fig. 2. Chromosomal localization of human infant thymus ESTs. The shadow boxes represent the locations of ESTs obtained by radiation hybrid mapping. Diseases linked to these locations are indicated on the left side of each chromosome. When an EST is located on two or more chromosomes, it is represented by an asterisk. Chromosome assignments using a somatic cell hybrid panel are indicated on the bottom of each chromosome with an italic letter.



Acknowledgements

the active cell cycling of thymocytes. The infant thymus ESTs can also be used as useful mapping markers for radiation hybrid mapping in addition to providing the expression information of active genes in the human infant thymus.

This work was supported by grants from the Ministry of Education, the Ministry of Science and Technology of Korea, and the Canadian Genome Analysis and Technology Program. L.-C.Tsui is a Senior Scientist of the Medical Re-

search Council of Canada, holds the Sellers Chair in Cystic Fibrosis Research, and is a Howard Hughes International Scholar. We would like to thank J.H. Lee, H.C. Lee, J.P. Jeon, J.C. Choi, D.S. Park, and W.K. Cho for their assistance in sequencing.

References

- Adams, M.D., Soares, M.B., Kerlavage, A.R., Fields, C., and Venter, J.C. 1993a. Rapid cDNA sequencing (expressed sequence tags) from a directionally cloned human infant brain cDNA library. *Nature Genet.* **4**: 373–380.
- Adams, M.D., Kerlavage, A.R., Fields, C., and Venter J.C. 1993b. 3,400 new expressed sequence tags identify diversity of transcripts in human brain. *Nature Genet.* **4**: 256–267.
- Adams, M.D., Kerlavage, A.R., Fleischmann, R.D., Fuldner, R.A., Bult, C.J., Lee, N.H., Kirkness, E.F., Weinstock, K.G., Gocayne, J.D., White, O., Sutton, G., Blake, J.A., Brandon, R.C., Man-Wai, C., Clayton, R.A., Cline, T.R., Cotton, M.D., Earle-Hughes, J., Fine, L.D., Fitzgerald, L.M., Fitzhugh, W.M., Fritchman, J.L., Geoghagen, N.S., Glodek, A., Gnehm, C.L., Hanna, M.C., Hedblom, E., Hinkle, P.S., Kelley, J.M. Jr., Kelley, J.C., Liu, L.-I., Marmaros, S.M., Merrick, J.M., Moreno-Palanques, R.F., McDonald, L.A., Nguyen, D.T., Pelligrino, S.M., Phillips, C.A., Ryder, S.E., Scott, J.L., Saudek, D.M., Shirley, R., Small, K.V., Spriggs, T.A., Utterback, T.R., Weidman, J.F., Li, Y., Bednarik, D.P., Cao, L., Cepeda, M.A., Coleman, T.A., Collins, E.J., Dimke, D., Feng, D.-F., Ferrie, A., Fischer, C., Hastings, G.A., He, W.W., Hu, J.S., Greene, J.M., Gruber, J., Hudson, P., Kim, A.K., Kozak, D.L., Kunsch, C., Hungjun, J., Li, H., Meissner, P.S., Olsen, H., Raymond, L., Wei, Y.F., Wing, J., Xu, C., Yu, G.L., Ruben, S.M., Dillion, P.J., Fannon, M.R., Rosen, C.A., Haseltine, W.A., Fields, C., Fraser, C.M., and Venter, J.C. 1995. Initial assessment of human gene diversity and expression patterns based upon 83 million nucleotides of cDNA sequence. *Nature*, **377**: 3–17.
- Affara, N.A., Bentley, E., Davey, P., Pelmeur, A., and Jones, M.H. 1994. The identification of novel gene sequences of the human adult testis. *Genomics*, **22**: 205–210.
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. 1990. Basic local alignment search tool. *J. Mol. Biol.* **215**: 403–410.
- Berry, R., Stevens, T.J., Walter, N.A.R., Wilcox, A.S., Rubano, T., Hopkins, J.A., Weber, J., Goold, R., Soares, M., and Sikela, J.M. 1995. Gene-based sequence-tagged sites (STSs) as the basis for a human gene map. *Nature Genet.* **10**: 415–423.
- Boguski, M.S., and Schuler, G.D. 1995. ESTablishing a human transcript map. *Nature Genet.* **10**: 319–371.
- Chan, Y.L., Diaz, J.J., Denoroy, L., Madjar, J.J., and Wool, I.G. 1996. The primary structure of rat ribosomal protein L10: Relationship to a jun binding protein and to a putative Wilms' tumor suppressor. *Biochem. Biophys. Res. Commun.* **225**: 952–956.
- Choi, S.S., Yun, J.W., Choi, E.K., Cho, Y.G., Sung, Y.C., and Shin, H.S. 1995. Construction of a gene expression profile of a human fetal liver by single-pass cDNA sequencing. *Mamm. Genome*, **6**: 653–657.
- Collins, F.S. 1995. Positional cloning moves from perdictional to traditional. *Nature Genet.* **9**: 347–350.
- Drwanga, H.L., Toji, L.H., Kim, C.H., Greene, A.E., and Mullivor, R.A. 1993. NIGMS human/rodent somatic cell hybrid mapping panel 1 and 2. *Genomics*, **16**: 311–314.
- Frigerio, J.M., Berthezene, P., Garrido, P., Ortiz, E., Barthelmy, S., Vasseur, S., Sastre, B., Seleznieff, I., Dagorn, J.C., and Iovanna, J.L. 1995. Analysis of 2166 clones from a human colorectal cancer cDNA library by partial sequencing. *Hum. Mol. Genet.* **4**: 37–43.
- Hillier, L., Lennon, G., Becker, M., Bonaldo, M.F., Chiapelli, B., Chissoe, S., Dietrich, N., DuBuque, T., Favello, A., Gish, W., Hawkins, M., Hultman, M., Kucaba, T., Lacy, M., Le, M., Le, N., Mardis, E., Moore, B., Morris, M., Parsons, J., Prange, C., Riffkin, L., Rohlfing, T., Schellenberg, K., Soares, M.B., Tan, F., Thierry-Meg, J., Trevaskis, E., Underwood, K., Wohldman, P., Waterston, R., Wilson, R., and Marra, M. 1996. Generation and Analysis of 280,000 Human expressed sequence tags. *Genome Res.* **6**: 807–828.
- Hwang, D.M., Fung, Y.W., Wang, R.X., Laurensen, C.M., Ng, S.H., Lam, W.Y., Tsui, K.W., Fung, K.P., Waye, M., Lee, C.Y., and Liew, C.C. 1995. Analysis of expressed sequence tags from a fetal human heart cDNA library. *Genomics*, **30**: 293–298.
- Kawamoto, S., Matsumoto, Y., Mizuno, K., Okubo, K., and Matsubara, K. 1996. Expression profiles of active genes in human and mouse livers. *Gene*, **174**: 151–158.
- Kirkness, E.F. 1996. Assessment of human gene diversity and expression pattern using expressed sequence tags. *Essays in Biochem.* **31**: 1–9.
- Lamerdin, J.E., Athwal, R.S., Kansara, M.S., Sandhu, A.R., Patanjali, S.R., Weissman, S.M., and Carrano, A.V. 1995. Chromosomal localization and expressed sequence tag generation of clones from a normalized human adult thymus cDNA library. *Genome Res.* **5**: 359–367.
- Lee, N.H., Weinstock, K.G., Kirkness, E.F., Earle-Hughes, J.A., Fuldner, R.A., Marmaros, S., Glodek, A., Gocayne, J.D., Adams, M.D., Kerlavage, A.R., Fraser, C.M., and Venter, J.C. 1995. Comparative expressed-sequence-tag analysis of differential gene expressed profiles in PC-12 cells before and after nerve growth factor treatment. *Proc. Natl. Acad. Sci. U.S.A.* **92**: 8303–8307.
- Liew, C.C., Hwang, D.M., Fung, Y.W., Laurensen, C., Cukerman, E., Tsui, S., and Lee, C.Y. 1994. A catalogue of genes in the cardiovascular system as identified by expressed sequence tags. *Proc. Natl. Acad. Sci. U.S.A.* **91**: 10 645 – 10 649.
- Okubo, K., Itoh, K., Fukushima, A., Yoshii, J., and Matsubara, K. 1995. Monitoring cell physiology by expression profiles and discovering cell type-specific genes by compiled expression profiles. *Genomics*, **30**: 178–186.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. 1989. *Molecular cloning: A laboratory manual*. 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Sudo, K., Chinen, K., and Nakamura, Y. 1994. 2058 expressed sequence tags (ESTs) from a human fetal lung cDNA library. *Genomics*, **24**: 276–279.
- Sutton, G.G., White, O., Adams, M.D., and Kerlavage, A.R. 1995. TIGR Assembler: A new tool for assembling large shotgun sequencing projects. *Genome Sci. & Tech.* **1**: 9–19.
- Tanaka, T., Ogiwara, A., Uchiyama, I., Takegi, T., Yazaki, Y., and Nakamura, Y. 1996. Construction of a normalized directionally clones cDNA library from adult heart and analysis of 3040 clones by partial sequencing. *Genomics*, **35**: 231–235.
- Wool, I.G. 1996. Extraribosomal functions of ribosomal proteins. *Trends Biochem. Sci.* **21**: 164–165.

Appendix 1. Infant thymus ESTs matched to known genes in the database.

Cell Division (6.8%)	Parathyroid hormone	Copper transport protein HAH1
BTG1 gene (2)	PI-3 kinase	Csa-19 (4)
BTG2 gene	Plexin 2 (3)	dnaJ like protein
Cellular oncogene c-fos	Pre-B cell stimulating factor homologue (2)	Ferritin heavy chain (2)
Cyclin D3 (2)	Protein kinase (rho-associated)	Ferritin light chain
Cyclin protein	Protein kinase 63 kDa related to rat ERK3	Heat shock cognate 70 kDa protein
Deoxynucleotidyltransferase	Protein kinase, TTK	Heat shock cognate 89 kDa protein (2)
DNA synthesis inhibitor	Protein tyrosine phosphatase (3)	Heat shock protein 90
Histone H1B, H2A, H2B, H4 genes	Rap1B mRNA	HLA classI locus C
Histone H2A, F (2)	Ras-related C3 botulinum toxin substrate(rac)	HLA classII SB beta-chain
Histone H2A.2	Renin	HLA-A2 classI antigen (3)
Histone H3.3 (4)	Rho GDP dissociation inhibitor 2	HLA-B27
Histone H3.3, B	Rod outer segment membrane protein 1 (Rom1)	HLA-C 1
HMG (High mobility group) 2	Serine/threonine kinase receptor 2	HLA-DQA2
Inhibitor of apoptosis protein 2	Stathmin (2)	HLA-DR, alpha, heavy chain
MA-3(apoptosis-related gene) (6)	Thymosin beta 10 (2)	HLA-Dw12 DQ beta
Nonhistone protein HMG1 mRNA	Thymosin beta 4 (13)	HLA-E
Notch3	Tumor necrosis factor (4)	HLA-SB beta
Nucleosome assemble protein 2 (2)		HLA-xrp gene
Oncogene ets-2	Cell Structure/Mobility (7.1%)	hMSH6 gene, exon 6-10
Oncogene PTI-1	Actin, beta (5)	Ig associated invariant gamma chain
Prothymosin alpha (6)	Actin, cytoplasmic	Ig gamma 1 chain C region
Proto oncogene c-sis	Actin, gamma (2)	Ig J chain gene
Proto oncogene Ets-1	Ankyrin binding glycoprotein-1 (4)	Ig kappa chain VK-1
SET nuclear phosphoprotein (3)	Capping protein, alpha subunit 1 (4)	Ig kappa light chain
Tis11d	CENP-F kinetochore protein mRNA	Ig lambda light chain (3)
TRE5 sequence (tre oncogene)	Clathrin coat assembly protein-like	Ig mu heavy chain C-region
	Cofilin (2)	RAG-1 (7)
Cell Signaling/Communication (11.4%)	COP(36 kDa coatomer), epsilon	RAG-2 (2)
14-3-3 eta chain	Cytokeratin	RAG-3
14-3-3, beta	Dynamitin (2)	rGSTK1-1
14-3-3, epsilon	Elastin (4)	RING3 gene
ADP-ribosylation factor 1 (2)	Fibrinogen gamma chain	T-cell receptor beta chain (D-J-C) (2)
Androgen receptor associated protein 24	Heterochromatin protein	T-cell receptor beta chain loci
Aurora-related kinase 2 (ARK 2)	Keratin (2)	T-cell receptor beta-chain (C region) (3)
B-cell growth factor	Kid (kinesin-like DNA binding protein)	T-cell receptor, T3 epsilon
Cadherin	Kinectin (motor kinesin binding protein)	Transferrin receptor
Calcyclin/PRA	Laminin B1	VH4-34 (isotype switch)
Calmodulin (CALM1) gene	Matrin 3	
Calmodulin dependent protein phosphatase	Moesin-related mRNA sequence	Gene/Protein Expression (34.6%)
Calmodulin homologue (NB-1)	Moitotic kinesin like protein	Acidic ribosomal phosphoprotein P0
Calmodulin-I mRNA (2)	Myosin light chain, non-muscle, alkali	Acidic ribosomal phosphoprotein P1
Calmodulin-like pseudogene	Profilin 1	Acidic ribosomal phosphoprotein P2
cdc2/CDC28 like protein kinase	RIG-like 14-1 mRNA	Antisecretory factor 1
Coupling protein G(s) alpha subunit	SPARC/osteonectin	Bat2 gene
Cytokine receptor CRFB4	Tubulin alpha (3)	CAAT box binding transcription factor (CTF)
ERK activator kinase	Tubulin beta (2)	Calpain (calcium activated neutral protease)
Glucocorticoid receptor		Cathepsin B
gpISG20	Cell/Organism Defense (11.8%)	Cathepsin L (2)
Grb3-3	27 kDa heat shock protein (2)	CCAAT/enhancer binding protein delta
GTP binding protein beta subunit-like protein	28 kDa heat shock protein	Cystein-rich sequence binding protein
GTP-binding protein (rhoA) (2)	APEX nuclease	DNA binding protein(neurodap 1)
hTGR	Beta-2 microglobulin (2)	DNA binding protein, KET
ICLN protein	CAMPATH-1 (3)	DNA binding protein, TAXREB107
IL2 receptor gamma	CD1 antigen R2, MHC-related (6)	eEF-1 alpha (20)
Inerferone-gamma IEF	Chitinase precursor	eEF-1 beta (2)
Laminin receptor (4)	Coagulation factor IX gene	eEF-1 gamma related protein
MAGHO mRNA	Complement C2 precursor	Enhancer of split m9/m10
<u>Oxytocin receptor</u>	Complement C3	FKBP (FK506-binding protein)

Appendix 1 (continued).

G17 gene	Ribosomal protein S27 (2)	ATPase, H ⁺ transport, vacuolar (2)
Helicase (218kD Mi-2 protein) (2)	Ribosomal protein S28, yeast homolog	ATPase, NA ⁺ / K ⁺ transport, alpha subunit
Helix-loop-helix-leucine zipper (SREBP-1)	Ribosomal protein S29	ATPase, proton subunit D, vacuolar (3)
Heterogenous ribonucleoprotein homolog (3)	Ribosomal protein S2a	Co-beta glucosidase(proactivator)
Hlark mRNA	Ribosomal protein S3a (6)	CYP3B6 gene cryptic exon (cytochrom p450)
hnRNP core protein A1	Ribosomal protein S4 (2)	Cytochrome C oxidase polypeptide I (COX I)
hnRNP D (2)	Ribosomal protein S5	Cytochrome C oxidase subunit IV (COX 4)
hnRNP-C	Ribosomal protein S6	Cytochrome c oxidase subunit VIa
hnRNP-C2	Ribosomal protein S7	Cytochrome C oxidase subunit Vlb
hnRNP-E1 (2)	Ribosomal protein S8 (2)	Cytochrome C oxidase subunit VIIB
hnRNP-H	Ribosomal protein S9	Deoxycytidine kinase (3)
hnRNP-K	Ribosomal protein YL30	Enyol CoA dehydrogenase
MAD-3 encoding Ikb-like acitivity	RNA binding protein, HU-K4	Glycealdehyde-3-phosphate dehydrogenase
Myocyte specific enhancer factor (2)	RNA helicase #46, ATP-dependent	GTP cyclohydrolase I
Nuclear p68 protein (3)	RNA polymerase I	Iduronate sulphate sulphatase (IDS)
Plasminogen activator inhibitor-1	Seb4	Importin beta subunit
Polyadenylation specificity factor	Sec61(protein transport protein)	Inosine monophosphate dehydrogenase type II
Prolyl 4 hydroxylase, alpha subunit (4)	SOX-4 (Sry-like HMG box protein)	Lactate dehydrogenase A
Proteasome component C9	Splicing factor mRNA	Lactate dehydrogenase B
Proteasome related gene (LMP 2)	Splicing factor SRp75	Liver phosphatase 2A (2)
Proteasome subunit HC8 (2)	Splicing factor, SF1-HL1 isoform	Lysophosphatidic acid acyltransferase-alpha (2)
Proteasome subunit p55	SW1/SNF complex 155 kDa subunit (BAF155)	Malate dehydrogenase
Proteasome-like subunit	TCF-1 (T-cell specific transcription factor)	mGLT(glutamate transporter)-1 mRNA
putative RNA binding protein	TFIID 55 kDa subunit	Monocarboxylate transporter 1 (2)
Ribosomal protein L1	TFIIB mRNA	Muscle glycogen phosphorylase
Ribosomal protein L10 (3)	Transcription factor (Staf 50)	N-acetylglucosaminyltransferase III DNA,
Ribosomal protein L17	Transcription factor AF-1p	NAD(P) ⁺ dependent malic enzyme mRNA (2)
Ribosomal protein L21 (2)	Transcription factor, ASF	NADH:ubiquitin oxidoreductase
Ribosomal protein L23 (5)	Transglutaminase type I	Nm23 protein (nucleoside diphosphate kinase)
Ribosomal protein L24	Translation factor Sui1 homolog (3)	Oligoadenylate synthetase
Ribosomal protein L26 (4)	Translation initiation factor p40 subunit	Phospholipase A2
Ribosomal protein L27 (2)	Translation initiation factor 3	Phosphotyrosyl phosphatase activator
Ribosomal protein L28 (2)	Translation initiation factor, Eif4g2	Purin 5'-nucleotidase
Ribosomal protein L3 (2)	Translation repressor NAT1 mRNA (7)	r-aminobutylaldehyde dehydrogenase
Ribosomal protein L30 (7)	U21.1 hnRNP	Ribonuclease 6 precursor
Ribosomal protein L32	Ubiquitin (2)	Ribonucleoside diphosphate reductase M1
Ribosomal protein L35 (4)	Ubiquitin conjugating enzyme (ubc4)	Ribonucleotide reductase M2
Ribosomal protein L36	Ubiquitin conjugating enzyme E2 (2)	Secretory carrier membrane protein
Ribosomal protein L37 (3)	USF-2(upstream stimulatory factor-2)	sterol carrier protein (3)
Ribosomal protein L37a	Wilm's tumor-related protein (6)	Ubiquinal-cytochrome C reductase
Ribosomal protein L38	Zinc finger protein RING (RZF)	
Ribosomal protein L4	Zinc finger protein, HPP1 (3)	
Ribosomal protein L41, yeast homologue	Zinc finger protein, ZNF	
Ribosomal protein L44, yeast homologue		
Ribosomal protein L5 (3)		
Ribosomal protein L6		
Ribosomal protein L7 (2)		
Ribosomal protein L7a (3)		
Ribosomal protein L9 (4)		
Ribosomal protein S10 (3)		
Ribosomal protein S11 (6)		
Ribosomal protein S14 (3)		
Ribosomal protein S16		
Ribosomal protein S17 (2)		
Ribosomal protein S18 (4)		
Ribosomal protein S20 (4)		
Ribosomal protein S24 (2)		
Ribosomal protein S25 (2)		
Ribosomal protein S26 (2)		
	Metabolism (10.4%)	
	2' oxidoglutarate dehydrogenase	
	Adenosin Deaminase (ADA)	
	Adenosin monophosphate deaminase1	
	Adenosine triphosphatase	
	Aldehyde dehydrogenase	
	Alpha enolase gene (2)	
	Apolipoprotein A-IV gene	
	ATP synthase, beta subunit (2)	
	ATP synthase, gamma subunit (2)	
	ATP synthase, subunit a	
	ATP synthase, subunit c (2)	
	ATP synthase, subunit d	
	ATP synthase, subunit f	
	ATP synthase, subunit g	
		Unclassified (17.9%)
		23KD highly basic protein(Ig gamma)
		AICL(activation induced c-type lectin)
		Ataxia telagietcusa (ATM) gene
		Autoantigen DFS70
		BBC1(breast basic conserved) protein
		Beta amyloid protein(APP) gene
		Brain 0-44 mRNA
		BRCA2(breast cancer susceptibility)
		CG1 protein
		CLE7 (<i>G.gallus</i>) (3)
		COX7RP
		CpG DNA (5)
		Dead box, x isoform(DBX) alternative transcript 2
		Down syndrome region
		DT1P1A10, CAG repeat
		DT1P1A2, CAG repeat
		Duffy blood group antigen Fya-b+

Appendix 1 (concluded).

E25 mRNA (2)	Hypothetical protein, KIAA0161 gene	Nuclear Factor IV
EB virus small RNA associated protein (EAP)	Hypothetical protein, KIAA0217 gene	Nucleolar phosphoprotein B23(nucleophomin)
EBI1-ligand chemokine (ELC)	Hypothetical protein, KIAA0220 gene	Nucleolar protein P120(ribonucleoprotein)
GP91-PHOX gene promoter region	Hypothetical protein, KIAA0252 gene	Ocular melanoma associated antigen
GRSF-1(G-rich sequence factor-1)	Hypothetical protein, KIAA0261 gene	ORF mRNA
GT233 mRNA	Hypothetical protein, KIAA0323 gene	p38-2G4
HE5 (CDw52-like epididymal protein) (5)	IEF7422	Proliferation-associated protein pag
Heat shock protein hsp70	Interferon inducible gene (3)	Prostatein C3 subunit gene
Hepatocyte nuclear factor 1 promoter	JKA10 mRNA induced upon T-cell activation	putative nuclear pore complex protein(Npap60)
HepG2 3' region cDNA	Leucine rich protein	PxF protein (3)
HLA class III containing NOTCH4 gene (2)	LLReps3	Retinoblastoma product binding protein (RBQ-1)
Housekeeping	LTG9/MLLT3, C-terminal	Retroposon SINE-R11 (2)
Human familial Alzheimer's disease gene	Lymphoid restricted membrane proein, Jaw1	S153 clone mRNA
Human mRNA for 3-7 gene product	Lysosomal-associated membrane glycoprotein1	Serine-rich neutrophil protein
Huntington's disease region	Lysosome associated membrane protein2	Split hand/split foot 1 (DSS1) (3)
Hypothetical protein, KIAA0034 gene (3)	Melanoma associated antigen	T-cell surface glycoprotein E2
Hypothetical protein, KIAA0037 gene	Metallopanstimulin 1	Tera mRNA
Hypothetical protein, KIAA0040 gene	MGC-24 (PNA-binding protein)	Thymocyte antigen CD1b
Hypothetical protein, KIAA0040 gene	MHC class I HLA-C 1 gene	Transformation related protein (2)
Hypothetical protein, KIAA0045 gene	MHC class I HLA-C 1 locus C heavy chain	Transmembrane protein (4)
Hypothetical protein, KIAA0083 gene	Na ⁺ /H ⁺ exchange regulatory co-factor	Unknown antigen
Hypothetical protein, KIAA0101 gene	NAC, alpha (4)	Unknown protein within p53 intron 1
Hypothetical protein, KIAA0107 gene	NifU-like protein (hNifU) mRNA	X11gene
Hypothetical protein, KIAA0128 gene	Ninein (centromal protein)	XG blood group
Hypothetical protein, KIAA0159 gene	NK-tumor recognition molecule-related protein	ZFM 1 protein alternatively spliced product

Note: Human infant ESTs matching to 416 distinct known genes were divided into 7 groups according to the putative function (Adams et al. 1995). Numbers in parentheses indicate the frequency of the ESTs. In the case of non-human matched ESTs, the organism is indicated in parentheses.