



Title	Picosecond time-resolved resonance Raman observation of the iso-CH₂Cl-I and iso-CH₂I-Cl photoproducts from the "photoisomerization" reactions of CH₂Cl in the solution phase
Author(s)	Kwok, WM; Ma, C; Parker, AW; Phillips, D; Towrie, M; Matousek, P; Zheng, X; Phillips, DL
Citation	Journal Of Chemical Physics, 2001, v. 114 n. 17, p. 7536-7543
Issued Date	2001
URL	http://hdl.handle.net/10722/42355
Rights	Creative Commons: Attribution 3.0 Hong Kong License

Picosecond time-resolved resonance Raman observation of the iso-CH₂Cl-I and iso-CH₂I-Cl photoproducts from the "photoisomerization" reactions of CH₂I-Cl in the solution phase

Wai Ming Kwok and Chensheng Ma

Department of Chemistry, Imperial College of Science, Technology and Medicine, London SW7 2AY, United Kingdom

Anthony W. Parker^{a)}

Central Laser Facility, CLRC Rutherford Appleton Laboratory, Chilton, Didcot, Oxfordshire OX11 0QX, United Kingdom

David Phillips^{a)}

Department of Chemistry, Imperial College of Science, Technology and Medicine, London SW7 2AY, United Kingdom

Michael Towrie and Pavel Matousek

Central Laser Facility, CLRC Rutherford Appleton Laboratory, Chilton, Didcot, Oxfordshire OX11 0QX, United Kingdom

Xuming Zheng

Department of Applied Chemistry, Zhejiang Institute of Science and Technology, Hangzhou 310033, People's Republic of China

David Lee Phillips^{a),b)}

Department of Chemistry, University of Hong Kong, Hong Kong S.A.R., People's Republic of China

(Received 13 December 2000; accepted 15 February 2001)

We report a preliminary picosecond Stokes time-resolved resonance Raman investigation of the initial formation and subsequent decay of the photoproduct produced following 267 nm excitation of CH₂I-Cl in acetonitrile solution. Density-functional theory computations were done for several probable photoproduct species. Comparison of these computational results and results from a recent femtosecond transient absorption study to our present picosecond resonance Raman spectra indicate that the iso-CH₂Cl-I species is mainly produced and associated with the ~460 nm transient absorption band. The iso-CH₂Cl-I species appears to decay and form appreciable amounts of the more stable iso-CH₂I-Cl species that is associated with a ~370 nm transient absorption band after a few hundred ps. © 2001 American Institute of Physics. [DOI: 10.1063/1.1362178]

I. INTRODUCTION

Polyhalomethane molecules are of chemical interest from several viewpoints. They are used as reagents in a number of reactions in synthetic chemistry like cyclopropanation of olefins and diiodomethylation of carbonyl compounds.¹⁻⁹ For instance, ultraviolet photolysis of CH₂I₂ or activation of CH₂I₂ by a Zn(Cu) couple in the Simmons-Smith reaction has found utility to produce cyclopropanated products from olefins with high stereospecificity and little competition from C-H insertion.¹⁻⁵ Polyhalomethanes like CH₂I₂, CH₂Br₂, CH₂BrI, CH₂ClI, CHBr₃, and CHBr₂Cl have been observed in the troposphere in measurable quantities and are probably important sources of reactive halogens in the atmosphere.¹⁰⁻¹⁶ This has led to increasing interest in the atmospheric photochemistry and chemistry of polyhalomethane molecules.

Polyhalomethanes are also of interest for fundamental investigations of gas and condensed phase direct photodissociation reactions.¹⁷⁻⁵⁵

Ultraviolet excitation of gas-phase polyhalomethanes usually leads to a direct carbon-halogen bond breaking reaction(s).¹⁷⁻²⁹ Molecular beam anisotropy measurements show these primary reaction(s) typically occur in a time much less than the rotational period of the parent molecule^{17,19-22} and photofragment translational spectroscopy experiments for CH₂I₂,¹⁸ CH₂BrI,²² and CF₂I₂,^{23,24} indicate that the polyatomic photofragment typically receives large amounts of internal excitation of their rotational and vibrational degrees of freedom. Gas and solution phase resonance Raman investigations for several polyhalomethane molecules showed that the direct photodissociation reaction(s) have multidimensional reaction coordinates and Franck-Condon region dynamics that appear qualitatively consistent with a semirigid radical impulsive description of the photodissociation dynamics.³⁰⁻³⁹

Ultraviolet excitation, direct photoionization and radiolysis of CH₂I₂ in condensed phase environments leads to

^{a)} Authors to whom correspondence should be addressed.

^{b)} Electronic mail: phillips@hkucc.hku.hk

formation of characteristic absorption bands ~ 385 nm (strong intensity) and ~ 570 nm (moderate intensity)^{40–47} that have been assigned to several different probable photoproduct species like trapped electrons,⁴⁰ the CH₂I₂⁺ cation^{45,46} or the iso-CH₂I–I species.^{43,44} Several femtosecond transient absorption experiments have been done to follow the formation and decay of the photoproduct species.^{48–50} Although the three different femtosecond studies exhibited similar results with a fast rise followed by a fast decay and then by a slower rise, three different interpretations were given because of their differing assignments for the photoproduct species responsible for the 385 and 570 nm transient absorption bands.^{48–50} A recent nanosecond transient resonance Raman and density functional theory investigation demonstrated that the iso-CH₂I–I species is mostly responsible for the intense ~ 385 nm transient absorption band.⁵¹ Further picosecond time-resolved resonance Raman spectroscopy experiments (with ~ 1 ps resolution) showed that the iso-CH₂I–I species is produced vibrationally hot within several picoseconds and then subsequently vibrationally cools on the 4–50 ps time scale.⁵² We proposed a qualitative geminate and/or near geminate recombination mechanism of the initially produced hot CH₂I fragment and I fragment within the solvent cage to give the hot iso-CH₂I–I photoproduct.⁵²

Many polyhalomethane molecules exhibit characteristic transient absorption bands after ultraviolet excitation in condensed phase environments.^{40–47} A combination of nanosecond transient resonance Raman experiments and density-functional theory computations has begun to be used to clearly identify and characterize the photoproduct species that give rise to these transient absorption bands for a number of polyhalomethanes containing iodine and bromine atoms.^{53–56} Excitation of either A-band or B-band CHI₃ and CH₂BrI produces the same iso-CHI₃ and iso-CH₂I–Br species, respectively.^{53,54} This suggests that the production of iso-polyhalomethane species does not occur for a particular transition and probably happens generally after ultraviolet excitation of $n \rightarrow \sigma^*$ transitions localized on C–X bonds in the condensed phase. The lifetime or stability of the iso-polyhalomethane species produced can vary considerably and this will likely affect the chemistry of these species. We note that we could not detect any isomer species associated with CH₂ClI in room temperature solutions on the nanosecond time scale⁵³ although the iso-CH₂Cl–I species was readily observed in low temperature (12 K) matrices.^{43,44} A recent femtosecond transient absorption study for the ultraviolet photolysis of CH₂ClI in room temperature acetonitrile solutions has been reported.⁵⁷ This study observed fast formation of a strong band at 460 nm and a weaker band ~ 710 nm that decayed with time constant of ~ 100 ps. This was followed by formation of a new species that had a strong band ~ 370 nm (strong) and ~ 725 nm (weak). The first transient absorption bands (at 460 and 710 nm) were attributed to formation of the iso-CH₂Cl–I species which has a lifetime ~ 100 ps and the second transient absorption (at ~ 370 nm) at longer times was tentatively assigned to an ICl[–] species.⁵⁷

In this paper, we report picosecond time-resolved resonance Raman experiments to investigate the identity and formation of the two photoproduct species observed in the re-

cent femtosecond transient absorption study of the ultraviolet photolysis of CH₂ClI in room temperature solution. We utilize the results of density functional theory computations for several proposed photoproduct species and compare them to the experimental results to identify the two photoproduct species as iso-CH₂Cl–I and iso-CH₂I–Cl. Our ps experimental results indicate the iso-CH₂Cl–I species is formed first within several ps via recombination within the solvent cage of the initially produced CH₂Cl and I photofragments. The iso-CH₂Cl–I species then decays to form either the more stable iso-CH₂I–Cl species or parent molecule via isomerization reaction(s).

II. EXPERIMENT AND CALCULATIONS

The experiments were carried out using a ps-TR³ apparatus based on optical parametric amplifiers (OPAs) described in detail elsewhere.⁵⁸ Briefly, a 800 nm pulse generated from a femtosecond Ti:Sapphire oscillator is amplified from 2 to 3 mJ at 1 kHz in a regenerative amplifier. The output from the amplifier was frequency doubled in a 2 mm type I BBO (β -barium borate) crystal to generate the 400 nm probe pulses which was also used to pump the OPA for generating the 480 nm probe pulses for TR³ spectroscopy. The 267 nm pump wavelength was the third harmonic of the regenerative amplifier. Typical pump and probe pulse energies at the sample were ~ 5 – 15 μ J. The time resolution was ~ 1 ps as determined by the duration of the laser pulses (~ 1 ps FWHM). The ground and excited states of trans-stilbene absorb in the region of the 267 nm pump and the 400 nm (and 480 nm) probe laser beams, respectively, and were used to determine the time zero delay between the pump and probe laser beams in the TR³ experiments. The time zero was established by adjusting the optical delay between the pump and probe beams to a position where the depletion of the stilbene fluorescence was halfway to the maximum fluorescence depletion by the probe laser. The time zero accuracy was estimated to be ± 0.5 ps. In order to use the laser beams more effectively in the TR³ experiments and considering that the rotational reorientation dynamics are much faster than the dynamics investigated in this study, parallel polarization of the pump and probe laser beams was used rather than the magic angle polarization. The beams were focused to a spot size of around 100 μ m in a jet with diameter ~ 500 μ m. Scattered photons were collected at 90° using a parabolic aluminum mirror ($f^{\#}=0.8$, $f=4$ cm), dispersed in a triple stage spectrograph and detected by a liquid nitrogen cooled charge coupled device (CCD). Each spectrum presented here was subtracted from scaled probe-before-pump and scaled net solvent measurements in order to eliminate CH₂ClI ground-state Raman peaks and residual solvent Raman bands, respectively. Solvent Raman bands were used to calibrate the spectra with an estimated accuracy of ± 10 cm^{–1} and ± 20 cm^{–1} in absolute frequency for the 400 and 480 nm probes, respectively. The total acquisition time for each spectrum at each time delay was about 100 min.

CH₂ClI and spectroscopic grade acetonitrile solvent were obtained commercially and used without further purification. 1 liter of CH₂ClI (5×10^{-2} mol dm^{–3}) samples were prepared in acetonitrile. Into this solution was added several




FIG. 1. Absorption spectrum of CH_2ClI in acetonitrile solution with the pump and probe excitation wavelengths (in nm) for the picosecond time-resolved resonance Raman experiments shown above the spectra.

strands of copper wire to absorb the photoproduct iodine. During the experimental run the samples exhibited less than a few percent degradation as indicated by the ultraviolet (UV) absorption spectra recorded before and after the TR³ measurement.

The GAUSSIAN program suite (G98W)⁵⁹ was used for all of the density-functional theory computations presented here and the Sadlej-PVTZ (Sadlej triple- ζ plus valence polarization) basis sets were used.⁶⁰ B3LYP (Becke three parameter with Lee, Yang, Parr functional) calculations were done to find the optimized geometry and vibrational frequencies of the species examined. Time-dependent density-functional theory at the random phase approximation computations [TD(RPA)] were used to find estimates of the electronic transition energies and oscillator strengths of the species investigated.⁶¹

III. RESULTS AND DISCUSSION

Figure 1 presents the ultraviolet absorption spectrum of CH_2ClI in acetonitrile solution. Figures 2 and 3 show the Stokes picosecond time-resolved resonance Raman obtained for the photoproduct(s) following 267 nm photoexcitation of CH_2ClI in acetonitrile solution using 400 and 480 nm probe wavelengths, respectively. The Raman spectra appear to be composed of the fundamentals, overtones, and combination bands of several Franck–Condon active modes. Inspection of Fig. 2 shows that a Raman band $\sim 174\text{ cm}^{-1}$ appears first and as it decays (between 3 and 500 ps) a new Raman band $\sim 207\text{ cm}^{-1}$ begins to appear at later times (between 300 and 2000 ps). This suggests that a second photoproduct species is formed following decay of the first photoproduct species. The 480 nm spectrum shown in Fig. 3 appears to be mainly due to the first photoproduct species since most of the Raman band intensity decays by 300 ps and no Raman band is observed $\sim 207\text{ cm}^{-1}$. Table I lists the vibrational frequencies of the larger Raman bands as a function of pump–probe delay time. The femtosecond transient absorption study done by Akesson and co-workers⁵⁷ on the photolysis of CH_2ClI in acetonitrile solution indicates the initial photoproduct species




FIG. 2. Stokes picosecond time-resolved resonance Raman spectra of the photoproduct species produced from ultraviolet excitation of CH_2ClI in acetonitrile solution. Spectra were obtained at varying pump (267 nm) and probe (400 nm) time delays (as indicated to the right of each spectrum). Asterisks “*” mark parts of the spectra where solvent subtraction artifacts are present.

is likely formed vibrationally hot. Therefore, vibrational relaxation will probably affect the Raman band intensities observed in the spectra of Figs. 2 and 3. Our previous investigation of the closely related photolysis of CH_2I_2 in the solution phase showed that the vibrational relaxation of the initially produced $\text{iso-CH}_2\text{I}_2$ photoproduct was essentially complete after 50–100 ps depending on the solvent used.⁵² Thus we expect that vibrational relaxation will have minimal effects on the Raman band intensities observed after 100 ps in Figs. 2 and 3 where the decrease of the first photoproduct Raman band intensity and the increase of the second photoproduct Raman band intensity occurs.

In order to identify the photoproduct species we performed density-functional theory calculations to find the optimized geometry, vibrational frequencies and electronic absorption transitions for the $\text{iso-CH}_2\text{Cl-I}$, $\text{iso-CH}_2\text{I-Cl}$, CH_2ClI^+ cation, and CH_2Cl radical species that might be produced after 267 nm excitation of the sample. Table II lists the optimized geometry of these species found from B3LYP/Sadlej-PVTZ computations. Table III compares the B3LYP/Sadlej-PVTZ computed vibrational frequencies for the prob-




FIG. 3. Stokes picosecond time-resolved resonance Raman spectra of the photoproduct species produced from ultraviolet excitation of CH₂ICI in acetonitrile solution. Spectra were obtained at varying pump (267 nm) and probe (480 nm) time delays (as indicated to the right of each spectrum).

able photoproduct species to those of the experimental spectra for the fundamental bands. Table IV shows the B3LYP/Sadlej-PVTZ computed singlet electronic transition energies and oscillator strengths for the species given in Tables II and III.

The first photoproduct species that appears after several ps in the 400 and 480 nm probe spectra has four fundamental bands at ~ 141 , 174, 376, and 724 cm⁻¹ (values from the 30 ps spectrum in Fig. 2). These bands appear to have overtones and/or combination bands with each other. The four fundamental Raman bands exhibit good agreement with the computed vibrational frequencies for iso-CH₂Cl-I but not the other probable photoproduct species shown in Table III. For instance, the ~ 141 , 174, 376, and 724 cm⁻¹ experimental vibrational frequencies agree better with the values computed for iso-CH₂Cl-I (at 136, 194, 420, and 760 cm⁻¹, respectively) compared to those for iso-CH₂I-Cl (at 113, 236, 494, and 686 cm⁻¹, respectively). The first photoproduct species clearly shows two low-frequencies Raman bands at ~ 141 and 174 cm⁻¹ that have a combination band ~ 303 cm⁻¹. However, the CH₂CI⁺ cation has only one low frequency mode below 200 cm⁻¹ at ~ 191 cm⁻¹ (see Table III) and can be ruled out as the species responsible for the first photoproduct resonance Raman spectra shown in Figs. 2 and

3. Similarly, the CH₂Cl radical has no low-frequency modes below 200 cm⁻¹ and can also be ruled out as the identity of the first photoproduct species. Inspection of Table IV shows that the computed electronic transition energies and oscillator strengths are consistent with the assignment of the first photoproduct species to the iso-CH₂Cl-I species but not the other species examined. A broad electronic transition around the 453 nm position and 0.4932 oscillator strength computed for the iso-CH₂Cl-I species is consistent with our experimental observation of the first photoproduct species using both the 480 and 400 nm probe wavelengths (albeit with differing intensities). However, the computed electronic transition energy for the iso-CH₂I-Cl species is considerably blue shifted to ~ 332 nm and is not likely to be observed in the resonance Raman spectra obtained using 480 nm probe wavelength. Similarly, the computed electronic absorption transition energy for the CH₂CI⁺ cation is ~ 489 nm with a small oscillator strength of 0.0018 and is not likely to be associated with the first photoproduct species that is more clearly observable at 400 nm than at 480 nm. The CH₂Cl radical has no singlet electronic absorption transition above 250 nm and can also be ruled out as the identity of the first photoproduct species. Comparison of our experimental results for both the vibrational frequencies and the electronic absorption transition observed in a recent femtosecond study at ~ 460 nm⁵⁷ clearly demonstrate that the first photoproduct species is the iso-CH₂Cl-I species. This is consistent with the proposed assignment of Åkesson and co-workers.⁵⁷ Thus, we assign the vibrational fundamentals of the first photoproduct species (iso-CH₂Cl-I) as follows: The 141 cm⁻¹ fundamental to the ν_6 C-Cl-I bend mode, the 174 cm⁻¹ fundamental to the ν_5 Cl-I stretch mode, the 376 cm⁻¹ fundamental to the ν_9 CH₂ twist mode, and the 724 cm⁻¹ fundamental to the ν_4 CH₂ wag mode.

The second photoproduct species that begins to appear as the first photoproduct species disappears in the 300–500 ps region also has at least two low-frequency fundamental modes. The CH₂Cl radical can thus be ruled out as the second photoproduct species. Since there appears to be at least two fundamental vibrational modes, this also rules out the possibilities of diatomic photoproduct species like the ICl⁻ ion or ICl molecule. The CH₂CI⁺ cation does not seem too likely for the following reasons: The computed 379 cm⁻¹ mode is fairly far away from the experimentally observed 312 cm⁻¹ Raman band, the computed electronic absorption transitions are also far from the experimental ~ 370 nm transient absorption band observed for the second photoproduct species in a recent femtosecond study,⁵⁷ and the oscillator strengths for the computed electronic transitions are relatively weak. The most likely candidate for the second photoproduct is the iso-CH₂I-Cl species if we assume the 312 cm⁻¹ Raman band in the experimental is a combination band of two lower frequency modes as was observed in the iso-CH₂Cl-I species picosecond resonance Raman spectra. This would give a ~ 105 cm⁻¹ fundamental frequency which forms a combination band with the strong 207 cm⁻¹ fundamental. The 105 cm⁻¹ fundamental can be assigned to the iso-CH₂I-Cl computed 113 cm⁻¹ ν_6 C-I-Cl bend mode and

TABLE I. Stokes resonance Raman band vibrational frequencies (in cm⁻¹) observed for the picosecond resonance Raman spectra shown in Figs. 2 (400 nm probe wavelength) and 3 (480 nm probe wavelength). The solvent was acetonitrile for all of the spectra.

Tentative vibrational assignment	(a) Iso-CH ₂ Cl-I and Iso-CH ₂ I-Cl at 400 nm probe wavelength									
	Raman shift (in cm ⁻¹)									
	3 ps	10 ps	30 ps	70 ps	150 ps	300 ps	500 ps	700 ps	1000 ps	2000 ps
Iso-CH ₂ Cl-I										
ν_6	152	141	141	140	150					
ν_5	172	171	174	170	176	172				
$\nu_5 + \nu_6/2\nu_5$	287	299	303	311	301	314				
ν_9		396	376	383	389					
ν_4		721	724	725	722	725				
$\nu_4 + \nu_5$		879	886	885	884	877				
$\nu_4 + \nu_5 + \nu_6$		1059	1062	1057	1063					
$\nu_4 + 2\nu_5 + 2\nu_6$			1373							
$2\nu_4$			1468							
Iso-CH ₂ I-Cl										
ν_5						211	207	206	211	213
$\nu_5 + \nu_6$						314	312	316	323	322
$2\nu_5$						422	428	420	427	426
	(b) Iso-CH ₂ Cl-I at 480 nm probe wavelength									
	Raman shift (in cm ⁻¹)									
Tentative vibrational assignment	3 ps	10 ps	30 ps	70 ps	150 ps	300 ps				
	Iso-CH ₂ Cl-I									
ν_6	162	151	145	146	146	156				
ν_5	192	177	177	180	178	188				
$\nu_5\nu_6/2\nu_5$		306	304	315	311					
ν_9		342	340	363	363					
ν_4		717	719	720	725					
$\nu_4 + \nu_5$		884	899	898	898					
$\nu_4 + \nu_5 + \nu_6$			1039	1050	1056					
$\nu_4 + 2\nu_5 + 2\nu_6$			1398	1377	1369					
$2\nu_4$			1446	1490	1477					

the experimental 207 cm⁻¹ Raman band could be assigned to the computed 236 cm⁻¹ ν_5 I-Cl stretch mode. The iso-CH₂I-Cl computed electronic transition position at 332 nm is reasonably close to the experimental transient absorption band \sim 370 nm observed in the femtosecond experiments⁵⁷ and the calculated oscillator strength (0.4008) for this transition is large. The B3LYP/Sadlej-PVTZ density-functional theory computational results indicate the iso-CH₂I-Cl species is about 4.54 kcal/mol more stable than the iso-CH₂Cl-I species. This is consistent with our assignment of the first photoproduct (which has a lifetime \sim 100 ps) to the iso-CH₂Cl-I species and the second photoproduct species (which has a longer lifetime on the order of hundreds of ps to nanoseconds) to the iso-CH₂I-Cl species. We note that this situation is similar to that found for the iso-CH₂Br-I and iso-CH₂I-Br species. The iso-CH₂I-Br species was computed to be more stable by \sim 4.1 kcal/mol and was the only species with a sufficiently long lifetime to be observed in nanosecond transient resonance Raman study.⁵⁴ We note that the assignment of the second photoproduct species to iso-CH₂I-Cl is not as clear cut as the first photoproduct species assignment to iso-CH₂Cl-I. However, the assignment of the second photoproduct to iso-CH₂I-Cl is consistent with the experimental and computational data available at this time.

The assignment of the second photoproduct species to iso-CH₂I-Cl gives rise to some interesting questions. How is iso-CH₂I-Cl formed? How is the C-Cl bond broken? There are two different probable scenarios for formation of the iso-CH₂I-Cl species. First, the initial 267 nm excitation of CH₂CI could give rise to both photodissociation of the C-I bond and the C-Cl bond reaction channels (to give CH₂Cl+I and CH₂I+Cl fragments, respectively) which then undergo recombination to produce the iso-CH₂Cl-I and iso-CH₂I-Cl photoproduct species. Second, the 267 nm excitation of CH₂CI results in cleavage of the C-I bond to give CH₂Cl+I fragments that then recombine to produce the iso-CH₂Cl-I species that then isomerize later to form either the more stable iso-CH₂I-Cl or parent CH₂CI molecules. Laser flash photolysis⁶² studies of the A-band absorption (\sim 270 nm) of CH₂CI in the gas phase showed that C-I bond cleavage is predominantly the primary reaction channel similar to A-band photolysis of CH₃I. A solution phase resonance Raman study of the A-band short-time photodissociation dynamics of CH₂CI indicated that there is mainly motion along the C-I stretch coordinate accompanied by smaller dynamics along other coordinates and this is consistent with photodissociation of the C-I bond.³⁴ Inspection of the Stokes picosecond time-resolved spectra in Fig. 2 shows that there does not appear to be any Raman bands clearly attributable to the

TABLE II. Parameters for the optimized geometry computed from the B3LYP/Sadlej-PVTZ density-functional theory computations for the iso-CH₂Cl-I, iso-CH₂I-Cl, CH₂ClI⁺, and CH₂Cl species. Bond lengths are in Å and bond angles are in degrees.

Parameter	B3LYP/Sadlej-PVTZ
iso-CH ₂ Cl-I	
C-Cl	1.6404
C-H	1.0914
Cl-I	2.8606
Cl-C-H	117.66
H-C-H	122.62
C-Cl-I	123.34
D(H-C-Cl-I)	±82.05
iso-CH ₂ I-Cl	
C-I	1.9547
C-H	1.0929
I-Cl	2.6149
I-C-H	118.52
H-C-H	119.59
C-I-Cl	120.78
D(H-C-I-Cl)	±79.61
CH ₂ ClI ⁺ cation	
C-Cl	1.7341
C-I	2.1887
C-H	1.1057
Cl-C-I	117.27
Cl-C-H	111.84
I-C-H	102.51
H-C-H	110.12
D(Cl-C-I-H)	±122.89
CH ₂ Cl radical	
C-Cl	1.7198
C-H	1.0884
Cl-C-H	117.01
H-C-H	125.43
D(Cl-C-H-H)	±171.23

TABLE IV. Electronic absorption transition energies and oscillator strengths (in parentheses) found from the density-functional theory computations (URPA/UB3LYP/Sadlej-PVTZ) for the iso-CH₂Cl-I, iso-CH₂I-Cl, CH₂ClI⁺, and CH₂Cl species.

Molecule	URPA/UB3LYP/Sadlej-PVTZ Singlet Transition Energies
Iso-CH ₂ I-Cl	339 nm (0.0020) 332 nm (0.4008) 270 nm (0.0067) 237 nm (0.0281) 232 nm (0.0008) 185 nm (0.0803)
Iso-CH ₂ Cl-I	551 nm (0.0000) 538 nm (0.0197) 453 nm (0.4932) 241 nm (0.0006) 234 nm (0.0206) 200 nm (0.0040)
CH ₂ ClI ⁺ cation	6849 nm (0.0000) 489 nm (0.0018) 274 nm (0.0000) 247 nm (0.0013) 244 nm (0.0001) 200 nm (0.0070)
CH ₂ Cl radical	224 nm (0.0024) 188 nm (0.0023) 180 nm (0.0000)

TABLE III. Comparison of the experimental fundamental vibrational frequencies (in cm⁻¹) found from the time-resolved resonance Raman picosecond spectra (this work) to the calculated B3LYP/Sadlej-PVTZ density functional theory vibrational frequencies.

Ultraviolet excitation of CH ₂ ClI		B3LYP Calc.		B3LYP Calc.		B3LYP Calc.	
400 nm ps-TR ³	400 nm ps-TR ³	iso-CH ₂ Cl-I		iso-CH ₂ I-Cl		CH ₂ ClI ⁺ cation	
30 ps in acetonitrile	500 ps in acetonitrile	Sadlej-PVTZ		Sadlej-PVTZ		Sadlej-PVTZ	
Experiment	Experiment						
		A' ν ₁ , sym. CH str.	3119	A ₁ ν ₁ , sym. CH str.	3114	A' ν ₁ , sym. CH str.	3002
		ν ₂ , CH ₂ scissor	1409	ν ₂ , CH ₂ scissor	1361	ν ₂ , CH ₂ def. or sciss.	1321
		ν ₃ , C-Cl stretch	975	ν ₃ , C-I stretch	794	ν ₃ , CH ₂ wag	1103
724		ν ₄ , CH ₂ wag	760	ν ₄ , CH ₂ wag	686	ν ₄ , C-Cl stretch	783
174	207	ν ₅ , Cl-I stretch	194	ν ₅ , I-Cl stretch	236	ν ₅ , C-I stretch	379
141	105 ^a	ν ₆ , C-Cl-I bend	136	ν ₆ , C-I-Cl bend	113	ν ₆ , I-C-Cl bend	191
		A'' ν ₇ , asym. CH str.	3276	A'' ν ₇ , asym. CH str.	3254	A'' ν ₇ , asym. CH str.	3050
		ν ₈ , CH ₂ rock	1020	ν ₈ , CH ₂ rock	869	ν ₈ , CH ₂ twist	1050
376		ν ₉ , CH ₂ twist	420	ν ₉ , CH ₂ twist	494	ν ₉ , CH ₂ rock	527
		CH ₂ Cl					
		Sadlej-PVTZ					
		A' ν ₁ , CH sym. str.	3134				
		ν ₂ , CH ₂ def. or scissor	1361				
		ν ₃ , C-Cl str.	827				
		ν ₄ , CH ₂ wag	242				
		A'' ν ₅ , CH asym. str.	3307				
		ν ₆ , CH ₂ rock	978				

^aValue obtained from subtraction of vibrational frequency for ν₅ from the combination band ν₅ + ν₆ in the 500 ps 400 nm Raman spectrum. str.=stretch; sym.=symmetric; asym.=asymmetric; def.=deformation.

iso-CH₂I-Cl species until after 150 ps when the iso-CH₂Cl-I species initially formed begins to show a noticeable intensity decrease. This is consistent with recent femtosecond transient absorption results⁵⁷ which show the first photoproduct species (associated with the strong ~460 nm and weaker ~710 nm absorption bands) decays with a lifetime ~100 ps and the second photoproduct species (associated with the intense ~370 nm and weak ~725 nm absorption bands) begins to appear as the first photoproduct species decays appreciably. There seems to be a correlation between the disappearance of the first photoproduct species and the appearance of the second photoproduct species. This and the fact that only the C-I bond cleavage channel is predominantly observed following A-band excitation of CH₂ClI indicates that the second mechanism for formation of the iso-CH₂I-Cl photoproduct is most likely. Our results suggest that the iso-CH₂Cl-I photoproduct is first produced from initially formed CH₂Cl and I fragments via recombination within the solvent cage and then subsequently isomerize to give either the more stable iso-CH₂I-Cl or parent CH₂ClI molecules. The iso-CH₂I-Cl probably then decays back to the parent CH₂ClI molecule. Since the 267 nm photon does not have enough energy to break the C-Cl bond by itself, the isomerization of the iso-CH₂Cl-I product to the iso-CH₂I-Cl product probably proceeds via a transition state that involves concerted formation of the C-I bond, strengthening of the I-Cl bond and cleavage of the C-Cl bond. It is conceivable that the cooperative formation of the C-I bond and strengthening of the I-Cl bond as the C-Cl bond breaks would make it easier to break the C-Cl bond for the isomerisation reaction. We note that our results for CH₂ClI are somewhat similar to recent results reported by Reid and coworkers for OCIO.⁶³ They photoexcited OCIO and observed fast geminate recombination of the ClO and O fragments to produce a vibrationally hot OCIO photoproduct that subsequently formed some ClOO with a 27.9 ± 4.5 ps time constant appearance time that subsequently decayed with a time constant of 398 ± 50 ps.⁶³

It is interesting that we observed a much smaller transient resonance Raman signal for the isomer photoproducts from ultraviolet excitation of CH₂ClI than previously found for isomer photoproducts from excitation of CH₂I₂, CH₂Br₂, and CHI₃ in the solution phase.^{52,64} This could be due to more efficient solvent cage escape of the CH₂Cl fragment initially produced by ultraviolet excitation of CH₂ClI in comparison to the larger and more massive CH₂I, CH₂Br, and CHI₂ fragments initially formed from ultraviolet excitation of CH₂I₂, CH₂Br₂, and CHI₃ in condensed phase environments. Molecular beam studies on ultraviolet C-X photodissociation reactions in haloalkanes and dihaloalkanes generally exhibit the trend of partitioning more of the available energy into populating modes associated with the internal degrees of freedom of the photofragments as the parent molecule becomes more massive and/or branched in structure.⁶⁵⁻⁷¹ Thus, the CH₂Cl fragment formed from CH₂ClI ultraviolet photodissociation would be expected to have much more translational energy than the larger and more massive CH₂I, CH₂Br, and CHI₂ fragments produced from corresponding photodissociation reactions of CH₂I₂,

CH₂Br₂, and CHI₃. This would be consistent with more fragments escaping the solvent cage for the initially produced CH₂Cl fragment which leads to our observing a lower signal level for the time-resolved resonance Raman spectra of the isomer species formed from solvent induced recombination. Escape of the photofragments from the solvent cage would likely produce some I-solvent contact charge transfer complexes.^{72,73} The transient absorption associated with iodine atom-alkane (or cycloalkane) solvents appears to have a very broad absorption in the 300-400 nm region that has an extinction coefficient of ~1700 M⁻¹ cm⁻¹ in *n*-octane solvent⁷² (Ref. 72 cautions that this value may be somewhat too high). The iodine atom-acetonitrile solvent contact charge transfer absorption band probably has a similar moderate extinction coefficient. The transient absorption bands associated with the two isomers of chloriodomethane are very strong and on the order of 10 000 M⁻¹ cm⁻¹ or greater (see Ref. 44) and have computed oscillator strengths on the order of 0.4-0.5 (this work). Thus, one would expect the I-solvent absorption may be more difficult to observe clearly compared to the isomer-chloriodomethane that have much stronger transitions in the 350-400 nm region. This may be part of the reason that the I-solvent transient absorption was not clearly seen in the recently reported femtosecond transient absorption study.⁵⁷ An alternative possibility is that there is not much difference in the solvent cage escape for the faster CH₂Cl fragment compared to CH₂I and CH₂Br fragments, but the faster CH₂Cl fragment has a lower quantum yield for formation of the isomer dihalomethane species than the slower and heavier fragments from photolysis of CH₂I₂ and CH₂BrI. This would also be consistent with the low-Raman signal observed for the iso-chloriodomethane species here and the lack of observation of the I-solvent complex transient absorption band in the femtosecond transient absorption study.⁵⁷ It is not clear how the quantum yield for formation of the isomer species depends on the translational energy of the fragments. Another possibility to be considered (though not mutually exclusive with respect to the probability of solvent cage escape taking place), is that the fast CH₂Cl fragment undergoes reaction with the solvent more easily than the CH₂I, CH₂Br and CHI₂ fragments produced in the corresponding photodissociation reactions of CH₂I₂, CH₂Br₂, and CHI₃. We note that halogen atom-solvent complexes absorption band position and intensity and stability of the complex can vary noticeably depending on the type of halogen atom and type of solvent.⁷²⁻⁷⁶ Further work is needed to better understand how the quantum yield for solvent cage escape versus geminate (or near-geminate) recombination vary as a function of the polyhalomethane mass and substituents.

ACKNOWLEDGMENTS

D.L.P. would like to thank the Research Grants Council (RGC) of Hong Kong and the Committee on Research and Conference Grants (CRCG) from the University of Hong Kong for financial support of this research. A.W.P. and D.P. are grateful to the EPSRC for financial support. W.M.K. gratefully acknowledges financial support from the Croucher

foundation, Hong Kong. The picosecond resonance Raman experiments were carried out within the Central Laser Facility, CLRC Rutherford Appleton Laboratory.

- ¹H. E. Simmons and R. D. Smith, *J. Am. Chem. Soc.* **81**, 4256 (1959).
- ²D. C. Blomstrom, K. Herbig, and H. E. Simmons, *J. Org. Chem.* **30**, 959 (1965).
- ³N. J. Pienta and P. J. Kropp, *J. Am. Chem. Soc.* **100**, 655 (1978).
- ⁴P. J. Kropp, N. J. Pienta, J. A. Sawyer, and R. P. Polniaszek, *Tetrahedron* **37**, 3229 (1981).
- ⁵P. J. Kropp, *Acc. Chem. Res.* **17**, 131 (1984).
- ⁶E. C. Friedrich, J. M. Domek, and R. Y. Pong, *J. Org. Chem.* **50**, 4640 (1985).
- ⁷E. C. Friedrich, S. E. Lunetta, and E. J. Lewis, *J. Org. Chem.* **54**, 2388 (1989).
- ⁸S. Durandetti, S. Sibille, and J. Pérchon, *J. Org. Chem.* **56**, 3255 (1991).
- ⁹J. M. Concellón, P. L. Bernad, and J. A. Pérez-Andrés, *Tetrahedron Lett.* **39**, 1409 (1998).
- ¹⁰Th. Class and K. Ballschmiter, *J. Atmos. Chem.* **6**, 35 (1988).
- ¹¹S. Klick and K. Abrahamsson, *J. Geophys. Res. [Oceans]* **97**, 12683 (1992).
- ¹²K. G. Heumann, *Anal. Chim. Acta* **283**, 230 (1993).
- ¹³R. M. Moore, M. Webb, R. Tokarczyk, and R. Wever, *J. Geophys. Res., [Oceans]* **101**, No. 9, 20899 (1996).
- ¹⁴J. C. Mössigner, D. E. Shallcross, and R. A. Cox, *J. Chem. Soc., Faraday Trans.* **94**, 1391 (1998).
- ¹⁵L. J. Carpenter, W. T. Sturges, S. A. Penkett, P. S. Liss, B. Alicke, K. Hebestreir, and U. Platt, *J. Geophys. Res. [Oceans]* **104**, 1679 (1999).
- ¹⁶B. Alicke, K. Hebestreir, J. Stutz, and U. Platt, *Nature (London)* **397**, 572 (1999).
- ¹⁷M. Kawasaki, S. J. Lee, and R. Bersohn, *J. Chem. Phys.* **63**, 809 (1975).
- ¹⁸G. Schmitt and F. J. Comes, *J. Photochem.* **14**, 107 (1980).
- ¹⁹P. M. Kroger, P. C. Demou, and S. J. Riley, *J. Chem. Phys.* **65**, 1823 (1976).
- ²⁰S. J. Lee and R. Bersohn, *J. Phys. Chem.* **86**, 728 (1982).
- ²¹L. J. Butler, E. J. Hints, and Y. T. Lee, *J. Chem. Phys.* **84**, 4104 (1986).
- ²²L. J. Butler, E. J. Hints, and Y. T. Lee, *J. Chem. Phys.* **86**, 2051 (1987).
- ²³G. Baum, P. Felder, and J. R. Huber, *J. Chem. Phys.* **98**, 1999 (1993).
- ²⁴E. A. J. Wannemacher, P. Felder, and J. R. Huber, *J. Chem. Phys.* **95**, 986 (1991).
- ²⁵W. Radloff, P. Farmanara, V. Stert, E. Schreiber, and J. R. Huber, *Chem. Phys. Lett.* **291**, 173 (1998).
- ²⁶S. R. Cain, R. Hoffman, and R. Grant, *J. Phys. Chem.* **85**, 4046 (1981).
- ²⁷K.-W. Jung, T. S. Ahmadi, and M. A. El-Sayed, *Bull. Korean Chem. Soc.* **18**, 1274 (1997).
- ²⁸S. L. Baughcum, H. Hafmann, S. R. Leone, and D. Nesbitt, *Faraday Discuss. Chem. Soc.* **67**, 306 (1979).
- ²⁹S. L. Baughcum and S. R. Leone, *J. Chem. Phys.* **72**, 6531 (1980).
- ³⁰J. Zhang and D. G. Imre, *J. Chem. Phys.* **89**, 309 (1988).
- ³¹J. Zhang, E. J. Heller, D. Huber, D. G. Imre, and D. Tannor, *J. Chem. Phys.* **89**, 3602 (1988).
- ³²W. M. Kwok and D. L. Phillips, *Chem. Phys. Lett.* **235**, 260 (1995).
- ³³W. M. Kwok and D. L. Phillips, *J. Chem. Phys.* **104**, 2529 (1996).
- ³⁴W. M. Kwok and D. L. Phillips, *J. Chem. Phys.* **104**, 9816 (1996).
- ³⁵S. Q. Man, W. M. Kwok, and D. L. Phillips, *J. Phys. Chem.* **99**, 15705 (1995).
- ³⁶S. Q. Man, W. M. Kwok, A. E. Johnson, and D. L. Phillips, *J. Chem. Phys.* **105**, 5842 (1996).
- ³⁷F. Duschek, M. Schmitt, P. Vogt, A. Materny, and W. Kiefer, *J. Raman Spectrosc.* **28**, 445 (1997).
- ³⁸M. Braun, A. Materny, M. Schmitt, W. Kiefer, and V. Engel, *Chem. Phys. Lett.* **284**, 39 (1998).
- ³⁹X. Zheng and D. L. Phillips, *Chem. Phys. Lett.* **313**, 467 (1999).
- ⁴⁰J. P. Simons and P. E. R. Tatham, *J. Chem. Soc. A* **1966**, 854.
- ⁴¹G. P. Brown and J. P. Simons, *Trans. Faraday Soc.* **65**, 3245 (1969).
- ⁴²H. Mohan, K. N. Rao, and R. M. Iyer, *Radiat. Phys. Chem.* **23**, 505 (1984).
- ⁴³G. Maier and H. P. Reisenauer, *Angew. Chem. Int. Ed. Engl.* **25**, 819 (1986).
- ⁴⁴G. Maier, H. P. Reisenauer, J. Lu, L. J. Schaad, and B. A. Hess, Jr., *J. Am. Chem. Soc.* **112**, 5117 (1990).
- ⁴⁵L. Andrews, F. T. Prochaska, and B. S. Ault, *J. Am. Chem. Soc.* **101**, 9 (1979).
- ⁴⁶H. Mohan and R. M. Iyer, *Radiat. Eff.* **39**, 97 (1978).
- ⁴⁷H. Mohan and P. N. Moorthy, *J. Chem. Soc., Perkin Trans.* **2**, 277 (1990).
- ⁴⁸B. J. Schwartz, J. C. King, J. Z. Zhang, and C. B. Harris, *Chem. Phys. Lett.* **203**, 503 (1993).
- ⁴⁹K. Saitow, Y. Naitoh, K. Tominaga, and Y. Yoshihara, *Chem. Phys. Lett.* **262**, 621 (1996).
- ⁵⁰A. N. Tarnovsky, J.-L. Alvarez, A. P. Yartsev, V. Sündstrom, and E. Åkesson, *Chem. Phys. Lett.* **312**, 121 (1999).
- ⁵¹X. Zheng and D. L. Phillips, *J. Phys. Chem. A* **104**, 6880 (2000).
- ⁵²W. M. Kwok, C. Ma, A. W. Parker, D. Phillips, M. Towrie, P. Matousek, and D. L. Phillips, *J. Chem. Phys.* **113**, 7471 (2000).
- ⁵³X. Zheng and D. L. Phillips, *Chem. Phys. Lett.* **324**, 175 (2000).
- ⁵⁴X. Zheng and D. L. Phillips, *J. Chem. Phys.* **113**, 3194 (2000).
- ⁵⁵X. Zheng, W. M. Kwok, and D. L. Phillips, *J. Phys. Chem. A* **104**, 10464 (2000).
- ⁵⁶X. Zheng, W.-H. Fang, and D. L. Phillips, *J. Chem. Phys.* **113**, 10934 (2000).
- ⁵⁷A. N. Tarnovsky, M. Wall, M. Rasmusson, T. Pascher, and E. Åkesson, *J. Chin. Chem. Soc. (Taipei)* **47**, 769 (2000).
- ⁵⁸M. Towrie, P. Matousek, A. W. Parker, and W. Shaikh, *Meas. Sci. Technol.* **9**, 816 (1998).
- ⁵⁹M. J. Frisch, G. W. Trucks, H. B. Schlegel *et al.* GAUSSIAN 98, Revision A.7, Gaussian, Inc., Pittsburgh, PA, 1998.
- ⁶⁰A. J. Sadlej, *Theor. Chim. Acta* **81**, 339 (1992).
- ⁶¹R. Bauernschmitt and R. Ahlrichs, *Chem. Phys. Lett.* **256**, 454 (1996).
- ⁶²G. Schmitt and F. J. Comes, *J. Photochem. Photobiol., A* **41**, 13 (1987).
- ⁶³C. L. Thomsen, M. P. Philpott, S. C. Hayes, and P. J. Reid, *J. Chem. Phys.* **112**, 505 (2000).
- ⁶⁴W. M. Kwok, C. Ma, A. W. Parker, D. Phillips, M. Towrie, P. Matousek, and D. L. Phillips, *J. Chem. Phys.* (submitted).
- ⁶⁵S. J. Riley and K. R. Wilson, *Faraday Discuss. Chem. Soc.* **53**, 132 (1972).
- ⁶⁶C. Paterson, F. G. Godwin, and P. A. Gorry, *Mol. Phys.* **60**, 729 (1987).
- ⁶⁷Q. Zhu, J. R. Cao, Y. Wen, J. Zhang, Y. Huang, W. Fang, and X. Wu, *Chem. Phys. Lett.* **144**, 486 (1988).
- ⁶⁸W. K. Kang, K. W. Jung, D. C. Kim, K.-H. Jung, and H. S. Kim, *Chem. Phys.* **196**, 363 (1995).
- ⁶⁹F. G. Godwin, C. Paterson, and P. A. Gorry, *Mol. Phys.* **61**, 827 (1987).
- ⁷⁰T. K. Minton, P. Felder, R. J. Brudzynski, and Y. T. Lee, *J. Chem. Phys.* **81**, 1759 (1984).
- ⁷¹G. M. Nathanson, T. K. Minton, S. F. Shane, and Y. T. Lee, *J. Chem. Phys.* **90**, 6157 (1990).
- ⁷²S. R. Logan, R. Bonneau, J. Jousset-Dubien, and P. Fornier De Violet, *J. Chem. Soc., Perkin Trans. 1* **71**, 2148 (1975).
- ⁷³P. Fornier De Violet, R. Bonneau, and J. Jousset-Dubien, *J. Chim. Phys. Phys.-Chim. Biol.* **70**, 1404 (1973).
- ⁷⁴K. D. Raner, J. Luszytky, and K. U. Ingold, *J. Phys. Chem.* **93**, 564 (1990).
- ⁷⁵Z. B. Alfassi, R. E. Huie, J. P. Mittal, P. Neta, and L. C. T. Shoute, *J. Phys. Chem.* **97**, 9120 (1993).
- ⁷⁶M. P. Philpott, S. C. Hayes, C. L. Thomsen, and P. J. Reid, *Chem. Phys.* **263**, 389 (2001).