



<b>Title</b>	<b>Are Self-Similar States in Fibonacci Systems Transparent?</b>
<b>Author(s)</b>	<b>Jin, GJ; Wang, Z</b>
<b>Citation</b>	<b>Physical Review Letters, 1997, v. 79 n. 26, p. 5298</b>
<b>Issued Date</b>	<b>1997</b>
<b>URL</b>	<b><a href="http://hdl.handle.net/10722/42440">http://hdl.handle.net/10722/42440</a></b>
<b>Rights</b>	<b>Creative Commons: Attribution 3.0 Hong Kong License</b>

## Are Self-Similar States in Fibonacci Systems Transparent?

In a recent Letter [1] Maciá and Domínguez-Adame intended to address the physical nature of critical wave functions in a generalized Fibonacci system. Apart from several interesting results presented, one main conclusion reached is that “*self-similar wave functions are those exhibiting higher transmission coefficients in a finite Fibonacci system.*” Although there exist extended or transparent states in many aperiodic systems under some special conditions, the conclusion itself is unfortunately incorrect and misleading, because it is based on a serious miscalculation of the transmission coefficient. In this Comment, we wish to clarify this point and present a correct calculation.

For the Hamiltonian considered in Ref. [1], four original transfer matrices ( $X, Y, Z, W$ ) can be cast into two new matrices  $R_A(=ZYX)$  and  $R_B(=WX)$ .  $R_A$  and  $R_B$  are arranged in a Fibonacci sequence. In the context of the formulation of Ref. [1], there always exists one energy  $E$  satisfying a relation

$$E = \alpha \frac{1 + \gamma^2}{1 - \gamma^2}, \quad (1)$$

where  $\alpha$  represents the on-site energy,  $\gamma$  the transfer integral. For these energies,  $[R_A, R_B] = 0$ . Based on

Eq. (1), the global transfer matrix  $M(N)$  in Ref. [1] is obtained. Using the  $M(N)$ , the authors further derive the transmission coefficient as

$$\tau(N) = \frac{1}{1 + [(1 - \gamma^2)^2 / (4 - E^2)\gamma^2] \sin^2(N\phi)}, \quad (2)$$

where  $\phi$  is a function of  $E$ ,  $\alpha$ , and  $\gamma$ . Then they suggest that the transparent condition be fulfilled when  $\sin(N\phi) = 0$ . However, we must note that the condition  $\sin(N\phi) = 0$  may not be consistent with Eq. (1), particularly in self-similar states. Therefore, under the condition  $\sin(N\phi) = 0$  in the self-similar states, Eq. (2) cannot be used and thus  $\tau(E) = 1$  should not be expected in the self-similar states. In the Letter, they take  $N = F_{17}$ ,  $\gamma = 2$ ,  $\alpha = 0.1$ , and  $E = -\sqrt{\alpha^2 + 4 \cos(1160\pi/N)} = 0.3348\dots$  [correspond to  $\sin(N\phi) = 0$ ] to plot Fig. 2 in Ref. [1], which statistically exhibits self-similar features and is claimed to be at a transparent state  $\tau(E) = 1$ . Unfortunately, the above three energy parameters ( $E$ ,  $\alpha$ , and  $\gamma$ ) do not satisfy Eq. (1), so their claim that this self-similar state corresponds to a transparent state does not make sense.

Actually, if we consider  $E$ ,  $\alpha$ , and  $\gamma$  as three independent parameters and denote the global transfer matrix as  $T(N)$  with matrix elements  $t_{i,j}$ s ( $i, j = 1, 2$ ), the transmission coefficient can be obtained as

$$\tau(N) = \frac{4 - E^2}{[t_{21} - t_{12} + (t_{22} - t_{11})E/2]^2 + (t_{22} + t_{11})^2(1 - E^2/4)}, \quad (3)$$

where  $t_{i,j}$  is a function of the three energy parameters. Note that Eq. (3) is more general and  $T_N \neq M_N$  in a general case. Once  $T(N) = M(N)$  (i.e., Eq. (1) holds), it is straightforward to show that Eq. (3) is equivalent to Eq. (2). From Eq. (3), we can check that when Eq. (1) is satisfied, (i) if  $\gamma = 2$ ,  $\alpha = 0.75$ ,  $E = -1.25$ , then  $\tau(F_{16}) = 0.5909\dots$ ; (ii) if  $\gamma = 2$ ,  $\alpha = 0.5$ ,  $E = -5/6$ , then  $\tau = 0.7425\dots$ , as obtained in Ref. [1]; the corresponding states are not self-similar. However, if we substitute  $\gamma = 2$ ,  $\alpha = 0.1$ ,  $E = -\sqrt{0.1^2 + 4 \cos^2(1160\pi/F_{17})}$  (corresponding to a self-similar state) into Eq. (3),  $\tau(N) = 0.2298\dots$ , instead of the result  $\tau = 1$  in Ref. [1].

On the other hand, when Eq. (1) is fulfilled, we can further show that  $R_A = R_C^3$ ,  $R_B = R_C^2$ , and  $M(N) = R_C^N$ , where

$$R_C = \begin{pmatrix} \gamma^{-1}(E - \alpha) & -\gamma \\ \gamma^{-1} & 0 \end{pmatrix}. \quad (4)$$

Then in terms of Chebyshev polynomials of the second order,  $M(N)$  can be obtained more easily.

Finally, we wish to pinpoint that if  $R_A$  and  $R_B$  commute, which corresponds to a zero-invariant case, one may find a higher transmission coefficient more easily. For example, (i)  $\gamma = 2$ ,  $\alpha = 0.1$ ,  $E = -1/6$ ,  $\tau(F_{17}) = 0.9232\dots$ ; (ii)  $\gamma = 1.5$ ,  $\alpha = 0.1$ ,  $E =$

$-13/50$ ,  $\tau(F_{17}) = 0.9938\dots$ ; (iii)  $\gamma = 2$ ,  $\alpha = 1.5$ , and  $E = -2.5$ ,  $\tau = 1$ . Notice that these states are not self-similar. It is well known that, in the on-site or in the transfer model, the self-similar states correspond to nonzero invariants [2]. In the mixing model, can the states with zero invariant be self-similar? From both analytical and numerical calculations, we find actually that if  $R_A$  and  $R_B$  commute, there exist only extended states rather than *self-similar states*, in sharp contrary to the conclusion reached in Ref. [1].

G.J. Jin<sup>1</sup> and Z.D. Wang<sup>2</sup>

<sup>1</sup>Department of Physics and National Laboratory of Solid State Microstructures, Nanjing University  
Nanjing 210093, People's Republic of China

<sup>2</sup>Department of Physics, The University of Hong Kong  
Pokfulam Road, Hong Kong, People's Republic of China

Received 7 October 1996 [S0031-9007(97)04854-0]  
PACS numbers: 71.23.Ft, 61.44.-n

[1] E. Maciá and F. Domínguez-Adame, Phys. Rev. Lett. **76**, 2957 (1996).

[2] M. Kohmoto, B. Sutherland, and C. Tang, Phys. Rev. B **35**, 1020 (1987).