



<b>Title</b>	<b>Enzymatic activity of endophytic fungi of six native seedling species from Doi Suthep-Pui National Park, Thailand</b>
<b>Author(s)</b>	<b>Lumyong, S; Lumyong, P; McKenzie, EHC; Hyde, KD</b>
<b>Citation</b>	<b>Canadian Journal of Microbiology, 2002, v. 48 n. 12, p. 1109-1112</b>
<b>Issued Date</b>	<b>2002</b>
<b>URL</b>	<b><a href="http://hdl.handle.net/10722/42683">http://hdl.handle.net/10722/42683</a></b>
<b>Rights</b>	<b>Creative Commons: Attribution 3.0 Hong Kong License</b>

NOTE

## Enzymatic activity of endophytic fungi of six native seedling species from Doi Suthep-Pui National Park, Thailand

Saisamorn Lumyong, Pipob Lumyong, Eric H.C. McKenzie, and Kevin D. Hyde

**Abstract:** Endophytic fungi were isolated from the stems, petioles, midribs, and leaves of seedlings of six native tree species collected from Doi Suthep-Pui National Park, Thailand. Endophytes were isolated from all tissue samples investigated, and taxa included five ascomycetes, eight anamorphic taxa, and numerous sterile mycelia. Twenty-six strains were tested for their ability to produce cellulase, mannanase, proteinase, and xylanase. The ability to produce these enzymes was distributed amongst the strains tested. Rainforest seedlings supported a diverse array of endophytes that have a wide range of enzymatic activities. The implication of enzyme production in relation to lifestyle abilities of the endophytes is discussed.

*Key words:* cellulase, hydrolytic enzyme, mannanase, protease, xylanase.

**Résumé :** Des champignons endophytes ont été isolés des tiges, des pétioles, des nervures médianes et des feuilles des plants de six espèces d'arbres indigènes recueillis dans le Parc National Doi Suthep-Pui, Thaïlande. Des endophytes ont été isolés à partir de tous les échantillons de tissus analysés et les taxons comprenaient cinq ascomycètes, huit taxons anamorphiques et de nombreux mycéliums stériles. Nous avons évalué la capacité de vingt-six souches à produire de la cellulase, de la mannanase, de la protéinase et de la xylanase. La capacité de produire ces enzymes était répartie parmi les souches analysées. Les plants issus de forêts tropicales ont soutenu le développement d'un éventail varié d'endophytes ayant une large gamme d'activités enzymatiques. L'incidence de la production des enzymes sur les habitudes de vie des endophytes est débattue.

*Mots clés :* cellulase, enzymes hydrolytiques, mannanase, protéase, xylanase.

[Traduit par la Rédaction]

Numerous authors have reported the presence of endophytes in temperate plants (e.g., Petrini et al. 1982; Petrini and Fisher 1988), but there has been less emphasis on tropical species (Rodrigues and Petrini 1997). Recent studies on endophytes of tropical plants include bamboo (Umali et al. 1999; Lumyong et al. 2000), banana (Brown et al. 1998; Photita et al. 2001), *Cuscuta reflexa* (Suryanarayanan et al. 2000), *Parthenium hysterophorus* (Romero et al. 2001), palms (Rodrigues and Samuels 1990; Rodrigues 1994; Guo et al. 1998; Taylor et al. 1999; Fröhlich et al. 2000), and wild Zingiberaceae (Bussaban et al. 2001).

The role of endophytes in plants has been the subject of much debate. It may range from the protection of the plants against herbivory to improvement in mineral absorption and drought tolerance (Bacon and Hill 1996). Some latent pathogens live endophytically, while others have been shown to be the initial saprobes that colonize dead plant material (Guo et al. 1998). Endophytes are important producers of secondary metabolites, and there has been biotechnological interest in their potential to produce novel metabolites (Dreyfuss and Chapela 1995; Strobel et al. 1996a, 1996b; Karim et al. 1997; Peláez et al. 1998).

Received 22 May 2002. Revision received 15 November 2002. Accepted 11 December 2002. Published on the NRC Research Press Web site at <http://cjfm.nrc.ca> on 11 January 2003.

**S. Lumyong.**<sup>1</sup> Department of Biology, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand.

**P. Lumyong.** Department of Plant Pathology, Faculty of Agriculture, Chiang Mai University, Chiang Mai 50200, Thailand.

**E.H.C. McKenzie.** Landcare Research, Private Bag 92170, Auckland, New Zealand.

**K.D. Hyde.** Centre for Research in Fungal Diversity, Department of Ecology and Biodiversity, The University of Hong Kong, Pokfulam Road, Hong Kong.

<sup>1</sup>Corresponding author (e-mail: [schoi009@chiangmai.ac.th](mailto:schoi009@chiangmai.ac.th)).

**Table 1.** Fungal endophytes isolated from rainforest seedlings.

Endophyte	Host					
	<i>Camellia sinensis</i> var. <i>assamensis</i>	<i>Cinnamomum</i> <i>inerts</i>	<i>Garcinia</i> <i>cowa</i>	<i>Litsea</i> <i>salicifolia</i>	<i>Manglietia</i> <i>garrettii</i>	<i>Trichilla</i> <i>connaroides</i>
<i>Colletotrichum</i> spp.	8	11	31	–	36	13
<i>Curvularia</i> sp.	–	–	–	–	1	–
<i>Didymella</i> sp.	–	–	1	–	–	–
<i>Glomerella cingulata</i>	3	12	28	5	12	1
<i>Guignardia cocoicola</i>	1	2	3	3	1	5
<i>Helminthosporium</i> sp.	–	–	–	–	4	–
<i>Nigrospora</i> sp.	1	2	1	–	2	2
<i>Pestalotiopsis</i> sp.	–	1	–	–	1	1
<i>Phoma</i> sp.	3	2	2	6	–	10
<i>Phomopsis</i> sp.	4	1	2	8	–	–
<i>Sporormia</i> sp.	1	–	–	–	–	–
<i>Xylaria</i> sp.	2	–	–	–	2	–
<i>Mycelia sterilia</i>	9	23	7	1	7	18
Unidentified sp.*	7	1	–	1	–	–

Note: Data are percent colonization rate.

\*Unidentified anamorphic fungi.

To establish the functional role of endophytes, it would be useful to determine their patterns of substrate utilization and enzyme production (Carroll and Petrini 1981). We report on endophytes isolated from seedlings of six native rainforest trees in northern Thailand, one of the few reports of endophytes from seedlings of native tropical plants (Rodrigues 1994). We tested the ability of selected isolates to produce cellulase, mannanase, protease, and xylanase to elucidate their possible roles.

Six seedling species (Table 1), all large, native rainforest tree species, were selected for this study in an area of Doi Suthep-Pui National Park at 1000 m above sea level. The seedlings were grown in a nursery that was situated within deciduous forest. Five healthy seedlings of each species, approx. 50 cm high, were randomly selected, returned to the laboratory, and processed within 24 h.

From each plant, five old and five young leaves were selected, and all stems were cut from the seedlings and treated (6 seedlings × (i) 5 specimens (5 old leaves, 5 young leaves), (ii) 5 petioles, and (iii) 1 stem = 336 samples) for isolation of endophytic fungi following the method of Photita et al. (2001). Colonization rate is expressed as a percentage.

Colonization rate =

$$\frac{\text{Total number of samples yielding } \geq 1 \text{ isolate} \times 100}{\text{Total number of samples in that trial}}$$

Twenty-six fungal isolates were tested for cellulase, mannanase, and xylanase production using 1% carboxymethyl cellulose, 1% locust bean gum (Difco Laboratories, Detroit, Mich.), and 1% xylan as carbon source, respectively (Downie et al. 1994; Pointing 1999). A modified agar diffusion method incorporating Congo red dye (Downie et al. 1994) was used as a qualitative assay. Protease activity was assayed using casein hydrolysis medium, which contained 1% skimmed milk. After incubation at 30°C, the diameter of the clear zone was measured.

The 336 tissue samples yielded five ascomycetes, eight anamorphic taxa, and numerous sterile mycelia (Table 1). Endophytes were isolated from all types of tissue. The sterile mycelia were not separated into morphospecies, and no basidiomycetes were isolated.

Most of the endophytes were isolated from several plant species, and all plant species hosted at least six endophytic taxa (Table 1). *Colletotrichum* spp., *Glomerella cingulata*, and sterile mycelia were the most common taxa isolated from most hosts. Some fungi may be specific to certain host seedlings. *Helminthosporium* sp. and *Curvularia* sp. were isolated only from *Manglietia garrettii* Craib, *Didymella* sp. only from *Garcinia cowa* Roxb., and *Sporormia* sp. only from *Camellia sinensis* (L.) var. *assamica* (Mast.) Kitamura. *Glomerella cingulata*, *Guignardia cocoicola*, and sterile mycelia occurred in all hosts. The colonization rate from each host and each plant tissue type is presented in Table 2. Most taxa were isolated from the stem, while the number of taxa isolated from other plant tissues varied within a low range. Fewer taxa were isolated from the young leaves.

The endophytes found were similar to those found in mature plants of other species at Doi Suthep-Pui National Park (Bussaban et al. 2001; Photita et al. 2001) and in other tropical plants (e.g., Suryanarayanan et al. 2000). Mature plants, however, tend to support a larger fungal assemblage (e.g., Taylor et al. 1999). The current data supports the hypothesis that endophytic fungi colonize plant tissues via inoculum from the surrounding environment (Fröhlich et al. 2000). As seedling leaves age, more spores will fall onto them, and thus, older leaves can be expected to have a larger and more diverse endophyte assemblage.

There was considerable variation in enzyme production between the endophytes tested (Table 3). The knowledge of enzyme production by fungi provides some idea of their biotechnological potential and may give insight into the lifestyles of endophytes. There were some interesting trends in the results that hint at possible roles for some of the endophytes isolated from the various seedlings. Most strains of

**Table 2.** Comparison of colonization rate in different seedling species and tissue types.

Seedling species	Colonization rate of each taxa per tissue type												
	Old tissue					Young tissue							
	Stem	Petiole	Top midrib	Between vein	Bottom midrib	Top intervein	Bottom intervein	Top midrib	Between vein	Bottom midrib	Top intervein	Bottom intervein	Top vein
<i>Camellia sinensis</i> var. <i>assamica</i>	9	5.6	2	2.3	4	1	2	1.3	1	1.5	1	1	0
<i>Cinnamomum iners</i>	6	6	6.6	4.6	3.6	3	2.6	5.3	6	4.6	2.3	2.3	2.6
<i>Garcinia cowa</i>	9.8	4.4	6.8	7.4	5.2	6.8	6.4	5.2	4.8	5.2	5.6	5.6	4
<i>Listea salicifolia</i>	12.6	4.4	1.6	5	2.8	2.8	2.2	0	0	1	1	1	1.4
<i>Manglietia garrettii</i>	6.6	4.6	6.4	4.8	5.4	3.8	4	5.6	3.6	3.6	3.2	3.2	5.5
<i>Trichillia connaroides</i>	8.8	10	6.6	3.8	1.6	4.8	4.6	2	6.6	2.2	0	0	1.6

**Table 3.** Production of enzymes by endophytes.

Endophyte	Host	Enzyme production ratio											
		Cellulase	Mannanase	Protease	Xylanase								
<i>Colletotrichum</i> sp.	Mg	1	1	+	0								
	Cs	3	4	+	2								
	Ci	3	3	+	4								
<i>Pestalotiopsis</i> sp.	Mg	3	4	-	0								
	Gc	0	0	+	0								
	Cs	0	4	+	0								
<i>Phoma</i> sp.	Ls	0	3	-	2								
	Mg	2	2	+	0								
	Gc	4	4	+	0								
	Tc	2	0	+	2								
<i>Phomopsis</i> sp.	Cs	4	0	-	0								
	Ls	4	0	-	0								
	Mg	2	0	+	3								
	Gc	4	4	-	4								
<i>Xylaria</i> sp.	Tc	4	4	-	4								
	Ls	3	2	+	1								
	Ci	4	4	-	4								
	Mg	4	3	+	1								
<i>Mycelia sterilia</i>	Gc	3	0	+	0								
	Tc	4	4	+	4								
	Cs	4	4	-	3								
	Ls	4	0	-	4								
<i>Mycelia sterilia</i>	Mg	2	0	-	1								
	Gc	2	3	-	1								
	Tc	3	4	-	4								
	Cs	2	2	-	2								

**Note:** Enzyme production ratio = the ratio of clear zone diameter to that of colony diameter. The extracellular enzymatic reactions were classified into the following four types: (i) strong reaction, the extracellular enzyme ratio was higher than or equal to 2; (ii) medium reaction, the extracellular enzyme ratio was less than 2 but more than 1; (iii) weak reaction, the extracellular enzyme ratio was equal to or less than 1; and (iv) no reaction, there is no reaction at all. In the case of protease, the reaction was small and could only be measured as + or -. Ci, *Cinnamomum iners* Reinw. ex Bl.; Cs, *Camellia sinensis* var. *assamica*; Gc, *Garcinia cowa*; Ls, *Listea salicifolia* Nees ex Roxb.; Mg, *Manglietia garrettii*; and Tc, *Trichillia connaroides* (Wight & Arn.) Benth.

*Pestalotiopsis* did not produce any cellulases or xylanases but showed some evidence of mannanase production. *Pestalotiopsis* species are mostly weak pathogens and often develop on leaf spots produced by other pathogens (Guba 1961). Most *Phoma* strains tested did not produce any mannanases or xylanases but did produce cellulases. These fungi are similar to the second group of Carroll and Petrini (1981), which are probably incapable of penetrating living cells. Such endophytes may require simple carbon sources for active growth, a nutritional feature that is common in symbiotic fungi (Carroll and Petrini 1981).

The production of cellulases, mannanases, and xylanases by *Phomopsis*, *Xylaria*, and sterile mycelia strains is interesting. Carroll and Petrini (1981) suggested that such fungi may be latent pathogens or vigorous decomposers after plant death. These taxa may be dormant in the leaves until plant death, and the evidence here indicates that they may then adopt a saprobic lifestyle and degrade dead leaves and possibly wood. *Diaporthe* species (anamorphs of *Phomopsis*) are commonly found on leaves and on wood (Wehmeyer 1933).

*Xylaria* species are also commonly found on degrading wood and many species have the unusual ability amongst ascomycetes to degrade lignins (Urairuj et al. 2003).

The ability of the sterile mycelia to produce cellulases, mannanases, and xylanases may also reflect their role in nature in degrading dead leaves or stems. Guo et al. (1998, 2000) have shown that many endophytic sterile mycelia are also saprobic ascomycetes (e.g., *Astrosphaeriella bakeriana* and *Myelosperma tumidum*) that can be found on decaying plant material. This study indicates that the endophytes within the rainforest seedlings may also have the ability to become saprobes following plant death.

## Acknowledgements

This research was supported by the Thailand Research Fund (BB/17/2539) and NRCT (Thailand-Japan Cooperation Program 2540-2540). K.D. Hyde thanks the Institute of Science and Technology Development of Chiang Mai University for funding his visit to Chiang Mai University.

## References

- Bacon, C.W., and Hill, N.S. 1996. Symptomless grass endophytes: products of coevolutionary symbiosis and their role in the ecological adaptations of grasses. In *Endophytic fungi in grasses and woody plants: systematics, ecology and evolution*. Edited by S.C. Redlin and L.M. Carris. APS Press, St. Paul, Minnesota. pp. 155–178.
- Brown, K.B., Hyde, K.D., and Guest, D.I. 1998. Preliminary studies on endophytic fungal communities of *Musa acuminata* species complex in Hong Kong and Australia. *Fung. Divers.* **1**: 27–51.
- Bussaban, B., Lumyong, S., Lumyong, P., McKenzie, E.H.C., and Hyde, K.D. 2001. Endophytic fungi from *Amomum siamense*. *Can. J. Microbiol.* **47**: 943–948.
- Carroll, G., and Petrini, O. 1981. Patterns of substrate utilization by some fungal endophytes from coniferous foliage. *Mycologia*, **75**: 53–63.
- Downie, B., Hilhorst, H.W.M., and Bewley, J.D. 1994. A new assay for quantifying endo- $\beta$ -mannanase activity using congo red dye. *Phytochemistry*, **36**: 829–835.
- Dreyfuss, M.H., and Chapela, I.H. 1995. Potential of fungi in the discovery of novel, low molecular weight pharmaceuticals. In *The discovery of natural products with therapeutic potential*. Edited by V.P. Gullo. Butterworth-Heinemann, Oxford, U.K. pp. 49–80.
- Fröhlich, J., Hyde, K.D., and Petrini, O. 2000. Endophytic fungi associated with palms. *Mycol. Res.* **104**: 1202–1212.
- Guba, E.F. 1961. Monograph of *Pestalotia* and *Monochaetia*. Harvard University Press, Cambridge, Massachusetts, U.S.A.
- Guo, L.D., Hyde, K.D., and Liew, A.C.Y. 1998. A method to promote sporulation in palm endophytic fungi. *Fung. Divers.* **1**: 109–113.
- Guo, L.D., Hyde, K.D., and Liew, A.C.Y. 2000. Identification of endophytic fungi from *Livistona chinensis* based on morphology and rDNA sequences. *New Phytol.* **147**: 617–630.
- Karim, M.I.A., Tomita, F., Koshimizu, K., Ali, A.M., Hassan, Z., Ariff, A., Radu, S., Mohtar, M., Shaari, K., Ohta, Y., Kawasu, K., Suto, M., and Akita, M. 1997. Screening and isolation of bioactive compounds from microorganisms and higher plants. In *Conservation and sustainable use of biodiversity*. Edited by A.H. Zakri and K. Mat-Salleh. Research Management Unit, UKM, Bangi, Malaysia. pp. 57–70.
- Lumyong, S., Thongantha, S., Lumyong, P., and Tomita, F. 2000. Endophytic fungi from 13 bamboo species in Thailand. *Biotechnology for Sustainable Utilization of Biological Resources in the Tropics*, **14**: 96–101.
- Peláez, F., Collado, J., Arenal, F., Basilio, A., Cabello, A., Díez Matas, M.T., García, J.B., del val González, A., González, V., Gorrochategui, J., Hernández, P., Martín, I., Platas, G., and Vicente, F. 1998. Endophytic fungi from plants living on gypsum soils as a source of secondary metabolites with antimicrobial activity. *Mycol. Res.* **102**: 755–761.
- Petrini, O., and Fisher, P.J. 1988. A comparative study of fungal endophytes in xylem and whole stem of *Pinus sylvestris* and *Fagus sylvatica*. *Trans. Br. Mycol. Soc.* **91**: 233–238.
- Petrini, O., Stone, J., and Carroll, F.E. 1982. Endophytic fungi in evergreen shrubs in western Oregon: a preliminary study. *Can. J. Bot.* **60**: 789–790.
- Photita, W., Lumyong, S., Lumyong, P., and Hyde, K.D. 2001. Endophytic fungi of wild banana (*Musa acuminata*) at Doi Suthep-Pui National Park, Thailand. *Mycol. Res.* **105**: 567–574.
- Pointing, S.B. 1999. Qualitative methods for determination of lignocellulolytic enzyme production by tropical fungi. *Fung. Divers.* **2**: 17–33.
- Rodrigues, K.F. 1994. The foliar fungal endophytes of the Amazonian palm *Euterpe oleracea*. *Mycologia*, **86**: 376–385.
- Rodrigues, K.F., and Petrini, O. 1997. Biodiversity of endophytic fungi in tropical regions. In *Biodiversity of tropical microfungi*. Edited by K.D. Hyde. Hong Kong University Press, Hong Kong.
- Rodrigues, K.F., and Samuels, G.J. 1990. Preliminary study of endophytic fungi in a tropical palm. *Mycol. Res.* **94**: 827–830.
- Romero, A., Carrión, G., and Rico-Gray, V. 2001. Fungal latent pathogens and endophytes from leaves of *Parthenium hysterophorus* (Asteraceae). *Fung. Divers.* **7**: 81–87.
- Strobel, G.A., Hess, W.M., Ford, E.J., Sidhu, R.S., and Yang, X. 1996a. Taxol from fungal endophytes and the issue of biodiversity. *J. Ind. Microbiol.* **17**: 417–423.
- Strobel, G.A., Yang, X., Sears, J., Kramer, R., Sidhu, R.S., and Hess, W.M. 1996b. Taxol from *Pestalotiopsis microspora*, an endophytic fungus of *Taxus wallichiana*. *Microbiology (Reading, U.K.)*, **142**: 435–440.
- Suryanarayanan, T.S., Senthilarasu, G., and Muruganandam, V. 2000. Endophytic fungi from *Cuscuta reflexa* and its host plants. *Fung. Divers.* **4**: 117–123.
- Taylor, J.E., Hyde, K.D., and Jones, E.B.G. 1999. Endophytic fungi associated with the temperate palm *Trachycarpus fortunei* both within and outside of its natural geographic range. *New Phytol.* **142**: 335–346.
- Umali, T.E., Quimio, T.H., and Hyde, K.D. 1999. Endophytic fungi in leaves of *Bambusa tuldoidea*. *Fung. Sci.* **14**: 11–18.
- Urairuj, C., Khanongnuch, C., and Lumyong, S. 2003. Ligninolytic enzymes from new sources: tropical endophytic *Xylaria* species. *Fung. Divers.* **13**: In press.
- Wehmeyer, L.E. 1933. The genus *Diaporthe* Nitschke and its segregates. *Univ. Mich. Stud.* **9**: 1–349.