



<b>Title</b>	<b>The dynamics of inter-Sertoli (SC) tight junctions (TJ) are regulated by transforming growth factor-beta 3 (TGF-beta 3) via the p38 mitogen-activated protein (MAP) kinase signaling pathway</b>
<b>Author(s)</b>	<b>Lui, WY; Lee, WM; Cheng, CY</b>
<b>Citation</b>	<b>The 41st Annual Meeting of the American Society for Cell Biology, Washington, DC., 8-12 December 2001. In Molecular Biology of the Cell, 2001, v. 12 suppl., p. 117A-118A</b>
<b>Issued Date</b>	<b>2001</b>
<b>URL</b>	<b><a href="http://hdl.handle.net/10722/47059">http://hdl.handle.net/10722/47059</a></b>
<b>Rights</b>	<b>Molecular Biology of the Cell. Copyright © American Society for Cell Biology.</b>

Conclusion: The intra-acrosomal alkalinization took place during incubation conducive for sperm capacitation and induced to alter the nature of acrosomal contents to be ready for release at acrosome exocytosis.

635

#### A 32kDa Protein (IAM 32) is the Major Integral Protein of the Sperm Inner Acrosomal Membrane

Alma Rosa Barajas-Espinosa, Yang Yu, Richard Oko, Anatomy and Cell Biology, Queen's University, Queen's University, Botterell Hall, Kingston, ON K7L-3N6 Canada

The inner acrosome membrane (IAM) participates in binding to the egg's zona pellucida and may also be involved in subsequent zona-penetration, yet its composition remains unknown. Our objective, therefore, was to obtain information on the integral protein constituents of the IAM. For this purpose we utilized sonicated and isolated bull sperm heads (SSPH) whose plasmalemma and acrosomal contents were missing but whose IAM remained bound to the SDS-insoluble perinuclear theca. A Triton X-100 extract of the SSPH fraction revealed a major protein of 32kDa (IAM 32) and a less abundant 38kDa protein (SP 38) which could be eliminated by prior high salt extraction of the SSPH fraction. Anti-serum raised against and specific to bull IAM 32 cross-reacted with co-migrating proteins of rat sperm and immunogold labeled the IAM of sonicated and whole bull and murid sperm, and of acrosome reacted rodent sperm. The amino terminal sequence of the isolated bull IAM 32 protein had no similarity to proteins in the NCBI protein data bases. In conclusion, we have uncovered the major integral membrane protein of the IAM that may contribute to the organization of the IAM matrix which, along with IAM 32, remains peripherally attached to the IAM after the acrosome reaction. (Supported by the National Science and Engineering Research Council of Canada (NSERC)).

636

#### Characterization of Rat Sperm Capacitation *In Vitro* and its Effect on Changes in Lipid and Protein Markers

Joseph A Wamstad<sup>1</sup>, Kenneth P Roberts<sup>2</sup>, Kathy M Ensrud<sup>1</sup>, David W Hamilton<sup>1</sup>, <sup>1</sup>Genetics, Cell Biology and Development, University of Minnesota, 321 Church St SE, Minneapolis, MN 55455, <sup>2</sup>Urologic Surgery, University of Minnesota

Capacitation of rat sperm is not well-defined in the literature. Here we characterize rat sperm capacitation *in vitro* through analysis of tyrosine phosphorylation kinetics, cholesterol removal kinetics (with BSA and methyl- $\beta$ -cyclodextrin [MBCD]) and motility. Using defined capacitation conditions, we also report effects on lipid raft (using GM<sub>1</sub> as a marker) and protein (CRISP-1) changes in the membrane. Quantification of cholesterol removal in cauda epididymal rat sperm capacitated *in vitro* shows that the amount of cholesterol extracted from the membrane increases directly with the concentration of MBCD; cholesterol extraction occurs rapidly, reaching peak levels within 15 to 30 minutes. Following removal of cholesterol with MBCD or BSA, tyrosine phosphorylation of specific proteins increases, as determined by western blot analysis, with the rate dependent on the concentration and type of cholesterol binding agent. Concurrently, treatment with MBCD or BSA results in the initiation of hyperactive motility. Following capacitation, there is an increase in sperm membrane fluidity as measured by changes in the localization of GM<sub>1</sub>, a known marker of lipid rafts. Using cholera toxin to label GM<sub>1</sub> there is a shift from sharp, intense staining in the post-acrosomal sheath and equatorial band region of the head in sperm incubated in non-capacitating conditions to a diffuse, unrestricted staining across the entire head in sperm incubated in capacitating conditions. Western blot analysis and immunocytochemistry show that the amount of CRISP-1 on sperm capacitated with MBCD declines slightly during capacitation, but the majority remains on sperm and does not move within the plane of the membrane. (Research supported by USPH grant HD-11962)

637

#### Peripheral Bound Membrane Proteins are Involved in the Maintenance of Boar Sperm Viability by Oviductal Apical Plasma Membrane Preparations *in vitro*

Roslyn Marie Ann Elliott<sup>1</sup>, Anne Duncan<sup>2</sup>, Paul Frederick Watson<sup>2</sup>, William Vincent Holt<sup>1</sup>, Alireza Fazeli<sup>1</sup>, <sup>1</sup>Institute of Zoology, Regent's Park, London, NW1 4RY United Kingdom, <sup>2</sup>Royal Veterinary College, United Kingdom

We have previously shown that oviductal apical plasma membrane (APM) preparations enhance boar sperm longevity *in vitro*. In the current investigation we studied the nature and location of active factors in APM. Oviductal epithelium was scraped, homogenised and purified to prepare APM vesicles. A sample of APM was heat treated to destroy protein activity. A further sample of APM was incubated at 40°C in 1 M NaCl for 30 minutes, followed by separation of peripheral and integral membrane bound proteins by centrifugation at 100,000g for 1 hour. Washed swim-up boar spermatozoa ( $25 \times 10^6$  sperm/ml) diluted with Tyrode's medium (25  $\mu$ l) were added to 25  $\mu$ l experimental APM with a final protein concentration of 200  $\mu$ g/ml and to medium only (control). Treatments included heat-treated vs non heat-treated APM and peripheral vs integral membrane bound proteins, alongside whole APM. These were incubated at 39°C and 5% CO<sub>2</sub> for 24 hours. Sperm viability was determined with Ethidium Homodimer-1 and SYBR 14. Results were expressed as

percentage of the original viability (viability index $\pm$ SEM). The heat treatment significantly ( $P < 0.05$ ) reduced the viability enhancing ability of APM (59 $\pm$ 5, 78 $\pm$ 9 and 65 $\pm$ 5 for heat-treated, non heat-treated and control respectively,  $n=8$ ). There was a significant increase ( $P < 0.05$ ) in the proportion of viable spermatozoa incubated with the peripheral fraction in comparison to all other treatments (97 $\pm$ 2, 75 $\pm$ 6, 85 $\pm$ 2 and 58 $\pm$ 5 for peripheral, integral, original APM fractions and control respectively,  $n=12$ ). We suggest that active factor(s) responsible for the maintenance of boar sperm viability is/are of protein nature and peripherally bound to the APM of porcine oviductal epithelial cells. *This study was funded by the Department for Environment, Food and Rural Affairs, UK.*

638

#### SP 38: the Major Peripheral Protein of the Sperm Inner Acrosomal Membrane

Yang Yu, Ronald Tovich, Alma Rosa Barajas-Espinosa, Mirabotalib Kazemie, Richard Oko, Anatomy and Cell Biology, Queen's University, Queen's University, Botterell Hall, Kingston, ON K7L-3N6 Canada

The consequence of the acrosome reaction is the exposure of the inner acrosomal membrane (IAM) as the leading surface of the sperm head, which then binds secondarily to the egg's zona pellucida, a prerequisite for zona-penetration. However, the IAM proteins involved have not been identified probably because the protein composition of the IAM was unknown. Our objective, therefore, was to obtain direct information on the peripheral protein constituents of the IAM. For this purpose, we devised a fractionation procedure to isolate apical tips of the rat sperm heads, which consisted solely of the IAM bound to the SDS-insoluble perforatorium. High salt incubation of this tip fraction extracted a major protein of 38 kDa coincident with partial removal of an electron dense layer attached to the IAM. A prominent 38kDa was also salt extractable from sonicated and isolated bull sperm heads that had retained their IAM. Antiserum raised against and specific to this bull protein cross-reacted with the co-migrating 38kDa protein obtained from the rat tips and immunogold labeled the IAM of these tips, of sonicated and whole bull and murid sperm, and of acrosome reacted murid sperm. Cloning and sequencing of the bull 38kDa protein revealed its identity as SP 38 (Dev. Biol. 168:575-583, 1995), an acrosome protein with zona/ZP 2 binding capability whose precise location within the acrosome was unknown. In conclusion, this mammalian sperm protein contributes to an electron dense layer or matrix that is peripherally attached to the IAM after the acrosome reaction. This suggests SP 38 is a strong candidate for secondary zona-binding. (Supported by NSERC).

639

#### Do D-3 Phosphoinositides Signal Actin Polymerization during Ascidian Sperm Activation?

Lamar Blackwell, Dave Bolger, Mohammad Hanizavareh, Emily Zebadua, Robert A Koch, Biological Science, Cal State Fullerton

Sperm activation in the sea squirt *Ascidia ceratodes* is characterized by mitochondrial translocation (MTL), an actin:myosin-dependent movement known to require elevation of both intracellular pH (pHi) and free calcium ion concentration ([Ca<sup>2+</sup>]<sub>i</sub>). Previously, we have shown that myosin activation requires a G protein-mediated pathway involving inositol 1,4,5-trisphosphate-mediated internal Ca release and a PKC-dependent internal alkalization that precedes external Ca entry. Here, we explore signaling elements that are involved in triggering actin polymerization. In MTL assays, the actin polymerization inhibitor latrunculin (10 $\mu$ M) completely blocked high pH artificial sea water (ASW)-induced sperm activation (positive control), and the actin polymerization inducer jasplakinolide (7.4 $\mu$ M) stimulated sperm activation equal to positive controls, an action blocked by latrunculin. Dual labeling with fluorescently tagged phalloidin and DNaseI revealed that filamentous actin was distributed most heavily on the mitochondrion whereas monomeric actin was also found along the length of the tail. Sperm activation appears to increase filamentous actin on the mitochondrion. In MTL assays, the phosphatidylinositol 3-kinase (PI3K) inhibitor LY294002 (50 $\mu$ M) blocked sperm activation induced by pH 9.4 ASW but not that induced by the G protein activator mas7 (3.5 $\mu$ M) or the PKC activator OAG (50 $\mu$ M), agents shown to be part of the myosin activation pathway. Liposomes that incorporated phosphatidylinositol 3,4,5-trisphosphate (PIP<sub>3</sub>) stimulated levels of sperm activation similar to positive controls. Indirect immunofluorescence using anti-profilin antibodies showed profilin to be present on the mitochondrion, providing a possible connection between PI3K-induced PIP<sub>3</sub> production and actin polymerization. (Funded by CSUF University Student Research Initiative grant to MH; NIH R25-GM56820 to RAK for LB, DB & EZ; NIH R15HD36500 to RAK.)

640

#### The Dynamics of Inter-Sertoli (SC) Tight Junctions (TJ) are Regulated by Transforming Growth Factor- $\beta$ 3 (TGF- $\beta$ 3) via the p38 Mitogen-Activated Protein (MAP) Kinase Signaling Pathway

Wing-yei Lui<sup>1</sup>, Will M Lee<sup>2</sup>, C. Yan Cheng<sup>1</sup>, <sup>1</sup>Population Council, 1230 York Avenue, New York, NY 10021, <sup>2</sup>Department of Zoology, University of Hong Kong, Hong Kong

Earlier studies have shown that the inter-Sertoli TJ dynamics are regulated, at least in part, by TGF- $\beta$ 3, possibly via its effects on occludin/claudin-11/ZO-1 (Lui et al., *Endocrinology* 142:1865-1877, 2001). We now report that this TGF- $\beta$ 3 regulatory effect on TJ-functionality is mediated via the MEK/ERK/p38-MAP kinase signal

transduction pathway. Using SCs cultured *in vitro* to allow the assembly of TJs when the TJ-permeability barrier was monitored by the transepithelial electrical resistance (TER) across the SC epithelium, we have examined if the TGF- $\beta$ -induced TJ disruption is mediated via one of the four upstream signal transducers, namely MEKs, Smad2/Smad3, Cdc42/Rac, and Ras. Cellular distribution studies using RT-PCR have shown that both SC and germ cells (GC) express almost similar levels of mRNA encoding for MEK2, Smad2, and small GTPases, such as Cdc42, Rac and N-Ras. A TGF- $\beta$ -induced transient increase in MEK2 expression, but not Smad2, Cdc42/Rac2, or N-Ras, was detected in SC during the assembly of the TJ-barrier. The TGF- $\beta$ -mediated (3 ng/ml) inhibitory effect on the assembly of TJs could be reversed dose-dependently by SB202190 at 0.1 nM-1 mM, a specific p38-MAP kinase inhibitor. We next investigated the protein expression of p-p38-MAP kinase (activated phosphorylated form) *versus* total p38-MAP kinase (nonphosphorylated inactive form) using SC lysates by immunoblottings and specific antibodies against p-p38- and p38-MAP kinase with a chemiluminescence-based detection system. It was found that the presence of TGF- $\beta$  indeed regulated the production of p-p38 MAP kinase protein during TJ assembly. In summary: the TGF- $\beta$ -mediated effects on the inter-Sertoli TJ dynamics and the blood-testis-barrier functionality are regulated via the p38-MAP kinase pathway. [Supported in part by grants from CONRAD (CICCR96-05-A to CYC), and HKRGC (HKU7245/00M to WML/CYC)].

641

### T3-Regulated Expression of a Novel Attachment Factor, PB-cadherin, may be Critical for Development of Neonatal Testicular Stem Cells

Ji Wu, William F. Jester, Jr., Joanne M. Orth, Dept. of Anatomy & Cell Biology, Temple University School of Medicine, 3400 N. Broad St, Philadelphia, PA 19140

In the rodent testis, contact-mediated interactions between gonocytes, or neonatal stem cells, and Sertoli cells are critical for development. Previously, we showed that Thyroid Hormone (T3) regulates expression in neonates of at least one Sertoli cell-gonocyte attachment factor, NCAM. We subsequently used a rat cDNA microarray and detected expression of another factor, short-type PB-cadherin (STPB-C) in neonatal Sertoli cell-gonocyte co-cultures. PB-cadherin is a novel cadherin mainly expressed in pituitary gland and brain of adults which is involved in development by regulating Ca<sup>2+</sup>-dependent cell-cell adhesion. Therefore, our present aims were (1) to explore expression of STPB-C *in vivo*, (2) to localize STPB-C mRNA in co-cultures with *in situ* hybridization, and (3) to determine if expression of STPB-C is regulated by T3. RT-PCR was used to generate cDNA for STPB-C from total RNA isolated from cocultures, cDNA was cloned into pPCR-Script™ Amp SK(+) cloning vector, and plasmid DNA was isolated and sequenced to confirm the fidelity of the STPB-C cDNA portion of the plasmid. In subsequent Northern analysis of testicular RNA, expression of STPB-C was strong on day 1, then diminished appreciably by day 3, became barely detectable by day 15, and disappeared in testes of adults. When neonatal cocultures were treated with T3 (10 nM, 24 hr) or vehicle, STPB-C mRNA was strongly expressed by neonatal stem cells and weakly by Sertoli cells *in situ*, while Northern analysis indicated that expression of STPB-C was down-regulated by T3 *in vitro*. Thus, regulation of PB-cadherin by T3 during the early neonatal period may be critical in development of the stem cell population from which all maturing germ cells subsequently arise. (Supported by NIH HD-15563.)

642

### Regulation of Catenins in The Rat Epididymis

Daniel G Cyr, Sophie DeBellefeuille, Human Health Research Centre, INRS-institut Armand Frappier, 245 Hymus boulevard, Pointe Claire, PQ H9R1G6 Canada

Adhering junctions are essential for the formation and regulation of tight junctions. In the epididymis, tight junctions between adjacent principal cells form the blood-epididymal barrier which creates a specific environment within the lumen of the epididymis necessary for sperm maturation. Adhering junctions in the epididymis are composed of a transmembrane protein, cadherin, and catenins (alpha, beta and P120). The objective of this study was to determine the effects of testis and testicular androgens on the immunolocalization of catenins (alpha, beta and P120) in the epididymis. In intact control adult rat epididymis, each of the three catenins were localized along the lateral plasma membranes of adjacent principal cells as well as between principal and both clear and basal cells. Twenty one days following orchidectomy there was a marked increase in the cytosolic staining of both alpha- and beta-catenin, particularly in the corpus and cauda epididymidis, suggesting a loss in the integrity of the adhering junctions. Interestingly, immunostaining for P120 appeared to be unaltered by orchidectomy. In orchidectomized rats that had been given testosterone implants at the time of orchidectomy, the immunolocalization of alpha- and beta-catenin was maintained along the lateral plasma membrane of epididymal principal cells. These data suggest that androgens can maintain the integrity of adhering junctions in the epididymis and may represent a mechanism by which androgens can regulate tight junctions and the blood-epididymal barrier in adult rats.

643

### Regulation of Sertoli(SC)-Germ(GC) Cell Anchoring Junction (AJ) Dynamics by RhoB GTPase

Wing-yei Lui, C. Yan Cheng, Population Council, 1230 York Avenue, New York, NY 10021

During spermatogenesis, there are extensive AJ restructuring, however, the mechanistic pathway that regulates AJ dynamics is not known. Rho GTPases have been implicated in the actin organization and cytoskeletal control. For instance, Rho GTPases are known to regulate AJ functionality by redistributing cadherins during AJ assembly. Using RT-PCR, both SC and GC were found to express RhoB. Moreover, the assembly of SC-GC AJs, but not SC TJs, was associated with a transient induction of RhoB expression. These changes were confirmed with a monospecific RhoB antibody for immunoblottings using cell lysates and a chemiluminescence-based detection system. Disruption of AJs in SC-GC cocultures by hypotonic treatment also induced a surge in RhoB expression, which became visible within 5-min. Moreover, when SC-GC AJs *in vivo* was disrupted by treatment of rats with a single dose of 1-(2,4-dichlorobenzyl)-indazole-3-carbohydrazide (DCIC, 300 mg/kg b.w. by gavage), a surge in RhoB expression by ~4-fold was also detected within 1-hr. This is long before the depletion of germ cells from epithelium become visible, which required ~10-day. When GCs were added to the SC epithelium (GC:SC, 1:1), which had been cultured for 5-day at  $0.5 \times 10^6$  cells/cm<sup>2</sup> on Matrigel-coated dishes, and cultured for 2-day to allow the assembly of AJs; inclusion of DCIC at 250 ng/ml to the cocultures also induced a surge in RhoB expression ~5 min-1 hr. These results thus illustrate that RhoB is activated during the assembly of AJ, and prior to the actual disruption of AJs. These changes are possibly needed because RhoB regulates redistribution of AJ-proteins, such as cadherins. In summary: (i) RhoB is an important signaling molecule that regulates AJ dynamics in the testis; and (ii) the DCIC-induced GC loss from the epithelium is mediated via the Rho GTPase signaling pathway

644

### The dynamics of Sertoli (SC)-germ cell (GC) anchoring junctions (AJs) are regulated by E-cadherin, N-cadherin and Src

Nikki Pui Yue Lee<sup>1</sup>, Will M Lee<sup>2</sup>, C Yan Cheng<sup>1</sup>, <sup>1</sup>Population Council, 1230 York Avenue, New York, NY 10021, <sup>2</sup>Department of Zoology, The University of Hong Kong, Hong Kong

Earlier studies have shown that the assembly of AJs between SC is associated with a transient induction in N-cadherin/ $\beta$ -catenin expression. Moreover, the junction dynamics in the testis are regulated by the interplay of phosphatases and kinases that regulates the intracellular phosphoprotein content. To expand these earlier studies, we have assessed the role of E-cadherin, N-cadherin and Src (an AJ-associated signaling molecule) in AJ dynamics. When SC were cultured at  $0.5 \times 10^6$  cells/cm<sup>2</sup> or SC-GC were cocultured (SC:GC ratio at 1:1, SC at  $0.5 \times 10^6$  cells/cm<sup>2</sup>) *in vitro* on Matrigel-coated dishes to allow the assembly of AJs, it was associated with a transient induction in the expression of E-cadherin, N-cadherin and Src. Similar changes were detected when cell lysates were prepared from these samples for immunoblottings using the corresponding antibodies and a chemiluminescence-based detection system. Cellular distribution studies by semi-quantitative RT-PCR revealed that while SC expressed almost twice as much N-cadherin when compared to GC, GC expressed almost 3-times as much E-cadherin as SC, suggesting GC play an important role contributing to the AJ-associated protein pool in the testis. Also, the expression of SC E-cadherin and N-cadherin were stimulated by testosterone and DHT by ~ 3-10 fold at  $10^{-9}$  V  $10^{-7}$  M, suggesting androgens may also regulate AJ functionality via their effects on AJ-associated proteins. Work is now in progress to assess whether there are changes in the phosphorylation status of these proteins during AJ assembly. In summary, these results demonstrate that the dynamics of AJs are regulated, at least in part, by N-cadherin, E-cadherin, Src and androgens. [Supported in part by grants from the CONRAD Program (CICCR96-05-A to CYC) and Hong Kong Research Grant Council (HKU7245/00M to WML/CYC)]

645

### The role of inducible nitric oxide synthase (iNOS) and endothelial nitric oxide synthase (eNOS) in the dynamics of tight (TJ) and anchoring junctions (AJ)

Nikki Pui Yue Lee, C Yan Cheng, Population Council, 1230 York Avenue, New York, NY 10021

NOS catalyzes the enzymatic oxidation of L-arginine to nitric oxide (NO), which plays a critical role in a variety of bioregulatory functions including junction assembly. Also iNOS and eNOS are present in the testis and are implicated in the regulation of spermatogenesis. However, their role(s) in junction dynamics in the testis has not been explored. When SC were cultured at  $0.5 \times 10^6$  cells/cm<sup>2</sup> on Matrigel-coated dishes for up to 7-days, there was a transient induction in eNOS expression, but not iNOS, coinciding with the assembly of inter-Sertoli TJ-permeability barrier. When SCs were cultured at  $0.5 \times 10^6$  cells/cm<sup>2</sup> on Matrigel-coated dishes for 5-days to allow the assembly of both TJs and AJs, freshly isolated GCs were then added onto the SC epithelium at a SC:GC ratio of 1:1 to initiate SC-GC AJ assembly; there was an increase in iNOS expression, but not eNOS, coinciding with the assembly of SC-GC AJs. To further explore the involvement of iNOS in SC-GC AJ dynamics, an *in vivo* model was used in which 1-(2,4-dichlorobenzyl)-indazole-3-carbohydrazide (DCIC) was used to induce GC depletion from the epithelium. A significant increase in iNOS, but not eNOS,