



Title	Transgenic plant-derived siRNAs can suppress propagation of influenza virus in mammalian cells
Author(s)	Zhou, Y; Chan, JH; Chan, AY; Chak, RKF; Wong, EYL; Chye, ML; Peiris, JSM; Poon, LLM; Lam, E
Citation	Febs Letters, 2004, v. 577 n. 3, p. 345-350
Issued Date	2004
URL	http://hdl.handle.net/10722/48507
Rights	Creative Commons: Attribution 3.0 Hong Kong License

Transgenic plant-derived siRNAs can suppress propagation of influenza virus in mammalian cells

Yuanxiang Zhou^{a,*}, Jack Y. Chan^b, Annie Y. Chan^b, Regina K.F. Chak^a, Elaine Y.L. Wong^a, Mee-Len Chye^a, Joseph S.M. Peiris^b, Leo L.M. Poon^b, Eric Lam^{a,1}

^aDepartment of Botany, University of Hong Kong, Pokfulam Road, Hong Kong Special Administrative Region, PR China

^bDepartment of Microbiology, Queen Mary Hospital, University of Hong Kong, Pokfulam Road, Hong Kong Special Administrative Region, PR China

Received 6 August 2004; revised 5 October 2004; accepted 11 October 2004

Available online

Edited by Hans-Dieter Klenk

Abstract As an example of the cost-effective large-scale generation of small-interfering RNA (siRNAs), we have created transgenic tobacco plants that produce siRNAs targeted to the mRNA of the non-structural protein NS1 from the influenza A virus subtype H1N1. We have investigated if these siRNAs, specifically targeted to the 5'-portion of the NS1 transcripts (*5mNS1*), would suppress viral propagation in mammalian cells. Agroinfiltration of transgenic tobacco with an *Agrobacterium* strain harboring a *5mNS1*-expressing binary vector caused a reduction in *5mNS1* transcripts in the siRNA-accumulating transgenic plants. Further, H1N1 infection of siRNA-transfected mammalian cells resulted in significant suppression of viral replication. These results demonstrate that plant-derived siRNAs can inhibit viral propagation through RNA interference and could potentially be applied in control of viral-borne diseases. © 2004 Published by Elsevier B.V. on behalf of the Federation of European Biochemical Societies.

Keywords: Transgenic plant; Small-interfering RNA; Gene encoding the non-structural protein NS1; Influenza virus; Mammalian cells; Anti-viral

1. Introduction

RNA interference (RNAi) is an ancient and evolutionarily conserved activity in eukaryotes. It results in RNA-mediated RNA degradation in a sequence-specific manner. Originally described in plants as a concerted inactivation of host genes and transgenes transcribing the same or similar sequences [1], it has been confirmed to occur in many different organisms. Examples include quelling in *Neurospora crassa* [2], and RNAi in *Caenorhabditis elegans* [3], *Drosophila* [4] and mammals [5]. In all these cases, RNAi is achieved through several closely coordinated steps: (1) an endonuclease Dicer with RNase III activity cleaves the dsRNA into 21–23 bp small interfering RNAs (siRNAs); (2) the siRNAs interact with a multicom-

ponent nuclease to form an RNA-induced silencing complex (RISC); (3) the siRNA in the RISC directs the complex to the target RNA through sequence complementarity; (4) RNA polymerization begins from the siRNA to form dsRNA; and (5) the dsRNA is cleaved into siRNAs [6,7]. The resulting siRNAs would then initiate another round of RNA cleavage.

Studies using synthetic [8], in vitro transcribed [9,10] and in vivo transcribed [11,12] siRNAs, as well as viral-mediated siRNA delivery [13], have demonstrated that well-designed siRNAs can effectively suppress target gene expression. Hence, RNAi technology could eventually be applied in the therapeutics of human and animal viral diseases of which the molecular components, e.g., viral sequences, are known, and in the case of infectious diseases, of which the relevant pathogens have been identified. In plants, viral-resistance has already been achieved through a plant RNAi pathway termed post-transcriptional gene silencing (PTGS) [14].

Although some understanding on siRNA inhibition of viral propagation [8,11] has been achieved, the local folding of the target RNAs that reduces siRNA accessibility within a transcript [15] makes it necessary to test out many different siRNAs before optimal transcript degradation can be attained [8,16]. For example [8], 20 siRNA oligos were screened before identification of one that could satisfactorily suppress replication of the influenza virus in mammalian cells. Also, siRNA-mediated gene suppression in mammals requires the dsRNA to be smaller than 30 bp to ensure specificity [17], as long dsRNA can provoke non-specific degradation of RNA transcripts and a general shutdown of protein translation [18]. Therefore, it is impossible to transfect mammalian cells with long-dsRNA-producing constructs essential for making multiple siRNAs. The high cost in RNA oligo synthesis and the toxic effects of long dsRNA in mammalian cells could be ameliorated by cost-effective techniques in simultaneous large-quantity production of different siRNAs to achieve a satisfactory level for RNAi-mediated gene suppression.

Here, we demonstrate that tobacco (*Nicotiana tabacum*) can be engineered by *Agrobacterium*-mediated transformation to produce siRNAs targeting the mRNA for the non-structural NS1 protein of the influenza virus A/WSN/33, subtype H1N1. The transgenic plants could effectively accumulate siRNAs that specifically target gene encoding the non-structural protein NS1 (*NS1*) transcripts. Transfection of mammalian cells with plant-derived siRNAs followed by infection of the influenza virus revealed significant reduction in viral propagation.

* Corresponding author. Fax: +852-28583477.
E-mail address: seanzhou@hkucc.hku.hk (Y. Zhou).

¹ Present address: Biotech Center, Rutgers University, 59 Dudley Road, New Brunswick, NJ 08901-8520, USA.
Abbreviations: NS1, gene encoding the non-structural protein NS1; MDCK, Madin–Darby canine kidney; siRNA, small-interfering RNA; RNAi, RNA interference; PTGS, post-transcriptional gene silencing; *EYFP*, gene encoding the enhanced yellow fluorescent protein

90 Our data demonstrate that plants can be used as an econom-
91 ical and sustainable source for large-scale production of di-
92 versified siRNAs.

93 **2. Materials and methods**

94 **2.1. Construction of hairpin RNA vector and generation of transgenic**
95 **tobacco plants**

96 A 0.4-kb fragment representing the 5'-portion of the *NSI* mRNA
97 (*5mNSI*) from the influenza virus strain A/WSN/33 subtype H1N1
98 (Fig. 1A and B) was amplified by reverse-transcriptase polymerase
99 chain reaction (RT-PCR) using forward primer 5'-
100 ggcgccgcgggatccatggacccaacactgtg-3' with *NotI* (in italics) and
101 *BamHI* (in bold) sites incorporated at its 5'-end, and reverse primer 5'-
102 caactagtatttcgcttcagtatga-3' with an added *SpeI* site (in italics). The
103 underlined nucleotides represent *NSI* sequences. The PCR product
104 was initially cloned in pGEM-T Easy vector (Promega) for verification
105 of DNA sequence. Subsequently, the pGEM-T Easy derivative was
106 digested with either *BamHI* and *SpeI* or *NotI* and *SpeI*. The 0.4-kb
107 *BamHI-SpeI 5mNSI* fragment was cloned into corresponding sites in a
108 pBluescript SKII(-) derivative that contains the *Arabidopsis TGA1*
109 intron [19] inserted at its *SpeI-XbaI* site. Next, the 0.4-kb *NotI-SpeI*
110 *5mNSI* fragment from the pGEM-T Easy derivative was cloned in the
111 *NotI-XbaI* site of the pBluescript SK(II)(-) derivative containing the
112 DNA fusion of "sense *5mNSI-TGA1* intron", to generate a dsRNA
113 cassette "sense *5mNSI-TGA1* intron-antisense *5mNSI*". This cassette
114 was then released by *NotI* and *BamHI* digestion, and, with the help of
115 a *NotI/XbaI* adaptor (upper strand, 5'-GGCCGAGTTGTTA-3'; lower
116 strand, 5'-CTAGTAACAAC-3'), was cloned in the *BamHI-XbaI*
117 site between the *CaMV* 35S promoter and the *nos* terminator, in an-
118 other pBluescript SKII(-) derivative. The resulting vector therefore
119 contains a cassette of "35S-s *5mNSI-TGA1* intron-as *5mNSI nos*"
120 (Fig. 1C). This cassette was further digested with *NotI* and *KpnI*, and
121 was cloned into corresponding sites within the T-DNA in a pBI101
122 backbone plasmid derivative (Clontech, Palo Alto, USA). The binary
123 vector was then mobilized into *Agrobacterium tumefaciens* strain
124 GV3101/MP90 for transformation of tobacco cultivar Samsun NN by
125 the leaf-disk procedure [20].

126 **2.2. siRNA detection**

127 Total RNA samples were extracted from tobacco leaves using
128 TRIzol (Invitrogen). Twenty micrograms of total RNA was separated
129 on a 15% polyacrylamide gel containing 7 M urea and was electroblot-
130 tted onto a nitrocellulose membrane (GeneScreen Plus®, PerkinEl-
131 mer Life Sciences, Inc.). The blot was then hybridized overnight at 42

°C to [³²P]UTP-labeled *5mNSI* riboprobes generated using the Ri-
boprobe® in vitro Transcription Systems (Promega), in a solution of
50% (v/v) formamide, 250 mM NaCl, 7% SDS and 125 mM phosphate
buffer, pH 7.0. After hybridization, the blot was washed twice with 2×
SSC plus 0.5% SDS and was then analyzed using a phospho-imager.
The volumes of the synthetic siRNA and of the siRNA from transgenic
plants were measured using an ImageQuant software (Molecular Dy-
namics), and the amount of siRNA in the plant RNA sample was
calculated based on its volume relative to that of synthetic, known
amount of RNA oligos.

2.3. Transient expression assay by agroinfiltration

Agrobacterium cells containing the *5mNSI*-expressing binary vector
and those containing an *EYFP-T2m* (*EYFP*, gene encoding the en-
hanced yellow fluorescent protein) expressing binary vector [21] were
inoculated in an induction solution containing 1 g/l NH₄Cl, 0.3 g/l
MgSO₄·7H₂O, 0.15 g/l KCl, 0.01 g/l CaCl₂, 0.0025 g/l FeSO₄·7H₂O, 2
mM phosphate, 1% glucose, 20 mM 2-(*N*-morpholino)ethanesulfonic
acid (MES, pH 5.5), 100 μM acetosyringone, 50 μg/ml kanamycin and
50 μg/ml gentamycin. The *EYFP-T2m* contains *EYFP* fused in-frame
to a mutant version of the *Arabidopsis TGA2* gene (*T2m*) and is used as
an expression reference after agroinfiltration. Following overnight
culture at 28 °C, the cells were collected by centrifugation at 3000 × *g*
for 15 min, and then resuspended in an infiltration solution containing
10 mM MES (pH 5.5), 10 mM MgSO₄ and 100 μM acetosyringone.
The resuspended *Agrobacterium* cells were adjusted to an OD₆₀₀ of 0.8
with the same solution before infiltration of tobacco leaves using a 1 ml
syringe. After two days, total RNA was extracted from the infiltrated
leaf areas for Northern blot analysis.

2.4. Northern blot analysis

Five micrograms of total RNA, extracted from the agroinfiltrated
and non-infiltrated leaf areas, were separated on a 1.2% agarose gel,
blotted with 20× SSC onto a nitrocellulose membrane, and hybridized
to [³²P]dCTP-labeled DNA probes generated from *5mNSI* and *EYFP*
DNA fragments using a Rediprime™ II Random Prime Labelling
System (Amersham, UK). Hybridization was performed at 65 °C
overnight in a buffer containing 250 mM NaCl, 7% SDS and 125 mM
phosphate, pH 7.0. After hybridization, the blot was washed twice at
room temperature in 2× SSC plus 0.5% SDS, then at 65 °C for 15 min
in 0.2× SSC plus 0.1% SDS. The blot was analyzed using a phospho-
imager.

2.5. Transfection of mammalian cells followed by infection with
influenza virus

Confluent Madin-Darby canine kidney (MDCK) cells grown in a T-
175 flask were washed twice with phosphate-buffered saline (PBS) and
trypsinized for 10 min in 10 ml trypsin solution at 37 °C. After ter-




Fig. 1. The *5mNSI* sequence and the hairpin RNA construct used in producing *5mNSI* siRNAs in tobacco. (A) Schematic representation of *NSI* mRNA. The cap and poly(A) tail structures are shown, and location of the 0.4-kb *5mNSI* fragment beginning from the first codon (atg) is indicated. (B) cDNA sequence of the *5mNSI*, with the sequence of the synthetic siRNA NS-128 used by Ge et al. [8] underlined. (C) Diagram showing RNAi cassette in a binary vector. The sense (S) and antisense (AS) *5mNSI* fragments are separated by the *Arabidopsis TGA1* intron (I-*TGA1*), and are under the control of the *CaMV* 35S promoter.

132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176

177 mination of trypsinization with 20 ml PBS, the cells were collected by
178 centrifugation for 5 min at 15 000 rpm, and were washed twice in 30 ml
179 cold PBS, followed by one wash in 30 ml cold RPMI1640 medium
180 (Gibco), before resuspension in cold RPMI1640 to a density of 1×10^7
181 cells/ml. Subsequently, 500 μ l of resuspended cells was transferred into
182 a 0.4 cm pre-chilled cuvette, and was mixed with 10 μ l water, 10 μ l
183 water with 42 ng NS-128, 10 μ l wild-type RNA sample or 10 μ l RNA
184 sample containing 42 ng siRNAs from transgenic plant. Equal
185 amounts of total RNA from wild-type or transgenic plants were used.
186 The cuvette was kept on ice for 10 min, before electroporation at 0.4
187 kV and 960 μ F using a gene pulser system (Bio-Rad). Cells were then
188 transferred into 5.6 ml of pre-warmed MDCK medium (MEM, 10%
189 cow serum, 1% penicillin and 1% streptomycin). Three milliliters was
190 transferred into a 6-well plate and incubated at 37 °C for 24 h before
191 infection with the influenza virus.

192 2.6. Virus infection and hemagglutination (HA) titer test

193 Twenty-four hours after transfection, cells in each well were washed
194 twice with PBS, and 300 μ l diluted influenza virus strain A/WSN/33
195 (MOI=0.001 in PBS) was added into the well. After shaking the
196 mixture for 1 h, the viruses in the supernatant were discarded, and 2 ml
197 infection medium [0.5 μ g/ml TPCK-trypsin (Sigma), 0.5% FCS (Gibco),
198 and 1% PS with MEM (Gibco)] was added into the well. The cells
199 were then incubated at 37 °C. Supernatants were collected at different
200 post-infection time points for the HA titer test as described [8].

201 3. Results and discussion

202 Influenza A viruses are medically important viral pathogens
203 that cause significant mortality and morbidity throughout the
204 world. Their easy transmission, antigenic shift and drift have
205 made current methodology of vaccination and therapy limited
206 in efficacy [22]. Inhibitors of the anti-M2 ion channel and
207 neuraminidase are common drugs for influenza, but both have
208 their drawbacks. The anti-M2 ion channel inhibitors (e.g.,
209 amantadine) induce viruses to develop drug-resistant muta-
210 tions, while the neuraminidase inhibitors (e.g., Tamiflu),
211 though very potent, are effective only at early disease onset. To
212 investigate if plant-derived siRNAs against the influenza virus
213 could inhibit viral replication, we selected a 0.4-kb fragment
214 representing the 5'-portion of the *NS1* gene in strain A/WSN/
215 33, subtype H1N1. The NS virion RNA (vRNA) consists of
216 about 890 nucleotides and encodes two non-structural pro-
217 teins, NS1 and NS2. The sequence of this vRNA is highly
218 conserved among different subtypes of influenza viruses [23].
219 The NS1 protein has not only been proposed to regulate viral
220 replication cycle, splicing and translation of mRNAs [24], but
221 also been shown to have inhibitory effect on cellular mRNA
222 maturation and cellular anti-viral response [25]. Thus, the NS1
223 gene plays an important role in virus replication and virus-
224 host interactions. The chosen 0.4-kb fragment was amplified
225 by PCR, and then sequentially cloned in sense and antisense
226 orientations, on either side of the *Arabidopsis TGAI* intron.
227 The resulting cassette of "sense-intron-antisense" was ex-
228 pressed from the *CaMV* 35S promoter in a binary vector
229 (Fig. 1C). Hence, transgenic plants obtained in *Agrobacterium*-
230 mediated plant transformation from this binary vector should
231 produce hairpin dsRNA, which would subsequently be pro-
232 cesses into siRNAs by the PTGS machinery.

233 Reports have shown that "sense-antisense" cassettes can be
234 transcribed to produce siRNAs after transfection of host cells
235 [11,26,27]. To investigate if the construct generated in this
236 study (Fig. 1C) could produce siRNA in transgenic tobacco,
237 RNAs from leaves of primary transformants was separated on
238 a gel of 15% polyacrylamide and 7 M urea, blotted onto ni-




Fig. 2. Accumulation of the *5mNSI* siRNA in selected primary transformants and in wild-type tobacco (WT). (A) Twenty micrograms of total RNA from leaves of transgenic tobacco was separated on a 15% polyacrylamide gel containing 7 M urea, blotted and hybridized to [32 P]UTP-labeled *5mNSI* riboprobes. (B) Normalization of RNA loading was based on the separation of 6 μ g of total RNA on a 1.2% agarose gel.

trocellulose membrane and hybridized to [32 P]UTP-labeled
5mNSI riboprobes. Of 21 independent transformants
screened, 13 showed obvious siRNA production. The levels of
siRNA accumulation in different lines varied, some produced
obvious signals after an overnight exposure using a phospho-
imager, while others barely yielded visible signals (data not
shown). The siRNA signals in selected transgenic lines are
shown in Fig. 2. Transgenic lines 1 and 2 had apparent accu-
mulation of *5mNSI* siRNAs, while lines 8 and 9 produced
much lower levels of the same siRNAs. In line 10, the siRNAs
were barely detectable.

A variation in siRNA levels may be due to several reasons.
First, T-DNA location in the genome could affect expression.
In *Agrobacterium*-mediated plant transformation, T-DNA is
transferred from the bacterium to the eukaryotic host cell and
further integrated into the host genome [28]. If the transgene
were inserted in the genome where active transcription occurs,
the transgene would be active. Otherwise, it would be less
active or even silent. Second, the copy number of the transgene
may be a contributing factor in expression levels, although in
some cases, transgene activity may not be directly proportional
to its copy number due to co-suppression. Third, methylation
of transgene may occur, especially at or near promoter if it is
considered foreign. As a safeguard, the host generally has a
mechanism to methylate and inactivate the transgene. This has
been reported with foreign DNA expressing dsRNA in PTGS
[29,30] and is supported by a requirement of DNA methylase
in initiating RNA-dependent DNA methylation [31].

As revealed by an increasing number of reports, siRNA is
the hallmark in triggering RNAi. Therefore, the accumulated
5mNSI siRNAs in the transgenic plants should initiate deg-
radation of *NS1* transcripts or endogenous tobacco transcripts
with sequences complementary to *5mNSI*. A BLAST analysis
was performed with *5mNSI* as query sequence for such com-
plementation in transcripts of tobacco or species evolutionarily
close to tobacco, but no match was identified. Northern blot
analysis of tobacco total RNA with the *5mNSI* probe also did
not yield any obvious bands. Therefore, *5mNSI* does not seem
to share homology to any tobacco transcripts and would not
cause unintended degradation of RNA transcribed from en-
dogenous genes. This is consistent with the fact that no ab-
normal phenotypes were observed in all the transgenic lines

281 (data not shown). To test if the plant-derived *5mNSI* siRNAs
 282 were functional in degrading *NSI* transcripts specifically,
 283 *5mNSI*- and *EYFP-T2m*-expressing binary vectors were in-
 284 troduced into *Agrobacterium* cells which were used to co-in-
 285 filtrate leaves of wild-type tobacco and those of transgenic
 286 tobacco lines expressing *5mNSI* siRNAs. As shown in Fig. 3B,
 287 all the three transgenic lines 1, 9 and 19, representing high,
 288 middle and low accumulation of *5mNSI* siRNA, respectively,
 289 had reduced *5mNSI* RNA levels, indicating that plant-derived
 290 *5mNSI* siRNAs indeed triggered PTGS of *NSI* in vivo. A
 291 negative correlation was observed between the levels of *5mNSI*
 292 siRNAs and *5mNSI* transcripts in infiltrated tobacco leaves.
 293 To obtain a percentage of the *5mNSI* transcript level in the

294 transgenic lines relative to that of wild-type, volumes of each
 295 *5mNSI* band and of the reference *EYFP* band were determined
 296 using the ImageQuant software, and percentage was calculated
 297 using the formula described in Fig. 3 legend. In transgenic
 298 tobacco line 1, which had the highest level of *5mNSI* siRNA
 299 accumulation, the percentage was only 0.4%, demonstrating
 300 high efficiency of this line in *5mNSI*-specific RNA degradation
 301 (Fig. 3C).

302 In RNAi studies, synthetic or in vitro expressed siRNAs
 303 have been used in transfection of target cells [26], and injection
 304 of worms [32] and animals [33], for evaluation of siRNA effi-
 305 cacy. To test if the *5mNSI* siRNAs produced in transgenic
 306 tobacco could be potentially used in suppressing viral propa-




Fig. 3. Suppression of *5mNSI* transcript accumulation in siRNA-expressing lines. Wild-type tobacco and transgenic plants expressing different levels of *5mNSI* siRNAs were co-infiltrated with two binary vectors separately expressing *5mNSI* and a fusion fragment of *EYFP T2m*. After two days, leaf samples were collected for RNA analysis by Northern blot analysis. (A) Part of the T-DNA in the two binary vectors. (B) Northern blot analysis showing levels of *EYFP-T2m* and *5mNSI* transcripts in the different infiltrated samples. The 28S rRNA was stained with ethidium bromide. The *5mNSI* levels are lowered in transgenic plants when compared to levels in WT. (C) The *5mNSI* transcript level, as a percentage of the wild-type, was calculated with data from three separate infiltrations. Calculation was performed according to the formula of:

$$\text{5mNSI level (\% of WT)} = \frac{(100) \times (\text{5 mNSI volume of transgenic line}) \times (\text{EYFP volume of wild-type})}{(\text{5mNSI volume of wild type}) \times (\text{EYFP volume of transgenic line})}$$




Fig. 4. Plant-derived *5mNSI* siRNAs can suppress replication of the influenza virus A/WSN/33 in mammalian cells. (A) Total RNA (10 μ g) from primary transformants 1 and 2 was separated on a 15% polyacrylamide gel, blotted onto a Nylon membrane and probed with [³²P]UTP-labeled *5mNSI* RNA probes. Quantity of siRNAs in the RNA samples was calculated based on its relative volume to that of known amount of synthetic siRNA oligo. These RNA preparations were then used for transfection of MDCK cells. (B) Normalization of RNA loading was based on the separation of 6 μ g of total RNA on a 1.2% agarose gel. (C) Suppression of viral replication as revealed in three independent HA titer assays. MDCK cells were first transfected with water (mock), a siRNA oligo NS-128 used by Ge et al. [8], and RNA from wild-type tobacco (control) or from two transgenic lines (line 1 and line 2) expressing siRNAs, and were then infected by influenza virus strain A/WSN/33 24 h post-transfection. HA titer was determined at 24 (blue boxes), 36 (red boxes) and 48 h (yellow boxes) post-infection.

307 gation in mammalian cells infected with the influenza virus,
 308 *5mNSI* siRNAs were harvested from the leaves of transgenic
 309 plants for transfection of MDCK cells. The amount of siRNA
 310 in total RNA was quantified by siRNA analysis (Fig. 4A). As a
 311 positive control, we used siRNA oligo NS-128 (5'-
 312 CGGCUUCGCCGAGAUCAGAdAdT-3'), since it has been
 313 proven best of three *NSI*-targeting siRNA oligos [8]. Cells
 314 transfected with RNA from non-transformed plants were the
 315 negative control and cells transfected with water constituted
 316 the mock transfection. Twelve hours after transfection, cells
 317 were infected by the influenza virus strain A/WSN/33
 318 (MOI = 0.001). The HA titer, which is an indicator of viral
 319 replication, was determined at 12, 24 and 36 h post-infection.
 320 The mock-transfected and the negative control cells showed
 321 similar HA titer, indicating that RNA from wild-type tobacco
 322 plants did not suppress viral replication. Though the HA titer
 323 values varied in three separate sets of transfection and infec-
 324 tion studies, a phenomenon unavoidably associated with
 325 conditions of the cells, e.g., passage history, both plant-derived
 326 and synthetic siRNAs significantly reduced H1N1 viral repli-
 327 cation. The anti-viral effect of siRNA was most prominent at
 328 36 h post-infection (Fig. 4B). In one set of experiments, plant-
 329 derived siRNA proved superior to the NS-128 oligo (Fig. 4B,
 330 experiment A).

331 These results strongly support our hypothesis that *5mNSI*
 332 siRNA from transgenic plants can effectively suppress repli-

333 cation of the influenza virus in mammalian cells. In addition,
 334 plant siRNAs showed similar suppression ability as the syn-
 335 thetic siRNA NS-128, demonstrating that plant-derived siR-
 336 NAs confer the same efficacy. Given the fact that transgenic
 337 plants can generate siRNAs targeting different areas of the
 338 *5mNSI* transcript, and that *NSI* sequences are highly con-
 339 served among influenza viruses [23], *5mNSI* siRNAs from
 340 transgenic plants should suppress the replication of a broad
 341 range of influenza viral subtypes with sequences homologous
 342 to the *5mNSI*.

343 While our results clearly indicate anti-viral effects of plant-
 344 derived *5mNSI* siRNAs, this study is primarily focusing on
 345 developing a strategy for economical and sustainable produc-
 346 tion of siRNAs. Besides using transgenic technology described
 347 in this study, a pool of siRNAs can also be generated with
 348 Dicer-dependent kits. When compared with the transgenic
 349 approach, the latter method is much more expensive, since it
 350 involves expensive reagents (i.e., dNTP, Dicer, and RNA
 351 polymerase), complicated steps (i.e., in vitro transcription, in
 352 vitro cleavage of dsRNA, and clean-up) and experienced re-
 353 searcher. The high cost not only limits production scale, but
 354 also requires repetition of the production process if the siR-
 355 NAs are to be used over and over again. Therefore, our proof-
 356 of-concept study demonstrates that transgenic plants are su-
 357 perior to the commercial kits for siRNA production and the
 358 time taken for generating them would be well compensated.

359 Though the *NSI*-targeting siRNAs possess anti-viral effects,
360 those against the NP genes would be more potent in sup-
361 pressing viral replication, as revealed by studies using mam-
362 malian cells [8] and animals [34]. In both studies, one of the
363 NP-targeting siRNAs, Np-1496, significantly reduced the virus
364 titers. These observations indicate that mRNA of the NP gene
365 might be a better target of siRNA, if positional effects on
366 siRNA accessibility could be faithfully addressed. In our fu-
367 ture study of using transgenic plant-derived siRNAs for viral
368 suppression, generating NP-targeting siRNAs would be a
369 more practical practice.

370 In conclusion, *5mNSI* siRNAs capable of activating RNAi
371 in mammalian cells against *NSI* were produced in transgenic
372 tobacco plants. The efficacy of the plant-derived siRNAs was
373 tested in vivo by agroinfiltration of the *5mNSI*-expressing
374 construct in leaves of transgenic tobacco and in vitro by ap-
375 plication of these siRNAs in mammalian cells to inhibit in-
376 fluenza viral replication. This cost-effective technique in
377 utilizing transgenic plants for large-scale siRNA production
378 could have advantages over current methods involving the use
379 of synthetic RNA oligos, the expression of short hairpin RNA
380 in *Escherichia coli* [35,36] and the transfection of mammalian
381 cells with short dsRNA. In addition, plant cells can apparently
382 tolerate expression of long dsRNAs, enabling the length of the
383 target gene fragment to be easily manipulated for optimal
384 suppression. Moreover, fragments producing siRNAs target-
385 ing multiple sites of the viral genome can be fused together so
386 that one transgenic plant can produce siRNAs for simulta-
387 neous silencing of multiple genes. This could provide a more
388 robust and sustained viral protection minimizing the likeli-
389 hood of the virus developing resistance to the siRNA through
390 mutation of the target sequence.

391 *Acknowledgements:* This study is in part supported by a Hong Kong
392 Research Grant Council Grant (7342/03M) to E. Lam and Y.X. Zhou,
393 a Hong Kong Research Grant Council Grant (7356/03M) to L.L.M.
394 Poon, and funds from the Department of Botany, University of Hong
395 Kong to M.L. Chye.

396 References

397 [1] Napoli, C., Lemieux, C. and Jorgensen, R. (1990) *Plant Cell* 2,
398 279–289.
399 [2] Cogoni, C. and Macino, G. (1997) *Proc. Natl. Acad. Sci. USA* 94,
400 10233–10238.
401 [3] Fire, A., Xu, S., Montgomery, M.K., Kostas, S.A., Drivers, S.E.
402 and Mello, C.C. (1998) *Nature* 391, 806–811.
403 [4] Zamore, P.D., Tuschl, T., Sharp, P.A. and Bartel, D.P. (2000) *Cell*
404 101, 25–33.
405 [5] Wianny, F. and Zernicka-Goetz, M. (2000) *Nat. Cell Biol.* 2, 70–75.
406 [6] Hannon, G.J. (2002) *Nature* 418, 244–251.

[7] Denli, A.M. and Hannon, G.J. (2003) *Trends Biochem. Sci.* 28,
196–201. 407
[8] Ge, Q., McManus, M.T., Nguyen, T., Shen, C.H., Sharp, P.A.,
Eisen, H.N. and Chen, J. (2003) *Proc. Natl. Acad. Sci. USA* 100,
2718–2723. 409
[9] Sohail, M., Doran, G., Riedemann, J., Macaulay, V. and
Southern, E.M. (2003) *Nucleic Acids Res.* 31. 410
[10] Myers, J.W., Jones, J.T., Meyer, T. and Ferrel Jr., J.E. (2003) *Nat.*
Biotechnol. 21, 324–328. 411
[11] Lee, N.S., Dohjima, T., Bauer, G., Li, H., Li, M.J., Ehsani,
A., Salvaterra, P. and Rossi, J. (2002) *Nat. Biotechnol.* 19,
500–505. 412
[12] Kaykas, A. and Moon, R.T. (2004) *BMC Cell Biol.* 5, 16–26. 413
[13] Xia, H.B., Mao, Q.W., Paulson, H.L. and Davidson, B.L. (2002)
Nat. Biotechnol. 20, 1006–1010. 414
[14] Tenllado, F., Llave, C. and Diaz-Ruiz, R. (2004) *Virus Res.* 102,
85–96. 415
[15] McManus, M.T. and Sharp, P.A. (2002) *Nat. Rev. Genet.* 3, 737–
747. 416
[16] Holen, T., Amarzguioui, M., Wiiger, M.T., Babaie, E. and Prydz,
H. (2002) *Nucleic Acids Res.* 30, 1757–1766. 417
[17] Gil, J. and Esteban, M. (2000) *Apoptosis* 5, 107–114. 418
[18] Stark, G.R., Kerr, I.M., Williams, B.R.G., Silverman, R.H. and
Schreiber, R.D. (1998) *Annu. Rev. Biochem.* 67, 264–277. 419
[19] Kotani, H., Nakamura, Y., Sato, S., Asamizu, E., Kaneko, T. and
Miyajima, N. (1998) *DNA Res.* 5, 203–216. 420
[20] Gallois, P. and Marinho, P. (1995) *Methods Mol. Biol.* 49, 39–
48. 421
[21] Pontier, D., Miao, Z.H. and Lam, E. (2001) *Plant J.* 27, 529–538. 422
[22] Webby, R.J. and Webster, R.G. (2001) *Philos. Trans. R. Soc.*
London 356, 1817–1828. 423
[23] Suarez, D.L. and Perdue, M.L. (1998) *Virus Res.* 54, 59–69. 424
[24] Garcia-Sastre, A., Egorov, A., Matasovoy, D., Brandt, S., Lery,
D.E. and Durbin, J.E., et al. (1998) *Virology* 252, 324–330. 425
[25] Krug, R.M., Yuan, W., Noah, D.L. and Latham, A.G. (2003)
Virology 309, 181–189. 426
[26] Sui, G.C., Soohoo, C., Affar, E.B., Gay, F., Shi, Y., Forrester,
W.C. and Shi, Y. (2002) *Proc. Natl. Acad. Sci. USA* 99, 5515–
5520. 427
[27] Klahre, U., Crete, P., Leuenberger, S.A., Lglesias, V.A. and Meins
Jr., F. (2002) *Proc. Natl. Acad. Sci. USA* 99, 11981–11986. 428
[28] Tzfira, T., Li, J., Lacroix, B. and Citovsky, V. (2004) *Trends*
Genet. 20, 375–383. 429
[29] Chan, S.W.L., Zilberman, D., Xie, Z., Johansen, L.K., Carrington,
J.C. and Jacobsen, S.E. (2004) *Science* 303, 1336. 430
[30] Aufsatz, W., Mette, M.F., van der Winden, J., Mztzke, A.J. and
Matzke, M. (2002) *Proc. Natl. Acad. Sci. USA* 99 (Supp 14),
16499–16506. 431
[31] Cao, X., Sufsatz, W., Zilberman, D., Mette, M.F., Huang, M.S.,
Matzke, M. and Jacobsen, S.E. (2003) *Curr. Biol.* 13, 2212–2217. 432
[32] Newmark, P.A., Reddien, P.W., Cebria, F. and Sanchez Alva-
rado, A. (2003) *Proc. Natl. Acad. Sci. USA* 100 (Supp 11), 11861–
11865. 433
[33] Sorensen, D.R., Leirdal, M. and Sioud, M. (2003) *J. Mol. Biol.*
327, 761–766. 434
[34] Tompkins, S.M., Lo, C.Y., Tumpey, T.M. and Epstein, S.L.
(2004) *Proc. Natl. Acad. Sci. USA* 101, 8682–8686. 435
[35] Timmons, L. and Fire, A. (1998) *Nature* 395, 854. 436
[36] Ketting, R.F., Fischer, S.E.J., Bernstein, E., Sijen, T., Hannon,
G.J. and Plasterk, R.H.A. (2001) *Genes Dev.* 15, 2654–2659. 437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466