



Title	P. gingivalis LPS enhances human α-defensins expression in gingival epithelia
Author(s)	Jin, LJ; Lu, Q; Darveau, RP; Samaranayake, LP
Citation	6th Annual Scientific Meeting of the International Association for Dental Research (Chinese Division), Shanghai, China, 24-25 October 2005. In Journal of Dental Research, 2005 v. 84 Sp Iss B, p. 192, abstract no. O-192
Issued Date	2005
URL	http://hdl.handle.net/10722/53667
Rights	Creative Commons: Attribution 3.0 Hong Kong License

***P. gingivalis* LPS enhances human β -defensins expression in gingival epithelia**

L. JIN¹, Q. LU¹, R. DARVEAU², and L.P. SAMARANAYAKE¹, ¹University of Hong Kong, Hong Kong, ²University of Washington, Seattle, USA

Objectives: Human β -defensins (hBDs) are a group of small, broad-spectrum cationic antimicrobial peptides. Our recent studies showed that the expression levels and pattern of hBDs were associated with periodontal conditions. The present study aimed to determine the effect of *P. gingivalis* LPS on the mRNA expression of hBDs as well as pattern recognition receptors and related adaptive molecules in reconstituted human gingival epithelia (RHGE) model. **Methods:** In the dose-dependant study, RHGE were treated with *E. coli* 055:B5 LPS or *P. gingivalis* LPS at various concentrations (1 ng/ml - 10 μ g/ml) for 24 hours. While in the time-dependant study, RHGE were treated with 10 ng/ml of *E. coli* LPS or 10 μ g/ml of *P. gingivalis* LPS for 2, 6, 12, 18, 24, 36, 48, 72 and 96 hours, respectively. The mRNA expression of hBD-1, 2, and 3, CD14, LBP, TLR-2 and -4, MD-2 and MyD88 was detected by RT-PCR. **Results:** hBD-1, 2, and 3 mRNAs were basally expressed by RHGE. Overall, *P. gingivalis* LPS (10 μ g/ml) markedly enhanced the expression of hBDs at 24 hours with a concurrent expression of TLR-2; whereas *E. coli* LPS (1 or 10 ng/ml) slightly upregulated the message with a concurrent expression of TLR-4. *P. gingivalis* LPS enhanced expression of hBD-2 and 3 mRNA was observed at 2 hours. The expression of CD14, MD-2 and MyD88 was independent of the stimulation of *P. gingivalis* or *E. coli* LPS. **Conclusions:** *P. gingivalis* LPS may enhance the *in vitro* expression of hBDs 1-3 mRNAs in the reconstituted human gingival epithelia, which might be associated with TLR-2 signaling pathway. Supported by Hong Kong Research Grants Council (RGC HKU 7310/00M).

[Periodontal Research - Pathogenesis](#)

[The Preliminary Program for 6th Annual Scientific Meeting of the IADR Chinese Division \(October 24-25, 2005\)](#)