



Title	Plaque, gingivitis and periodontal diseases in relation to diagnosis of general diseases
Author(s)	Soder, B; Airila-Mansson, S; Jin, L; Klinge, B; Soder, PO
Citation	80th General Session and Exhibition of the International Association for Dental Research, San Diego, California, USA, 6-9 March 2002, v. 81 n. Sp A, p. A-450
Issued Date	2002
URL	http://hdl.handle.net/10722/53796
Rights	Creative Commons: Attribution 3.0 Hong Kong License

3670 Plaque, Gingivitis and Periodontal Diseases in relation to Diagnosis of General Diseases

B. SODER¹, S. AIRILA-MANSSON¹, L. JIN², B. KLINGE¹, and P.-O. SODER¹,
¹Karolinska Institutet, Huddinge, Sweden, ²University of Hong Kong, China

Objectives: The aim was to study the amount of plaque, gingival inflammation and periodontitis in subjects in relation to general diseases diagnosed at hospital and proceeding death, in a 13 year longitudinal study.
Methods: The participants consisted of 1664 subjects randomly selected 30 to 40 years of age, 842 females, 35.7 (+- 2.9 SD) yr. and 822 males 35.8 (+- 2.8) yr. Their hospital care and death was followed during 13 years. Oral examinations were carried out at baseline with determination of plaque (PLI), gingivitis (GI) and number of teeth with pockets 5 mm and deeper. After 13 years the subjects were divided in two groups, HM-group subjects who had been treated at hospitals before they passed away, 10 females and 13 males and HH-group, 1653, healthy or treated at hospital. The International Classification of Diseases, 9th Revision (ICD-9) has been used for the diagnosis of diseases and death. Analysis of Variance was the statistical method used. **Results:** During the 13 years 95.7 % (22 /23) of the subjects in the HM-group had visited hospital in mean 9 (+- 8.2 SD) times. Neoplasms (ICD 140-239), Mental Disorders (ICD 290-319) and Diseases of the Circulatory System (ICD 390-459) were the most frequent diagnosis found during their stay at hospital. There was statistically significant difference in PLI at baseline ($p < 0.05$) between the HM-group and HH-group 0.94 (+-1.4 SE) and 0.70 (+-0.01 SE) respectively. Statistically significant difference ($p = 0.01$) was also found for GI at baseline between the HM-group and HH-group, 1.55 (+-0.12 SE) and 1.27 (+-0.01 SE) respectively. **Conclusions:** Amount of plaque and gingival inflammation might be marker for development of certain diseases which can be the cause death. The study was supported by the Karolinska Institutet

[Seq #330 - Periodontal Medicine](#)

11:00 AM-12:15 PM, Saturday, 9 March 2002 San Diego Convention Center Exhibit Hall C

[Back to the Periodontal Research - Diagnosis / Epidemiology Program](#)
[Back to the IADR/AADR/CADR 80th General Session \(March 6-9, 2002\)](#)