



<b>Title</b>	<b>Why are we still promoting breast self-examination?</b>
<b>Author(s)</b>	<b>Tarrant, M</b>
<b>Citation</b>	<b>International Journal Of Nursing Studies, 2006, v. 43 n. 4, p. 519-520</b>
<b>Issued Date</b>	<b>2006</b>
<b>URL</b>	<b><a href="http://hdl.handle.net/10722/54326">http://hdl.handle.net/10722/54326</a></b>
<b>Rights</b>	<b>Creative Commons: Attribution 3.0 Hong Kong License</b>



Available online at [www.sciencedirect.com](http://www.sciencedirect.com)



INTERNATIONAL JOURNAL OF  
**NURSING  
STUDIES**

International Journal of Nursing Studies ■ (■■■■) ■■■-■■■

[www.elsevier.com/locate/ijnurstu](http://www.elsevier.com/locate/ijnurstu)

Commentary

# Why are we still promoting breast self-examination?

Marie Tarrant\*

*Department of Nursing Studies, Faculty of Medicine, University of Hong Kong, 4/F, William M.W. Mong Block, 21 Sassoon Road, Hong Kong, China*

Received 2 December 2005; accepted 9 December 2005

**Keywords:** Breast self-examination; Breast cancer screening

Nurses have long been advocates of breast self-examination (BSE), believing that not only were they promoting a practice that could be life-saving but that they were also empowering women to take greater control over their health. There is an abundance of research in the nursing literature reporting the measurement of women's BSE practices, the psychometric correlates of BSE practice and strategies attempting to increase BSE practice among women (Champion and Menon, 1997; Chouliara et al., 2004; Petro-Nustus and Mikhaïl, 2002; Reis et al., 2004; Secginli and Nahcivan, 2004). In their study to be published in forthcoming issue of IJNS, Secginli and Nahcivan (in press) sought to identify variables correlated with the breast cancer screening behaviours of BSE and mammography in Turkish women, presumably so that rates of both these practices could be increased. While I do not take issue with the methodology or results presented in this paper, it is still, nonetheless, fundamentally flawed. The authors clearly delineate the rates of participation in both screening activities and the psychometric correlates of both BSE and mammography separately. There is a tendency, however, when discussing the benefits of screening to combine both BSE and mammography as if they were interconnected. BSE and mammography are two discrete procedures and should be discussed as such. Furthermore, all studies examining breast screening practices are designed on the premise that through early detection of breast lumps, breast cancer mortality can be

reduced and lives can be saved. In the introduction and literature review sections of their paper, Secginli and Nahcivan present no evidence on the effectiveness of either BSE or mammography in detecting breast cancer and decreasing mortality. Perhaps, this is because although the benefits of mammography are still currently being debated (Goodman, 2002; Olsen and Gotzsche, 2005; U.S. Preventive Services Task Force, 2002), a preponderance of evidence has now clearly shown that BSE does not save lives and offers no benefit to women.

Results from two large randomized controlled trials (RCTS) involving almost 400,000 women in Russia and China have shown that BSE is not effective in reducing mortality from breast cancer, and does not improve the probability of survival after breast cancer diagnosis (Semiglazov et al., 1999; Thomas et al., 2002). Moreover, both studies also demonstrated that regularly practicing BSE was significantly more likely to cause harm by way of increased biopsies for benign breast lumps. In a systematic review of the benefits of BSE, the Cochrane group has concluded that "screening by breast self-examination cannot be recommended" (Kosters and Gotzsche, 2005) and most experts no longer recommend BSE (Baxter and Canadian Task Force on Preventive Health, 2001; Elmore et al., 2005; Harris and Kinsinger, 2002). Furthermore, even in countries such as Turkey, where mammography screening may not be widely available, because of its lack of demonstrable benefits, promoting BSE is not a prudent use of the limited funds available for preventive services (Thomas et al., 2002).

\*Tel.: +852 2819 2643; fax: +852 2872 6079.  
E-mail address: [tarrantm@hku.hk](mailto:tarrantm@hku.hk).

57  
59  
61  
63  
65  
67  
69  
71  
73  
75  
77  
79  
81  
83  
85  
87  
89  
91

1 Contrary to the recommendation of Secginli and  
 3 Nahcivan that more longitudinal studies on the factors  
 5 influencing the use of BSE are required, the evidence  
 7 against BSE is sufficiently compelling that the Cochrane  
 9 group has also concluded that “it is unlikely that  
 11 additional trials investigating breast-self examination as  
 13 a single general screening method would be worthwhile”  
 15 (Kosters and Gotzsche, 2005).

17 What does all of this mean for nursing? Firstly, it  
 19 means that promoting BSE at a population level and  
 21 investigating factors which can increase performance of  
 23 BSE are not worthy of valuable time and money.  
 25 Resources should be focused on promoting and  
 27 investigating screening practices with proven benefits,  
 29 or on more accurate measurement of the benefits of  
 31 other screening practices currently in use, such as  
 mammography. What these findings do not mean,  
 however, is that we should teach women to ignore their  
 breasts. Education on BSE should be replaced by breast  
 awareness education, where women are taught the  
 cardinal sign of breast cancer, a painless lump, and the  
 necessity of seeking prompt medical evaluation of that  
 lump (Harris and Kinsinger, 2002). Additionally, if  
 women choose to continue to regularly perform BSE,  
 they should be informed that the benefits are unproven  
 and that it may result in unnecessary biopsies for benign  
 breast lumps (Thomas et al., 2002). Finally, women  
 should continue to participate in mammography screen-  
 ing programs and receive annual clinical breast exams as  
 indicated by the national or regional guidelines for  
 where they reside.

33 The time has come, therefore, to say good-bye to BSE.  
 35 There is no evidence to support the practice, and the best  
 37 available evidence tells us that it does more harm than  
 39 good. It is natural that nurses would not willingly give  
 41 up promoting a practice that they have strongly believed  
 43 in and have invested in considerably. However, if we are  
 45 fully embracing an evidence-based practice we have to  
 47 go with the evidence, even if we do not like what it is  
 49 telling us.

## References

45 Baxter, N., Canadian Task Force on Preventive Health, C.,  
 47 2001. Preventive health care, 2001 update: should women be  
 49 routinely taught breast self-examination to screen for breast  
 cancer? *Canadian Medical Association Journal* 164 (13),  
 1837–1846.

- Champion, V., Menon, U., 1997. Predicting mammography  
 and breast self-examination in African American women.  
*Cancer Nursing* 20 (5), 315–322. 51
- Chouliara, Z., Papadioti-Athanasidou, V., Power, K.G., Swanson,  
 V., 2004. Practice of and attitudes toward breast self-  
 examination (BSE): a cross-cultural comparison between  
 younger women in Scotland and Greece. *Health Care for  
 Women International* 25 (4), 311–333. 55
- Elmore, J.G., Armstrong, K., Lehman, C.D., Fletcher, S.W.,  
 2005. Screening for breast cancer. *Journal of the American  
 Medical Association* 293 (10), 1245–1256. 57
- Goodman, S.N., 2002. The mammography dilemma: a crisis for  
 evidence-based medicine? *Annals of Internal Medicine* 137  
 (5), 363–365. 59
- Harris, R., Kinsinger, L.S., 2002. Routinely teaching breast  
 self-examination is dead. What does this mean? *Journal of  
 the National Cancer Institute* 94 (19), 1420–1421. 63
- Kosters, J.P., Gotzsche, P.C., 2005. Regular self-examination  
 or clinical examination for early detection of breast cancer  
 [systematic review]. *Cochrane Database of Systematic  
 Reviews* 4. 65
- Olsen, O., Gotzsche, P.C., 2005. Screening for breast cancer  
 with mammography. *Cochrane Database of Systematic  
 Reviews* 4. 67
- Petro-Nustus, W., Mikhail, B.I., 2002. Factors associated with  
 breast self-examination among Jordanian women. *Public  
 Health Nursing* 19 (4), 263–271. 69
- Reis, J., Trockel, M., King, T., Remmert, D., 2004. Computer-  
 ized training in breast self-examination: a test in a  
 community health center. *Cancer Nursing* 27 (2), 162–168. 71
- Secginli, S., Nahcivan, N.O., 2004. Reliability and validity of  
 the breast cancer screening belief scale among Turkish  
 women. *Cancer Nursing* 27 (4), 287–294. 73
- Secginli, S., Nahcivan, N.O. Factors associated with breast  
 cancer screening behaviours in a sample of Turkish women:  
 a questionnaire survey. *International Journal of Nursing  
 Studies*, in press. 75
- Semiglazov, V.F., Moiseenko, V.M., Manikhas, A.G., Prot-  
 senko, S.A., Kharikova, R.S., Ivanov, V.G., Barash, N.I.,  
 Seleznev, I.K., Migmanova, N.S., Ivanova, O.A., Orlov,  
 A.A., Popova, R.T., Chagunava, O.L., 1999. Role of breast  
 self-examination in early detection of breast cancer: Russia/  
 WHO prospective randomized trial in St. Petersburg.  
*Cancer Strategy* 1, 145–151. 77
- Thomas, D.B., Gao, D.L., Ray, R.M., Wang, W.W., Allison,  
 C.J., Chen, F.L., Porter, P., Hu, Y.W., Zhao, G.L., Pan,  
 L.D., Li, W., Wu, C., Coriaty, Z., Evans, I., Lin, M.G.,  
 Stalsberg, H., Self, S.G., 2002. Randomized trial of breast  
 self-examination in Shanghai: final results. *Journal of the  
 National Cancer Institute* 94 (19), 1445–1457. 79
- U.S. Preventive Services Task Force, 2002. Screening for breast  
 cancer: recommendations and rationale. *Annals of Internal  
 Medicine* 137 (5), 344–346. 81