



Title	Suppression of hypoxia inducible factor-1 (HIF-1) by YC-1 is dependent on murine double minute 2 (Mdm2)
Author(s)	Lau, CK; Yang, ZF; Lam, CT; Tam, KH; Poon, RTP; Fan, ST
Citation	Biochemical And Biophysical Research Communications, 2006, v. 348 n. 4, p. 1443-1448
Issued Date	2006
URL	http://hdl.handle.net/10722/54335
Rights	Creative Commons: Attribution 3.0 Hong Kong License



ELSEVIER

Available online at www.sciencedirect.com



BBRC

Biochemical and Biophysical Research Communications xxx (2006) xxx–xxx

www.elsevier.com/locate/ybbrc

Suppression of hypoxia inducible factor-1 α (HIF-1 α) by YC-1 is dependent on murine double minute 2 (Mdm2) [☆]

Chi Keung Lau, Zhen Fan Yang ^{*}, Chi Tat Lam, Ka Ho Tam,
Ronnie Tung Ping Poon, Sheung Tat Fan

Center for the Study of Liver Disease and Department of Surgery, The University of Hong Kong, Pokfulam, Hong Kong, China

Received 29 July 2006

Abstract

Inhibition of HIF-1 α activity provides an important strategy for the treatment of cancer. Recently, 3-(5'-hydroxymethyl-2'-furyl)-1-benzyl indazole (YC-1) has been identified as an anti-HIF-1 α drug in cancer therapy with unclear molecular mechanism. In the present study, we aimed to investigate the effect and mechanism of YC-1 on HIF-1 α in a hepatocellular carcinoma cell line under hypoxic condition, which was generated by incubating cells with 0.1% O₂. The phenotypic and molecular changes of cells were determined by cell proliferation assay, apoptosis assay, luciferase promoter assay, and Western blot analysis. YC-1 arrested tumor cell growth in a dose-dependent manner, whereas it did not induce cell apoptosis. Hypoxia-induced upregulation of HIF-1 α was suppressed by YC-1 administration. YC-1 inhibited HIF-1 α protein synthesis under normoxia and affected protein stability under hypoxia. YC-1 suppressed the expression of total and phosphorylated forms of murine double minute 2 (Mdm2), whereas this inhibitory effect was blocked by overexpression of Mdm2. In conclusion, YC-1 suppressed both protein synthesis and stability of HIF-1 α in HCC cells, and its inhibitory effects on HIF-1 α were dependent on Mdm2.

© 2006 Elsevier Inc. All rights reserved.

Keywords: Hypoxia inducible factor-1 α ; YC-1; Murine double minute 2; Hepatocellular carcinoma

Hepatocellular carcinoma (HCC) is one of the five most common malignancies in the world, with an increasing incidence in both Asian and Western countries [1]. Only a small proportion of patients are suitable candidates for liver transplantation, surgical resection or other surgical treatments due to the advanced stage of tumor or poor hepatic functional reserve. Transarterial chemoembolization is one of the major alternatives for the treatment of HCC patients with an advanced stage [2,3]. However, the long-term survival is unsatisfactory and the role of hypoxia

in stimulating cancer growth is thought to be one of the reasons that lead to treatment failure [4].

Hypoxia is a common phenomenon in solid tumors, as oxygen supply usually does not meet the demand of tumor cells during progression [5]. The reduced oxygen levels in tumor tissues induce serial changes of hypoxia-related molecules that promote angiogenesis, among which hypoxia inducible factor-1 α (HIF-1 α) is the most predominant one [6,7]. Overexpression of HIF-1 α was associated with angiogenesis, tumor invasion, and poor prognosis of various types of cancers [8–12]. In HCC, it was reported that activation of HIF-1 α promoted upregulation of VEGF, a key player during angiogenesis [13,14]. In addition to hypoxic condition, HIF-1 α could be upregulated by some therapeutic approaches, such as transarterial chemoembolization, resulting in treatment failure and poor outcomes [15]. Due to the importance of HIF-1 α in tumor progression and angiogenesis, 50

[☆] Financial support: The study was supported by the Small Project Funding of the University of Hong Kong, and the grant from the Research Grant Council of the Hong Kong Special Administrative Region (Project No. HKU 7621/06M).

^{*} Corresponding author. Fax: +852 2818 9249.

E-mail address: zfyang@hkucc.hku.hk (Z.F. Yang).

51 targeting HIF-1 α becomes a potential approach of cancer
52 therapy that has attracted great interest [12,16–18].

53 A number of chemicals and drugs have been discovered
54 in recent years for targeting HIF-1 α , one of which is 3-(5'-
55 hydroxymethyl-2'-furyl)-1-benzyl indazole (YC-1). YC-1
56 was first identified as an activator of platelet guanylate
57 cyclase in 1994 and was used as a vessel dilator in circula-
58 tion disorders [19]. Under hypoxic condition, YC-1 exhib-
59 ited anticancer effects through inhibition of HIF-1 α activity
60 [20]. However, little is known about the possible mecha-
61 nism of YC-1-mediated HIF-1 α suppression. As the rela-
62 tionship between murine double minute 2 (Mdm2) and
63 HIF-1 α has been demonstrated by some studies, we
64 designed the present study to investigate the potential role
65 of Mdm2 in YC-1-mediated HIF-1 α suppression.

66 Materials and methods

67 *Cell lines.* HepG2 human HCC cell line was purchased from the
68 American Type Culture Collection (Manassas, VA). Cells were main-
69 tained as monolayer culture in Dulbecco's modified Eagle's medium
70 (DMEM) with 10% fetal bovine serum (FBS) and 1% penicillin (Life
71 Technologies, Carlsbad, CA) at 37 °C in a humidified atmosphere of 5%
72 CO₂ in air.

73 *Cell proliferation assay.* Cell proliferation was determined by 3,[4,5-
74 dimethylthiazol-2-yl]-2,5-diphenyl-tetrazolium bromide (MTT) assay. The
75 HepG2 cells (1×10^4) were inoculated into 96-well plates, and treated with
76 1% dimethylsulfoxide (DMSO) in 10% FBS-DMEM or different doses (1,
77 5, and 10 μ M) of YC-1 (dissolved in 1% DMSO-10% FBS-DMEM),
78 respectively, for 12 h before incubating in a humidified atmosphere of 95%
79 N₂/5% CO₂ (the final oxygen content estimated to be 0.1%) for 24 h. MTT
80 was then added into each well and the cells were incubated for another 4 h.
81 The reaction was stopped with 0.04 M hydrochloride (in isopropanol) and
82 measured at λ 570–630 nm in a V_{\max} kinetic microplate reader (Molecular
83 Devices Corporation, Sunnyvale, CA). The cell proliferation index was
84 expressed as means \pm SD.

85 *Cytofluorometric apoptosis analysis.* The HepG2 cells (5×10^5) were
86 inoculated into each well of a six-well plate, and treated with 1%
87 DMSO in 10% FBS-DMEM and different doses (1, 5, and 10 μ M) of
88 YC-1, respectively, in a hypoxic condition for 24 h. The cells were then
89 labeled with Annexin V-FITC (BD Biosciences Pharmingen, San Diego,
90 CA), and detected in a FACS Calibur (Becton Dickinson Immunocytometry
91 Systems, San Jose, CA). Unstained cells were used as a nega-
92 tive control.

93 *Terminal deoxynucleotidyl transferase-mediated dUTP nick end labeling*
94 (*TUNEL*) assay. The TUNEL technique was performed to detect apop-
95 totic cells using the in situ cell death detection kit (Roche Diagnostics,
96 Indianapolis, IN). Briefly, the HepG2 cells were cultured on cover slides
97 with different treatments. After 24-h incubation, cover slides were fixed
98 with 4% paraformaldehyde for 1 h and permeabilized by 0.1% Triton X-
99 100 at 4 °C for 2 min. The slides were then incubated with TUNEL
100 reaction mixture for 1 h at 37 °C. After washing, the slides were incubated
101 with horse-radish peroxidase-conjugated anti-fluorescein antibody for
102 30 min at 37 °C. After substrate reaction, slides were counterstained with
103 hematoxylin, and the number of apoptotic nuclei was examined under a
104 light microscope with the magnification of 400.

105 *Western blot.* The HepG2 cells (5×10^5) were inoculated into each well
106 of a 6-well plate, and treated with 1% DMSO in 10% FBS-DMEM and
107 10 μ M of YC-1, respectively, for different time intervals under hypoxic
108 condition according to the experimental design. After exposure of cells to
109 the indicated agents and time courses, reactions were terminated by
110 addition of lysis buffer (Cell Signaling Technology, Beverly, MA). The cell
111 lysates were electrophoresized on 8–12% SDS-PAGE. The primary anti-
112 bodies were anti-HIF-1 α (Calbiochem, San Diego, CA), anti- β -actin

(Santa Cruz Biotechnology, Santa Cruz, CA), anti-Mdm2 and anti-
phosphorylated Mdm2 (P-Mdm2) (Cell Signaling Technology). The rela-
tive protein level was expressed by a ratio to β -actin.

116 *HIF-1 α protein synthesis and protein stability.* In the protein synthesis
117 experiment, to determine the optimal doses and time intervals of protea-
118 some inhibitor, MG132 (Sigma–Aldrich, St. Louis, MO), at different
119 doses, was added into the cell line, and incubated for different time peri-
120 ods, respectively. The expression of HIF-1 α was examined by Western
121 blot. Based on the findings of the above protocols, the dose of 40 μ M
122 MG132 and incubation time of 4 and 6 h was chosen for the following
123 experiments. The HepG2 cells were pre-treated with 10 μ M YC-1 for 12 h
124 before adding 40 μ M MG132 and incubated for 4 and 6 h, respectively,
125 and the expression of HIF-1 α was determined by Western blot. In the
126 protein stability experiment, the HepG2 cells were incubated under hypox-
127 ic condition (0.1% O₂) for 4 h before administration of 100 μ M protein
128 synthesis inhibitor, cycloheximide (Sigma–Aldrich) with or without 10 μ M
129 YC-1, and incubated for another 30 and 60 min, respectively. Cells were
130 lysed and protein was extracted for Western blot analysis of HIF-1 α
131 expression.

132 *Cell transfection.* Cytomegalovirus (CMV)-Mdm2 plasmid (a gift from
133 Dr. Bert Vogelstein) [21] and empty vector were transfected for 24 h before
134 being treated with 5 μ M YC-1 under hypoxic condition. The levels of
135 HIF-1 α , Mdm2 and P-Mdm2 were also detected by the standard Western
136 blot protocol.

137 *Transfections and luciferase reporter assay.* The HepG2 cells (1×10^5)
138 were transfected with 1 μ g of pGL3-Mdm2 reporter plasmid (a gift from
139 Dr. Jason M. Shohet) [22] and 1 μ g of pRL-TK (*Renilla* luciferase, Pro-
140 mega, Madison, WI) as a normalization control. Cell transfection was
141 achieved by using Fugene 6 transfection reagent (Roche Diagnostics,
142 Indianapolis, IN). The luciferase activities were measured by luminometer
143 using the Dual-Luciferase Reporter Assay System according to the man-
144 ufacturer's instruction (Promega).

145 Results

146 Under hypoxic condition, YC-1 exerted a dose-depend-
147 ent inhibition of cell growth in the HepG2 cells with
148 IC₅₀ of 5 μ M (Fig. 1A). To further examine whether the
149 effect of YC-1 on tumor cells was cytostatic or cytotoxic,
150 cytofluorometric apoptosis assay was performed. Under
151 the same experimental conditions, YC-1 exhibited no sig-
152 nificant effect on tumor cell death even with a concentra-
153 tion of 10 μ M in a 24-h treatment (Fig. 1B). Similar to
154 the results of Annexin-V staining, TUNEL assay did not
155 identify any difference in the number of apoptotic cells
156 between the groups with and without YC-1 treatment in
157 the HepG2 cells, even with the highest dose tested
158 (10 μ M) (Fig. 1C).

159 When the tumor cells were pre-treated with 10 μ M YC-1
160 for 12 h before incubating in 0.1% O₂ for another 4 h, the
161 protein expression of HIF-1 α was significantly decreased in
162 the HepG2 cells, compared with that without YC-1 treat-
163 ment (data not shown).

164 As HIF-1 α protein is subjected to rapid degradation
165 under normoxia by the process of pVHL-mediated ubiqui-
166 tin-proteasome pathway, whereas the hypoxic condition
167 blocks the effect of degradation and leads to accumulation
168 of HIF-1 α protein. A proteasome inhibitor, MG132, was
169 used to prevent proteasome-mediated HIF-1 α protein deg-
170 radation under normoxia and the effect of YC-1 on HIF-1 α
171 protein synthesis was determined by measuring the accu-
172 mulation of protein at certain time points using Western




Fig. 1. YC-1 inhibited tumor cell growth under hypoxic condition. (A) The HepG2 cells were treated with different doses (1, 5, and 10 μM) of YC-1 for 12 h before incubating under 0.1% O_2 for another 24 h. The cell viabilities were assayed using MTT as described in the Materials and methods. The number of apoptotic cells was determined by (B) cytofluorometric apoptosis assay (Annexin V-FITC labeling) and (C) TUNEL assay. Under the conditions with or without YC-1 treatment, no significant difference in the number of apoptotic cells was detected by both assays. The percentage of Annexin V-FITC positive cells was expressed as means \pm SD. Arrows pointed to the apoptotic nuclei. DMSO, dimethyl sulfoxide.




Fig. 2. (A) YC-1 inhibited HIF-1 α protein synthesis under normoxic condition. To inhibit the HIF-1 α protein degradation, a proteasome inhibitor, MG132, was used. (a) and (b) The HepG2 cells were treated with different doses (10, 20 or 40 μM) of MG132 for 4 h, or incubated for different time periods (1, 2 or 4 h) before determination of HIF-1 α protein levels using Western blot. MG132 exhibited a dose and time dependent suppression of HIF-1 α protein degradation. (c) After treated with 10 μM YC-1 and MG132 (40 μM) for 4- or 6-h, a downregulation of HIF-1 α was detected. (B) YC-1 inhibited HIF-1 α protein stability under hypoxic condition. The HepG2 cells were pre-treated with 0.1% O_2 for 4 h before cycloheximide (100 μM) was added with or without YC-1 (10 μM), and incubated for 30 or 60 min. Cells were harvested and the HIF-1 α protein levels were detected using Western blot. DMSO, dimethylsulfoxide. Representative of three independent experiments.

173 blot. The effect of MG132 on proteasome inhibition was in
174 a dose and time dependent manner (Fig. 2A-a). As MG132
175 at the dose of 40 μM (Fig. 2A-a) and with the incubation
176 time of 4 h (Fig. 2A-b) had the most significant inhibitory
177 effect (with no obvious morphological changes of the cells),
178 these dose and time point were chosen for the YC-1 exper-
179 iment. Compared to the control groups, the protein synthe-
180 sis of HIF-1 α in the HepG2 cells was affected by YC-1 and
181 a significant inhibitory effect was observed at the 6-h time
182 point (Fig. 2A-c).

183 In addition to the effect of YC-1 on HIF-1 α protein syn-
184 thesis, its effect on protein stability was also tested. After
185 incubating the cells under hypoxic condition for 4 h, a pro-
186 tein synthesis inhibitor, cycloheximide, was added into the
187 culture medium with or without YC-1 treatment. It was

found that the expression of HIF-1 α protein in the DMSO 188
control group was much higher than that in the YC-1 treated 189
HepG2 cells (Fig. 2B). 190

As both the HIF-1 α protein synthesis and stability could 191
be affected by YC-1 in the HepG2 cells and Mdm2 was a 192
potential upstream molecule that regulated HIF-1 α expres- 193
sion, the possible link between Mdm2 and YC-1-mediated 194
HIF-1 α suppression was investigated. The HepG2 cells 195
were treated with 10 μM YC-1 under hypoxia for 1, 2, 196
and 4 h, respectively, and the expression of HIF-1 α , total 197
Mdm2, and P-Mdm2 was detected by Western blot. A 198
concurrent downregulation of HIF-1 α , total Mdm2, and 199

200 P-Mdm2 was detected with YC-1 treatment for 2 and 4 h
201 under hypoxic condition (Fig. 3A).

202 In order to further examine whether YC-1 mediated its
203 effect on HIF-1 α expression through suppression of




Fig. 3. (A) YC-1 suppressed the expression of HIF-1 α , total and phosphorylated forms of Mdm2 under hypoxic condition in a time dependent manner. The HepG2 cells were treated with 10 μ M YC-1 under hypoxia for different time intervals (1, 2 or 4 h). Cells were then harvested for the detection of HIF-1 α , Mdm2, and P-Mdm2 expression using Western blot. (B) Upregulation of Mdm2 by transfection reversed YC-1-mediated HIF-1 α suppression. The HepG2 cells were transfected with either empty vector or CMV-Mdm2 for 24 h. After transfection, the cells were treated with DMSO or 10 μ M YC-1 and incubated under 0.1% O₂ for 4 h before determination of HIF-1 α , Mdm2, and P-Mdm2 expression using Western blot. (C) YC-1 suppressed the promoter activity of Mdm2 in the HepG2 cells. Cells were co-transfected with 1 μ g of pGL3-Mdm2 reporter plasmid and 1 μ g of pRL-TK as a normalization control. The luciferase activity or *Renilla* luciferase activity was measured by luminometer using Dual-Luciferase Reporter Assay System according to the manufacturer's instruction. DMSO: dimethyl sulfoxide. The Firefly luciferase activity was normalized with *Renilla* luciferase activity. * $P < 0.05$, compared with DMSO control under hypoxia (Student's *t* test). Representative of three independent experiments.

Mdm2, under hypoxia, cells were transfected with CMV-Mdm2 plasmid for 24 h before DMSO or YC-1 was added. The transfection of Mdm2 induced a significant increase in the expression of total Mdm2 and P-Mdm2. In addition, the enhanced expression of Mdm2 by transfection could increase HIF-1 α level despite the presence or absence of YC-1 treatment in the HepG2 cells (Fig. 3B).

The previous experiments revealed that YC-1 might mediate its inhibitory effect on HIF-1 α expression by downregulation of Mdm2 protein. It was of interest to know whether YC-1 affected Mdm2 expression at the transcriptional level or protein level. Therefore, wild type Mdm2 promoter constructed in luciferase reporter plasmid was transfected before YC-1 administration. It was found that 10 μ M YC-1 significantly suppressed Mdm2 transcription in hypoxic HepG2 cells by an average of 2-fold compared with DMSO control (Fig. 3C).

Discussion

In the present study, we demonstrated that YC-1 inhibited the growth of HCC cells. This was consistent with the study of Wang et al., [23], which suggested that YC-1 exhibited an anti-proliferative effect by arresting the cell cycle in the G0-G1 phase in HCC cells. Similar effect was also found in endothelial cells and mesangial cells [24,25]. However, our data did not support a previous finding in prostate cancer that YC-1 could induce apoptosis of tumor cells [26]. Even with the dose of 10 μ M, YC-1 exhibited no effect on induction of cell apoptosis examined by both TUNEL assay and cytofluorometric apoptosis assay, suggesting that YC-1 inhibited the activity of HCC cells through a cytostatic pathway rather than a cytotoxic one.

Although the anti-HIF-1 α effect of YC-1 has been well demonstrated in several studies, the molecular basis of YC-1-mediated HIF-1 α suppression remains largely unclear. The present study revealed that YC-1 could affect both protein synthesis and protein stability of HIF-1 α , suggesting dual effects of YC-1 on suppressing HIF-1 α expression. To further explore the suppressive effect of YC-1 on protein synthesis, we performed another set of experiments to investigate whether this inhibitory effect was related to the mammalian target of rapamycin (mTOR) signaling pathway, as several downstream molecules of mTOR, such as ribosomal S6 kinase and eukaryote initiation factor 4E binding protein 1, were key regulators in protein translation and synthesis [27,28]. However, we did not detect any changes of these molecules after YC-1 treatment (data not shown), implying that YC-1-mediated inhibition of protein synthesis was independent of mTOR signaling pathway. Therefore, further studies are needed to explore other pathways that are related to protein synthesis.

Based on some studies demonstrating that Mdm2 might play a potential role in HIF-1 α protein stability [29,30], we investigated the relationship among YC-1, HIF-1 α , and

259 Mdm2 in the present study. With the downregulation of
 260 HIF-1 α , the protein level of Mdm2 was significantly
 261 decreased with YC-1 administration in a time dependent
 262 manner, indicating that Mdm2 might be involved in YC-
 263 1-mediated HIF-1 α suppression. To further prove this
 264 hypothesis, we induced upregulation of Mdm2 in the
 265 HepG2 cells by transfection before DMSO or YC-1 admin-
 266 istration, and found that the increased expression of Mdm2
 267 could reverse the inhibitory effect of YC-1 on HIF-1 α
 268 expression, suggesting that YC-1 regulated HIF-1 α expres-
 269 sion was Mdm2 dependent. To further explore whether
 270 YC-1 functioned on Mdm2 at a transcriptional level, we
 271 measured the promoter activity of Mdm2 under the condi-
 272 tions with or without YC-1 treatment, and found that YC-
 273 1 could decrease the promoter activity of Mdm2, suggest-
 274 ing that YC-1 might act on the transcriptional level of
 275 Mdm2. In addition, by detecting a downregulation of Fli-
 276 1, an upstream transcriptional regulator of Mdm2 [31], this
 277 study suggested that YC-1 functioned on the transcription-
 278 al level of Mdm2 in the cells with endogenous Mdm2.

279 In conclusion, YC-1 retarded cell growth and exhibited
 280 a cytostatic effect in the HCC cells under hypoxic condi-
 281 tion. YC-1 downregulated HIF-1 α expression by affecting
 282 both protein synthesis and stability, and the inhibitory
 283 effects of YC-1 on HIF-1 α were dependent on Mdm2.

284 Acknowledgments

285 The CMV-Mdm2 plasmid was kindly provided by Dr.
 286 Bert Vogelstein, the Howard Hughes Medical Institute
 287 and Sidney Kimmel Comprehensive Cancer Center, the
 288 Johns Hopkins Oncology Center, MD. The pGL3-Mdm2
 289 reporter was kindly provided by Dr. Jason M. Shohet,
 290 Center for Cell and Gene Therapy, Texas Children's Can-
 291 cer Center, Baylor College of Medicine, Houston, TX. We
 292 thank Dr. Julian J. Lum, Abramson Family Cancer Re-
 293 search Institute, Department of Cancer Biology, University
 294 of Pennsylvania School of Medicine, Philadelphia, PA for
 295 his valuable comments.

296 References

297 [1] H.B. El Serag, Epidemiology of hepatocellular carcinoma, Clin. Liver
 298 Dis. 15 (2001) 87–107.
 299 [2] J.M. Llovet, M.I. Real, X. Montana, R. Planas, S. Coll, J. Aponte, C.
 300 Ayuso, M. Sala, J. Muchart, R. Sola, J. Rodes, J. Bruix, Barcelona
 301 Liver Cancer Group. Arterial embolisation or chemoembolisation
 302 versus symptomatic treatment in patients with unresectable hepato-
 303 cellular carcinoma: a randomised controlled trial, Lancet. 359 (2002)
 304 1734–1739.
 305 [3] C.M. Lo, H. Ngan, W.K. Tso, C.L. Liu, C.M. Lam, R.T. Poon, S.T.
 306 Fan, J. Wong, Randomized controlled trial of transarterial lipiodol
 307 chemoembolization for unresectable hepatocellular carcinoma, Hepa-
 308 tology 35 (2002) 1164–1171.
 309 [4] C.B. O'Suilleabhain, R.T. Poon, J.L. Yong, G.C. Ooi, W.K. Tso, S.T.
 310 Fan, Factors predictive of 5-year survival after transarterial chemo-
 311 embolization for inoperable hepatocellular carcinoma, Br. J. Surg. 90
 312 (2003) 325–331.
 313 [5] C. Menon, D.L. Fraker, Tumor oxygenation status as a prognostic
 314 marker, Cancer Lett. 221 (2005) 225–235.

[6] T. Schmid, J. Zhou, B. Brune, HIF-1 and p53: communication of
 315 transcription factors under hypoxia, J. Cell. Mol. Med. 8 (2004) 423–
 316 431.
 317 [7] M. Wartenberg, F.C. Ling, M. Muschen, F. Klein, H. Acker, M.
 318 Gassmann, K. Petrat, V. Putz, J. Hescheler, H. Sauer, Regulation of
 319 the multidrug resistance transporter *P*-glycoprotein in multicellular
 320 tumor spheroids by hypoxia-inducible factor (HIF-1) and reactive
 321 oxygen species, FASEB J. 17 (2003) 503–505.
 322 [8] R. Bos, P.J. van Diest, J.S. de Jong, G.P. van der, V. van d, W.E. van
 323 der, Hypoxia-inducible factor-1 α is associated with angiogenesis,
 324 and expression of bFGF, PDGF-BB, and EGFR in invasive breast
 325 cancer, Histopathology 46 (2005) 31–36.
 326 [9] K. Nakanishi, S. Hiroi, S. Tominaga, S. Aida, H. Kasamatsu, S.
 327 Matsuyama, T. Matsuyama, T. Kawai, Expression of hypoxia-
 328 inducible factor-1 α protein predicts survival in patients with
 329 transitional cell carcinoma of the upper urinary tract, Clin. Cancer
 330 Res. 11 (2005) 2583–2590.
 331 [10] V.E. Theodoropoulos, A.Ch. Lazaris, F. Sofras, I. Gerzelis, V.
 332 Tsoukala, I. Ghikonti, K. Manikas, I. Kastriotis, Hypoxia-inducible
 333 factor 1 α expression correlates with angiogenesis and unfavor-
 334 able prognosis in bladder cancer, Eur. Urol. 46 (2004) 200–208.
 335 [11] D. Zagzag, H. Zhong, J.M. Scalzitti, E. Laughner, J.W. Simons, G.L.
 336 Semenza, Expression of hypoxia-inducible factor 1 α in brain
 337 tumors: association with angiogenesis, invasion, and progression,
 338 Cancer 88 (2000) 2606–2618.
 339 [12] G.L. Semenza, Targeting HIF-1 for cancer therapy, Nat. Rev. Cancer
 340 3 (2003) 721–732.
 341 [13] E. Dupuy, P. Hainaud, A. Villemain, E. Bodevin-Phedre, J.P.
 342 Brouland, P. Briand, G. Tobelem, Tumoral angiogenesis and tissue
 343 factor expression during hepatocellular carcinoma progression in a
 344 transgenic mouse model, J. Hepatol. 38 (2003) 793–802.
 345 [14] S. Yasuda, S. Arii, A. Mori, N. Isobe, W. Yang, H. Oe, A. Fujimoto,
 346 Y. Yonenaga, H. Sakashita, M. Imamura, Hexokinase II and VEGF
 347 expression in liver tumors: correlation with hypoxia-inducible factor 1
 348 α and its significance, J. Hepatol. 40 (2004) 117–123.
 349 [15] Z.F. Yang, R.T.P. Poon, J. To, D.W. Ho, S.T. Fan, The potential
 350 role of hypoxia inducible factor 1 α in tumor progression after
 351 hypoxia and chemotherapy in hepatocellular carcinoma, Cancer Res.
 352 64 (2004) 5496–5503.
 353 [16] J.M. Brown, W.R. Wilson, Exploiting tumour hypoxia in cancer
 354 treatment, Nat. Rev. Cancer 4 (2004) 437–447.
 355 [17] E.J. Yeo, Y.S. Chun, J.W. Park, New anticancer strategies targeting
 356 HIF-1, Biochem. Pharmacol. 68 (2004) 1061–1069.
 357 [18] O. Stoeltzing, M.F. McCarty, J.S. Wey, F. Fan, W. Liu, A. Belcheva,
 358 C.D. Bucana, G.L. Semenza, L.M. Ellis, Role of hypoxia-inducible
 359 factor 1 α in gastric cancer cell growth, angiogenesis, and vessel
 360 maturation, J. Natl. Cancer Inst. 96 (2004) 946–956.
 361 [19] F.N. Ko, C.C. Wu, S.C. Kuo, F.Y. Lee, C.M. Teng, YC-1, a novel
 362 activator of platelet guanylate cyclase, Blood 84 (1994) 4226–4233.
 363 [20] E.J. Yeo, Y.S. Chun, Y.S. Cho, J. Kim, J.C. Lee, M.S. Kim, J.W.
 364 Park, YC-1: a potential anticancer drug targeting hypoxia-inducible
 365 factor 1, J. Natl. Cancer Inst. 95 (2003) 516–525.
 366 [21] J.D. Oliner, K.W. Kinzler, P.S. Meltzer, D.L. George, B. Vogelstein,
 367 Amplification of a gene encoding a P53-associated protein in human
 368 sarcomas, Nature 358 (1992) 80–83.
 369 [22] A. Slack, Z. Chen, R. Tonelli, M. Pule, L. Hunt, A. Pession, J.M.
 370 Shohet, The P53 regulatory gene MDM2 is a direct transcriptional
 371 target of MYCN in neuroblastoma, Proc. Natl. Acad. Sci. USA 102
 372 (2005) 731–736.
 373 [23] S.W. Wang, S.L. PanL, J.H. Guh, H.L. Chen, D.M. Huang, Y.L.
 374 Chang, S.C. Kuo, F.Y. Lee, C.M. Teng, YC-1 [3-(5'-Hydroxymethyl-
 375 2'-furyl)-1-benzyl Indazole] exhibits a novel antiproliferative effect
 376 and arrests the cell cycle in G0-G1 in human hepatocellular
 377 carcinoma cells, J. Pharmacol. Exp. Ther. 312 (2005) 917–925.
 378 [24] H.K. Hsu, S.H. Juan, P.Y. Ho, Y.C. Liang, C.H. Lin, C.M. Teng,
 379 W.S. Lee, YC-1 inhibits proliferation of human vascular endothelial
 380 cells through a cyclic GMP-independent pathway, Biochem. Phar-
 381 macol. 66 (2003) 263–271.
 382

- 383 [25] W.C. Chiang, C.M. Teng, S.L. Lin, Y.M. Chen, T.J. Tsai, B.S.
384 Hsieh, YC-1-inhibited proliferation of rat mesangial cells through
385 suppression of cyclin D1-independent of cGMP pathway and
386 partially reversed by p38 MAPK inhibitor, *Eur. J. Pharmacol.* 517
387 (2005) 1–10.
- 388 [26] Y.T. Huang, S.L. Pan, J.H. Guh, Y.L. Chang, F.Y. Lee, S.C. Kuo,
389 C.M. Teng, YC-1 suppresses constitutive nuclear factor-kappaB
390 activation and induces apoptosis in human prostate cancer cells, *Mol.*
391 *Cancer Ther.* 4 (2005) 1628–1635.
- 392 [27] K. Inoki, M.N. Corradetti, K.L. Guan, Dysregulation of the TSC-
393 mTOR pathway in human disease, *Nat. Genet.* 37 (2005) 19–24.
- [28] D.A. Guertin, D.M. Sabatini, An expanding role for mTOR in
394 cancer, *Trends Mol. Med.* 11 (2005) 353–361. 395
- [29] D. Chen, M. Li, J. Luo, W. Gu, Direct interactions between HIF-1
396 alpha and Mdm2 modulate p53 function, *J. Biol. Chem.* 278 (2003)
397 13595–13598. 398
- [30] R. Ravi, B. Mookerjee, Z.M. Bhujwala, C.H. Sutter, D. Artemov, Q.
399 Zeng, L.E. Dillehay, A. Madan, G.L. Semenza, A. Bedi, Regulation
400 of tumor angiogenesis by p53-induced degradation of hypoxia-
401 inducible factor 1alpha, *Genes Dev.* 14 (2000) 34–44. 402
- [31] A.H. Truong, D. Cervi, J. Lee, Y. Ben-David, Direct transcriptional
403 regulation of MDM2 by Fli-1, *Oncogene* 24 (2005) 962–969. 404
405

UNCORRECTED PROOF