

Title	Assimilation of phytate-phosphorus by the extracellular phytase activity of tobacco (<i>Nicotiana tabacum</i>) is affected by the availability of soluble phytate
Author(s)	Lung, SC; Lim, BL
Citation	Plant And Soil, 2006, v. 279 n. 1-2, p. 187-199
Issued Date	2006
URL	http://hdl.handle.net/10722/54340
Rights	Creative Commons: Attribution 3.0 Hong Kong License

Editorial Manager(tm) for Plant and Soil
Manuscript Draft

Manuscript Number: PLSO1724R1

Title: Assimilation of phytate-phosphorus by the extracellular phytase activity of tobacco (*Nicotiana tabacum*) is affected by the availability of soluble phytate.

Article Type: Manuscript

Section/Category:

Keywords: inositol hexakisphosphate; phosphate; phosphorus; phytase; phytate; tobacco.

Corresponding Author: Dr. Boon Leong Lim, PhD

Corresponding Author's Institution: The University of Hong Kong

First Author: Shiu-Cheung Lung

Order of Authors: Shiu-Cheung Lung; Boon Leong Lim, PhD

Manuscript Region of Origin:

Abstract: Phytate, the major organic phosphorus in soil, is not readily available to plants as a source of phosphorus (P). It is either complexed with cations or adsorbed to various soil components. The present study was carried out to investigate the extracellular phytase activities of tobacco (*Nicotiana tabacum* variety GeXin No.1) and its ability to assimilate external phytate-P. Whereas phytase activities in roots, shoots and growth media of Pi-fed 14-day-old seedlings were only 1.3-4.9% of total acid phosphatase (APase) activities, P starvation triggered an increase in phytase secretion up to 914.9 mU mg⁻¹ protein, equivalent to 18.2% of total APase activities. Much of the extracellular phytase activities were found to be root-associated than root-released. The plants were not able to utilize phytate adsorbed to sand, except when insoluble phytate salts were preformed with Mg²⁺ and Ca²⁺ ions for supplementation. Tobacco grew better in sand supplemented with Mg-phytate salts (31.9 mg dry weight plant⁻¹; 0.68% w/w P concentration) than that with Ca-phytate salts (9.5 mg plant⁻¹; 0.42%), presumably due to its higher solubility. We conclude that insolubility of soil

phytate is the major constrain for its assimilation. Improving solubility of soil phytate, for example, by enhancement of citrate secretion, may be a feasible approach to improve soil phytate assimilation.

Manuscript: PLSO 1724

Authors(s): Lung S.C. and Lim B.L.

Title: Assimilation of phytate-phosphorus by the extracellular phytase activity of tobacco (*Nicotiana tabacum*) is affected by the availability of soluble phytate

Revision according to the referees' comments:

Editor's comments

1. Authority of the species has been included in abstract and materials in the revised manuscript (p.2, line 6).
2. We verified the numbers carefully and amended the numbers on p.2, line 14 and p.15, line 7.
3. The abbreviation of liter (L) has been changed accordingly.

Referee 2

1. The focus of this study is to investigate the ability of tobacco plants to assimilate external IHP, and to study whether the lack of a secretory phytase and/or the solubility of IHP are the factors that constrain IHP-P assimilation. While the adsorption of IHP to goethite (Ognalaga, et al, 1994, Soil Sci. Soc. Am. J. 58, 332-337) and Al/Fe precipitates (Shang et al, 1992, Geoderma, 53, 1-14) have been characterized, no report on how IHP is sorbed to sand is available from the literature. Our data confirmed the sorption of IHP to sands but it is beyond our capacity to speculate how it was absorbed in this study.
2. Referee 2 raised an interesting question of why plants express phytase despite the "inaccessibility" of soil phytate in an evolutionary point of view. First of all, while our data indicated absorbed phytate is not readily available to plant, it does not necessarily mean that soil IHP-P is absolutely not utilizable. In fact, as cited by Referee 2 (Page 21, Lines 1-10), small amount of IHP can be detected in soil solution (Espinosa et al., 1999). This compound might be phytase-labile. In addition, the present study also demonstrated considerable plant growth on Ca-IHP and Mg-IHP salts, indicating that the precipitation reaction limits IHP solubility but does not completely make it unavailable to plant phytase.

Furthermore, secretory plant phytases induced during P deprivation are not necessarily monospecific towards phytate. They could exhibit activities towards a broad range of P-monoesters and diesters.

3. The words “for the existence” in Page 3 Line 20 have been deleted in the revised manuscript.
4. As suggested by the reviewer, the erroneous statement in Page 4 Line 9 “as the existence of IHP in soil solution has never been reported (Hayes et al., 2000b)” has been removed. A statement “While in soil leachate, 77% of the P compounds is orthophosphate, and IHP constitutes only 3.3% of the total P (Espinosa et al., 1999).” was included in the discussion section in Page 21 Line 6. This article directly detected various P species, including IHP, using anion exchange column and HPLC. Another study (Turner et al., 2002) cited by Referee 2 employed an indirect method of phosphatase hydrolysis to characterize soil organic P. We think that the study of Turner et al. (2002) is less conclusive because the fungal phytase employed in the study was not monospecific towards IHP. In addition, no explanation has been provided to account for the absence of phytase-specific P compounds in moist soil extracts. Hence, we decide not to cite this reference.
5. Persulfate digestion is commonly used for the determination of IHP content in a number of publications. Our calculation was based on an IHP standard curve, where various concentrations of IHP were processed by the same procedure. Therefore, even if the persulfate method does not recover IHP completely, the determination of IHP should be accurate.

Referee 3

1. The research gap of the present study has been stated in page 5 lines 11-15 of the revised manuscript.
2. The recipe of Murashige and Skoog (1962) medium suggested that approximately 1 mM P_i is optimal for growth of tobacco plants. Our preliminary experiments showed that ten-fold reduction of this P supply resulted in P_i deficiency symptoms of plants including reduced shoot biomass and lower shoot-to-root ratio. Thus, 0.1 mM P_i , in addition to no-P, was chosen as a mean to study phytase activities under P_i deficiency. The situation of sterile hydroponic culture appears to be different from the field conditions, where acid phosphatase-labile organic P compounds can

be a P source to support plant growth, even though the existence of P_i in soil solution is in micromolar range. In fact, Anderson (1980) has well documented the continual introduction of organic P compounds into soil.

3. The reference to Microcal Origin software has been provided in Page 10 Line 7 in the revised manuscript.

1 Title: Assimilation of phytate-phosphorus by the extracellular phytase activity of tobacco
2 (*Nicotiana tabacum*) is affected by the availability of soluble phytate.

3

4 Shiu-Cheung Lung and Boon L. Lim*

5 *Department of Zoology, The University of Hong Kong, Pokfulam Road, Hong Kong, China*

6

7 * Corresponding Author

8 Address: Department of Zoology, The University of Hong Kong, Pokfulam Road, Hong Kong,

9 China

10 Tel: 852-22990826

11 Fax: 852-25599114

12 E-mail: bllim@hkucc.hku.hk

13

14 Number of text pages: 30

15 Number of tables: 2

16 Number of figures: 5

17

18 Key words: inositol hexakisphosphate, phosphate, phosphorus, phytase, phytate, tobacco

1 **Abstract**

2

3 Phytate, the major organic phosphorus in soil, is not readily available to plants as a source
4 of phosphorus (P). It is either complexed with cations or adsorbed to various soil components.
5 The present study was carried out to investigate the extracellular phytase activities of tobacco
6 (*Nicotiana tabacum* variety GeXin No.1) and its ability to assimilate external phytate-P.
7 Whereas phytase activities in roots, shoots and growth media of P₁-fed 14-day-old seedlings
8 were only 1.3-4.9% of total acid phosphatase (APase) activities, P starvation triggered an
9 increase in phytase secretion up to 914.9 mU mg⁻¹ protein, equivalent to 18.2% of total APase
10 activities. Much of the extracellular phytase activities were found to be root-associated than
11 root-released. The plants were not able to utilize phytate adsorbed to sand, except when
12 insoluble phytate salts were preformed with Mg²⁺ and Ca²⁺ ions for supplementation. Tobacco
13 grew better in sand supplemented with Mg-phytate salts (31.9 mg dry weight plant⁻¹; 0.68%
14 w/w P concentration) than that with Ca-phytate salts (9.5 mg plant⁻¹; 0.42%), presumably due
15 to its higher solubility. We conclude that insolubility of soil phytate is the major constrain for
16 its assimilation. Improving solubility of soil phytate, for example, by enhancement of citrate
17 secretion, may be a feasible approach to improve soil phytate assimilation.

1 **Introduction**

2

3 Phosphorus (P) is a macronutrient that constitutes vital molecules such as nucleic acids,
4 phospholipids and sugar phosphates in all living organisms. Terrestrial plants generally meet
5 their P requirement by the uptake of soil P in inorganic form (P_i ; Marschner, 1995). However,
6 majority of the soil P (50-80% of the total) exists as organic compounds (Turner et al., 2002),
7 which are unavailable to plants unless mineralization takes place. Plants have adapted to
8 mineralize organic P compounds, primarily by the secretion of acid phosphatases (APases),
9 which are a broad classification of hydrolytic enzymes catalyzing the breakdown of
10 P-monoesters with acid pH optima (Vincent et al, 1992). Despite the active APase secretion,
11 particularly in response to P_i starvation (Duff et al., 1994), earlier report (Anderson, 1980) has
12 indicated the relatively rare occurrence of APase-labile P species in soil, including nucleic
13 acids (up to 2.4% of organic P) and phospholipids (less than 5% of organic P). By comparison,
14 inositol hexakisphosphates (IP_6), a stable compound highly invulnerable to chemical and
15 enzymatic degradation, is the predominant soil organic P that can constitute more than 50% of
16 organic P and 25% of total P (Anderson, 1980).

17

18 In nature, IP_6 mainly exists in *myo*- form (phytate; IHP) as a product of a common
19 sequestrating activity in plants that generate a metabolic pool of P storage in seeds and grains,
20 whereas the origin of other stereoisomers (*scyllo*-, *neo*- and *chiro*- forms) in soil remains

1 controversial (Turner et al., 2002). Despite the abundance of IHP in soil, whether it contributes
2 to plant P nutrition is still in question. Earlier studies demonstrated that IHP is of poor
3 availability to a wide range of plants, including maize, wheat, legumes, grasses and pasture
4 species, cultivated on sterile agar, sand or sand-vermiculate media (Findenegg and Nelemans,
5 1993; Hayes et al., 2000; Richardson et al., 2000). Adams and Pate (1992) is the only group
6 that has demonstrated plant growth with IHP comparable to the P_i -fed controls. Nonetheless,
7 one of the critical requirements for the assimilation of IHP-P by plants is considered to be its
8 availability in soil (Adams and Pate, 1992). In fact, the dominance of IHP over other forms of
9 organic compounds in soil takes into account their strengthened stabilities through adsorption
10 and precipitation reactions (Turner et al., 2002).

11
12 Hydrolytic enzymes that catalyze the cleavage of phosphomonoester bonds in IHP are
13 collectively known as phytases. In plant kingdom, the roles of phytases have not yet been well
14 understood. Currently purified and characterized plant phytases are APases that possess high
15 specific activities against IHP. According to the sequence homologies, they are either classified
16 as histidine acid phosphatases (HAP) or purple acid phosphatases (PAP), primarily discovered
17 in maize (Hegeman and Grabau, 2001) and soybean (Maugenest et al., 1997), respectively. The
18 temporal expression of both phytases at the early stage of germination may be of relevance for
19 the mobilization of stored IHP-P to nourish plant growth, in agreement with earlier reports that
20 demonstrated increasing phytase activities in growing seedlings and germinating pollens (Goel

1 and Sharma, 1979; Lin and Dickinson, 1985; Mandal et al., 1974; Walker, 1974). Many plant
2 phytases have also been described in roots (Asmar, 1997; Hayes et al., 1999; Hübel and Beck,
3 1996; Li et al., 1997a, b; Richardson et al., 2000). Whilst maize root phytase was confined to
4 the endodermis (Hübel and Beck, 1996), P_i starvation elicited phytase expression to higher
5 levels in roots of some other plant species (Hayes et al., 1999; Li et al., 1997a, b), suggesting
6 that these enzymes may be pertinent to external P acquisition. Extracellular phytase activities
7 have also been reported in several plant species (Asmar, 1997; Hayes et al., 1999; Li et al.,
8 1997a; Richardson et al., 2000).

9
10 Although adsorption and precipitation reaction of IHP in soil environment have long been
11 hypothesized to be the reasons that limited IHP-P assimilation by plants, solid evidence is still
12 scarce. Thus, the present study aimed to investigate the ability of tobacco plants to assimilate
13 adsorbed and precipitated IHP, and to study whether the lack of a secretory phytase and/or the
14 insolubility of IHP are the limiting factors that constrain IHP-P assimilation by tobacco plants.

1 **Materials and methods**

2

3 *Plant cultivation*

4 Tobacco (*Nicotiana tabacum*) variety 'GeXin No1' was obtained from the Shanghai
5 Institute of Plant Physiology and Ecology. Seeds were surface-sterilized in 20% (v/v) Clorox
6 for 20 min, followed by several rinses with sterile, deionized water. For sand culture, sterilized
7 seeds were germinated on P-deficient MS agar for 10 days. The seedlings (1 plant pot⁻¹ in
8 triplicates) were transplanted onto 250 g quartzsand or riversand (0.1-1.0 mm sieved size),
9 which had been acid-washed and rinsed thoroughly with deionized water, oven-dried (110 °C;
10 72 h) and mixed with 60 mL of sterile nutrient solution: 1.5 mM KNO₃, 1.2 mM Ca(NO₃)₂, 0.4
11 mM NH₄NO₃, 0.5 mM MgSO₄, 0.3 mM K₂SO₄, 0.3 mM (NH₄)₂SO₄, 1.5 μM MnSO₄, 1.5 μM
12 ZnSO₄, 1.5 μM (NH₄)₆Mo₇O₂₄, 0.5 μM CuSO₄, 0.5 μM NaB₄O₇, 27 μM Fe-EDTA(Na) and 5
13 mM MES, pH 6.0. The original water content of the sand was maintained by irrigation with
14 P-deficient nutrient solution every other day. For sterile hydroponic culture, sterilized seeds
15 were sown on 20 mL of liquid MS medium (about 50 seeds plate⁻¹) and maintained aseptically.
16 IHP used in plant culture was purchased from Sigma, St. Louis, USA in the form of
17 dodecasodium salt (cat no. P3168). All plants were cultivated in controlled environment
18 cabinet with a day/night regime of 16/8 h, 22/16 °C, and photon flux density of 100 μmol m⁻²
19 s⁻¹.

20

1 *Phytase and acid phosphatase assays*

2 Plants were grown in sterile hydroponic culture with modified P_i concentrations (0, 0.1 or
3 1 mM) for 14 days. Phytase and APase activities were measured in (i) crude shoot, root extracts
4 and growth media, (ii) intact roots (root-associated activities), and (iii) root-bath solutions
5 (root-released activities), following the same procedure described previously (Asmar, 1997;
6 Richardson et al., 2000) with minor modifications. Phytase and APase activities were measured
7 in MES/Ca buffer (15 mM MES, 0.5 mM $CaCl_2$, pH 5.5) at room temperature using either 1
8 mM Na-IHP or 10 mM *p*-nitrophenyl phosphate (*p*NPP) as substrates, respectively. Phytase
9 activity assays were conducted over 1 h, unless otherwise specified, and terminated by equal
10 volume of 10% (w/v) TCA. The liberated P_i was quantified by the molybdenum-blue reaction
11 (Murphy and Riley, 1962). APase activities were assayed over 30 min and terminated by equal
12 volume of 0.25 M NaOH. P_i release was derived from the equimolar liberation of *p*-nitrophenol
13 (*p*NP) product, quantified spectrophotometrically at 405 nm over standard solutions. One
14 enzyme unit (1 U) is defined as the activity that releases 1 μ mol of phosphate per min under
15 the specified assay conditions. In experiment (i), shoot and root materials were ground in 15
16 mM MES, 1 mM PMSF, pH 5.5, and the crude extracts were centrifuged (20 min, 14000 rpm).
17 The supernatant was dialyzed against MES/Ca buffer to minimize the intrinsic P_i background
18 for assays. Soluble proteins in the growth media were concentrated by ultrafiltration using the
19 centrifugal filter units (Millipore, Bedford, MA) prior to phytase and APase assays. The total
20 soluble protein content was determined by Bradford assay (Bradford, 1976). Experiments (ii)

1 and (iii) were conducted aseptically in a laminar flow hood. Prior to assays, residual P_i on the
2 root surface was eliminated by several rinses with MES/Ca buffer and the excess solution was
3 removed by blotting with sterile tissue paper. In experiment (ii), the roots portions of the intact
4 seedlings (50 plants plate⁻¹) were incubated in 20 mL of MES/Ca buffer that contained 1 mM
5 Na-IHP for 2 h and the liberated P_i was quantified in the incubated solution. Corrections were
6 made to account for P efflux/influx (Asmar, 1997; Richardson et al., 2000) by replacing
7 Na-IHP with different P_i concentrations (0-2.4 mg P L⁻¹) in the incubated solution. In
8 experiment (iii), root-released phytase was collected by incubating the root portions of the
9 intact seedlings (50 plants plate⁻¹) in 20 mL of MES/Ca buffer for 4 h. The incubated solution
10 was then filtered (0.45 μ m) to remove the sloughed-off cells (Asmar, 1997) and concentrated
11 by ultrafiltration using the centrifugal filter units (Millipore, Bedford, MA), prior to phytase
12 assays.

13

14 *Sorption experiments*

15 IHP-sorption capacities of the quartzsand and riversand used for plantation were estimated
16 by incubating 7.5 mL of 1.794 mM Na-IHP solution, with (pH 6 by HCl) or without (pH 11)
17 pH adjustment, in 25 g sand samples (i.e. 100 mg P kg⁻¹) at room temperature. At t = 0, 5 min,
18 20 min, 1 h, 3 h, 8 h and 24 h, 50 μ L of the soil solution was collected. The solid residue in the
19 soil solution was removed by centrifugation (5 min, 6,000 g) and the solution P concentration
20 was determined by persulfate digestion (US EPA, 1993).

1

2 *Sand culture*

3 Tobaccos were grown in sand culture for 20 days with (i) adsorbed IHP, (ii) supplemented
4 IHP salts, or (iii) supplemented NaH_2PO_4 as the sole P source. At harvest, shoot and root
5 materials were oven-dried at 80 °C for 72 h for dry weight (DW) determination. To measure
6 shoot P concentration, the dried shoot samples were subsequently ashed at 500 °C for 16 h and
7 the residues dissolved in 0.1 M HCl (8.75 mg sample mL^{-1}). The P_i concentration of the acid
8 solution was determined using the molybdenum-blue reaction (Murphy and Riley, 1962). For
9 experiment (i), 250 g quartzsand or riversand was incubated with Na-IHP solution (100 mg P
10 kg^{-1} sand) at room temperature for 24 h. The sand-IHP complex was extensively rinsed with
11 distilled water until no IHP was detected in the water extract by persulfate digestion (US EPA,
12 1993). The samples were made homoionic by equilibration with 3 changes of 0.1 M CaCl_2 or
13 MgCl_2 solutions, followed by extensive rinses with distilled water until no chloride was
14 detected in the water extract as indicated by the absence of white precipitate upon addition of
15 1% (w/v) AgNO_3 . The IHP-adsorbed sand samples were then air-dried and used for plantation.
16 For experiment (ii), insoluble Ca- and Mg-IHP salts were prepared by incubating 10 mM
17 Na-IHP solution with 50 mM CaCl_2 or MgCl_2 solution at room temperature, respectively. The
18 precipitated salts were allowed to settle and the supernatant decanted off. The solid was then
19 washed thrice with distilled water, resuspended in nutrient solution and added to the sand (100
20 mg P kg^{-1} sand) for plant growth. Experiment (iii) was carried out to estimate the plant growth

1 responses to different P_i dosage. NaH_2PO_4 was added from a 0.1 M stock solution to the
2 nutrient solution, which was subsequently added to the sand to provide 0, 12.5, 25, 50, 100,
3 200 or 400 mg P kg^{-1} sand for plant growth. The DW and shoot P concentration of plants
4 grown at each supplied P_i level were fitted to the Mitscherlich model (Ware et al., 1982) with
5 the equation $y = a + be^{-cx}$, such that y = shoot DW or P concentration, x = supplied P_i level, and
6 a , b , c = constants, where a represents the maximum yield, with the aid of analysis program
7 Microcal Origin version 7.5 (Microcal Software Inc., Northampton, USA).

8

9 *Assays with elevated cation levels*

10 Plants were grown in sterile hydroponic culture with modified P_i concentrations (0, 0.1 or
11 1 mM) for 14 days. Crude root extracts were obtained by the same procedure described above
12 and the intrinsic P_i was removed by dialysis against Ca^{2+} -free MES (15 mM, pH 5.5) buffer.
13 Ca^{2+} and Mg^{2+} ions were added to the samples in the form of chloride solutions and the phytase
14 activities were assayed as described above. The final divalent ion concentrations in the reaction
15 mixtures were 0, 0.5, 1, 2, 4 and 8 mM.

1 **Results**

2

3 *Phytase and acid phosphatase activities*

4 Phytase and APase activities of 14-day-old tobacco seedlings grown under different P_i
5 treatments (0, 0.1 or 1 mM) were assayed in the crude root and shoot extracts, as well as in the
6 growth media. The APase activities in shoots (1,029-1,449 mU mg⁻¹ protein) were generally
7 higher than that in roots (503-618 mU mg⁻¹ protein; Table 1). Comparable APase activity levels
8 were observed in the crude root extracts across different P_i treatments, whereas a 10-fold
9 reduction in P_i supply (i.e. 0.1 mM) significantly ($P<0.05$) increased the APase activities in
10 shoots by 40%, although similar difference between no-P plants and P_i -fed plants (1 mM) was
11 statistically insignificant ($P\geq 0.05$; Table 1). APase was believed to be one of the predominant
12 secretory proteins, by a record of 4,323-5,017 mU mg⁻¹ protein in the growth media. Subtle
13 (10-16%) but significant ($P<0.05$) increase in the activities was observed from the growth
14 media of no-P and low-P plants (0.1 mM), compared to the P_i -fed controls (Table 1). Specific
15 phytase activities of crude root and shoot extracts from P_i -fed plants were only 15.6 and 13.8
16 mU mg⁻¹ protein (Table 1). Whilst the reduction of P_i supply only increased the specific
17 phytase activities in roots by 47% (no-P plants) and 68% (low-P plants), the stimulatory
18 response to P deprivation was more prominent in shoot, where specific phytase activities were
19 boosted by 4.2-fold under low-P condition and 3.1-fold in no-P condition, concomitant with an
20 increase in secretory phytase activities by 1.9-fold and 3.4-fold, respectively (Table 1). Across

1 different P_i treatment, the increase in phytase/APase ratio was generally proportional to that in
2 specific phytase activities, resulting from the fact that changes in APase activities were
3 relatively insubstantial (Table 1).

4

5 *Root-associated and root-released phytase activities*

6 Root-associated activities were assayed by the incubation of intact roots with IHP solution
7 and subsequent quantification of P_i liberation, whereas *in vitro* assays for root-released
8 activities were conducted after collection of secretory proteins in root-bath solution (Table 2).

9 Confounded by the possible P_i efflux or uptake during the assay period, the presented
10 root-associated activities have been subjected to corrections made by the estimation of P_i
11 efflux/influx rate using a series of control plants incubated with different P_i concentrations over
12 the assay period. While P efflux from P_i -fed plants was evident (data not shown), a near
13 balance between P efflux and influx rates was observed in no-P and low-P plants, depending on
14 the P_i concentration of the incubated solution (data not shown). Extracellular phytase activities
15 of 14-day-old tobacco seedlings were primarily root-associated, ranging from 819 mU g⁻¹ root
16 DW in P_i -fed plants to 1,367 mU g⁻¹ root DW in no-P plants (Table 2). By comparison,
17 root-released activities were hardly detectable, even though secretory proteins have been
18 collected over an extended period of 4 h and extensively concentrated for *in vitro* assays (Table
19 2).

20

1 *IHP sorption capacities of sand media*

2 Time course experiments demonstrated that the riversand and quartzsand used for
3 plantation in the present study possessed contrasting IHP sorption capacities. Whereas no IHP
4 was lost in the soil solution when incubated with quartzsand for up to 24 h, IHP concentration
5 in the soil solution at the starting 100 mg P kg^{-1} level was immediately halved upon 10-min and
6 30-min incubation with riversand at pH 6 and 11, respectively (Figure 1). Near-equilibria of
7 IHP adsorption at 84 and 65 mg P kg^{-1} riversand were reached within 3-h incubation at pH 6
8 and 11, respectively (Figure 1).

9

10 *P assimilation in sand culture*

11 IHP-P assimilation by tobacco plants was investigated by the cultivation of 10-day-old
12 seedlings in sand culture with IHP supplied in various forms for 20 days. For comparison, plant
13 growth response to different P_i dosage, ranging from 0 to 400 mg P kg^{-1} sand in the form of
14 NaH_2PO_4 , is depicted in Figure 2. Results from all growth parameters, i.e. shoot and root DW,
15 and shoot P concentration, were generally consistent in response to different P_i dosage.
16 Maximum yield of quartzsand-grown plants was achieved by supplying 100 mg P kg^{-1} sand, at
17 which the shoot DW and P concentration were 55 mg plant^{-1} and 0.79% (w/w), respectively
18 (Figures 2A and 2C). Even though the shoot P concentration could be further boosted to 0.90%
19 (w/w) by the supply of 200 mg P kg^{-1} sand, the yield of dry biomass was notably inhibited at
20 this level, concomitant with a drastic decline in both DW and shoot P concentration of plants

1 grown with 400 mg P kg⁻¹ sand (Figures 2A and 2C). Plant growth under more than 100 mg P
2 kg⁻¹ sand dosage of NaH₂PO₄ is considered to be inhibited by high Na⁺ concentration,
3 corroborated by 35% decrease in shoot DW when doubling the Na⁺ concentration by the
4 replacement of NaH₂PO₄ with Na₂HPO₄ at a level of 100 mg P kg⁻¹ sand (data not shown). On
5 the contrary, same P_i dosage (0-400 mg P kg⁻¹ sand) for plant growth on riversand did not result
6 in any obvious inhibition by high Na⁺ concentration (Figures 2B and 2D). Maximum yield of
7 shoot DW (69 mg plant⁻¹) and P concentration [0.68% (w/w)] were achieved at 200 and 400
8 mg P kg⁻¹ sand, respectively (Figures 2B and 2D). Utilization of IHP-P by tobacco plants were
9 studied by the provision of IHP in either adsorbed or supplemented forms. Adsorbed IHP was
10 prepared by pre-incubation of the sand media with IHP solution, followed by extensive rinses,
11 with or without subsequent Ca²⁺ and Mg²⁺ homo-ionization. In agreement with the results from
12 sorption experiments that quartzsand possessed poor IHP-sorption capacity (Figure 1), DW and
13 shoot P concentrations of the plants grown on quartzsand pre-incubated with IHP was
14 indifferent from that of the no-P controls (Figures 3 and 4). Nonetheless, similar results were
15 obtained from the plants grown on the high IHP-sorbing riversand, indicating that adsorbed
16 IHP was a poor P source for plant growth (Figures 3 and 4). Alternatively, IHP was directly
17 supplemented for plant growth in the form of cationic complexes. As preliminary experiments
18 showed that supplementation of IHP in the forms of Mn²⁺, Al³⁺, Cu²⁺, Zn²⁺ and Fe³⁺ salts
19 demonstrated toxicity to the plants and thus unsuitable for use (data not shown), results from
20 the provision of Na⁺ (soluble), Ca²⁺ and Mg²⁺ (insoluble) salts was only reported here.

1 Riversand-grown plants did not respond notably to any of the cationic complexes, with only
2 subtle but significant ($P<0.05$) improvement in both shoot DW and P concentration by
3 supplementation of Mg-IHP, but not Na-IHP and Ca-IHP, compared to the no-P controls
4 (Figures 3 and 4). On the contrary, P nutrition of the quartzsand-grown plants was considerably
5 improved to different extents by the supplementation of either one of the 3 IHP salts (Figures 3
6 and 4). The utilization of insoluble Ca-IHP was as efficient as of soluble Na-IHP by the
7 quartzsand-grown plants, yielding approximately 9-11 mg shoot DW plant⁻¹ and 0.42% (w/w)
8 shoot P concentration (Figures 3 and 4; upper panels), equivalent to the P_i-fed plants at 5 and 23
9 mg P kg⁻¹ sand levels, respectively (Figures 3 and 4; lower panels). The effects of IHP
10 supplementation on quartzsand-grown plants were much more prominent in the form of Mg²⁺
11 salts, yielding 32 mg shoot DW plant⁻¹ and 0.68% (w/w) shoot P concentration (Figures 3 and
12 4; upper panels), equivalent to P_i-fed plants at 26 and 57 mg P kg⁻¹ sand levels, respectively
13 (Figures 3 and 4; lower panels). The retarded plant growth on Na-IHP, compared to Mg-IHP,
14 was believed to be the outcome of extremely high Na⁺ concentration from the commercial
15 preparation of IHP in the form of Na⁺ salts (12 Na:IHP molar ratio; Sigma-Aldrich P3168)., as
16 similar growth retardation was observed when NaH₂PO₄ standard was supplied in high
17 concentrations (Figure 2).

18

19 *Effects of divalent ions on root phytase activities*

20 Severe hindrance of plant phytase activities in the presence of elevated divalent ion

1 levels could not be observed from *in vitro* assays. Ca^{2+} and Mg^{2+} ions were found to possess
2 slightly enhance and inhibitory effects on the phytase activities in the crude root extracts
3 from 14-day-old tobacco seedlings in a concentration-dependent manner, respectively. Root
4 phytase activities were notably heightened by up to 4 mM CaCl_2 (Figure 5). In the presence of
5 1 mM CaCl_2 , the relative activities were peaked at 110%, 120% and 130% in P_i -fed plants,
6 low-P plants and no-P plants, respectively, although the activities were declined when IHP
7 started to precipitate with Ca^{2+} at 8 mM concentration (Figure 5). Increasing Mg^{2+}
8 concentrations resulted in a relatively steady declination in root phytase activities of seedlings
9 grown under all P_i treatments, down to 53-59% activities left at the precipitating level (8 mM;
10 Figure 5). The results indicated that tobacco plant phytases did possess ability to hydrolyze IHP
11 complexed with Ca^{2+} and Mg^{2+} , which was in line with our observation in sand culture.
12 However, while the insoluble Ca and Mg-IHP were in direct contact with the crude root
13 phytase preparations in *in vitro* assays, only solubilized IHP from Ca and Mg-complexes was
14 accessible to the root-associated phytases, which constitute predominant extracellular phytase
15 activities (Table 2). Thus, even though Mg^{2+} was found to be more inhibitory to plant phytase
16 than Ca^{2+} (Figure 5), plants grown with less stable Mg-IHP complex outperformed that with
17 Ca-IHP in sand culture (Figures 3 and 4).

1 Discussion

2

3 *Intrinsic plant acid phosphatases and phytases*

4 While tobacco plants (*Nicotiana tabacum*) have extensively been used in transgenic
5 phytase researches, from recombinant protein production (Ullah et al., 1999; Verwoerd et al.,
6 1995) to functional studies of microbial enzymes on plant growth promotion (George et al.,
7 2005; Lung et al., 2005; Yip et al., 2003), little is known about the intrinsic phytase in this
8 model plant. The present study is the first attempt to demonstrate low P_i-inducible extracellular
9 phytase activities in tobacco seedlings. It is generally perceived that plant phytases are APases
10 that are able to hydrolyze IHP as a member of their broad spectra of P-monoester substrates,
11 hence, phytase activities were compared with APase activities (against *p*NPP) in the present
12 study, and in other publications (Asmar, 1997; Hayes et al., 1999; Hübel and Beck, 1996; Li et
13 al., 1997a, b; Richardson et al., 2000). The ubiquitous APase functions in plant P metabolism
14 (Duff et al., 1994) account for the measured shoot APase in P_i-fed plants (1029 mU mg⁻¹
15 protein; Table 1), which was accompanied with considerable phytase activity (13.8 mU mg⁻¹
16 protein; Table 1). P_i limitation elicited subtle increase in shoot and secretory APase ($P < 0.05$;
17 Table 1), probably as an adaptation for P_i acquisition and recycling (Duff et al., 1994), which is
18 concomitant with more prominent increase in phytase activities (Table 1). The lesser extent of
19 increase in phytase activities of roots suggested that the elicited phytase might not be
20 accumulated intracellularly, but targeted to secretion pathway, corroborated by the 3 to 4-fold

1 increase in secretory phytase activity (Table 1). Even though precise comparison of phytase
2 activities among different plant species described elsewhere is confounded by the experimental
3 variations (plant age, assay pH and temperature, etc.), specific root phytase activities under P_i
4 deficiency were mostly ranged from tens to hundreds milli-unit mg^{-1} protein, i.e. 1 to 5 % of
5 total APase activities (Hayes et al., 1999; Hübel and Beck, 1996; Li et al., 1997b; Richardson
6 et al., 2000). The specific phytase activity of tobacco roots (26.2 mU mg^{-1} at 0.1 mM P_i supply;
7 Table 1) was comparable to that of wheat (26.7 mU mg^{-1} ; Richardson et al., 2000) and a bit
8 lower than that of maize (88 mU mg^{-1} ; Hübel and Beck, 1996), whereas the extreme abundance
9 of root phytase (440 mU mg^{-1} ; Li et al., 1997a) in tomato was a rare occurrence. Nonetheless,
10 the ultimate location of the phytase is one of the determining factors governing its level in
11 roots. While some phytases were confined intracellularly (Hübel and Beck, 1996), phytase
12 secretion has been described in many species (Asmar, 1997; Hayes et al., 1999; Li et al., 1997a;
13 Richardson et al., 2000). In tobacco, the extracellular phytase activities were found to be
14 mostly root-associated (Table 1). Whereas the possibility of cell wall-protein association
15 should not be ruled out, the abundance of secretory phytase in growth media suggested that the
16 negligible root-released activity levels might be due to inadequate time for secretory protein
17 accumulation, as protein release from the apoplastic space to the external medium is a passive
18 process.

19

20 *Adsorption and precipitation: constrains for IHP-P assimilation*

1 While many plant species are incapable of utilizing IHP-P in medium (Hayes et al, 2000;
2 Richardson et al, 2000; Richardson et al., 2001), the notable growth of other plant species on
3 IHP as the sole P source was found to be medium-dependent. For instance, *Trifolium*
4 *subterraneum* which had demonstrated substantial IHP utilization ability in agar medium was
5 found to exhibit poor P nutrition on IHP-supplemented sand-vermiculate medium (Hayes et al.,
6 2000). Similarly, IHP, an equally good source of P as KH_2PO_4 for the growth of lupins in
7 quartzsand, was found to be the poorest form of P supplementation to the soil, compared to P_i ,
8 glycerophosphate and ribonucleic acid (Adams and Pate, 1992). In some experiments that
9 utilized commercial phytase for stimulating plant growth, 10-fold higher IHP and phytase
10 dosages were required in soil than in quartzsand to produce significant effects (Findenegg and
11 Nelemans, 1993). Observations from the present study also clearly demonstrated the poor plant
12 growth on adsorbed IHP (Figures 3 and 4), in despite of the notable extracellular phytase
13 activities in hydroponic culture (Table 1 and 2). In fact, all these findings reflect the real
14 situation in soil environment that IHP rarely exists in enzyme-labile form. Due to the dense
15 negative charges, IHP is strongly adsorbed to insoluble salts such as Fe and Al oxides and
16 hydrous oxides (Shang et al., 1992), as well as clay minerals such as illite, kaolinite and
17 montmorillonite (Celi et al., 1999). Apart from adsorption, availability of IHP in soil is
18 believed to be severely limited by cationic precipitation, such as Fe- and Al-IHP in acid soils
19 and Ca-IHP in alkaline soils (Turner et al., 2002). Maenz et al. (1999) ranked the cation-IHP
20 complexes in descending order of stability: $\text{Fe}^{2+} \sim \text{Zn}^{2+} \sim \text{Fe}^{3+} > \text{Mn}^{2+} > \text{Ca}^{2+} > \text{Mg}^{2+}$, which is

1 in line with the present observation that Mg-IHP was a better source of P for plant growth than
2 Ca-IHP (Figures 3 and 4), assuming that plant phytases preferably act on solubilized and
3 dissociated IHP from the complexes. Instead of direct attack on the insoluble cation-IHP
4 complexes, IHP solubilization is believed to be a crucial step for plant phytase hydrolysis,
5 supported by the better assimilation of P from Mg and Ca-IHP by plants on quartzsand (Figures
6 3 and 4), compared to riversand that possessed a higher potency to bind soluble IHP (Figure 1).
7 It is believed that assimilation of P from Mg and Ca-IHP is an innate ability of plants, as the
8 IHP storage in seeds and grains was mostly associated with Mg^{2+} and Ca^{2+} ions to form
9 complexes commonly referred to as phytins. However, despite the remarkable plant growth on
10 Mg and Ca-IHP in the present study (Figures 3 and 4), these unstable complexes, once
11 introduced into soil from plant remains, could be immediately stabilized by the prevalent ions
12 (e.g. Fe^{3+} , Al^{3+} , Zn^{2+} , etc) in soil, rendering them invulnerable to enzymatic hydrolysis. In view
13 of the three-dimensional structures of the three characterized phytase families, beta-propeller
14 phytase (BPP; Shin et al., 2001), HAPs (Lim et al., 2000; Liu et al., 2004) and plant PAPs
15 (Strater et al., 1995; Schenk et al., 2005), the cleavage of phosphomonoester bonds requires the
16 precise coordination of the phosphate ions with the amino acid residues in the active site. For
17 instance, the substrate binding site of *E. coli* HAP, located in a deep indentation inside the
18 molecule, requires all six deprotonated phosphate for hydrolytic reaction (Lim et al., 2000).
19 Thus, the adsorption and precipitation reactions may render the IHP molecule inaccessible to
20 the substrate-binding pockets of phytases for mineralization.

1

2 *Implications on GM approaches for IHP-P assimilation*

3 From the present study, it is showed that tobacco plants do secrete phytase under P_i
4 starvation, but this adaptation may not satisfy plant P nutrition if the availability of soluble
5 phytate is limited. In soil, the majority of IHP is adsorbed to soil components or is precipitated
6 with cations. While in soil leachate, 77% of the P compounds is orthophosphate, and IHP
7 constitutes only 3.3% of the total soluble P (Espinosa et al., 1999). The minute amount of
8 soluble IHP in soil solution may still be a sparing source of P for plant, rationalizing phytase
9 secretion under P starvation. Some recent studies attempted to explore the usability of the IHP
10 pool in bulk soil. Transgenic strategies to improve IHP-P assimilation in plants have been
11 demonstrated by the extracellular secretion of recombinant phytases, including HAP (George et
12 al., 2005; Richardson et al., 2001; Zimmermann et al., 2003), BPP (Lung et al., 2005; Chan et
13 al., 2005) and PAP (Xiao et al., 2005). While IHP utilization ability was significantly improved
14 in all of these transgenic plants grown in sterile agar or hydroponic culture, no enhancement of
15 IHP assimilation by these transgenic plants has been reported in soil. Even though
16 Zimmermann et al. (2003) were able to demonstrate better P nutrition of transgenic plants in a
17 non-sterile quartz-loess-peat mixture by irrigating plants with 100 μ M IHP solution twice a
18 week, no assessment of the P-sorption capacities of this substrate has been performed. Perhaps,
19 supplementation of IHP solution was adequate to provide sufficient soluble IHP for
20 mineralization. In fact, George et al. (2005) have shown that transgenic plants secreting HAP

1 did not lead to an improved P nutrition from P-deficient soils without the amendment with IHP,
2 coincident with our observations that transgenic plants secreting BPP (Lung et al., 2005) did
3 not exhibit better growth performance in a high P-sorbing sand (unpublished data). Thus, in
4 agreement with the present study, both native and recombinant phytase exudation from plant
5 roots may not ensure significant IHP-P assimilation in the soil environment. To circumvent the
6 obstacle of phytate insolubility, phytase secretion should be supplemented by some means, for
7 instance, organic acid exudation, to desorb or solubilize the adsorbed or precipitated IHP.
8 Hence, co-exudation of recombinant phytase and organic anions (e.g. citrate, malate and
9 oxalate) may be a promising strategy to improve P acquisition from IHP in soil.

1 **Acknowledgement**

2

3 This research was supported by the Research Grants Council (project HKU 7335/04M) of the

4 Hong Kong Special Administrative Region Government, China.

1 **References**

- 2 Adams M A and Pate J S 1992 Availability of organic and inorganic forms of phosphorus to
3 lupins (*Lupinus* spp.). *Plant Soil* 145, 107-113.
- 4 Anderson G 1980 Assessing organic phosphorus in soils. In *The role of phosphorus in*
5 *agriculture*. Eds. F E Khasawneh, E C Sample and E J Kamprath. pp 411-432. American
6 Society of Agronomy.
- 7 Asmar F 1997 Variation in activity of root extracellular phytase between genotypes of barley.
8 *Plant Soil* 195, 61-64.
- 9 Bradford M M 1976 A rapid and sensitive method for the quantification of microgram
10 quantities of protein utilizing the principle of protein-dye banding. *Anal. Biochem.* 72,
11 248-254.
- 12 Celi L, Lamacchia S, Marsan F A and Barberis E 1999 Interaction of inositol hexaphosphate on
13 clays: adsorption and charging phenomena. *Soil Sci.* 164, 574-585.
- 14 Chan W L, Lung S C and Lim B L 2005 Properties of beta-propeller phytase expressed in
15 transgenic tobacco. *Protein Expr. Purif.* (in press)
- 16 Duff S M G, Sarath G and Plaxton W C 1994 The role of acid phosphatases in plant
17 phosphorus metabolism. *Physiol. Plant* 90, 791-800.

- 1 Espinosa M, Turner B L and Haygarth P M 1999 Preconcentration and separation of trace
2 phosphorus compounds in soil leachate. *J. Environ. Qual.* 28, 1497-1504.
- 3 Findenegg G R and Nelemans J A 1993 The effect of phytase on the availability of P from
4 *myo*-inositol hexaphosphate (phytate) for maize roots. *Plant Soil* 154, 189-196.
- 5 George T S, Simpson R J, Hadobas P A and Richardson A E 2005 Expression of a fungal
6 phytase gene in *Nicotiana tabacum* improves phosphorus nutrition of plants grown in
7 amended soils. *Plant Biotechnol. J.* 3, 129-140.
- 8 Goel M and Sharma C B 1979 Multiple forms of phytase in germination cotyledons of
9 *Cucurbita maxima*. *Phytochemistry* 18, 1939-1942.
- 10 Hübel F and Beck E 1996 Maize root phytase. Purification, characterization, and localization
11 of enzyme activity and its putative substrate. *Plant Physiol.* 112, 1429-1436.
- 12 Hayes J E, Richardson A E and Simpson R J 1999 Phytase and acid phosphatase activities in
13 extracts from roots of temperate pasture grass and legume seedlings. *Plant Physiol.* 26,
14 801-809.
- 15 Hayes J E, Richardson A E and Simpson R J 2000 Components of organic phosphorus in soil
16 extracts that are hydrolysed by phytase and acid phosphatase. *Biol. Fertil. Soils* 32,
17 279-286.

- 1 Hegeman C E and Grabau E 2001 A novel phytase with sequence similarity to purple acid
2 phosphatases is expressed in cotyledons of germinating soybean seedlings. *Plant Physiol.*
3 126, 1598-1608.
- 4 Li M, Osaki M, Honma M and Tadano T 1997a Purification and characterization of phytase
5 induced in tomato roots under phosphorus-deficient conditions. *Soil. Sci. Plant Nutr.* 43,
6 179-190.
- 7 Li M, Osaki M, Rao I M and Tadano T 1997b Secretion of phytase from the roots of several
8 plant species under phosphorus-deficient conditions. *Plant Soil* 195, 161-169.
- 9 Lim D, Golovan S, Forsberg C W and Jia Z 2000 Crystal structures of *Escherichia coli* phytase
10 and its complex with phytate. *Nat. Struct. Biol.* 7, 108-113.
- 11 Lin J J and Dickinson D B 1985 Studies of phytic acid and phytase in germinating lily pollen.
12 *Plant Physiol.* 77, S46.
- 13 Liu Q, Huang Q, Lei X G and Hao Q 2004 Crystallographic snapshots of *Aspergillus fumigatus*
14 phytase, revealing its enzymatic dynamics. *Structure (Camb.)* 12, 1575-1583.
- 15 Lung S C, Chan W L, Yip W, Wang L, Yeung E C and Lim B L 2005 Secretion of
16 beta-propeller phytase from tobacco and *Arabidopsis* roots enhances phosphorus
17 utilization. *Plant Sci.* 169, 341-349.

- 1 Maenz D D, Engele-Schaan C M, Newkirk R W and Classen H L 1999 The effects of minerals
2 and mineral chelators on the formation of phytase-resistant and phytase-susceptible forms
3 of phytic acid in solution and in a slurry of canola meal. *Anim. Feed Sci. Technol.* 81,
4 177-192.
- 5 Mandal N C, Burman S and Biswas B B 1974 Isolation, purification and characterization of
6 phytase from germinating mung beans. *Phytochemistry* 13, 1047-1051.
- 7 Marschner H 1995 Mineral nutrition of higher plants. Academic Press, London.
- 8 Maugenest S, Martinez I and Lescure A M 1997 Cloning and characterization of a cDNA
9 encoding a maize seedlings phytase. *Biochem. J.* 322, 151-157.
- 10 Murphy J and Riley J P 1962 A modified single solution method for the determination of
11 phosphate in natural waters. *Anal. Chim. Acta* 27, 31-36.
- 12 Richardson A E, Hadobas P A and Hayes J E 2000 Acid phosphomonoesterase and phytase
13 activities of wheat (*Triticum aestivum* L.) roots and utilization of organic phosphorus
14 substrates by seedlings grown in sterile culture. *Plant Cell Environ.* 23, 397-405.
- 15 Richardson A E, Hadobas P A and Hayes J E 2001 Extracellular secretion of *Aspergillus*
16 phytase from *Arabidopsis* roots enables plants to obtain phosphorus from phytate. *Plant J.*
17 25, 641-649.

- 1 Schenk G, Gahan L R, Carrington L E, Mitic N, Valizadeh M, Hamilton S E, de Jersey J and
2 Guddat L W 2005 Phosphate forms an unusual tripodal complex with the Fe-Mn center of
3 sweet potato purple acid phosphatase. Proc. Natl. Acad. Sci. U.S.A. 102, 273-278.
- 4 Shang C, Stewart J W B and Huang P M 1992 pH effect on kinetics of adsorption of organic
5 and inorganic phosphates by short-range ordered aluminum and iron precipitates.
6 Geoderma 53, 1-14.
- 7 Shin S, Ha N C, Oh B C, Oh T K and Oh B H 2001 Enzyme mechanism and catalytic property
8 of beta propeller phytase. Structure (Camb.) 9, 851-858.
- 9 Strater N, Klabunde T, Tucker P, Witzel H, Krebs B, Donegan K K, Palm C J, Fieland V J,
10 Porteous L A, Ganio L M, Schaller D L, Bucuo L Q and Seider R J 1995 Crystal structure
11 of a purple acid phosphatase containing a dinuclear Fe(III)-Zn(II) active site. Science 268,
12 1489-1492.
- 13 Turner B L, Papházy M J, Haygarth P M and McKelvie I D 2002 Inositol phosphates in the
14 environment. Phil. Trans. R. Soc. Lond. B 357, 449-469.
- 15 Ullah A H J, Sethumadhavan K, Mullaney E J, Ziegelhoffer T and Phillips S A 1999
16 Characterization of recombinant fungal phytase (*phyA*) expressed in tobacco leaves.
17 Biochem. Biophys. Res. Commun. 264, 201-206.
- 18 US EPA 1993 Methods of chemical analysis of water and wastes. Environ. Monit. Systems

- 1 Lab., Cincinnati, OH.
- 2 Verwoerd T C, van Paridon P A, van Ooyen A J J, van Lent J W M, Hoekema A and Pen J 1995
- 3 Stable accumulation of *Aspergillus niger* phytase in transgenic tobacco leaves. Plant
- 4 Physiol. 109, 1199-1205.
- 5 Vincent J B, Crowder M W and Averill B A 1992 Hydrolysis of phosphate monoesters: a
- 6 biological problem with multiple chemical solutions. Trends Biochem. Sci. 17, 105-110.
- 7 Walker K A 1974 Change in phytic acid and phytase during early development of *Phaseolus*
- 8 *vulgaris* L. Planta 116, 91-98.
- 9 Ware G O, Ohki K and Moon L C 1982 The Mitscherlich plant growth model for determining
- 10 critical nutrient deficiency levels. Agron. J. 74, 88-91
- 11 Xiao K, Harrison M J and Wang Z Y 2005 Transgenic expression of a novel *M. truncatula*
- 12 phytase gene results in improved acquisition of organic phosphorus by Arabidopsis. Planta,
- 13 in press.
- 14 Yip W, Wang L, Cheng C, Wu W, Lung S and Lim B L 2003 The introduction of a phytase
- 15 gene from *Bacillus subtilis* improved the growth performance of transgenic tobacco.
- 16 Biochem. Biophys. Res. Commun. 310, 1148-1154.
- 17 Zimmermann P, Zardi G I, Lehmann M, Zeder C, Amrhein M, Frossard E and Bucher M 2003

- 1 Engineering the root-soil interface via targeted expression of a synthetic phytase gene in
- 2 trichoblasts. *Plant Biotechnol. J.* 1, 353-360.

1 **FIGURE LEGENDS**

2

3 *Figure 1.* Time course of IHP adsorption on quartzsand (QS) and riversand (RS). Na-IHP
4 solution (pH 6 or 11) was incubated with sand sample at 100 mg P kg⁻¹. At different time
5 intervals, soil solution was collected for IHP quantification by persulfate digestion.

6

7 *Figure 2.* Growth parameters of plants with different P_i dosage in sand culture. Ten-day-old
8 tobacco seedlings were grown on quartzsand (QS) or riversand (RS) with different NaH₂PO₄
9 concentrations (mg P kg⁻¹ sand) for 20 days. Shoot/root dry weights (DW) and shoot P
10 concentration were determined. Shoot DW of plants grown in 0-100 mg P kg⁻¹ QS (A) and
11 0-400 mg P kg⁻¹ RS (B) were fitted to the Mitscherlich equation. For shoot P concentration,
12 data from 0-200 mg P kg⁻¹ QS (C) and 0-400 mg P kg⁻¹ RS (D) were fitted to the Mitscherlich
13 equation.

14

15 *Figure 3.* Dry weights of plants grown with different IHP treatments in sand culture. IHP was
16 supplied in either adsorbed or supplemented forms as the sole P source. Shoot and root dry
17 weights were shown in the upper panel. Equivalent soil P_i levels were estimated from the
18 corresponding Mitscherlich equations depicted in Figure 2A and 2B.


19


1 *Figure 4.* Shoot P concentration of plants grown with different IHP treatments in sand culture.
2 Shoot P concentrations were shown in upper panel. Equivalent soil P_i levels were estimated
3 from the corresponding Mitscherlich equations depicted in Figure 2C and 2D.



4


5 *Figure 5.* Effects of divalent ions on root phytase activities. Tobacco seeds were sown on liquid
6 MS medium with different P_i treatments (0, 0.1 or 1 mM) and grown for 14 days. Phytase
7 activities were assayed in crude root extracts in the presence of different Ca²⁺ or Mg²⁺
8 concentrations.

9









line figure 5

[Click here to download high resolution image](#)




Table 1. Phytase and acid phosphatase activities of crude root / shoot extracts and growth media from 14-day-old tobacco plants

P treatment	Phytase activity (mU mg ⁻¹ protein)			Acid phosphatase activity (mU mg ⁻¹ protein)			Phytase / acid phosphatase activity (%)		
	Root extracts	Shoot extracts	Growth media	Root extracts	Shoot extracts	Growth media	Root extracts	Shoot extracts	Growth media
1 mM P _i	15.6 ± 0.9 ^c	13.8 ± 2.5 ^c	210.0 ± 14.1 ^c	589 ± 62 ^{ab}	1029 ± 120 ^b	4323 ± 95 ^b	2.7 ± 0.3 ^b	1.3 ± 0.1 ^c	4.9 ± 0.4 ^c
0.1 mM P _i	26.2 ± 1.6 ^a	72.1 ± 9.8 ^a	600.9 ± 14.8 ^b	618 ± 75 ^a	1449 ± 216 ^a	4761 ± 186 ^a	4.3 ± 0.3 ^a	5.2 ± 0.3 ^a	12.6 ± 0.8 ^b
No-P	23.0 ± 2.0 ^b	56.3 ± 9.2 ^b	914.9 ± 18.6 ^a	503 ± 23 ^b	1232 ± 138 ^{ab}	5017 ± 118 ^a	4.6 ± 0.4 ^a	4.6 ± 0.4 ^b	18.2 ± 0.5 ^a

Each observation is the mean of at least 3 replicates ± 1 standard deviation.

Within each column, the values marked by different letters (a, b, c) are significantly different ($P < 0.05$).

Table 2. Extracellular phytase activities of 14-day-old tobacco plants

P treatment	Extracellular phytase activities	
	Root-associated activity (mU g ⁻¹ root DW)	Root-released activity (mU mg ⁻¹ root DW h ⁻¹)
1 mM P _i	819 ± 82 ^c	< 0.01
0.1 mM P _i	1,134 ± 55 ^b	< 0.01
No-P	1,367 ± 62 ^a	< 0.01

Each observation is the mean of at least 3 replicates ± 1 standard deviation.

Within each column, the values marked by different letters (a, b, c) are significantly different ($P < 0.05$).