



<b>Title</b>	<b>Effects of smoking on healing response to non-surgical periodontal therapy: A multilevel modelling analysis</b>
<b>Author(s)</b>	<b>Wan, CP; Leung, WK; Wong, MCM; Wong, RMS; Wan, P; Lo, ECM; Corbet, EF</b>
<b>Citation</b>	<b>Journal Of Clinical Periodontology, 2009, v. 36 n. 3, p. 229-239</b>
<b>Issued Date</b>	<b>2009</b>
<b>URL</b>	<b><a href="http://hdl.handle.net/10722/55450">http://hdl.handle.net/10722/55450</a></b>
<b>Rights</b>	<b>Creative Commons: Attribution 3.0 Hong Kong License</b>

1  
2  
3 **Effects of smoking on healing response to non-surgical periodontal**  
4  
5 **therapy: A multilevel modeling analysis**  
6  
7  
8  
9

10 Chi Pui Wan, W. Keung Leung\*, May C. M. Wong, Ruby M. S. Wong, Peng Wan,

11  
12 Edward C. M. Lo, Esmonde F. Corbet

13  
14  
15 Faculty of Dentistry, The University of Hong Kong, Hong Kong SAR, China  
16  
17  
18  
19

20 Running title: smokers' healing: multilevel analysis  
21  
22  
23  
24

25 Keywords: Periodontitis; Smoking; Models, Statistical;  
26  
27  
28  
29

30 **\*Corresponding author:**  
31

32 W. K. Leung, Room 3B39, 34 Hospital Road, Prince Philip Dental Hospital, Faculty of  
33

34 Dentistry, The University of Hong Kong, Hong Kong SAR, China. Telephone:  
35

36 +852-28590417; Fax: +852-28587874; E-mail: [ewkleung@hkucc.hku.hk](mailto:ewkleung@hkucc.hku.hk)  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Date of resubmission: November 21, 2008

### **Conflict of interest and sources of funding statement**

The authors declare that they have no conflict of interests.

The work described in this paper was partially supported by grants from the Research Grants Council of the Hong Kong Special Administrative Region, China (HKU 7331/00M) and Merck Sharp & Dohme (Asia) Limited (26004077).

For Peer Review

1  
2  
3 *Wan CP, Leung WK, Wong MCM, Wong RMS, Wan P, Lo ECM, Corbet EF. Effects of*  
4 *smoking on healing response to non-surgical periodontal therapy: A multilevel modeling*  
5 *analysis. J Clin Periodontol*  
6  
7  
8  
9

10  
11  
12 **Abstract**  
13

14  
15 **Aim:** To investigate factors predicting non-surgical periodontal treatment responses  
16  
17  
18 using multilevel multiple regression.  
19

20  
21 **Methods:** 40 men (mean 45.6 years) were recruited. 20 were smokers. 12-month  
22  
23  
24 reduction in probing pocket depth (PPD) and gain in probing attachment level (PAL) of  
25  
26  
27 5814 sites were analyzed with **594 being initially diseased sites (initial PPD  $\geq$**   
28  
29  
30 **5mm).**  
31

32  
33  
34 **Results:** Variance Component models showed site level variations contributed  
35  
36  
37 about 70-90% of the total variance. About 10% reduction of the total variations  
38  
39  
40 of PPD reduction in initially diseased sites was achieved with the inclusion of the  
41  
42  
43 **10 predictors in the multilevel multiple regression. Multilevel multiple regression**  
44  
45  
46 showed that three predictors - subject-level: non-smokers; tooth-level: anterior  
47  
48  
49 teeth; site-level: sites without plaque at baseline, were significantly associated  
50  
51  
52 with greater reduction in PPD in initially diseased sites over the 12 months study  
53  
54  
55 period. (p<0.05). No consistent predictor was found for PAL gain.  
56

57  
58 **Conclusion:** Multilevel analysis was applied on periodontal treatment response data.  
59

60  
Smokers showed less favorable PPD reduction at deep sites after non-surgical

1  
2  
3  
4 periodontal therapy.  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

1  
2  
3  
4 Smoking is considered as a well-established risk factor for periodontal diseases, a  
5  
6 chronic infectious disorder caused by bacterial plaque characterized by destruction of  
7  
8 tooth supporting tissue. Smokers have increased risks of experiencing periodontal  
9  
10 attachment loss (Grossi et al. 1994, Haffajee and Socransky 2001a, Susin et al. 2004,  
11  
12 Ng & Leung, 2006), radiographic bone loss (Grossi et al. 1995, Bergstrom 2004,  
13  
14 Baljoon et al. 2005) and tooth loss post-treatment (Leung et al. 2006, Matuliene et al.  
15  
16 2008). Smokers are found to harbor a higher prevalence of periodontal pathogens  
17  
18 (Haffajee & Socransky, 2001b, van Winkelhoff et al. 2001).  
19  
20  
21  
22  
23

24 Apart from alterations of the periodontal microflora, smoking has been shown to  
25  
26 adversely **affect** the host immune response in various respects, including impaired  
27  
28 neutrophil function (Mariggio et al. 2001, Güntsch et al. 2006), lowered  
29  
30 immunoglobulin production (Mooney et al. 2001, Apatzidou et al. 2005), reduced  
31  
32 fibroblast function (Raulin et al. 1988), altered inflammatory mediator production  
33  
34 (Boström et al. 1998, 1999; Giannopoulou et al., 2003) and vasoconstrictive effects of  
35  
36 tissue exposed to cigarette smoke (Mirbod et al. 2001).  
37  
38  
39

40 Non-surgical mechanical periodontal therapy, including oral hygiene instruction,  
41  
42 scaling and root planing, is an effective treatment modality for periodontal disease  
43  
44 (Van der Weijden & Timmerman 2002, **Sanz & Teughels, 2008**); however, numerous  
45  
46 studies have indicated that smokers generally undergo less favorable improvements in  
47  
48 response to non-surgical therapy (Preber & Bergstrom, 1986; Preber et al., 1995;  
49  
50 Renvert et al., 1998; Jin et al., 2000). A systematic review evaluating the effect of  
51  
52 smoking on non-surgical periodontal therapy (Labriola et al., 2005) found that the  
53  
54 mean difference in probing pocket depth (PPD) reduction with an initial probing depth  
55  
56 of 5mm or more would be 0.433mm favoring non-smokers. On the other hand, the  
57  
58 same meta-analysis showed that there was no evidence of a difference observable in  
59  
60

1  
2  
3 clinical attachment level gain between smokers and non-smokers after non-surgical  
4  
5 periodontal therapy, although a review of clinical evidence (Heasman et al. 2006)  
6  
7 suggests that the majority of studies do show that smokers gain less clinical attachment  
8  
9 gain in response to periodontal therapy. It is agreed that achieving optimal treatment  
10  
11 responses to non-surgical periodontal therapy in smokers is a challenging task and that  
12  
13 the treatment outcome of the therapy may vary from patient to patient and also vary  
14  
15 among different teeth and tooth sites. It would be beneficial to understand factors at  
16  
17 patient, tooth and site levels that may affect these variations in treatment response in  
18  
19 both smokers and non-smokers.  
20  
21  
22  
23

24 Since the early 1990s, researchers have questioned the utility of single level  
25  
26 statistical analysis **of site-level or tooth-level data** in periodontal clinical trials  
27  
28 **because the correlations among sites and/or teeth within subjects invalidates**  
29  
30 **these methods. In applying single level statistical analysis to periodontal data,**  
31  
32 **many** earlier publications **chose to** present average sites' measurements generated on a  
33  
34 subject level. **However, such an approach** may not explicitly reflect the site-specific  
35  
36 nature of periodontal disease (Albandar & Goldstein 1992, Gilthorpe et al. 2000a,  
37  
38 Gilthorpe et al. 2000b, Gilthorpe et al. 2001). Application of multilevel modeling  
39  
40 analysis, which takes the clustering effect of periodontal research data into  
41  
42 consideration, may provide a more accurate explanation of the natural hierarchical  
43  
44 structure of clinical findings of periodontitis and the healing responses after  
45  
46 periodontal therapy. Two reports lately, adopted such approach in their periodontal trial  
47  
48 data analysis (Tomasi et al. 2007, Matuliene et al. 2008).  
49  
50  
51  
52  
53

54  
55 In the present prospective study, clinical healing responses of two groups of male  
56  
57 Chinese subjects: smokers or non-smokers - matched according to age, pre-operative  
58  
59 oral hygiene levels and periodontal disease severity - were recorded after non-surgical  
60  
periodontal therapy. The aim of this study was to compare the 12-month healing

1  
2  
3 response of male Chinese smokers and non-smokers with chronic periodontitis after  
4 non-surgical mechanical periodontal therapy using multilevel modeling analysis. The  
5 clinical data would be analyzed at site level. The null hypothesis of this clinical trial is  
6 that there is no difference in healing responses after non-surgical mechanical  
7 periodontal therapy of periodontitis affected male Chinese smokers and non-smokers.  
8  
9  
10  
11  
12  
13  
14  
15  
16

## 17 **Materials and methods**

### 18 **Sample size determination**

19  
20 This clinical study targeted subjects with chronic periodontitis who were otherwise  
21 systemically healthy. Sample size for the study was computed as follows. In a study  
22 among the same local population, patients with chronic periodontitis showed 4.6mm of  
23 probing pocket depth (PPD) reduction at 12 months after non-surgical therapy, with  
24 standard deviation (SD) of 1.6mm (Tong et al. 2003). Assuming that the SD would be  
25 the same for smokers and an expected difference of PPD reduction at the initially  
26 diseased sites between smokers and non-smokers of 2 mm, 20 subjects in each group  
27 were required to enable such a difference to be detected.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

### 43 **Patient selection and screening**

44  
45 New male patients attending the Reception Clinic of the Prince Philip Dental Hospital,  
46 Faculty of Dentistry, The University of Hong Kong and satisfying the inclusion criteria  
47 were recruited to participate in the study. The target sample size was at least 22  
48 subjects for each group, to allow for retention of 20 subjects in each group at 12  
49 months. For inclusion, patients had to be free of systemic disease, not undergoing  
50 orthodontic treatment, and displaying the following features:  
51  
52  
53  
54  
55  
56  
57  
58

- 59 1. 35- to 64-years-old ethnic Chinese with untreated chronic periodontitis  
60



2. Smokers with a smoking habit of  $\geq 10$  cigarettes per day for at least 10 years and expressing no interest in quitting smoking in the coming 12 months
3. Non-smokers with a smoking history of never having smoked
4. At least 16 standing teeth, with at least 1 tooth having PPD  $\geq 5$  mm in each quadrant, excluding the third molars.

Subjects were excluded if the patient interview revealed:

1. Known systemic diseases
2. History of taking systemic antibiotics in the preceding 30 days
3. History of dental treatment, other than oral hygiene instructions, in the preceding 30 days

The target sample size for each group was secured six months after the commencement of recruitment.

### **Patient management and non-surgical mechanical periodontal treatment**

The clinical study was carried out in the Periodontology Clinic, Prince Philip Dental Hospital, Faculty of Dentistry, The University of Hong Kong. Emergency treatment such as extraction, caries stabilization, initial endodontic therapy, if necessary, was completed before the non-surgical periodontal treatment. Six tooth-sites (mesio-buccal, mid-buccal, disto-buccal, mesio-lingual, mid-lingual, and disto-lingual) of each standing tooth were included in this study. One member of the research team (W.K.L.) checked the eligibility of all subjects and that all necessary **pre-treatment** preparations had been carried out. Receptionists of the Periodontology Clinic were then instructed to arrange the non-surgical periodontal treatment appointments (4-6 visits) under local anesthesia for all subjects to be delivered by a group of six

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

experienced dental hygienists within an 8-week period. Both smokers and non-smokers received **the** same non-surgical periodontal treatment, namely oral hygiene instruction regarding brushing and interdental cleaning, followed by quadrant-wise debridement under local anesthesia. Two research group members (P.W. and R.M.S.W.) at the end of the last dental hygienist treatment appointment independently clinically assessed the quality of the hygienists' care to ensure the completeness of the non-surgical periodontal therapy.

Any residual periodontal problems at conclusion of the study at 12 months, namely any sites with residual PPD  $\geq$  5 mm, were followed-up and appropriate periodontal treatment e.g. re-root planing or surgical treatment was arranged and delivered without delay. Smoking subjects were again reminded of the deleterious effects resulting from their continued smoking.

### Clinical examination

This was a 12-month prospective clinical study. Clinical parameters were obtained from the patients at baseline, and at 3, 6 and 12 months after completion of non-surgical therapy. All clinical examinations were performed by one examiner (C.P.W.).

Probing pocket depth (PPD) and probing attachment level (PAL) were measured and recorded for six sites of each tooth, excluding third molars. Custom-made poly-ethylene occlusal stents were made for each patient as reference guides for reproducibility of probing sites and for measurement of probing attachment level throughout the study. Except for initial baseline PAL data, which was collected using manual periodontal probe (PCP-UNC 15, Hu-Friedy probe®, Chicago, IL), each site was probed with an automated controlled-force periodontal probe, Florida Probe® (Florida Probe Co.). Probe tips were 0.45 mm in diameter and manufactured from

1  
2  
3 implant grade titanium. The resolution of 0.2 mm could be detected with controlled  
4  
5 force of 15g. Presence of plaque was recorded dichotomously as presence or absence  
6  
7 of plaque according to detection of plaque deposits determined by running the tip of a  
8  
9 periodontal probe along the tooth surface at the gingival margin of each site. Bleeding  
10  
11 on probing (BOP) was designated as positive if bleeding occurred within 10 seconds  
12  
13 after periodontal probing using the electronic probe.  
14  
15  
16  
17  
18

## 19 20 **Ethics**

21  
22 The research protocol was approved by the Ethics Committee, Faculty of Dentistry,  
23  
24 The University of Hong Kong. Written informed consent was obtained from all  
25  
26 participants before the commencement of the study.  
27  
28  
29  
30

## 31 32 **Data analysis**

### 33 34 **Routine statistical analysis**

35  
36 The data collected was entered into a computer and analyzed using the statistical  
37  
38 software package (SPSS). For comparing the difference in healing response between  
39  
40 smokers and non-smokers at the subject level, the primary efficacy measure was  
41  
42 **change in PPD and** change in PAL and the secondary efficacy measures included  
43  
44 PI%, BOP% and percentage of sites  $\geq 5.0$  mm. The significance level was set at  $p <$   
45  
46 0.0017 for multiple comparisons at the 3-, 6- and 12-month recalls within groups or  
47  
48 between groups. Differences between groups and between different time-points within  
49  
50 groups were tested by Mann-Whitney U test and Wilcoxon signed rank test  
51  
52 respectively.  
53  
54  
55  
56  
57  
58  
59  
60

## Multilevel analysis

In order to account for the hierarchical structure of periodontal disease measurements, site measurements clustered around individual teeth and then teeth clustered within subjects, analysis using a multilevel approach was adopted in this study (Gilthorpe et al. 2000b). PPD reductions at site level at 3 months, 6 months and 12 months (compared to baseline PPD) were analyzed using multilevel multiple regressions. A 3-level **random intercept regression** model was constructed: site at level 1, tooth at level 2 and subject at level 3. Variance Components models (with no independent variables included) were obtained initially to investigate the variance of the PPD reductions across all the 3 levels. **At different levels the random effects were assumed to be uncorrelated and followed normal distributions. Subsequently, ten independent variables with five on the subject-level, two on the tooth-level and three on the site-level were included in the multilevel multiple regression model. The five subject-level variables were: smoking (non-smoker vs. smoker), age (in years), number of missing teeth at baseline, % sites with plaque at baseline and % sites with BOP at baseline. The two tooth-level variables considered in the regression model were: the tooth position (posterior [premolars and molars] vs. anterior [incisors and canines]) and arch (lower vs. upper). The three site-level variables were: presence or absence of plaque at baseline, presence or absence of BOP at baseline, and surface (lingual vs. buccal). All the continuous variables were centered (subtracted from the mean) before the analysis. The analyses of the gain in PAL at 3, 6 and 12 months were performed in a similar manner: 3-level regression models were considered with ten independent variables. All the analyses were performed using the software MLwiN 2.1 (Rasbash et al. 2000). The level of significance was set at 0.05.**

In order to focus on the factors affecting the change of PPD and PAL of initially

1  
2 diseased sites (sites with PPD  $\geq$  5.0 mm at baseline), above-mentioned multilevel  
3  
4 multiple regressions were repeated for initially diseased sites only. Again, the level of  
5  
6  
7 significance was set to be at 0.05.  
8  
9

## 10 11 **Results**

### 12 13 **Routine statistical analysis**

#### 14 15 *Change of PPD and PAL at all sites*

16  
17 In the present study, 23 non-smokers and 23 smokers were recruited. Forty of the  
18  
19 enrolled subjects completed the study, 3 subjects being lost to follow-up in both the  
20  
21 smoker and the non-smoker groups. One smoker and three non-smokers could not  
22  
23 attend the scheduled recalls due to contemporaneous conflict with their job time-tables.  
24  
25 Two smokers quit smoking, one for personal reasons and the other having been  
26  
27 diagnosed to be suffering from hypertension was successfully counseled to quit  
28  
29 smoking by his physician.  
30  
31  
32  
33  
34

35  
36 Mean ages of the smokers and non-smokers who completed the study were  $46.2 \pm$   
37  
38  $6.8$  and  $45.0 \pm 5.9$  years, respectively. Regarding the tobacco consumption of smokers,  
39  
40 6 were light smokers while the **remaining** 14 were moderate smokers (Grossi et al.  
41  
42 1994). Their smoking-pack-years were  $20.8 \pm 8.7$ , ranging from 10 to 30. Mean  
43  
44 number of missing teeth (excluding third molars) was  $3.9 \pm 2.9$  teeth for smokers and  
45  
46  $3.7 \pm 2.8$  teeth for non-smokers ( $P > 0.05$ ). Other clinical data are shown in Table 1.  
47  
48 There was no difference between non-smokers and smokers in percentage of plaque,  
49  
50 mean full-mouth PPD, mean full-mouth PAL and percentage of sites with PPD  $\geq$  5mm  
51  
52 at baseline. Both groups showed poor oral hygiene and a high percentage of sites  
53  
54 with BOP at baseline, while smokers exhibited significantly less bleeding compared  
55  
56 with non-smokers ( $p = 0.003$ ).  
57  
58  
59  
60

Table 1 shows the change of subject level clinical parameters over the study

1  
2  
3 period. Throughout the course of the study, both non-smokers and smokers achieved  
4  
5 favorable improvements in their plaque control. This was demonstrated by significant  
6  
7 reductions of PI% at 3, 6 and 12 months compared to baseline in both groups. By 12  
8  
9 months, the mean PI% was reduced to less than 34%.

10  
11  
12 In addition, in response to non-surgical mechanical periodontal therapy, both  
13  
14 groups showed significant reductions in mean full-mouth BOP% compared to baseline.  
15  
16 By 12 months, the mean BOP% was reduced to less than 27%.

17  
18  
19 During the 12-month study period, full-mouth mean PPD in both groups was  
20  
21 found to be significantly reduced when compared to the baseline. Moreover, both  
22  
23 groups showed PAL gains compared to baseline. However, there was no significant  
24  
25 difference in mean full-mouth PPD reduction and mean full-mouth PAL gain between  
26  
27 non-smokers and smokers. Also, the proportion of sites with PPD  $\geq 5.0$  mm was  
28  
29 significantly reduced after the non-surgical periodontal therapy in both smokers and  
30  
31 non-smokers. However at 12 months, smokers showed less favorable results in terms  
32  
33 of significantly higher percentage residual pockets (PPD  $\geq 5.0$  mm) than non-smokers  
34  
35 (Table 1).  
36  
37  
38  
39  
40  
41  
42

#### 43 *Change of PPD and PAL at initially diseased sites*

44  
45 **For the 594 sites with initial PPD  $\geq 5.0$  mm, the mean PPD at these initially**  
46  
47 **diseased sites in smokers was  $5.85 \pm 0.48$  mm and in non-smokers was  $5.94 \pm 0.47$**   
48  
49 **mm. Both smokers and non-smokers showed significant reductions of probing**  
50  
51 **pocket depth at 3, 6 and 12 months when compared to baseline ( $p < 0.001$ ) (Table**  
52  
53 **1). In smokers, the PPD at initially diseased sites reduced from  $5.85 \pm 0.48$  mm at**  
54  
55 **baseline to  $3.00 \pm 0.80$  mm at 12 months. In non-smokers, the corresponding PPD**  
56  
57 **change was from  $5.94 \pm 0.47$  mm at baseline to  $2.49 \pm 0.50$  mm at 12 months**  
58  
59 **(Table 1). When comparing the two groups, non-smokers showed significantly**  
60

1  
2  
3 **greater PPD reduction at 6 and 12 months ( $p<0.01$ ) (Fig.1).**  
4

5 The change in PAL at initially diseased sites of the two groups is shown in Fig 2.  
6  
7 No significant difference between smokers and non-smokers was detected at any time  
8  
9 point.  
10

### 11 **Multilevel statistical analysis**

#### 12 *Change of PPD at all sites*

13  
14 Altogether, 5814 sites distributed on 969 teeth in these 40 subjects were included for  
15  
16 the analysis of reduction in PPD at 3, 6 and 12 months.  
17

18  
19 The overall mean reductions in PPD at 3, 6 and 12 months were 0.85 mm, 0.95  
20  
21 mm and 1.00 mm respectively (Table 2). The Variance Component models showed that  
22  
23 significant variations existed at all three levels of the multilevel structure (all 95%  
24  
25 confidence intervals did not cover the value of 0). Site-level variation contributed  
26  
27 about 80% of the total variation in reduction in PPD at 3, 6 and 12 months.  
28  
29

30  
31 **Ten independent variables were included in the multilevel multiple**  
32  
33 **regression and the random intercept models with significant variables only are**  
34  
35 **shown in Table 3. The intercept in the model for the reduction in PPD 3-month**  
36  
37 **was 0.62 mm. This indicates that the mean reduction in PPD at 3 months was**  
38  
39 **0.62 mm for buccal sites from lower anterior teeth with absence of plaque and**  
40  
41 **BOP at baseline in smokers with mean age of 45.58 years, with a mean 3.78**  
42  
43 **missing teeth and a mean 63.89% sites with BOP and 77.11% with plaque at**  
44  
45 **baseline.**  
46  
47

48  
49 **From the random intercept models for all sites there was no statistically**  
50  
51 **significant difference in PPD reduction between non-smokers and smokers**  
52  
53 **throughout the study period ( $p<0.05$ ).**  
54  
55

56  
57 **Consistently, sites on incisors and canines, on lingual aspects, sites with**  
58  
59  
60

1  
2  
3 presence of plaque and BOP at baseline, as well as sites from subjects with higher  
4  
5 percentages of sites with BOP showed significantly greater reduction in PPD at 3,  
6  
7 6 and 12 months.  
8

9  
10 The variances at each level were reduced by the inclusion of the ten  
11  
12 variables. The total variances of the models were reduced by 7%, 8% and 9%  
13  
14 respectively for reduction in PPD at 3, 6 and 12 months when compared to the  
15  
16 corresponding Variance Components models.  
17  
18

### 21 *Change of PAL at all sites*

22  
23 Again, 5814 sites distributed on 969 teeth in all the 40 subjects were included for the  
24  
25 analyses of gain in PAL at 3, 6 and 12 months.  
26  
27

28  
29 The overall mean gains in PAL at 3, 6 and 12 months were 0.24 mm, 0.30 mm and  
30  
31 0.37 mm respectively (Table 2). The Variance Component models showed that  
32  
33 significant variations existed at all three levels of the multilevel structure (all 95%  
34  
35 confidence intervals did not cover the value of 0) except for the tooth-level at 12  
36  
37 months. Site-level variation contributed from 80% to 90% of the total variation in gain  
38  
39 in PAL at 3, 6 and 12 months.  
40  
41

42  
43 From the regression models (Table 3), it was found that there was no  
44  
45 significant difference in the gain in PAL at 3, 6 and 12 months between the  
46  
47 smokers and non-smokers. Consistently, sites on lingual surfaces showed  
48  
49 significantly greater gains in PAL at 3, 6 and 12 months ( $p < 0.001$ ). Moreover,  
50  
51 sites on anterior teeth showed slightly greater PAL gain at 6 and 12 months ( $p$   
52  
53  $< 0.001$ ).  
54  
55

56  
57 The variations at the three levels were reduced by 0-30% with the inclusion  
58  
59 of the ten variables. The total variances of the models were reduced only by 2-4%  
60  
for the gain in PAL at 3, 6 and 12 months when compared to the corresponding



## Variance Components models.

### *Change in PPD at initially diseased sites*

Altogether, 594 sites with initial PPD  $\geq$  5mm, distributed on 324 teeth in these 40 subjects were included for the analyses of reduction in PPD of initially diseased sites at 3, 6 and 12-months.

The overall mean reductions in PPD of initially diseased sites at 3, 6 and 12 months were 2.55 mm, 2.77 mm and 3.16 mm respectively (Table 4). The Variance Component models showed that significant variations existed at all three levels of the multilevel structure (all 95% confidence intervals did not cover the value of 0) except for subject level at 3-month. Site-level variation contributed about 70% to 80% of the total variation in reduction in PPD at 3, 6 and 12 months.

**Similar to the analysis for all sites, 10 independent variables were included in the multilevel multiple regression, and the result of random intercept models are shown in Table 5. From the regression models, initially diseased sites of non-smokers consistently showed greater PPD reduction at 3, 6 and 12 months (0.41 mm, 0.79 mm and 0.68 mm respectively,  $p < 0.05$ ).**

**In accordance with analysis of all sites, initially diseased sites from anterior teeth were found to have undergone significantly greater reduction in PPD at 3, 6 and 12 months ( $p < 0.05$ ). Contrary to the results of the analysis of all sites, initially diseased sites on lingual aspects with presence of plaque at baseline showed less PPD reduction at 3, 6 and 12 months ( $p < 0.05$ ).**

**In the analysis for the initially diseased sites, the total variances of the models were reduced by only 9-13% respectively for reduction in PPD at 3, 6 and 12 months when compared to the corresponding Variance Components models.**

### *Change in PAL at initially diseased sites*

Those 594 initially diseased sites on 324 teeth in the 40 patients were included for the analyses of gain in PAL at 3, 6 and 12 months.

From the Variance Component models, the overall mean gains in PAL at 3, 6 and 12 months were 0.80 mm, 0.83 mm and 1.21 mm respectively (Table 4). Significant variations existed at tooth and sites levels but not subject level of the multilevel structure (all 95% confidence intervals did not cover the value of 0) at 3, 6 and 12 months. Site-level variation contributed most of the variation in gain in PAL at 3, 6 and 12 months, ranging from 75% to 80%.

**After the inclusion of the 10 variables, the total variances of the models were reduced by 2-5% for the gain in PAL at 3, 6 and 12 months when compared to the corresponding Variance Components models.**

From the regression models (Table 5), it was found that there was no significant difference between the smokers and non-smokers in the gain in PAL at 3, 6 and 12 months for initially diseased sites ( $p > 0.05$ ). Only subjects with higher percentage of sites with plaque at baseline showed slightly less PAL gain at 12 months ( $p < 0.05$ ).

For tooth level variables, only tooth position showed a significant effect on gain in PAL of initially diseased sites at 6 months. Sites from anterior teeth had significantly greater gain in PAL than sites on posterior teeth at 6 months ( $p < 0.05$ ).

For the site level, it was found that only sites with absence of plaque at baseline showed greater PAL gain at 3 and 6 months ( $p < 0.05$ ), while the effects of other variables were insignificant.

## Discussion

Previous studies have generally demonstrated that smokers have increased risk of periodontal destruction and less favorable healing in response to non-surgical periodontal therapy (Preber & Bergstrom 1986, Preber et al. 1995, Renvert et al. 1998, Jin et al. 2000). However, the factors affecting the variability of treatment outcomes among different smoking patients and at different sites within individual smokers are still not fully understood.

In much periodontal research statistical methods have been applied which generally ignore the fact that many observations are correlated, by combining all site observations into a mean value. Site level observations are not truly independent (Hujoel et al. 1990). Sites are clustered around a tooth and teeth are clustered in individuals. **It is therefore, inappropriate to analyse the site-level or subject level observations using single-level, univariate statistical methods since the correlation among sites and/or teeth within an individual invalidates these statistical methods.** Consequently, statistical analysis with assumption that the sites observations are independent would generate potentially misleading results (Tu et al., 2004).

Consequently, statistical analysis undertaken on the assumption that site observations are independent could generate potentially misleading interpretations of results (Tu et al. 2004).

A recent study employed a multilevel approach to investigate factors affecting the probability of “pocket closure” for diseased sites 3 months after two separate regimes of non-surgical periodontal therapy (Tomasi et al. 2007). However “pocket closure” is not the only healing response to non-surgical therapy. Therefore, the present study aimed, using multilevel modeling analysis, to investigate the possible factors affecting response of non-surgical periodontal therapy in male Chinese smokers and

1  
2  
3 non-smokers in terms of both PPD reduction and PAL gain.  
4

5 In the present study, results generated from traditional, routine statistical analysis  
6 are also presented. It was found that smokers showed less favorable responses after  
7 non-surgical therapy. At 12 months, smokers presented with a significantly higher  
8 percentage of residual pockets (Table 1). Additionally, smokers showed less PPD  
9 reduction in sites with initial PPD  $\geq$  5mm (Fig. 1). However, there was no statistically  
10 significant difference in the gain in PAL in initially diseased sites between smokers and  
11 non-smokers (Fig. 2). This is in agreement of a recent systematic review concerning  
12 effect of smoking on non-surgical therapy (Labriola et al. 2005), although a review of  
13 clinical evidence suggests that the majority of studies do show that clinical attachment  
14 gain in response to periodontal therapy is impaired in smokers (Heasman et al. 2006).  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

28 In order to account for the natural hierarchical structure of periodontal disease  
29 measurements, the present study adopted multilevel multiple regressions to analyze  
30 reductions in PPD and gains in PAL compared to baseline at 3, 6 and 12 months  
31 following non-surgical periodontal therapy. The Variance Component models of our  
32 study clearly showed that significant variation existed at most of the levels in the  
33 hierarchical structure at all time points (Tables 2 and 4). This indicates that subject,  
34 tooth and site level factors are all responsible for the outcome variations of PPD  
35 reduction and change in PAL in response to non-surgical periodontal therapy. In  
36 addition, this once more demonstrated that analysis which ignores the natural  
37 hierarchical structure of periodontal data might provide some inaccurate results.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

The advantage of a multilevel approach can be identified in the difference between routine subject level analysis, shown in Table 1, and the multilevel regression result, shown in Table 5. **Routine univariate statistical analysis showed the**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**difference of PPD reduction in initially diseased sites smokers and non-smokers to be significant only at 6 months ( $p<0.0017$ ) and marginally insignificant in 12-months ( $p=0.008$ ). On the other hand, the multilevel regression for initially diseased sites (Table 5) showed that sites from non-smokers achieved a significantly greater PPD reduction throughout the study period.**

Tables 2 and 4 demonstrate that the site level factors contributed around 70 to 80% of the total variance in healing outcomes, whereas tooth and subject levels only contributed the remaining 20% to 30%. This implies that most of the variations in outcomes to non-surgical periodontal therapy level result from factors acting at the site level. This is in agreement with a recent study also assessing the relative contribution of multilevel variation for the outcome of subgingival debridement (D'Aiuto et al. 2005) and with a report on both non-surgical and surgical therapy in single-rooted teeth (Kim et al. 2007), both of which found that site level factors had a much greater impact than subject level factors. Indeed, if tooth loss or tooth retention is the true outcome measure of significance after periodontal therapy, it is worth noting that tooth level factors have been shown to be more important than subject level factors in an analysis which factored in tooth and patient level features (Muzzi et al. 2006).

**In the multilevel multiple regression models (Tables 3 and 5), 10 independent variables were included. The percentage reduction in variance compared to Variance Component models indicates the amount of variation that could be explained by the 10 independent variables introduced. For PPD reduction, the independent variables used in the present study achieved about 10% reduction in variance at the 3-, 6- and 12-month re-examinations. Some variables such as presence of BOP at baseline and mean percentage of sites with BOP at baseline seem only to influence the variance for PPD reduction in general for all sites but do not influence the PPD reduction of initially diseased sites, which mostly**

1  
2  
3 exhibited BOP at baseline.

4  
5 Only 2-5% of variance reductions were obtained for gain in PAL in all sites  
6  
7 and in initially diseased sites using the same 10 independent variables (Table 3 &  
8  
9 5). It is rational to presume that factors affecting PPD reduction in response to  
10  
11 non-surgical periodontal therapy are different from those influencing PAL gain.  
12  
13 Further study involving further independent variables is warranted for investigating the  
14  
15 factors affecting gain in PAL after non-surgical periodontal therapy.  
16  
17

18  
19 By means of multilevel modeling analysis, apart from analyzing which variables  
20  
21 significantly affect the results of non-surgical periodontal therapy, an understanding of  
22  
23 the effects of these individual factors can be generated. **In the regression model,**  
24  
25 **utilizing data from 5814 sites of 969 teeth from 40 subjects for all sites (Table 3),**  
26  
27 **sites on anterior teeth, sites with presence of plaque and BOP at baseline, sites on**  
28  
29 **lingual aspects and sites from subjects with higher full-mouth mean BOP%**  
30  
31 **consistently showed greater PPD reduction.**  
32  
33

34  
35 From Table 3, it appears that the effect of percentage of sites with BOP at baseline  
36  
37 on PPD reduction is clinically insignificant (0.01mm). However, if a subject's baseline  
38  
39 BOP% were to be increased by 1%, the PPD reduction of sites in that subject would  
40  
41 have been 0.01mm greater. Hence if a subject presents with 50% higher BOP% at  
42  
43 baseline, the PPD reduction of sites in that subject would be all 0.5mm greater. Hence  
44  
45 greater reductions in PPD can be expected in those presenting with poorer plaque  
46  
47 control, and this may be of clinical importance.  
48  
49

50  
51  
52 **It is generally believed that deeper initial pockets show more PPD reduction.**  
53  
54 **However, researchers have questioned whether that correlation of PPD reduction**  
55  
56 **and baseline PPP measurement may only due to “mathematical coupling” (Tu et**  
57  
58 **al., 2002 and Tu et al., 2005). Since the objective of the present study was not**  
59  
60 **testing the relationship between change and initial value of PPD and PAL but**

1  
2  
3 focusing on the effect of smoking on response after non-surgical periodontal  
4  
5 therapy in terms of PPD reduction and PAL gain, the independent variables such  
6  
7 as initial PPD and PAL at baseline and full-mouth mean PPD and PAL at  
8  
9 baseline were not included in the analysis (Tu et al. 2004). Other multilevel  
10  
11 analysis strategies for investigating the relationship between change and initial  
12  
13 values are available to address this issue (Blance et al. 2005, Tu et al. 2005, Tu &  
14  
15 Gilthorpe, 2007).  
16  
17

18  
19 In treating patients with chronic periodontitis, it may be important to focus  
20  
21 attention on the response of diseased sites with periodontal pockets rather than  
22  
23 gingivitis sites or healthy sites with no increases in PPD. In the present study, a  
24  
25 separate set of multilevel multiple regressions was performed to investigate the effects  
26  
27 of variables on PPD reduction and PAL gain in sites with baseline PPD  $\geq$  5mm.  
28  
29 Non-smokers showed consistently greater PPD reduction at initially diseased sites  
30  
31 throughout the study (Table 5). **The differences were 0.41 mm, 0.79 mm and 0.68**  
32  
33 **mm at the 3-, 6- and 12-month recalls.** These results are in agreement with a  
34  
35 previous study demonstrating that smokers from the same population have generally  
36  
37 less favorable PPD reduction post-treatment (Jin et al. 2000) and implies that the effect  
38  
39 of smoking is to reduce the PPD reduction in sites with baseline PPD  $\geq$  5mm by **0.41**  
40  
41 **mm, 0.79 mm and 0.68 mm** at 3, 6 and 12 months post-therapy respectively. However,  
42  
43 it is important to note that the smoking status as a subject level variable was considered  
44  
45 in dichotomous fashion, i.e. if the patient is a current smoker or a never smoker. Future  
46  
47 studies could include a quantitative measurement such as pack-years and also include  
48  
49 former smokers in investigating any dose-related or residual effect of cigarette  
50  
51 smoking on periodontal healing.  
52  
53  
54  
55  
56  
57  
58

59 In addition, initially diseased sites from anterior teeth, diseased sites with absence  
60  
of plaque at baseline were found to undergo greater PPD reduction throughout the

1  
2  
3 course of the study in response to non-surgical periodontal therapy.  
4

5 **In the present study, we have applied the multilevel statistical analysis of the**  
6 **periodontal data derived from investigating treatment responses after**  
7 **non-surgical therapy in smokers and non-smokers. This approach has yielded**  
8 **new insights into and better understanding of the result of non-surgical**  
9 **periodontal treatment and has allowed a comparison of the treatment responses**  
10 **in Chinese male smokers and non-smokers.**  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

## 21 **Conclusion**

22  
23 The present study adds to the evidence that smokers generally show less favorable  
24 responses after non-surgical mechanical periodontal therapy in terms of pocket depth  
25 reduction. Utilizing multilevel modeling enabled an appreciation of the impact of tooth  
26 position and site level factors on healing responses to non-surgical periodontal therapy  
27 in both smokers and non-smokers. **Most of the variations were found to be**  
28 **associated with site level variables. On the basis of this study future studies with**  
29 **larger sample sizes and focusing on different site level variables are warranted.**  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## Reference

- 1  
2  
3  
4  
5 Albandar, J.M. & Goldstein, H. (1992) Multi-level statistical models in studies of  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- Albandar, J.M. & Goldstein, H. (1992) Multi-level statistical models in studies of periodontal diseases. *Journal of Periodontology* **63**, 690-695.
- Apatzidou, D.A., Riggio, M.P. & Kinane, D.F. (2005) Impact of smoking on the clinical, microbiological and immunological parameters of adult patients with periodontitis. *Journal of Clinical Periodontology* **32**, 973-983.
- Baljoon, M., Natto, S. & Bergström, J. (2005) Long-term effect of smoking on vertical periodontal bone loss. *Journal of Clinical Periodontology* **32**, 789-797.
- Bergström, J. (2004) Influence of tobacco smoking on periodontal bone height. Long-term observations and a hypothesis. *Journal of Clinical Periodontology* **31**, 260-266.
- Bergström, J. & Boström, L. (2001) Tobacco smoking and periodontal hemorrhagic responsiveness. *Journal of Clinical Periodontology* **28**, 680-685.
- Blance, A., Tu, Y.K. & Gilthorpe, M.S. (2005) A multilevel modelling solution to mathematical coupling. *Statistical Methods in Medical Research* **14**, 553-65.
- Boström, L., Linder, L.E. & Bergström J. (1998) Clinical expression of TNF-alpha in smoking-associated periodontal disease. *Journal of Clinical Periodontology* **25**, 767-773.
- Boström, L., Linder, L.E. & Bergström J. (1999) Smoking and cervicular fluid levels of IL-6 and TNF-alpha in periodontal disease. *Journal of Clinical Periodontology* **26**, 352-357.
- Clarke, N.G., Shephard, B.C. & Hirsch, R.S. (1981) The effects of intra-arterial epinephrine and nicotine on gingival circulation. *Oral Surgery, Oral Medicine, Oral Pathology* **52**, 577-582.
- D'Aiuto, F., Ready, D., Parkar, M. & Tonetti, M.S. (2005) Relative contribution of patient-, tooth-, and site-associated variability on the clinical outcomes of

1  
2 subgingival debridement. I. Probing depths. *Journal of Periodontology* **76**,  
3 398-405.  
4  
5

6  
7 Giannopoulou, C., Kamma, J.J. & Mombelli, A. (2003) Effect of inflammation,  
8 smoking and stress on gingival crevicular fluid cytokine level. *Journal of*  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

*Clinical Periodontology* **30**, 145-153.

Gilthorpe, M.S., Griffiths, G.S., Maddick, I.H. & Zamzuri, A.T. (2000a) The application of multilevel modelling to periodontal research data. *Community Dental Health* **17**, 227-235.

Gilthorpe, M.S., Griffiths, G.S., Maddick, I.H. & Zamzuri, A.T. (2001) An application of multilevel modelling to longitudinal periodontal research data. *Community Dental Health* **18**, 79-86.

Gilthorpe, M.S., Maddick, I.H. & Petrie, A. (2000b) Introduction to multilevel modelling in dental research. *Community Dental Health* **17**, 222-226.

Grossi, S.G., Genco, R.J., Machtei, E.E., Ho, A.W., Koch, G., Dunford, R., Zambon, J.J. & Hausmann, E. (1995) Assessment of risk for periodontal disease. II. Risk indicators for alveolar bone loss. *Journal of Periodontology* **66**, 23-29.

Grossi, S.G., Zambon, J.J., Ho, A.W., Koch, G., Dunford, R.G., Machtei, E.E., Norderyd, O.M. & Genco, R.J. (1994) Assessment of risk for periodontal disease. I. Risk indicators for attachment loss. *Journal of Periodontology* **65**, 260-267.

Güntsch, A., Erler, M., Preshaw, P.M., Sigusch, B.W., Klinger, G. & Glockmann, E. (2006) Effect of smoking on crevicular polymorphonuclear neutrophil function in periodontally healthy subjects. *Journal of Periodontal Research* **41**, 184-188.

Haffajee, A.D. & Socransky, S.S. (2001a) Relationship of cigarette smoking to attachment level profiles. *Journal of Clinical Periodontology* **28**, 283-295.

Haffajee, A.D. & Socransky, S.S. (2001b) Relationship of cigarette smoking to the

1  
2  
3 subgingival microbiota. *Journal of Clinical Periodontology* **28**, 377-388.

4  
5  
6 Heasman, L., Stacey, F., Preshaw, P.M., McCracken, G.I., Hepburn, S. & Heasman, P.A.  
7  
8 (2006) The effect of smoking on periodontal treatment response: a review of  
9  
10 clinical evidence. *Journal of Clinical Periodontology* **33**, 241-253.

11  
12  
13  
14 Hujoel, P.P., Loesche, W.J. & DeRouen, T.A. (1990) Assessment of relationships  
15  
16 between site-specific variables. *Journal of Periodontology* **61**, 368-372.

17  
18  
19  
20 Jin, L., Wong, K.Y., Leung, W.K. & Corbet, E.F. (2000) Comparison of treatment  
21  
22 response patterns following scaling and root planing in smokers and  
23  
24 non-smokers with untreated adult periodontitis. *Journal of Clinical Dentistry* **11**,  
25  
26 35-41.

27  
28  
29  
30 Johnson, J.D., Houchens, D.P., Kluwe, W.M., Craig, D.K. & Fisher, G.L. (1990)  
31  
32 Effects of mainstream and environmental tobacco smoke on the immune system  
33  
34 in animals and humans: a review. *Critical Reviews in Toxicology* **20**, 369-395.

35  
36  
37 Kaldahl, W.B., Kalkwarf, K.L., Patil, K.D., Dyer, J.K. & Bates, R.E. Jr. (1988)  
38  
39 Evaluation of four modalities of periodontal therapy. Mean probing depth,  
40  
41 probing attachment level and recession changes. *Journal of Periodontology* **59**;  
42  
43 783-793.

44  
45  
46 Kim, T.-S., Schenk, A., Lungeanu, D., Reitmeir, P. & Eickholz P. (2007) Nonsurgical  
47  
48 and surgical periodontal therapy in single-rooted teeth. *Clinical Oral*  
49  
50 *Investigations* **11**, 391-399.

51  
52  
53 Kraal, J.H. & Kenney E.B. (1979) The response of polymorphonuclear leukocytes to  
54  
55 chemotactic stimulation for smokers and non-smokers. *Journal of Periodontal*  
56  
57 *Research* **14**, 383-389.  
58  
59  
60

- 1  
2  
3 Labriola, A., Needleman, I. & Moles, D.R. (2005) Systematic review of the effect of  
4  
5 smoking on non-surgical periodontal therapy. *Periodontology 2000* **37**, 124-137.  
6  
7  
8 Leung, W. K., Ng, D.K.C., Jin, L. & Corbet, E.F. (2006) Tooth loss in treated  
9  
10 periodontitis patients responsible for their supportive care arrangements. *Journal*  
11  
12 *of Clinical Periodontology* **33**, 265-275.  
13  
14  
15 Lie, M.A., Timmerman, M.F., van der Velden, U. & van der Weijden, G.A. (1998)  
16  
17 Evaluation of 2 methods to assess gingival bleeding in smokers and non-smokers  
18  
19 in natural and experimental gingivitis. *Journal of Clinical Periodontology* **25**,  
20  
21 695-700.  
22  
23  
24  
25 Lindhe, J., Socransky, S.S., Nyman, S. & Westfelt, E. (1987) Dimensional alteration of  
26  
27 the periodontal tissues following therapy. *International Journal of Periodontics*  
28  
29 *and Restorative Dentistry* **7**(2), 9-21.  
30  
31  
32  
33 Mariggio, M.A., Guida, L., Laforgia, A., Santacroce, R., Curci, E., Montemurro, P. &  
34  
35 Fumarulo, R. (2001) Nicotine effects on polymorphonuclear cell apoptosis and  
36  
37 lipopolysaccharide-induced monocyte functions. A possible role in periodontal  
38  
39 disease? *Journal of Periodontal Research* **36**, 32-39.  
40  
41  
42  
43 Matuliene, G., Pjetursson, B.E., Salvi, G.E., Schmidlin, K., Brägger, U., Zwahlen, M.  
44  
45 & Lang, N.P. (2008) Influence of residual pockets on progression of periodontitis  
46  
47 and tooth loss: results after 11 years of maintenance. *Journal of Clinical*  
48  
49 *Periodontology* **35**, 685-695.  
50  
51  
52  
53 Mirbod, S.M., Ahing, S.I. & Pruthi, V.K. (2001) Immunohistochemical study of  
54  
55 vestibular gingival blood vessel density and internal circumference in smokers  
56  
57 and non-smokers. *Journal of Periodontology* **72**, 1318-1323.  
58  
59  
60  
60 Mooney, J., Hodge, P.J. & Kinane, D.F. (2001) Humoral immune response in  
early-onset periodontitis: influence of smoking. *Journal of Periodontal Research*

1  
2  
3 36, 227-232.  
4

5 Muzzi, L., Nieri, M., Cattabriga, M., Rotundo, R., Cairo, F. & Pini-Prato, G.P. (2006)

6  
7 The potential prognostic value of some periodontal factors for tooth loss: a  
8 retrospective multilevel analysis on periodontal patients treated and maintained  
9 over 10 years. *Journal of Periodontology* 77, 2084-2089.  
10  
11

12  
13  
14 Ng, S.K. & Leung, W.K. (2006) A community study on the relationship between stress,  
15 coping, affective dispositions and periodontal attachment loss. *Community*  
16  
17 *Dentistry and Oral Epidemiology* 34, 252-266.  
18  
19

20  
21 Page, R.C. & Kornman, K.S. (1997) The pathogenesis of human periodontitis: an  
22 introduction. *Periodontology 2000* 14, 9-11.  
23  
24

25  
26 Preber, H & Bergström, J. (1986) The effect of non-surgical treatment on periodontal  
27  
28 pockets in smokers and non-smokers. *Journal of Clinical Periodontology* 13,  
29  
30 319-323.  
31  
32

33  
34 Preber, H., Linder, L. & Bergström, J. (1995) Periodontal healing and periopathogenic  
35  
36 microflora in smokers and non-smokers. *Journal of Clinical Periodontology* 22,  
37  
38 946-952.  
39

40  
41 Ramfjord, S.P., Caffesse, R.G., Morrison, E.C., Hill, R.W., Kerry, G.J., Appleberry,  
42  
43 E.A., Nissle, R.R. & Stults, D.L. (1987) 4 modalities of periodontal treatment  
44  
45 compared over 5 years. *Journal of Clinical Periodontology* 14; 445-452.  
46

47  
48 Rasbash, J., Browne, W., Goldstein, H., Yang, M., Plewis, I., Healy, M., Woodhouse,  
49  
50 G., Draper, D., Langford, I. & Lewis, T. (2000). A user's guide to MLwiN.  
51  
52 Version 2.1. Multilevel Models Project, Institute of Education, University of  
53  
54 London.  
55

56  
57 Raulin, L.A., McPherson, J.C. 3rd, McQuade, M.J. & Hanson, B.S. (1998) The effect  
58  
59 of nicotine on the attachment of human fibroblasts to glass and human root  
60  
surfaces in vitro. *Journal of Periodontology* 59, 318-325.

- 1  
2  
3 Renvert, S., Dahlén, G. & Wikström, M. (1998) The clinical and microbiological  
4  
5 effects of non-surgical periodontal therapy in smokers and non-smokers. *Journal*  
6  
7 *of Clinical Periodontology* **25**, 153-157.  
8  
9  
10 Sanz, M. & Teughels, W. (2008) Innovations in non-surgical periodontal therapy:  
11  
12 Consensus Report of the Sixth European Workshop on Periodontology. *Journal*  
13  
14 *of Clinical Periodontology* **35**(s8), 3-7.  
15  
16  
17 Susin, C., Oppermann, R.V., Haugejorden, O. & Albandar, J.M. (2004) Periodontal  
18  
19 attachment loss attributable to cigarette smoking in an urban Brazilian population.  
20  
21 *Journal of Clinical Periodontology* **31**, 951-958.  
22  
23  
24 Tomasi, C., Leyland, A.H. & Wennström, J.L. (2007) Factors influencing the outcome  
25  
26 of non-surgical periodontal treatment: a multilevel approach. *Journal of Clinical*  
27  
28 *Periodontology* **34**, 682-690.  
29  
30  
31 Tong, K.S., Zee, K.-Y., Lee, D.H. & Corbet, E.F. (2003) Clinical responses to  
32  
33 mechanical periodontal treatment in Chinese chronic periodontitis patients with  
34  
35 and without *Actinobacillus actinomycetemcomitans*. *Journal of Periodontology*  
36  
37 **74**, 1582-1588.  
38  
39  
40 Tu, Y.K. & Gilthorpe, M.S. (2007) Revisiting the relation between change and initial  
41  
42 value: A review and evaluation. *Statistics in Medicine* **26**, 443-457.  
43  
44  
45 Tu, Y.K., Baelum, V. & Gilthorpe, M.S. (2005) The problem of analysing the  
46  
47 relationship between change and initial value in oral health research. *European*  
48  
49 *Journal of Oral Sciences*. **113**, 271-8.  
50  
51  
52 Tu, Y.K., Gilthorpe, M.S. & Griffiths, G.S. (2002) Is reduction of pocket probing depth  
53  
54 correlated with the baseline value or is it "mathematical coupling"? . *Journal of*  
55  
56 *Dental Research* **81**, 722-726.  
57  
58  
59 Tu, Y.K, Gilthorpe, M.S., Griffiths, G.S., Maddick, I.H., Eaton, K.A. & Johnson, N.W.  
60  
(2004) The application of multilevel modeling in the analysis of longitudinal

1  
2  
3 periodontal data--part II: changes in disease levels over time. *Journal of*  
4  
5 *Periodontology* **75**, 137-45.  
6

7 van Winkelhoff, A.J., Bosch-Tijhof, C.J., Winkel, E.G. & van der Reijden, W.A. (2001)

8  
9 Smoking affects the subgingival microflora in periodontitis. *Journal of*  
10  
11 *Periodontology* **72**, 666-671.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

**Legend**

*Fig. 1.* Change in probing pocket depth (PPD;  $\pm$  SD) of sites with PPD  $\geq$  5.0 mm at baseline. \*Statistically significant differences between groups after adjustment for multiple comparisons ( $p < 0.001$ ).

*Fig. 2.* Change in PAL ( $\pm$  SD) of sites with PPD  $\geq$  5.0 mm at baseline.

For Peer Review



## Clinical Relevance

*Scientific rationale:* It would be useful for clinicians to be able to predict outcomes of non-surgical mechanical periodontal therapy based on clinical data. The hierarchical structure of periodontal disease measurements, sites' measurements clustered around teeth and then teeth clustered within individuals applies to periodontal disease clinical findings and to outcomes of periodontal therapy, hence multilevel analysis approach is adopted in this study. *Practical implications:* Multilevel analysis revealed that **for diseased sites without plaque at baseline, from anterior teeth, in non-smokers were found to respond favorably throughout 12 month post-treatment.** Such analysis strategy could be applied to other periodontal treatment modalities.

Table 1. Subject level clinical parameters over study period

	Non-smokers (n = 20)				Smokers (n = 20)			
	Months post-treatment				Months post-treatment			
	Baseline	3	6	12	Baseline	3	6	12
Full mouth plaque %	75.45 ± 14.95	40.70 ± 17.21	32.81 ± 17.21	26.55 ± 14.19	77.36 ± 10.96	35.21 ± 23.50	26.36 ± 13.61	33.79 ± 15.07
Full mouth BOP %	73.45 ± 21.02	42.01 ± 15.53	37.95 ± 15.40	24.92 ± 10.44	<b>54.32 ± 13.68</b>	32.04 ± 11.73	<b>23.97 ± 9.65</b>	26.91 ± 10.85
Full mouth mean PPD (mm)	2.82 ± 0.73	1.95 ± 0.42	1.82 ± 0.31	1.71 ± 0.28	2.89 ± 0.52	2.06 ± 0.37	1.99 ± 0.34	2.01 ± 0.38
Full mouth mean PAL (mm)*	3.69 ± 0.97	---	---	---	3.71 ± 0.68	---	---	---
PPD reduction (mm)	---	0.88 ± 0.57	1.00 ± 0.55	1.11 ± 0.69	---	0.83 ± 0.28	0.91 ± 0.28	0.89 ± 0.32
PAL gain (mm)	---	0.18 ± 0.48	0.33 ± 0.54	0.50 ± 0.52	---	0.28 ± 0.18	0.26 ± 0.21	0.31 ± 0.42
% of pocket ≥ 5.0 mm	11.43 ± 12.14	1.98 ± 2.13	1.15 ± 1.52	0.80 ± 0.94	9.98 ± 9.69	2.76 ± 3.01	2.52 ± 2.68	<b>3.37 ± 3.24</b>
Diseased site mean PPD (mm)	5.94 ± 0.47	3.29 ± 0.57	2.89 ± 0.39	2.49 ± 0.50	5.85 ± 0.48	3.51 ± 0.71	<b>3.46 ± 0.54</b>	3.00 ± 0.80
Diseased site mean PAL (mm)*	6.86 ± 0.88	---	---	---	6.61 ± 0.64	---	---	---
Diseased site PPD reduction (mm)	---	2.65 ± 0.66	3.05 ± 0.61	3.45 ± 0.62	---	2.33 ± 0.50	<b>2.38 ± 0.57</b>	2.84 ± 0.75
Diseased site plaque %	86.48 ± 14.67	65.24 ± 26.59	50.22 ± 26.83	42.45 ± 25.57	92.53 ± 10.24	53.23 ± 29.31	45.70 ± 23.07	58.27 ± 21.86
Diseased site BOP%	90.25 ± 15.86	65.81 ± 21.94	55.74 ± 19.73	35.86 ± 22.49	<b>71.89 ± 19.54</b>	44.92 ± 22.47	36.68 ± 20.88	42.42 ± 21.54

**Bold fonts:** Statistically significance between groups regarding data at baseline (p<0.05)

**Bold and italic fonts:** Statistically significance between groups after adjustment for multiple comparison (p<0.0017)

\*Measured manually by PCP-UNC 15, Hu-Friedy probe, Chicago, IL (Cheng et al., 2008); all other measurements of PPD and PAL used Florida Probe®

Table 2. Variance Components models for reduction in PPD and gain in PAL for all sites

	Reduction in PPD			Gain in PAL		
	3-month	6-month	12-month	3-month	6-month	12-month
Mean (intercept)	0.85 (0.72, 0.99)	0.95 (0.82, 1.08)	1.00 (0.83, 0.16)	0.24 (0.13, 0.34)	0.30 (0.17, 0.42)	0.37 (0.23, 0.51)
Variance						
Subject (level-3)	0.18 (0.09, 0.26)	0.17 (0.09, 0.25)	0.27 (0.14, 0.39)	0.11 (0.06, 0.17)	0.14 (0.07, 0.21)	0.18 (0.10, 0.27)
Tooth (level-2)	0.15 (0.12, 0.18)	0.14 (0.10, 0.17)	0.15 (0.11, 0.19)	0.11 (0.08, 0.14)	0.13 (0.10, 0.16)	0.03 (0.00, 0.07)
Site (level-1)	1.15 (1.10, 1.20)	1.21 (1.16, 1.26)	1.53 (1.47, 1.59)	1.24 (1.19, 1.29)	1.22 (1.17, 1.27)	1.89 (1.82, 1.97)
Total variance	1.48	1.51	1.95	1.46	1.50	2.11
% total variance						
Subject (level-3)	12	11	14	8	10	9
Tooth (level-2)	10	9	8	7	9	1
Site (level-1)	78	80	78	85	81	90

95% confidence intervals in parenthesis.

Table 3. Random intercept models for reduction in PPD and gain in PAL for all sites

Variables	Reduction in PPD			Gain in PAL		
	3-month Estimate (SE)	6-month Estimate (SE)	12-month Estimate (SE)	3-month Estimate (SE)	6-month Estimate (SE)	12-month Estimate (SE)
Intercept	<b><i>0.62 ± 0.10</i></b>	<b><i>0.66 ± 0.09</i></b>	<b><i>0.60 ± 0.11</i></b>	0.13 ± 0.09	0.14 ± 0.10	<b><i>0.22 ± 0.10</i></b>
Subject-level						
Smoking (non-smoker vs smoker)	-0.11 ± 0.13	-0.10 ± 0.12	-0.01 ± 0.15	-0.15 ± 0.12	-0.04 ± 0.14	0.06 ± 0.14
Age at baseline	0.01 ± 0.01	0.01 ± 0.01	0.01 ± 0.01	0.01 ± 0.01	0.01 ± 0.01	<0.01 ± 0.01
Number of missing teeth	0.02 ± 0.02	0.03 ± 0.02	<b><i>0.06 ± 0.03</i></b>	0.01 ± 0.02	-0.01 ± 0.02	<0.01 ± 0.02
% of sites with plaque at baseline	<0.01 ± <0.01	<0.01 ± <0.01	-0.003 ± <0.01	<0.01 ± <0.01	-0.01 ± <0.01	-0.01 ± 0.01
% of sites with BOP at baseline	<b><i>&lt;0.01 ± &lt;0.01</i></b>	<b><i>&lt;0.01 ± &lt;0.01</i></b>	<b><i>0.01 ± &lt;0.01</i></b>	<0.01 ± <0.01	0.01 ± <0.01	<b><i>0.01 ± &lt;0.01</i></b>
Tooth-level						
Tooth position (post. vs. ant.)	<b><i>-0.10 ± 0.04</i></b>	<b><i>-0.11 ± 0.04</i></b>	<b><i>-0.140 ± 0.04</i></b>	<0.01 ± 0.04	<b><i>-0.13 ± 0.04</i></b>	<b><i>-0.15 ± 0.04</i></b>
Arch (lower vs. upper)	-0.06 ± 0.04	-0.04 ± 0.04	0.01 ± 0.04	-0.04 ± 0.04	0.02 ± 0.04	0.06 ± 0.04
	-0.11 ± 0.13	-0.10 ± 0.12	-0.01 ± 0.15	-0.15 ± 0.12	-0.04 ± 0.14	0.06 ± 0.14
Site-level						
Presence of plaque at baseline	0.01 ± 0.01	0.01 ± 0.01	0.01 ± 0.01	0.01 ± 0.01	0.01 ± 0.01	<0.01 ± 0.01
Presence of BOP at baseline	0.02 ± 0.02	0.03 ± 0.02	<b><i>0.06 ± 0.03</i></b>	0.01 ± 0.02	-0.01 ± 0.02	<0.01 ± 0.02
Surface (lingual vs buccal)	<0.01 ± <0.01	<0.01 ± <0.01	-0.003 ± <0.01	<0.01 ± <0.01	-0.01 ± <0.01	-0.01 ± 0.01
	<b><i>&lt;0.01 ± &lt;0.01</i></b>	<b><i>&lt;0.01 ± &lt;0.01</i></b>	<b><i>0.01 ± &lt;0.01</i></b>	<0.01 ± <0.01	0.01 ± <0.01	<b><i>0.01 ± &lt;0.01</i></b>
Variance						
Subject	0.11	0.10	0.14	0.10	0.13	0.13
Tooth	0.14	0.12	0.14	0.11	0.13	0.03
Site	1.12	1.17	1.49	1.22	1.20	1.87
Total variance	1.38	1.40	1.77	1.43	1.36	2.03
% reduction in variance (compared to Variance Component models in Table 2)						
Subject	34	41	46	11	9	30
Tooth	6	9	9	0	0	0
Site	2	3	3	1	2	1
Total variance	7	8	9	2	2	4

**Bold fonts:**  $p < 0.05$ ; **Bold and italic fonts:**  $p < 0.001$

Table 4. Variance Components models for reduction in PPD and gain in PAL for initially diseased sites\*

	Reduction in PPD			Gain in PAL		
	3-month	6-month	12-month	3-month	6-month	12-month
Mean (intercept)	2.55 (2.35, 2.74)	2.77 (2.55, 3.00)	3.16 (2.91, 3.42)	0.80 (0.63, 0.97)	0.83 (0.62, 1.03)	1.21 (0.98, 1.44)
Variance						
Subject (level-3)	0.15 (-0.01, 0.32)	0.24 (0.03, 0.45)	0.40 (0.11, 0.68)	0.00 (-0.10, 0.11)	0.17 (-0.01, 0.36)	0.16 (-0.06, 0.37)
Tooth (level-2)	0.35 (0.07, 0.63)	0.38 (0.10, 0.67)	0.43 (0.14, 0.71)	0.83 (0.41, 1.25)	0.39 (0.07, 0.70)	0.73 (0.28, 1.18)
Site (level-1)	2.07 (1.74, 2.39)	2.03 (1.71, 2.34)	1.90 (1.60, 2.20)	2.55 (2.15, 2.96)	2.32 (1.96, 2.68)	3.03 (2.55, 3.50)
Total variance	2.57	2.65	2.73	3.38	2.88	3.91
% total variance						
Subject (level-3)	6	9	14	0	6	4
Tooth (level-2)	14	15	16	25	14	19
Site (level-1)	80	76	70	75	80	77

95% confidence intervals in parenthesis.

\*Baseline PPD  $\geq$  5.0 mm

Table 5. Final multilevel multiple regression random intercept models for reduction in PPD and gain in PAL for initially diseased sites\*

Variables	Reduction in PPD			Gain in PAL		
	3-month Estimate (SE)	6-month Estimate (SE)	12-month Estimate (SE)	3-month Estimate (SE)	6-month Estimate (SE)	12-month Estimate (SE)
Intercept	<b>3.45 ± 0.27</b>	<b>3.26 ± 0.27</b>	<b>3.65 ± 0.28</b>	<b>1.71 ± 0.31</b>	<b>1.49 ± 0.29</b>	<b>2.12 ± 0.34</b>
Subject-level						
Smoking (non-smoker vs smoker)	<b>0.41 ± 0.20</b>	<b>0.79 ± 0.20</b>	<b>0.68 ± 0.24</b>	-0.19 ± 0.22	-0.09 ± 0.22	-0.13 ± 0.25
Age at baseline	0.01 ± 0.02	0.02 ± 0.02	<0.01 ± 0.02	0.02 ± 0.02	0.02 ± 0.02	<0.01 ± 0.02
Number of missing teeth	<0.01 ± 0.03	<0.01 ± 0.03	0.06 ± 0.04	-0.02 ± 0.03	-0.04 ± 0.04	<0.01 ± 0.04
% of sites with plaque at baseline	<0.01 ± <0.01	<0.01 ± <0.01	<b>-0.02 ± &lt;0.01</b>	-0.002 ± <0.01	-0.01 ± <0.01	<b>-0.02 ± &lt;0.01</b>
% of sites with BOP at baseline	<0.01 ± <0.01	<0.01 ± <0.01	<0.01 ± <0.01	<0.01 ± <0.01	<0.01 ± <0.01	<0.01 ± <0.01
Tooth-level						
Tooth position (post. vs. ant.)	<b>-0.35 ± 0.15</b>	<b>-0.48 ± 0.14</b>	<b>-0.35 ± 0.15</b>	-0.23 ± 0.18	<b>-0.31 ± 0.16</b>	-0.36 ± 0.19
Arch (lower vs. upper)	-0.02 ± 0.14	-0.02 ± 0.14	0.06 ± 0.14	-0.25 ± 0.17	-0.06 ± 0.15	-0.12 ± 0.18
Site-level						
Presence of plaque at baseline	<b>-0.55 ± 0.19</b>	<b>-0.44 ± 0.19</b>	<b>-0.45 ± 0.19</b>	<b>-0.48 ± 0.22</b>	<b>-0.45 ± 0.20</b>	-0.25 ± 0.24
Presence of BOP at baseline	-0.21 ± 0.20	0.10 ± 0.19	-0.16 ± 0.20	-0.14 ± 0.23	-0.03 ± 0.21	-0.41 ± 0.25
Surface (lingual vs buccal)	<b>-0.39 ± 0.13</b>	<b>-0.42 ± 0.13</b>	-0.20 ± 0.13	-0.07 ± 0.15	<0.01 ± 0.14	-0.09 ± 0.16
Variance						
Subject	0.11	0.05	0.16	0.00	0.07	0.06
Tooth	0.14	0.32	0.41	0.81	0.38	0.76
Site	1.12	1.94	1.85	2.50	2.30	2.97
Total variance	1.38	2.30	2.42	3.31	2.75	3.78
% reduction in variance (compared to Variance Component models in Table 4)						
Subject	79	80	60	100	57	64
Tooth	-8	17	4	3	1	-3
Site	6	4	3	2	1	2
Total variance	9	13	11	2	5	3

**Bold fonts:**  $p < 0.05$ ; **Bold and italic fonts:**  $p < 0.001$

\*Baseline PPD  $\geq 5.0$  mm




Fig. 1. Change in probing pocket depth (PPD;  $\pm$  SD) of sites with PPD  $\geq$  5.0 mm at baseline.  
\*Statistically significant differences between groups after adjustment for multiple comparisons  
( $p < 0.001$ ).

131x74mm (600 x 600 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Fig. 2. Change in PAL ( $\pm$  SD) of sites with PPD  $\geq$  5.0 mm at baseline.  
131x74mm (600 x 600 DPI)