



<b>Title</b>	<b>Molecular and cellular aspects of plasticity after neural injury</b>
<b>Author(s)</b>	<b>Chan, YS; Chen, LW; Yung, KKL; Shum, DKY</b>
<b>Citation</b>	<b>Journal of Medical Sciences, 2008, v. 1 n. 2, p. 73-79</b>
<b>Issued Date</b>	<b>2008</b>
<b>URL</b>	<b><a href="http://hdl.handle.net/10722/58284">http://hdl.handle.net/10722/58284</a></b>
<b>Rights</b>	<b>Creative Commons: Attribution 3.0 Hong Kong License</b>

# Molecular and cellular aspects of plasticity after neural injury

**Y S Chan<sup>a,c</sup>, L.W Chen<sup>d</sup>, K.K.L. Yunge, and D K Y Shum<sup>b,c</sup>**

*Departments of physiology<sup>a</sup> and Biochemistry<sup>b</sup>, Research Centre of Heart, Brain, Hormone and Healthy Aging<sup>c</sup>, Faculty of Medicine, The University of Hong Kong, Hong Kong, Institute of Neuroscience<sup>d</sup>, The Fourth Military Medical University, Xi'an, and Department of Biology<sup>e</sup>, Hong Kong Baptist University, Hong Kong, P R China*

This review focuses on our effort to address plasticity of the nervous system after neural injury. We have used different animal models to examine cellular mechanisms of plasticity underlying the pathological and repair processes. After severance of sensory input from one inner ear, topographic representations of space-centered coordinates in the brain undergo plastic changes. During vestibular compensation, tissue plasticity constitutes an important component for functional recovery of neuronal network. In Parkinsonian animals, modulation of signaling via glutamatergic synapses, neurotrophins and neurokinins contributes to the protection of basal ganglion neurons from degeneration, thereby delaying deterioration of motor functions. With the use of animal models of neural injury, we further

overcome the molecular restriction at the glial scar to enhance neural regrowth and remyelination, pointing to the possibility of developing new therapeutic strategies to stimulate neural plasticity and repair in the adult nervous system.

## *Modification of Space-centered Coordinates in the Brain after Unilateral Loss of Vestibular Function*

Vestibular information is centrally processed for the recognition of spatial orientation and for sensory-motor coordination.<sup>1,2</sup> It is well documented that vestibular nuclear (VN) neurons receive otolith inputs arising from the two sides.<sup>3-5</sup> Immediately after unilateral vestibular neurectomy or labyrinthectomy, patients exhibit oculomotor and postural disorders as well as a corresponding change in their spatial perceptual judgments.<sup>6,7</sup> However, it is unknown whether central vestibular neurons receiving inputs solely from the ipsilateral or contralateral otoliths can encode spatiotemporal information during natural otolithic stimulation after unilateral loss of vestibular input from the inner ear. In normal rats, otolith-related VN neurons exhibited a spectrum of spatiotemporal properties, spanning from narrowly to broadly tuned patterns.<sup>8-12</sup> The response vectors of these neurons were found to point in all horizontal directions,<sup>11</sup> indicating that all horizontal head orientations are encoded across an ensemble of VN neurons.<sup>13,14</sup> In hemi-labyrinthomized (HL) preparations in which the labyrinthine input from one side was eliminated,<sup>8</sup> a number of deranged neural patterns were observed. Besides a significant increase in the proportion of broadly spatiotemporal-tuned neurons, there was an imbalance in the distribution of response vectors<sup>9,15</sup> resulting in deranged spatial coding after HL. Another finding was the emergence of unilaterally sensitive neurons that were not found in labyrinth-intact animals.<sup>11,16</sup> The latter neurons

## Correspondence

Y S Chan

Department of Physiology, Faculty of Medicine, The University of Hong Kong, 21 Sassoon Road, Hong Kong, China,

Tel: (852) 2819 9263, Fax: (852) 2855 9730

Email: yschan@hkucc.hku.hk

showed large shifts in response vector with change in velocity,<sup>9</sup> indicating that signals of head orientations would be erroneously transmitted by these neurons during head movements after HL. In addition, asymmetries in spontaneous discharge, response dynamics and spatial coding behaviour were evidenced between neuronal subpopulations on the two sides,<sup>9,10,17</sup> suggesting a segregation of otolith signals reaching the ipsilateral and contralateral VN. These physiological correlates may contribute to both biased spatial coding and maladjusted otolith reflexes accompanying vestibular compensation. We therefore suggest within the VN, gating of spatial and temporal cues arising from the bilateral labyrinths are crucial for the recognition of head orientations in the normal state.

Progress has also been made to bridge the gap between remodeling of the perineuronal matrix and re-tuning of synaptic transmission after unilateral vestibular lesion. Perineuronal nets, lattice-like extracellular structures in the matrix surrounding neurons, were found to be disorganized with severance of peripheral inputs from one labyrinth but the nets resumed following adaptation to the change.<sup>18</sup> It is noteworthy that degradation of perineuronal net by treatment with chondroitinase ABC reactivated cortical plasticity and promoted functional recovery of the lesioned visual system in adult animals.<sup>19</sup> Since the perineuronal nets appear to be restrictive to tissue plasticity,<sup>20</sup> our finding therefore offers a structural correlate for freeing the perineuronal environment to synaptic modulations that lead to functional recovery of VN neurons in the course of vestibular compensation. How the biophysical properties of the matrix influence the interactions with cellular receptors and consequent signaling cascades remain challenging questions for further investigations. Given that neurotrophin-mediated signaling is important for promoting survival of neuronal populations in the central nervous system<sup>21</sup> and for the survival of the vestibular ganglia,<sup>22</sup> we examined the expression of high affinity tyrosine kinase receptors TrkA-C in the

VN. The vast majority of otolith-related neurons in various VN subnuclei of adult rats expressed TrkA, TrkB or TrkC receptors.<sup>23,24</sup> In hippocampal neurons, interaction of brain-derived neurotrophic factor (BDNF) / neurotrophin-4 with TrkB receptor altered the phosphorylation of NR125 and NR2B subunits,<sup>26</sup> thereby modulating the properties of N-methyl-D-aspartate (NMDA) channels.<sup>27,28</sup> Within the VN which receives signals of head movements via glutamatergic vestibular primary afferents,<sup>29,30</sup> otolith-related neurons highly expressed NMDA receptors.<sup>31,32</sup> Co-expression of NMDA and  $\alpha$ -amino-3-hydroxyl-5-methyl-4isoxazole-propionic acid (AMPA) receptor subunits was also observed in VN neurons,<sup>33</sup> implicating cross-modulation between these receptors. Pilot experiments using whole cell patch-clamp recording in VN neurons further demonstrated that changes in NMDA and AMPA receptor components are crucial for the differential demands of glutamatergic neurotransmission during the maturation of VN neurons.<sup>34</sup> That 90% of otolith-related VN neurons expressed either NR1 or NR231 and that a similar proportion expressed TrkB receptor<sup>35</sup> tempted us to infer that co-localization of TrkB and NR1/NR2 receptors in most otolith-related VN neurons contribute to the survival of central vestibular neurons during vestibular compensation.

#### *Role of Glutamatergic Synapses, Neurotrophins and Neurokinins in Neuroprotection*

Glutamate is also a key neurotransmitter in the basal ganglia.<sup>36</sup> One of the possible causes of cell death in the substantia nigra of Parkinsonian patients is thought to be glutamate-mediated excitotoxicity and over-activity of glutamatergic pathways.<sup>37,38</sup> We have conducted a series of studies to investigate the contribution of glutamate neurotransmission, neurotrophins and neurokinins in protecting basal ganglia neurons of Parkinsonian animals from degeneration. Subpopulations of striatal neurons showed stage-specific combinations of NMDA subunits during postnatal development.<sup>39,40</sup> The change in expression patterns of NR subunits may be related to functional maturation of neurons in the neostriatum. Similar observations have been

reported for the localization and expression of AMPA receptors,<sup>41-43</sup> metabotropic glutamate receptors (Lam et al. 2005)<sup>44</sup> and  $\gamma$ -aminobutyric acid (GABA) receptors<sup>45,46</sup> in the neostriatum. We have also found that the expression of NMDA and AMPA receptor subunits in the neostriatum changed after the induction of Parkinsonism in a 6-hydroxydopamine (6-OHDA)  $\alpha$ -lesioned rat model of the disease.<sup>47</sup> Other than the use of antagonists to block glutamatergic transmission,<sup>48</sup> we recently demonstrated that motor symptoms of Parkinson's disease in the rat model could be ameliorated after administration of an NR1 antisense oligonucleotide to the lesioned neostriatum.<sup>13</sup> These results thus shed new light on alternative treatments of motor problems associated with Parkinson's disease.

Several lines of evidence have shown that degeneration of dopaminergic neurons and onset of Parkinson's disease are closely related to the level of neurokinins (including substance P, neurokinin A and B) in the basal ganglia.<sup>49-51</sup> The biological functions of neurokinin as neurotransmitters or neuromodulators<sup>52</sup> are mediated by distinct G-protein coupled receptors. Neurons in the basal ganglia display distinct subclasses of neurokinin receptors, with a predominance of NK-3 receptor in the substantia nigra and NK-1 receptor in the neostriatum.<sup>53,54</sup> Administration of NK-3 receptor agonist to mice treated with 1-methyl-4-phenyl- $\alpha$ -1,2,3,6-tetrahydropyridine (MPTP) aggravated the motor symptoms of Parkinson's disease, indicating that NK-3 antagonism may improve the Parkinson's disease conditions after onset of the disease. In the ventral pallidum, cholinergic neurons expressed both receptors,<sup>55,56</sup> suggesting the differential roles of neurokinin receptors in the basal ganglia.

Substance P was also found to protect striatal cholinergic neurons from glutamate excitotoxicity, probably via modulating NMDA<sup>57</sup> and AMPA receptors.<sup>43</sup> In addition, the expression of substance P in striatal neurons was under the control of neurotrophic factors such as glial-derived neurotrophic factor (GDNF) and neurotrophin-

4/5,<sup>58</sup> thus promoting the survival of dopaminergic neurons and prevention of neuronal death within the basal ganglia circuitry.<sup>59</sup>

Experiments have also been conducted to examine the contributory role of reactive astrocytes in the recovery of brain functions in Parkinsonism. In MPTP-treated mice, significant re-expression of nestin protein, an intermediate filament that is expressed in migrating and proliferating cells during embryogenesis but restricted to areas of regeneration in adults,<sup>60</sup> was found in the caudate putamen while no obvious change was detected in the globus pallidus and substantia nigra.<sup>61,62</sup> In the caudate putamen, the majority of nestin-immunoreactive cells displayed astrocytic morphology and expressed glial fibrillary acid protein (GFAP), a marker for astrocytic glial cells.<sup>61</sup> In addition, these nestin-expressing glial cells exhibited proliferative (Ki-67) and neurotrophic (BDNF) properties.<sup>63</sup> Taken together, these results suggest that reactive astrocytes, through their neurotrophic functions and active interaction with dopaminergic neurons or progenitors, play important roles in the protection of surviving dopamine cells, thereby delaying deterioration of dopaminergic neurons in Parkinson's disease.<sup>64</sup>

#### *Neural Regeneration After Injury*

Structural plasticity of neural tissue plays a key role in the remodeling and regeneration of the adult nervous system after injury. The association of extracellular matrix molecules with neural regeneration or remodeling has been increasingly acknowledged.<sup>65-68</sup> For example, heparan sulfate proteoglycan is considered to play a permissive role while chondroitin sulfate proteoglycan has been conferred a restrictive role. Using a nerve bridge model, we demonstrated that supplementation of soluble heparan sulfates to the regenerative environment of sciatic nerves enhanced axonal reconnection of the severed nerve with the target muscle.<sup>69</sup> A full recovery of the nerve conduction velocity was also evidenced in the course of 20 weeks. Upregulation of chondroitin 6-sulphotransferase-1 was demonstrated to result in

enhanced mobility of Schwann cells that guided axonal regrowth in the injured sciatic nerve.<sup>70</sup> Within the spinal cord of adult rats, co-localization of syndecan-3, a transmembrane heparan sulfate proteoglycan, with heparanase was revealed in neurons and oligodendrocytes,<sup>35</sup> providing cellular basis for further study on the role of these components in enhancing regeneration and plasticity after spinal cord injury. Another extracellular matrix component, the chondroitin sulfate proteoglycans, were found to be up-regulated in regions of reactive gliosis after spinal cord injury.<sup>71</sup> We hypothesized that chondroitin sulfate proteoglycans deposited at the gliotic front constitute a molecular barrier to axonal growth into the transected spinal cord. Despite success in guiding axonal growth into the cellular graft that bridges across the transected cord, regeneration across the distal graft-host interface into the host spinal cord was limited.<sup>72,73</sup> We exploited activity of chondroitinase ABC *in vivo* to resolve restriction due to chondroitin sulfate moieties at the glial scar to enhance neural regrowth and remyelination in the host cord.<sup>74</sup> Our findings indicate that the regrowing axons could advance through the diminished chondroitin sulfate barrier at the interface, thus facilitating axonal regeneration into the caudal host spinal cord. Also, the prospect of the therapeutic use of the Schwann cell-seeded channel as a bridge to facilitate axonal regrowth across traumatic injury in the spinal cord is advanced.

### Acknowledgements

The authors are grateful to Simon S M Chan, Alice Y Y Lui and Kimmy F L Tsang of The University of Hong Kong for their excellent assistance in the experiments. This work was supported by grants from the Hong Kong Research Grants Council (HKU 7294/01M, 7270/02M, 7380/03M, N\_HKBU 202/02).

### References

1. Angelaki DE, Shaikh AG, Green AM, Dickman ID. Neurons compute internal models of the physical laws of motion. *Nature* 2004; 430: 560-564.
2. D.M. Merfeld, L.H. Zupan and R.J. Peterka Humans use internal models to estimate gravity and linear acceleration. *Nature* 1999; 398: 615-618.
3. Chan YS, Hwang JC, Cheung YM. Crossed sacculo-ocular pathway via the Deiters' nucleus in cats. *Brain Res Bull* 1977; 2: 1-6.
4. Uchino Y, Sato YH, Kushiro K, Zakir M, Imagawa MM, Ogawa Y, Katsuta M. Cross-striolar and commissural inhibition in the otolith system. *Ann N Y Acad Sci* 1999; 871: 162-172.
5. Wilson VJ, Gacek RR, Uchino Y, Susswein AJ. Properties of central vestibular neurons fired by stimulation of the saccular nerve. *Brain Res* 1978; 143: 251-261.
6. Curthoys IS. Vestibular compensation and substitution. *Curr Opin Neurol* 2000; 13: 27-30.
7. Seemungal BM, Gresty MA, Bronstein AM. The endocrine system, vertigo and balance. *Curr Opin Neurol* 2001; 14: 27-34.
8. Chan YS. Coding of head positions in neurons of bilateral vestibular nuclei of cats after unilateral labyrinthectomy: Response to off-vertical axis rotation. *Exp Brain Res* 1997; 114: 293-303.
9. Chan YS, Shum DK, Lai CH. Neuronal response sensitivity to bidirectional off vertical axis rotations: A dimension of imbalance in the bilateral vestibular nuclei of cats after unilateral labyrinthectomy. *Neuroscience* 1999; 94: 831-843.
10. Chan YS, Lai CH, Shum DK. Bilateral otolith contribution to spatial coding in the vestibular system. *J Biomed Sci* 2002; 9: 574-586.
11. Lai CH, Chan YS. Properties of otolith-related vestibular nuclear neurons in response to bidirectional off-vertical axis rotation of the rat. *Brain Res* 1995; 693: 39-50.
12. Lai CH, Chan YS. Spontaneous discharge and response characteristics of central otolith neurons of rats during postnatal development. *Neuroscience* 2001; 103: 275-288.
13. Lai CH, Tse YC, Shum DK, Yung KK, Chan YS. Postnatal expression of Fos in otolith-related brainstem neurons of rats following off-vertical axis rotation. *J Comp Neurol* 2004; 470: 282-296.
14. Lai SK, Lai CH, Yung KK, Shum DK, Chan YS. Maturation of otolith-related brainstem neurons in the detection of vertical linear acceleration in rats. *Eur J Neuroscience* 2006; 23(9): 2431-46.
15. Chan YS, Cheung YM. Response of otolith-related neurons in bilateral vestibular nucleus of acute hemilabyrinthectomized cats to off-vertical axis rotations. *Ann N Y Acad. Sci* 1992; 656: 755-765.

16. Chan YS, Cheung YM, Hwang JC. Response characteristics of neurons in the cat vestibular nuclei during slow and constant velocity off-vertical axes rotations in the clockwise and counterclockwise rotations. *Brain Res* 1987; 406: 294-301.
17. Lai CH, Chan YS. Spontaneous properties of otolith-related vestibular nuclear neurons in the decerebrate rats. *Brain Res* 1996; 739: 322-329.
18. Ma CW, Lai CH, Shum DK, Chan YS. Change in Perineuronal Net of Rat Vestibular Nuclear Neurons in Development and in Injury. *Neurosci Lett Suppl* 2006; P-13/25: 147.
19. T. Pizzorusso, P. Medini, N. Berardi, S. Chierzi, I.W. Fawcett, L. Maffei. Reactivation of ocular dominance plasticity in the adult visual cortex. *Science* 2002; 298: 1248-1251.
20. Celio MR, Spreafico De Biasi RS, Vitellaro-Zuccarello L. Perineuronal nets: past and present. *Trends Neurosci*, 1998; 21: 510-515.
21. Huang RJ, Reichardt LF. Neurotrophins: roles in neuronal development and function. *Ann Rev Neurosci* 2001; 24: 677-736.
22. Ernfors P, van de Water PT, Loring J, Jaenisch R. Complementary roles of BDNF and NT-3 in vestibular and auditory development. *Neuron* 1995; 14: 1153-1164.
23. Zhang FX, Lai CR, Lai SK, Yung KK, Shum DK, Chan YS. Neurotrophin receptor immunostaining in the vestibular nucleus of rats. *Neuroreport* 2003; 14: 851-855.
24. Zhang FX, Lai CH, Tse YC, Shum DK, Chan YS. Expression of Trk receptors in otolith-related neurons in the vestibular nucleus of rats. *Brain Res* 2005; 1062(1-2):92-100.
25. Suen PC, Wu K, Levine ES, Mount HT, Xu JL, Lin SY, et al. Brain derived neurotrophic factor rapidly enhances phosphorylation of the postsynaptic N-methyl-D-aspartate receptor subunit 1. *Proc. Natl. Acad. Sci* 1997; 94: 8191-8195.
26. Lin SY, Wu K, Levine ES, Mount HT, Suen PC, Black LB. BDNF acutely increases tyrosine phosphorylation of the BDNF acutely increases tyrosine phosphorylation of the NMDA receptor subunit 2B in cortical and hippocampal postsynaptic densities. *Mol Brain Res* 1998; 55: 20-27.
27. Pozzo-Miller LD, Gottschalk W, Zhang L, McDermott K, Du J, Gopalakrishnan R, et al. Impairments in high frequency transmission, synaptic vesicle docking and synaptic protein distribution in the hippocampus of BDNF knockout mice. *J Neurosci* 1999; 19: 4972-4983.
28. Schinder AF, Berninger B, Poo MM. Postsynaptic target specificity of neurotrophin-induced presynaptic potentiation. *Neuron* 2000; 25: 151-163.
29. Kinney GA, Peterson RW, Slater NT. The synaptic activation of N-methyl-D-aspartate receptors in the rat medial vestibular nucleus. *J Neurophysiol* 1994; 72: 1588-1595.
30. Lewis MR, Phelan KD, Shinnick-Gallagher P, Gallagher JP. Primary afferent excitatory transmission recorded intracellularly in vitro from rat medial vestibular neurons. *Synapse* 1989; 3: 149-153.
31. Chen LW, Lai CR, Law RY, Yung KK, Chan YS. Quantitative study of the coexpression of Fos and N-methyl-D-aspartate (NMDA) receptor subunits in otolith-related vestibular nuclear neurons of rats. *J Comp Neurol* 2003b; 406: 292-301.
32. Tse YC, Lai CH, Lai SK, Liu JX, Yung KK, Shum DK, et al. Developmental expression of NMDA receptor subunits in otolith neurons within the vestibular nucleus of rats. *J Comp Neurol* 2008; 508(2): 343-64.
33. Chen LW, Yung KK, Chan YS. Co-localization of NMDA receptors and AMPA receptors in neurons of the vestibular nucleus of rats. *Brain Res* 2000; 884: 87-97.
34. Lai SK, Yung WH, Chan YS. Developmental profile of ionotropic glutamate receptors in spinal vestibular neurons of rats. *Soc Neuroscience Abstract (U.S.A.)* 2005; 602: 12.
35. Zhang Y, Yeung MN, Liu J, Chau CH, Chan YS, Shum DK. Mapping heparanase expression in the spinal cord of adult rats. *J Comp Neurol* 2006; 494(2): 345-57.
36. Gerfen CR, Wilson CJ. The basal ganglia. In: L.W. Swanson (Ed.) *Handbook of Chemical Neuroanatomy, Vol. 12: Integrated Systems of the CNS, Part 3*. Elsevier, Amsterdam, 1996.
37. Beal MF. Excitotoxicity and nitric oxide in Parkinson's disease pathogenesis. *Ann Neurol* 1998; 44: S 110-114.
38. Blandini F, Nappi G, Greenamyre JT. Subthalamic infusion of an NMDA antagonist prevents basal ganglia metabolic changes and nigral degeneration in a rodent model of Parkinson's disease. *Ann Neurol* 2001; 49: 525-529.
39. Lau WK, Lui PW, Wong CK, Chan YS, Yung KK. Differential expression of N-methyl-D-aspartate receptor subunit messenger ribonucleic acids and immunoreactivity in the rat neostriatum during postnatal development. *Neurochemistry Int* 2003; 43: 47-65.
40. Lui PW, Suen KC, Chan YS, Yung WH, Yung KK. Striatal neurons but not nigral dopaminergic neurons in primary

- cell culture express endogenous functional N-methyl-D-aspartate receptors. *Mol. Brain Res* 2003; 120: 9-21.
41. Chen LW, Wei LC, Lang B, Ju G, Chan YS. Differential expression of AMPA receptor subunits in dopamine neurons of the rat brain: a double immunocytochemical study. *Neuroscience* 2001a; 106: 149-160.
  42. Chan WS, Chung EK, Lau WK, Chan YS, Yung KK. Differential expression of  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazole-propionate glutamate receptors in the rat neostriatum during postnatal development. *NeuroSignals* 2003; 12: 302-309.
  43. Hu HJ, Chen LW, Yung KK, Chan YS. Differential expression of AMPA receptor subunits GluRI, 2, 3, and 4 in substance P receptor-containing neurons of the neostriatal regions of rats. *Neuroscience Res* 2004; 49: 281-288.
  44. Lam RW, Chan YS, Yung KK. Differential localization of metabotropic glutamate receptors 1 and 2/3 in the rat striatum during early postnatal development. *Neuroembryology and Aging* 2005; 3: 36-41.
  45. Chen LW, Cao R, Liu HL, Ju G, Chan YS. The striatal GABA-ergic neurons expressing substance P receptors in the basal ganglia of mice. *Neuroscience* 2003a; 119: 919-925.
  46. Lau WK, Yung WH, Chan YS, Yung KK. Developmental assembly of GABA receptors in the rat substantia nigra. In: P.J. Magill and J.P. Bolam (Eds), *Basal Ganglia VIII*. Springer, New York, 2005, pp. 229-240.
  47. Lai SK, Tse YC, Yang MS, Wong CK, Chan YS, Yung KK. Gene expression of glutamate receptors GluRI and NRI is differentially modulated in striatal neurons in rats after 6-hydroxydopamine lesion. *Neurochemistry Int* 2003; 43: 639-653.
  48. Le DA, Lipton SA. Potential and current use of N-methyl-D-aspartate (NMDA) receptor antagonists in diseases of aging. *Drugs Aging* 2001; 18: 717-724.
  49. Benarroch EE, Schmeichel AM, Low PA, Parisi JE. Depletion of ventromedullary NK-1 receptor-immunoreactive neurons in multiple system atrophy. *Brain* 2003; 126: 2183-2190.
  50. Kowall NW, Quigley BJ Jr, Krause JE, Lu F, Kosofsky BE, Ferrante RJ. Substance P and substance P receptor histochemistry in human neurodegenerative diseases. *Regul Pept* 1993; 46: 174-185.
  51. Schiffmann SN, Dassel D, d'Alcantara P, Ledent C, Swillens S, Zoli M. A2A receptor and striatal cellular functions: regulation of gene expression, currents, and synaptic transmission. *Neurology* 2003; 61: S24-29.
  52. Diez-Guerra FJ, Richardson PJ, Emson PC. Subcellular distribution of mammalian tachykinins in rat basal ganglia. *J Neurochem* 1988; 50: 440-450.
  53. Kaneko S, Hikida T, Watanabe D, Ichinose H, Nagatsu T, Kreitman RJ, *et al.* Synaptic integration mediated by striatal cholinergic interneurons in basal ganglia function. *Science* 2000; 289: 633-637.
  54. Parent A, Cicchetti F, Beach TG. Striatal neurons displaying substance P (NK1) receptor immunoreactivity in human and non-human primates. *Neuroreport* 1995; 6: 721-724.
  55. Chen LW, Wei LC, Liu RL, Qiu Y, Chan YS. Cholinergic neurons expressing substance P receptor in the basal forebrain of the rat: a double immunocytochemical study. *Brain Res* 2001b; 904: 161-166.
  56. Chen LW, Wei LC, Liu HL, Rao ZR, Ju G, Chan YS. Cholinergic neurons expressing neuromedin K receptor (NK3) in the basal forebrain of the rat: a double immunofluorescence study. *Neuroscience* 2001c; 103: 413-422.
  57. Calvo N, Reiriz J, Pérez-Navarro E, Alberch J. Tachykinins protect cholinergic neurons from quinolinic acid excitotoxicity in striatal cultures. *Brain Res* 1996; 740: 323-328.
  58. Humpel C, Marksteiner J, Saria A. Glial-cell-line-derived neurotrophic factor enhances biosynthesis of substance P in striatal neurons in vitro. *Cell Tissue Res* 1996; 286: 249-255.
  59. Chen LW, Yung KK, Chan YS. Neurokinin peptides and neurokinin receptors as potential therapeutic intervention targets of basal ganglia in the prevention and treatment of Parkinson's disease. *Current Drug Targets*, 2004b; 5: 197-206.
  60. Wiese C, Rolletschek A, Kania G, Blyszczuk P, Tarasov KV, Tarasova Y, *et al.* Nestin expression - a property of multi-lineage progenitor cells? *Cell Mol Life Sci* 2004; 61: 2510-2522.
  61. Chen LW, Wei LC, Qiu Y, Liu RL, Rao ZR, Ju G, Chan YS. Significant up-regulation of nestin protein in the neostriatum of MPTP-treated mice. Are the striatal astrocytes regionally activated after systemic MPTP administration? *Brain Res* 2002; 925(1): 9-17.
  62. Wei LC, Shi M, Chen LW, Cao R, Zhang P, Chan YS. Nestin-containing cells express glial fibrillary acidic protein in the proliferative regions of central nervous system of postnatal developing and adult mice. *Dev Brain Res* 2002; 139: 9-17.
  63. Chen LW, Hu RJ, Liu HL, Yung KK, Chan YS. Identification

- of brain derived neurotrophic factor in nestin-expressing astroglial cells in the neostriatum of 1 methyl-4-phenyl-1,2,3,6-tetrahydropyridine-treated mice. *Neuroscience* 2004a; 126: 941-953.
64. Chen LW, Yung KK, Chan YS. Reactive astrocytes as potential manipulation targets in novel cell replacement therapy of Parkinson's disease. *Current Drug Targets* 2005; 6: 821-833.
65. Buss A, Brook GA, Kakulas B, Martin D, Franzen R, Schoenen J, *et al.* Gradual loss of myelin and formation of an astrocytic scar during Wallerian degeneration in the human spinal cord. *Brain* 2004; 127: 34-44.
66. Dityatev A, Schachner M. Extracellular matrix molecules and synaptic plasticity. *Nat Rev Neurosci* 2003; 4: 456-468.
67. Novikova LN, Novikov N, Kellerth JO. Biopolymers and biodegradable smart implants for tissue regeneration after spinal cord injury. *Curr Opin Neurol* 2003; 16(6): 711-5.
68. Sliver J, Miller JH. Regeneration beyond the glial scar. *Nat Rev Neurosci* 2004; 5: 146-156.
69. Chau CR, Shum DK, Chan YS, So KF. Heparan sulfates upregulate regeneration of transected sciatic nerve of adult guinea pigs. *Eur J Neuroscience* 1999; 11: 1914-1926.
70. Liu J, Chau CH, Liu H, Jang BR, Li X, Chan YS, *et al.* Upregulation of chondroitin 6-sulphotransferase-1 facilitates Schwann cell migration during axonal growth. *J Cell Sci* 2006; 119(Pt 5): 933-42.
71. Jones LL, Yamaguchi Y, Stallcup WB, Tuszynski MH. NG2 is a major chondroitin sulfate proteoglycan produced after spinal cord injury and is expressed by macrophages and oligodendrocyte progenitors. *J Neurosci* 2002; 22: 2792-2803.
72. Cheng H, Cao Y, Olson L. Spinal cord repair in adult paraplegic rats: partial restoration of hind limb function. *Science* 1998; 273: 510-513.
73. Li Y, Field PM, Raisman G. Repair of adult rat corticospinal tract by transplants of olfactory ensheathing cells. *Science* 1998; 277: 2000-2002.
74. Chau CH, Shum DK, Li H, Pei J, Lui YY, Wirthlin L, *et al.* Chondroitinase ABC enhances axonal regrowth through Schwann cell-seeded guidance channels after spinal cord injury. *FASEB J* 2004; 18: 194-196.

---

© Chan *et al.*: Licensee Bentham Open.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.