



Title	Triruthenium alkylidyne cluster complexes [H ₂ Ru ₃ (CO) ₉ (μ -3-CX)] (X=pyridine, 4-ethylpyridine)
Author(s)	Wong, WT; Wong, WY
Citation	Acta Crystallographica Section C: Crystal Structure Communications, 1994, v. 51 n. 1, p. 57-60
Issued Date	1994
URL	http://hdl.handle.net/10722/69285
Rights	Creative Commons: Attribution 3.0 Hong Kong License

high final *R* value is mainly due to the disorder of the polymethylene chain.

The authors thank the CNRS for financial support.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and complete geometry have been deposited with the IUCr (Reference: HR1021). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Baxter, L. A. M., Blake, A. J., Gould, R. O., Heath, G. A. & Stephenson, T. A. (1993). *Acta Cryst.* **C49**, 1311–1315.
- Diantouba, B.-A., Guiguemde, I., Tayeb, A., Goetz-Grandmont, G. J. & Brunette, J. P. (1994). *Solvent Extr. Ion Exch.* **12**, 325–347.
- Fair, C. K. (1990). *MOLÉN. An Interactive Intelligent System for Crystal Structure Analysis*. Enraf-Nonius, Delft, The Netherlands.
- Johnson, C. K. (1965). *ORTEP*. Report ORNL-3794. Oak Ridge National Laboratory, Tennessee, USA.
- Lakkis, M., Lakkis, Z., Goetz-Grandmont, G. J. & Brunette, J.-P. (1991). *Monatsh. Chem.* **122**, 9–15.
- Li, X.-J. & Yang, R.-D. (1992). *Polyhedron*, **11**, 1545–1550.
- Main, P., Fiske, S. J., Hull, S. E., Lessinger, L., Germain, G., Declercq, J.-P. & Woolfson, M. M. (1982). *MULTAN11/82. A System of Computer Programs for the Automatic Solution of Crystal Structures from X-ray Diffraction Data*. Univs. of York, England, and Louvain, Belgium.
- Maslen, H. S. & Waters, T. N. (1975). *Coord. Chem. Rev.* **17**, 137–175.
- Mehrotra, R. C., Bohra, R. & Gaur, D. P. (1978). *Metal β -Diketones and Allied Derivatives*. London: Academic Press.
- Miao, F.-M., Liu, X.-L. & Li, Y.-Q. (1991). *Wuji Huaxue Xuebao*, **7**(2), 129–132; *Chem. Abstr.* **116**, 245603r.
- Miyazaki, S., Mukai, H., Umetani, S., Kihara, S. & Matsui, M. (1989). *Inorg. Chem.* **28**, 3014–3017.
- Tobon-Cervantes, A., Toscano, R. A., Rosales, M. J., Fernandez-G., J. M. & Bernal-Uruchurtu, M. I. (1988). *Acta Cryst.* **C44**, 621–623.
- Zolotov, Yu. A. & Kuz'min, N. M. (1977). *The Extraction of Metals by Acylpyrazolones*. Moscow: Nauka. (In Russian.)

Acta Cryst. (1995). **C51**, 57–60

Triruthenium Alkylidyne Cluster Complexes [H₂Ru₃(CO)₉(μ_3 -CX)] (X = Pyridine, 4-Ethylpyridine)

WING-TAK WONG* AND WAI-YEUNG WONG

*Department of Chemistry, University of Hong Kong,
Pokfulam Road, Hong Kong*

(Received 27 September 1993; accepted 19 May 1994)

Abstract

Crystal structure analyses of nonacarbonyl-1 κ^3 C-, 2 κ^3 C, 3 κ^3 C-di- μ -hydrido-1:2 κ^2 H; 2:3 κ^2 H- μ_3 -(pyridin-1-

ylmethylidene)-triangulo-triruthenium, [Ru₃(μ -H)₂(μ_3 -C₆H₅N)(CO)₉], (I), and nonacarbonyl-1 κ^3 C-, 2 κ^3 C, 3 κ^3 C- μ_3 -[(4-ethylpyridin-1-yl)methylidene]-di- μ -hydrido-1:2 κ^2 H; 2:3 κ^2 H-triangulo-triruthenium, [Ru₃(μ -H)₂(μ_3 -C₈H₉N)(CO)₉], (II), reveal that both compounds consist of a triruthenium alkylidyne core with the nucleophiles [pyridine in (I) and 4-ethylpyridine in (II)] bonded to the μ_3 -bridging alkylidyne C atom.

Comment

This work is part of our structural and reactivity studies of triruthenium and triosmium alkylidyne carbonyl clusters. Several triosmium alkylidynes with the formula [H₂O₃(CO)₉(μ_3 -CY)] (Y = pyridine, quinoline or trimethyl phosphite) have been synthesized and fully characterized (Johnson, Lahoz, Lewis, Prior, Raithby & Wong, 1992). The title compounds, (I) and (II), have been prepared by a similar route.





Fig. 1. The molecular structure of [H₂Ru₃(CO)₉(μ_3 -Cpy)] (I).

The structures of (I) and (II), as shown by X-ray analyses, each consist of a triruthenium alkylidyne core with the nucleophile substituted at the apical C atom. Both molecules have approximate C_s symmetry. The hydride atoms of (I) and (II), evident from their ¹H NMR spectra, could not be located by X-ray analysis. However, potential-energy calculations (Orpen, 1980) suggested that one bridges the Ru1—Ru2 edge and the other the Ru2—Ru3 edge in both (I) and (II). These two edges [averages 2.844 Å for (I), 2.835 Å for (II)] are significantly longer than the unbridged Ru1—Ru3 bonds [2.739 (1) for (I), 2.729 (1) Å for (II)]. This is consistent with the general observation that *M—M* distances are increased when bridged by a hydride atom (Humphries & Kaesz, 1979). The N—C(alkylidyne) bond lengths in both (I) [1.46 (1) Å] and (II) [1.458 (9) Å] are comparable to those of the triosmium alkylidynes with other N-donors substituted on the apical C atom.




Fig. 2. The molecular structure of [H₂Ru₃(CO)₉(μ₃-C{4-Etpy})] (II).

Experimental

Compound (I)

Crystal data

[Ru₃H₂(C₆H₅N)(CO)₉]

M_r = 648.43

Monoclinic

Cc

a = 18.585 (3) Å

b = 9.562 (1) Å

c = 15.012 (4) Å

β = 131.83 (3)°

V = 1987.8 (12) Å³

Mo Kα radiation

λ = 0.71073 Å

Cell parameters from 25 reflections

θ = 10–12°

μ = 2.25 mm⁻¹

T = 293 K

Block

0.22 × 0.16 × 0.14 mm

Z = 4

D_x = 2.163 Mg m⁻³

Data collection

Enraf–Nonius CAD-4

diffractometer

ω–2θ scans

Absorption correction:

empirical

T_{min} = 0.901, *T_{max}* =

0.999

1456 measured reflections

1395 independent reflections

1295 observed reflections

[*F_o* > 3σ(*F_o*)]

Refinement

Refinement on *F*

R = 0.037

wR = 0.050

S = 1.876

1295 reflections

126 parameters

H-atom parameters not refined

w = 4*F_o*²/[σ²(*F_o*²)

+ (0.04*F_o*²)²]

Compound (II)

Crystal data

[Ru₃H₂(C₈H₉N)(CO)₉]

M_r = 676.49

Orthorhombic

*P*2₁2₁2₁

a = 10.971 (2) Å

b = 11.034 (2) Å

c = 18.100 (5) Å

V = 2191.0 (14) Å³

Z = 4

D_x = 2.051 Mg m⁻³

Orange

Crystal source: recrystallized from CH₂Cl₂/*n*-hexane

R_{int} = 0.022

θ_{max} = 22.5°

h = –20 – 20

k = 0 – 10

l = 0 – 16

3 standard reflections

frequency: 120 min

intensity variation: <2%

(Δ/σ)_{max} = 0.04

Δρ_{max} = 0.98 e Å⁻³

Δρ_{min} = –0.65 e Å⁻³

Extinction correction: none

Atomic scattering factors

from *International Tables for X-ray Crystallography* (1974, Vol. IV)

Mo Kα radiation

λ = 0.71073 Å

Cell parameters from 25 reflections

θ = 11–13°

μ = 2.05 mm⁻¹

T = 293 K

Block

0.32 × 0.22 × 0.20 mm

Orange

Crystal source: recrystallized from CH₂Cl₂/*n*-hexane

Data collection

Enraf–Nonius CAD-4

diffractometer

ω–2θ scans

Absorption correction:

empirical

T_{min} = 0.895, *T_{max}* =

0.999

2221 measured reflections

2221 independent reflections

2046 observed reflections

[*F_o* > 3σ(*F_o*)]

Refinement

Refinement on *F*

R = 0.028

wR = 0.040

θ_{max} = 25°

h = 0 – 13

k = 0 – 13

l = 0 – 21

3 standard reflections

frequency: 120 min

intensity variation: <3%

(Δ/σ)_{max} = 0.01

Δρ_{max} = 0.37 e Å⁻³

Δρ_{min} = –0.71 e Å⁻³

$S = 1.394$
 2046 reflections
 271 parameters
 H-atom parameters not refined
 $w = 4F_o^2 / [\sigma^2(F_o^2) + (0.04F_o^2)^2]$

Extinction correction: none
 Atomic scattering factors and anomalous-dispersion corrections from *International Tables for X-ray Crystallography* (1974, Vol. IV)

C31	0.1897 (7)	-0.0263 (7)	0.0835 (5)	4.7 (2)
C32	0.3525 (9)	-0.1655 (8)	0.1920 (5)	5.5 (2)
C33	0.3433 (8)	0.0856 (8)	0.1890 (4)	4.8 (2)
HA†	0.4864	-0.0214	-0.0477	5
HB†	0.3811	-0.1720	0.0592	5

† Coordinate fixed to define origin.

‡ These atoms were not located in the analyses. Their positions were deduced from potential-energy calculations (Orpen, 1980).

Table 1. Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

$B_{\text{eq}} = (8\pi^2/3) \sum_i \sum_j U_{ij} a_i^* a_j^* \mathbf{a}_i \cdot \mathbf{a}_j$ for Ru atoms in (I) and all non-H atoms in (II); B_{iso} for all others.

(I)	x	y	z	$B_{\text{iso}}/B_{\text{eq}}$
Ru1	0.646†	0.06082 (9)	0.107†	2.58 (2)
Ru2	0.44197 (5)	0.05228 (9)	-0.08242 (6)	2.50 (2)
Ru3	0.56793 (6)	-0.12780 (8)	-0.07440 (8)	2.76 (2)
O11	0.6638 (6)	0.328 (1)	0.2320 (7)	5.1 (2)
O12	0.8308 (8)	0.109 (2)	0.1591 (9)	7.3 (3)
O13	0.7431 (8)	-0.147 (2)	0.316 (1)	8.1 (4)
O21	0.3069 (7)	-0.054 (1)	-0.0450 (8)	5.9 (3)
O22	0.2944 (7)	0.052 (1)	-0.3519 (8)	5.6 (2)
O23	0.4194 (6)	0.359 (1)	-0.0558 (7)	5.0 (2)
O31	0.7477 (7)	-0.130 (1)	-0.0419 (9)	6.4 (3)
O32	0.4451 (8)	-0.237 (2)	-0.3263 (9)	7.5 (3)
O33	0.6159 (7)	-0.396 (2)	0.0713 (9)	6.4 (3)
N1	0.5553 (5)	0.193 (1)	-0.1453 (6)	2.6 (2)
C1	0.5564 (5)	0.086 (1)	-0.0753 (7)	1.9 (2)
C2	0.5185 (7)	0.168 (2)	-0.2559 (9)	3.8 (2)
C3	0.5112 (8)	0.271 (2)	-0.328 (1)	4.6 (3)
C4	0.5407 (7)	0.404 (1)	-0.2841 (9)	3.3 (2)
C5	0.5813 (8)	0.429 (1)	-0.170 (1)	3.9 (2)
C6	0.5875 (6)	0.325 (1)	-0.1021 (8)	3.0 (2)
C11	0.6582 (6)	0.226 (1)	0.1844 (8)	3.3 (2)
C12	0.7584 (8)	0.090 (2)	0.1354 (9)	4.2 (3)
C13	0.7039 (8)	-0.069 (2)	0.237 (1)	4.2 (3)
C21	0.3601 (7)	-0.017 (1)	-0.0550 (8)	3.3 (2)
C22	0.3470 (7)	0.055 (1)	-0.2497 (8)	3.3 (2)
C23	0.4258 (6)	0.243 (1)	-0.0664 (8)	3.1 (2)
C31	0.6789 (7)	-0.129 (1)	-0.0529 (9)	3.3 (2)
C32	0.4919 (7)	-0.194 (1)	-0.2314 (8)	3.8 (2)
C33	0.5953 (8)	-0.298 (2)	0.012 (1)	4.9 (3)
HA†	0.5329	0.0736	0.0749	5
HB†	0.4519	-0.1223	-0.1133	5
(II)				
Ru1	0.46060 (5)	0.10649 (4)	0.01346 (3)	2.873 (9)
Ru2	0.51780 (5)	-0.14357 (4)	0.01192 (3)	3.15 (1)
Ru3	0.35897 (5)	-0.03952 (5)	0.11844 (3)	3.47 (1)
O11	0.2242 (5)	0.1436 (5)	-0.0758 (4)	5.9 (1)
O12	0.6472 (5)	0.2137 (5)	-0.0903 (3)	5.5 (1)
O13	0.4423 (6)	0.3338 (5)	0.1077 (3)	6.4 (1)
O21	0.3920 (7)	-0.2921 (5)	-0.1087 (3)	6.4 (2)
O22	0.6052 (9)	-0.3539 (5)	0.1065 (3)	8.4 (2)
O23	0.7658 (6)	-0.0990 (6)	-0.0580 (4)	6.7 (2)
O31	0.0981 (5)	-0.0095 (6)	0.0612 (4)	6.8 (2)
O32	0.3460 (9)	-0.2397 (6)	0.2361 (4)	8.3 (2)
O33	0.3354 (8)	0.1612 (6)	0.2315 (3)	7.8 (2)
N1	0.6455 (5)	0.0084 (5)	0.1384 (3)	3.0 (1)
C1	0.5378 (7)	-0.0062 (5)	0.0921 (3)	3.2 (1)
C2	0.6499 (7)	-0.0434 (7)	0.2050 (4)	4.4 (2)
C3	0.753 (1)	-0.0316 (7)	0.2479 (4)	5.6 (2)
C4	0.8542 (8)	0.0341 (7)	0.2241 (4)	4.5 (2)
C5	0.8469 (7)	0.0842 (7)	0.1547 (4)	4.3 (2)
C6	0.7427 (7)	0.0729 (6)	0.1142 (4)	3.7 (1)
C7	0.964 (1)	0.050 (1)	0.2727 (5)	8.5 (3)
C8	1.049 (1)	-0.056 (2)	0.2694 (6)	12.7 (5)
C11	0.3127 (7)	0.1336 (6)	-0.0435 (4)	4.0 (1)
C12	0.5825 (7)	0.1724 (6)	-0.0517 (4)	3.4 (1)
C13	0.4503 (7)	0.2484 (7)	0.0738 (4)	4.1 (1)
C21	0.4404 (7)	-0.2409 (6)	-0.0665 (4)	4.0 (1)
C22	0.5750 (9)	-0.2752 (6)	0.0718 (4)	5.0 (2)
C23	0.6713 (7)	-0.1211 (6)	-0.0332 (4)	3.9 (1)

Table 2. Selected geometric parameters (\AA , °)

	(I)	(II)
Ru1—Ru2	2.8440 (6)	2.8297 (7)
Ru1—Ru3	2.739 (1)	2.7291 (8)
Ru2—Ru3	2.843 (2)	2.8409 (8)
Ru1—C1	2.060 (8)	2.071 (6)
Ru2—C1	2.08 (1)	2.109 (7)
Ru3—C1	2.06 (2)	2.052 (7)
N1—C1	1.46 (1)	1.458 (9)
N1—C2	1.33 (2)	1.335 (9)
N1—C6	1.35 (2)	1.35 (1)
C2—C3	1.40 (2)	1.38 (1)
C3—C4	1.36 (3)	1.39 (1)
C4—C5	1.36 (2)	1.37 (2)
C5—C6	1.38 (3)	1.51 (1)
C5—C6	—	1.36 (2)
C7—C8	—	1.50 (2)
Ru1—HA	1.82	1.82
Ru2—HA	1.78	1.76
Ru2—HB	1.78	1.76
Ru3—HB	1.82	1.83
Ru2—Ru1—Ru3	61.19 (4)	61.44 (2)
Ru1—Ru2—Ru3	57.59 (3)	57.54 (2)
Ru1—Ru3—Ru2	61.22 (3)	61.02 (2)
Ru1—C1—Ru2	86.7 (5)	85.3 (2)
Ru1—C1—Ru3	83.4 (3)	82.9 (3)
Ru2—C1—Ru3	86.7 (5)	86.1 (3)
C1—N1—C2	121 (1)	120.0 (6)
C1—N1—C6	122 (1)	120.7 (5)
C2—N1—C6	118 (2)	119.3 (6)
C4—C7—C8	—	112 (2)
Ru1—HA—Ru2	104.5	104.2
Ru2—HB—Ru3	104.6	104.0

The metal-atom positions were obtained by direct methods using *MULTAN*11/82 (Main, Fiske, Hull, Lessinger, Germain, Declercq & Woolfson, 1982). Each structure was refined by a full-matrix least-squares method with anisotropic displacement parameters for the Ru atoms in (I) and for all non-H atoms in (II). All calculations were performed using the *SDP* system (Enraf-Nonius, 1985) on a MicroVAX II computer. The absolute structures were assigned on the basis of the lowest R values after identical refinements.

We thank the Hong Kong Research Grants Council and the University of Hong Kong for support.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and bond distances and angles involving non-H atoms have been deposited with the IUCr (Reference: MU1090). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

Enraf-Nonius (1985). *Structure Determination Package. SDP/PDP User's Guide*. Version 3.0. Enraf-Nonius, Delft, The Netherlands.

- Humphries, A. P. & Kaesz, H. D. (1979). *Prog. Inorg. Chem.* **25**, 145–222.
- Johnson, B. F. G., Lahoz, F. J., Lewis, J., Prior, N. D., Raithby, P. R. & Wong, W.-T. (1992). *J. Chem. Soc. Dalton Trans.* pp. 1701–1708.
- Main, P., Fiske, S. J., Hull, S. E., Lessinger, L., Germain, G., Declercq, J.-P. & Woolfson, M. M. (1982). *MULTAN11/82. A System of Computer Programs for the Automatic Solution of Crystal Structures from X-ray Diffraction Data*. Univs. of York, England, and Louvain, Belgium.
- Orpen, A. G. (1980). *J. Chem. Soc. Dalton Trans.* pp. 2509–2516.

but there were discrepancies between the data sets reported. Besides, since photographic methods were used, both studies were not very precise. The value of R_f was not reported by Griffith (1943) but can be calculated from his published F_o and F_c values as 0.299 for 84 F_{hkl} . The value of R_f in the study of Kolesnikov & Baumer (1975) was 0.133 for 735 F_{hkl} . Since the precise data on the structure of $2Ag^+ \cdot C_2O_4^{2-}$, (I), are very important for understanding the mechanism of the photochemical and thermal decomposition of this salt (Boldyrev, 1993), we have undertaken a new structural study.

Acta Cryst. (1995). **C51**, 60–62

Silver Oxalate, $2Ag^+ \cdot C_2O_4^{2-}$

DMITRY YU. NAUMOV

*Novosibirsk State University, Pirogova Street 2,
Novosibirsk 630090, Russia*

ALEXANDER V. VIROVETS AND NINA V. PODBEREZSKAYA

*Institute of Inorganic Chemistry SD RAS,
Ak.Lavrent'eva Street 3, Novosibirsk 630090, Russia*

ELENA V. BOLDYREVA

*Institute of Solid State Chemistry SD RAS,
Derzhavina Street 18, Novosibirsk 630090, Russia*

(Received 6 April 1994; accepted 14 July 1994)


Abstract

The early data of Griffith [*J. Chem. Phys.* (1943), pp. 499–505] and of Kolesnikov & Baumer [*Vestn. Khar'k. Univ. Ser. Khim.* (1975), **127**, 38–41] on the structure of silver oxalate, $2Ag^+ \cdot C_2O_4^{2-}$, are revised and refined. In the structure, the oxalate anions form a framework with channels extended along [100] and Ag^+ cations are located in these channels, forming dimers with an $Ag—Ag$ bond distance of 2.945 (1) Å (close to that in metallic Ag). A noticeable elongation of C—C bond is observed.

Comment

Metal oxalates have been used in studies of the various aspects of solid-state reactivity since the early 1940's and are still attracting attention today (Brown, Dollimore & Galwey, 1980). The first structural studies of oxalates were reported 60 years ago (Hendricks, 1935); however, crystal structures of many metal oxalates remain unknown, mainly because of the experimental difficulties in obtaining single crystals.

Early structural studies of $2Ag^+ \cdot C_2O_4^{2-}$ were reported by Griffith (1943) and by Kolesnikov & Baumer (1975),



According to our data, the anion in the structure of silver oxalate is practically flat (deviations of C and O atoms from the common plane do not exceed 0.006 Å). The site symmetry is $\bar{1}$, but the geometry of the oxalate anion is very close to being idealized, with point symmetry mmm . The value of the C—C bond length (Table 2) somewhat exceeds the statistically averaged value [1.56 (2) Å] reported by Allen *et al.* (1987) for oxalate groups in various structures. The elongation of the C—C bond may be essential for the decomposition of silver oxalate, since, according to the model suggested, the primary step of the reaction is the cleaving of this bond (Boldyrev, 1993).

Each oxalate anion in the structure is coordinated to six Ag^+ cations. Two different types of $Ag—O$ bonds are observed (Table 2). Oxalate anions form a framework with the layers in the (200) planes and the channels extended along [100]. $Ag—Ag$ dimers are located in these channels; the $Ag—Ag$ distance of 2.945 (1) Å is close to that in metallic silver (2.889 Å; Guinier, 1956). Each Ag atom is coordinated to four O atoms of three oxalate anions (forming a flattened tetrahedron) and another Ag atom (Fig. 1). Ag atoms in the structure form strips of a 'stair type' extended along [100] at $y = \frac{1}{2}$ and $y = 0$ (corresponding values of z are ± 0.15 and $\frac{1}{2} \pm 0.15$). The width of a stair step is determined by the short $Ag—Ag$ distance; the height of a step is equal to the translation along x .

The results of our study allow revision of the data of Griffith (1943) and refinement of that of Kolesnikov & Baumer (1975). The values of the lattice parameters and the structure of the silver sub-lattice obtained in our study are in reasonable agreement with the data reported by Kolesnikov & Baumer (1975). They can be correlated also with the model of Griffith (1943) if one assumes that the value of the β angle is 104° (and not 76° , as stated in the text of the Griffith's paper). The structure of the oxalate sub-lattice in our