



Title	On opial-type integral inequalities
Author(s)	Cheung, WS; Zhao, CJ
Citation	Journal Of Inequalities And Applications, 2007, v. 2007, p. 38347
Issued Date	2007
URL	http://hdl.handle.net/10722/75449
Rights	Creative Commons: Attribution 3.0 Hong Kong License

Research Article

On Opial-Type Integral Inequalities

Wing-Sum Cheung and Chang-Jian Zhao

Received 22 January 2007; Accepted 4 April 2007

Recommended by Peter Yu Hin Pang

We establish some new Opial-type inequalities involving functions of two and many independent variables. Our results in special cases yield some of the recent results on Opial's inequality and also provide new estimates on inequalities of this type.

Copyright © 2007 W.-S. Cheung and C.-J. Zhao. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In the year 1960, Opial [1] established the following integral inequality.

THEOREM 1.1. *Suppose $f \in C^1[0, h]$ satisfies $f(0) = f(h) = 0$ and $f(x) > 0$ for all $x \in (0, h)$. Then the integral inequality holds*

$$\int_0^h |f(x)f'(x)| dx \leq \frac{h}{4} \int_0^h (f'(x))^2 dx, \quad (1.1)$$

where this constant $h/4$ is best possible.

Opial's inequality and its generalizations, extensions, and discretizations play a fundamental role in establishing the existence and uniqueness of initial and boundary value problems for ordinary and partial differential equations as well as difference equations [2–6]. The inequality (1.1) has received considerable attention and a large number of papers dealing with new proofs, extensions, generalizations, variants, and discrete analogs of Opial's inequality have appeared in some literature [7–22]. For an extensive survey on these inequalities, see [2, 6]. The main purpose of the present paper is to establish some new Opial-type inequalities involving functions of two and many independent variables. Our results in special cases yield some of the recent results on Opial's inequality and provide some new estimates on such types of inequalities.

2. Main results

Our main results are given in the following theorems.

THEOREM 2.1. *Let $u_i(s, t), v_i(s, t), i = 1, \dots, n$, be real-valued absolutely continuous functions defined on $[a, b] \times [c, d]$ and $a, b, c, d \in [0, \infty)$ with $u_i(s, c) = u_i(a, t) = u_i(a, c) = 0, v_i(s, c) = v_i(a, t) = v_i(a, c) = 0, i = 1, \dots, n$. Let F, G be real-valued nonnegative continuous and nondecreasing functions on $[0, \infty)^n$ with $F(0, \dots, 0) = 0, G(0, \dots, 0) = 0$ such that all their partial derivatives $\partial^2 F / \partial |u_i|^2, \partial F / \partial |u_i|, \partial^2 G / \partial |v_i|^2, \partial G / \partial |v_i|, i = 1, \dots, n$ are nonnegative continuous and nondecreasing functions on $[0, \infty)^n$. Let $\partial |u_i| / \partial s, \partial |u_i| / \partial t, \partial^2 |u_i| / \partial s \partial t, \partial |v_i| / \partial s, \partial |v_i| / \partial t, \partial^2 |v_i| / \partial s \partial t, i = 1, \dots, n$, be nonnegative continuous and nondecreasing functions on $[a, b] \times [c, d]$. Then*

$$\begin{aligned} & \int_a^b \int_c^d \left[F(|u_1(s, t)|, \dots, |u_n(s, t)|) \sum_{i=1}^n \left(\frac{\partial^2 G}{\partial |v_i|^2} \cdot \frac{\partial |v_i|}{\partial t} \cdot \frac{\partial |v_i|}{\partial s} + \frac{\partial G}{\partial |v_i|} \cdot \frac{\partial |v_i|}{\partial s \partial t} \right) \right. \\ & \quad + G(|v_1(s, t)|, \dots, |v_n(s, t)|) \sum_{i=1}^n \left(\frac{\partial^2 F}{\partial |u_i|^2} \cdot \frac{\partial |u_i|}{\partial t} \cdot \frac{\partial |u_i|}{\partial s} + \frac{\partial F}{\partial |u_i|} \cdot \frac{\partial |u_i|}{\partial s \partial t} \right) \\ & \quad \left. + S(s, t) \right] ds dt \\ & \leq F \left(\int_a^b \int_c^d \left| \frac{\partial^2 u_1}{\partial s \partial t} \right| ds dt, \dots, \int_a^b \int_c^d \left| \frac{\partial^2 u_n}{\partial s \partial t} \right| ds dt \right) \\ & \quad \cdot G \left(\int_a^b \int_c^d \left| \frac{\partial^2 v_1}{\partial s \partial t} \right| ds dt, \dots, \int_a^b \int_c^d \left| \frac{\partial^2 v_n}{\partial s \partial t} \right| ds dt \right), \end{aligned} \tag{2.1}$$

where

$$S(s, t) = \sum_{i=1}^n \frac{\partial F}{\partial |u_i|} \frac{\partial |u_i|}{\partial s} \cdot \sum_{i=1}^n \frac{\partial G}{\partial |v_i|} \frac{\partial |v_i|}{\partial t} + \sum_{i=1}^n \frac{\partial F}{\partial |u_i|} \frac{\partial |u_i|}{\partial t} \cdot \sum_{i=1}^n \frac{\partial G}{\partial |v_i|} \frac{\partial |v_i|}{\partial s}. \tag{2.2}$$

Proof. From the hypotheses on $u_i(s, t), v_i(s, t), i = 1, \dots, n$, we have

$$\begin{aligned} |u_i(s, t)| & \leq \int_a^s \int_c^t \left| \frac{\partial^2 u_i}{\partial \sigma \partial \tau}(\sigma, \tau) \right| d\sigma d\tau, \\ |v_i(s, t)| & \leq \int_a^s \int_c^t \left| \frac{\partial^2 v_i}{\partial \sigma \partial \tau}(\sigma, \tau) \right| d\sigma d\tau, \end{aligned} \tag{2.3}$$

for $s \in [a, b], t \in [c, d]$.

From (2.3) and in view of the hypotheses on all partial derivatives, and by letting

$$\begin{aligned} U_i(s, t) &= \int_a^s \int_c^t \left| \frac{\partial^2 u_i}{\partial \sigma \partial \tau}(\sigma, \tau) \right| d\sigma d\tau, \\ V_i(s, t) &= \int_a^s \int_c^t \left| \frac{\partial^2 v_i}{\partial \sigma \partial \tau}(\sigma, \tau) \right| d\sigma d\tau, \end{aligned} \quad (2.4)$$

we obtain

$$\begin{aligned} & \int_a^b \int_c^d \left[F(|u_1(s, t)|, \dots, |u_n(s, t)|) \sum_{i=1}^n \left(\frac{\partial^2 G}{\partial |v_i|^2} \cdot \frac{\partial |v_i|}{\partial t} \cdot \frac{\partial |v_i|}{\partial s} + \frac{\partial G}{\partial |v_i|} \cdot \frac{\partial^2 |v_i|}{\partial s \partial t} \right) \right. \\ & \quad + G(|v_1(s, t)|, \dots, |v_n(s, t)|) \sum_{i=1}^n \left(\frac{\partial^2 F}{\partial |u_i|^2} \cdot \frac{\partial |u_i|}{\partial t} \cdot \frac{\partial |u_i|}{\partial s} + \frac{\partial F}{\partial |u_i|} \cdot \frac{\partial^2 |u_i|}{\partial s \partial t} \right) \\ & \quad + \sum_{i=1}^n \frac{\partial F}{\partial |u_i|} \cdot \frac{\partial |u_i|}{\partial s} \cdot \sum_{i=1}^n \frac{\partial G}{\partial |v_i|} \cdot \frac{\partial |v_i|}{\partial t} \\ & \quad \left. + \sum_{i=1}^n \frac{\partial F}{\partial |u_i|} \cdot \frac{\partial |u_i|}{\partial t} \cdot \sum_{i=1}^n \frac{\partial G}{\partial |v_i|} \cdot \frac{\partial |v_i|}{\partial s} \right] ds dt \\ & \leq \int_a^b \int_c^d \left[F(U_1(s, t), \dots, U_n(s, t)) \cdot \sum_{i=1}^n \left(\frac{\partial^2 G}{\partial V_i^2} \cdot \frac{\partial V_i}{\partial t} \cdot \frac{\partial V_i}{\partial s} + \frac{\partial G}{\partial V_i} \cdot \frac{\partial^2 V_i}{\partial s \partial t} \right) \right. \\ & \quad + G(V_1(s, t), \dots, V_n(s, t)) \cdot \sum_{i=1}^n \left(\frac{\partial^2 F}{\partial U_i^2} \cdot \frac{\partial U_i}{\partial t} \cdot \frac{\partial U_i}{\partial s} + \frac{\partial F}{\partial U_i} \cdot \frac{\partial^2 U_i}{\partial s \partial t} \right) \\ & \quad \left. + \sum_{i=1}^n \frac{\partial F}{\partial U_i} \frac{\partial U_i}{\partial s} \cdot \sum_{i=1}^n \frac{\partial G}{\partial V_i} \frac{\partial V_i}{\partial t} + \sum_{i=1}^n \frac{\partial F}{\partial U_i} \frac{\partial U_i}{\partial t} \cdot \sum_{i=1}^n \frac{\partial G}{\partial V_i} \frac{\partial V_i}{\partial s} \right] ds dt \\ & = \int_a^b \int_c^d \frac{\partial^2}{\partial s \partial t} [F(U_1(s, t), \dots, U_n(s, t)) \cdot G(V_1(s, t), \dots, V_n(s, t))] ds dt \\ & = F(U_1(b, d), \dots, U_n(b, d)) \cdot G(V_1(b, d), \dots, V_n(b, d)) \\ & = F \left(\int_a^b \int_c^d \left| \frac{\partial^2 u_1}{\partial s \partial t} \right| ds dt, \dots, \int_a^b \int_c^d \left| \frac{\partial^2 u_n}{\partial s \partial t} \right| ds dt \right) \\ & \quad \cdot G \left(\int_a^b \int_c^d \left| \frac{\partial^2 v_1}{\partial s \partial t} \right| ds dt, \dots, \int_a^b \int_c^d \left| \frac{\partial^2 v_n}{\partial s \partial t} \right| ds dt \right). \end{aligned} \quad (2.5)$$

This completes the proof of inequality (2.1). \square

Remark 2.2. (i) Taking $G = 1$ in inequality (2.1), and in view of

$$\frac{\partial^2 G}{\partial |v_i|^2} \cdot \frac{\partial |v_i|}{\partial t} \cdot \frac{\partial |v_i|}{\partial s} + \frac{\partial G}{\partial |v_i|} \frac{\partial |v_i|}{\partial s \partial t} = 0, \quad S(s, t) = 0, \tag{2.6}$$

for $i = 1, \dots, n$, we have

$$\begin{aligned} & \int_a^b \int_c^d \left[\sum_{i=1}^n \left(\frac{\partial^2 F}{\partial |u_i|^2} \cdot \frac{\partial |u_i|}{\partial t} \cdot \frac{\partial |u_i|}{\partial s} + \frac{\partial F}{\partial |u_i|} \cdot \frac{\partial^2 |u_i|}{\partial s \partial t} \right) \right] ds dt \\ & \leq F \left(\int_a^b \int_c^d \left| \frac{\partial^2 u_1}{\partial s \partial t} \right| ds dt, \dots, \int_a^b \int_c^d \left| \frac{\partial^2 u_n}{\partial s \partial t} \right| ds dt \right), \end{aligned} \tag{2.7}$$

for $i = 1, \dots, n$.

Let $u_i(s, t)$ reduce to $u_i(t)$, where $i = 1, \dots, n$ and with suitable modifications, then (2.7) becomes the following inequality:

$$\int_a^b \left[\sum_{i=1}^n F'_i (|u_1(t)|, \dots, |u_n(t)|) |u'_i(t)| \right] dt \leq F \left(\int_a^b |u'_1(t)| dt, \dots, \int_a^b |u'_n(t)| dt \right). \tag{2.8}$$

This is a recent inequality which was given by Pečarić and Brnetić [18, 19].

Taking $n = 1$, inequality (2.7) reduces to

$$\int_a^b \int_c^d \left(\frac{\partial^2 F}{\partial |u|^2} \cdot \frac{\partial |u|}{\partial t} \cdot \frac{\partial |u|}{\partial s} + \frac{\partial F}{\partial |u|} \cdot \frac{\partial^2 |u|}{\partial s \partial t} \right) ds dt \leq F \left(\int_a^b \int_c^d \left| \frac{\partial^2 u}{\partial s \partial t} \right| ds dt \right). \tag{2.9}$$

Let $u(s, t)$ reduce to $u(t)$ and with suitable modifications, then the above inequality becomes the following inequality:

$$\int_a^b F'(|f(t)|) |f'(t)| dt \leq F \left(\int_a^b |f'(x)| dt \right). \tag{2.10}$$

This is an inequality which was given by Godunova and Levin [12].

(ii) Taking $G = F$ and $u_i(s, t) = v_i(s, t)$, $i = 1, \dots, n$, in inequality (2.1), we have

$$\begin{aligned} & \int_a^b \int_c^d \left[F(|u_1(s, t)|, \dots, |u_n(s, t)|) \sum_{i=1}^n \left(\frac{\partial^2 G}{\partial |u_i|^2} \cdot \frac{\partial |u_i|}{\partial t} \cdot \frac{\partial |u_i|}{\partial s} + \frac{\partial G}{\partial |u_i|} \frac{\partial |u_i|}{\partial s \partial t} \right) \right. \\ & \quad \left. + \sum_{i=1}^n \frac{\partial F}{\partial u_i} \frac{\partial u_i}{\partial s} \cdot \sum_{i=1}^n \frac{\partial F}{\partial u_i} \frac{\partial u_i}{\partial t} \right] ds dt \\ & \leq \frac{1}{2} \cdot F^2 \left(\int_a^b \int_c^d \left| \frac{\partial^2 u_1}{\partial s \partial t} \right| ds dt, \dots, \int_a^b \int_c^d \left| \frac{\partial^2 u_n}{\partial s \partial t} \right| ds dt \right). \end{aligned} \tag{2.11}$$

Taking $n = 1$, (2.11) reduces to

$$\int_a^b \int_c^d \left[F(|u(s,t)|) \left(\frac{\partial^2 G}{\partial |u|^2} \cdot \frac{\partial |u|}{\partial t} \cdot \frac{\partial |u|}{\partial s} + \frac{\partial G}{\partial |u|} \frac{\partial |u|}{\partial s \partial t} \right) + \frac{\partial F}{\partial u} \frac{\partial u}{\partial s} \frac{\partial F}{\partial u} \frac{\partial u}{\partial t} \right] ds dt \tag{2.12}$$

$$\leq \frac{1}{2} \cdot F^2 \left(\int_a^b \int_c^d \left| \frac{\partial^2 u}{\partial s \partial t} \right| ds dt \right).$$

Let $u(s,t)$ reduce to $u(t)$ and with suitable modifications, then (2.12) becomes the following inequality:

$$\int_a^b [F(|u(t)|) \cdot F'(|u(t)|) \cdot |u'(t)|] dt \leq \frac{1}{2} F^2 \left(\int_a^b |u'(t)| dt \right). \tag{2.13}$$

This is an inequality given by Pachpatte in [15].

Inequality (2.12) is also a similar form of the following inequality which was given by Yang [22]:

$$\int_{a_1}^{b_1} \int_{a_2}^{b_2} \left| f(t_1, t_2) \frac{\partial^2 f}{\partial t_1 \partial t_2} \right| dt_1 dt_2 \leq \frac{(b_1 - a_1)(b_2 - a_2)}{8} \int_{a_1}^{b_1} \int_{a_2}^{b_2} \left(\frac{\partial^2 f}{\partial t_1 \partial t_2}(t_1, t_2) \right)^2 dt_1 dt_2. \tag{2.14}$$

(iii) Let $u_i(s,t)$ and $v_i(s,t)$ reduce to $u_i(s)$ and $v_i(s)$, respectively, and with suitable modifications (where $i = 1, \dots, n$), then inequality (2.1) changes to the following inequality:

$$\int_a^b \left[F(|u_1(t)|, \dots, |u_n(t)|) \sum_{i=1}^n G_i(|v_1(t)|, \dots, |v_n(t)|) |v'_i(t)| \right. \\ \left. + G(|v_1(t)|, \dots, |v_n(t)|) \sum_{i=1}^n F'_i(|u_1(t)|, \dots, |u_n(t)|) |u'_i(t)| \right] dt \\ \leq F \left(\int_a^b |u'_1(t)| dt, \dots, \int_a^b |u'_n(t)| dt \right) \cdot G \left(\int_a^b |v'_1(t)| dt, \dots, \int_a^b |v'_n(t)| dt \right). \tag{2.15}$$

This is an inequality given by Agarwal and Pang in [2].

Taking $n = 1$, $G = 1$, $F(u) = u^2$, (2.15) changes to

$$\int_a^b |u(t)| |u'(t)| dt \leq \frac{1}{2} (b - a) \int_a^b |u'(t)|^2 dt. \tag{2.16}$$

This is another version of the Opial's inequality, (see [13]).

(iv) Taking $G = 1$, $F = (|u_1|, \dots, |u_n|) = \prod_{i=1}^n f_i(|u_i|)$, $i = 1, \dots, n$, in (2.1), (2.1) changes to a general form of the inequality which was given by Pachpatte [16], where the functions f_i must satisfy some suitable conditions, (see [16]).

THEOREM 2.3. Let $u_i(s, t)$, $v_i(s, t)$, F , G , $\partial^2 F/\partial|u_i|^2$, $\partial F/\partial|u_i|$, $\partial|u_i|/\partial s$, $\partial|u_i|/\partial t$, $\partial^2|u_i|/\partial s\partial t$, $\partial^2 G/\partial|v_i|^2$, $\partial G/\partial|v_i|$, $\partial|v_i|/\partial s$, $\partial|v_i|/\partial t$, $\partial^2|v_i|/\partial s\partial t$, $i = 1, \dots, n$, be as in Theorem 2.1. Let $p_i(s, t)$, $q_i(s, t)$, $i = 1, \dots, n$, be real-valued positive functions defined on $[a, b] \times [c, d]$ satisfying

$$\int_a^b \int_c^d p_i(s, t) ds dt = 1, \quad \int_a^b \int_c^d q_i(s, t) ds dt = 1 \quad (i = 1, \dots, n). \tag{2.17}$$

Let h_i , w_i , $i = 1, \dots, n$, be real-valued positive convex and increasing functions on $(0, \infty)^2$. Then the following integral inequality holds:

$$\begin{aligned} & \int_a^b \int_c^d \left[F(|u_1(s, t)|, \dots, |u_n(s, t)|) \sum_{i=1}^n \left(\frac{\partial^2 G}{\partial|v_i|^2} \cdot \frac{\partial|v_i|}{\partial t} \cdot \frac{\partial|v_i|}{\partial s} + \frac{\partial G}{\partial|v_i|} \frac{\partial^2|v_i|}{\partial s\partial t} \right) \right. \\ & \quad + G(|v_1(s, t)|, \dots, |v_n(s, t)|) \sum_{i=1}^n \left(\frac{\partial^2 F}{\partial|u_i|^2} \cdot \frac{\partial|u_i|}{\partial t} \cdot \frac{\partial|u_i|}{\partial s} + \frac{\partial F}{\partial|u_i|} \frac{\partial^2|u_i|}{\partial s\partial t} \right) \\ & \quad \left. + S(s, t) \right] ds dt \\ & \leq F \left[h_1^{-1} \left(\int_a^b \int_c^d p_1(s, t) h_1 \left(\frac{|\partial^2 u_1/\partial s\partial t|}{p_1(s, t)} \right) ds dt \right), \dots, \right. \\ & \quad \left. h_n^{-1} \left(\int_a^b \int_c^d p_n(s, t) h_n \left(\frac{|\partial^2 u_n/\partial s\partial t|}{p_n(s, t)} \right) ds dt \right) \right] \\ & \quad \cdot G \left[w_1^{-1} \left(\int_a^b \int_c^d q_1(s, t) w_1 \left(\frac{|\partial^2 v_1/\partial s\partial t|}{q_1(s, t)} \right) ds dt \right), \dots, \right. \\ & \quad \left. w_n^{-1} \left(\int_a^b \int_c^d q_n(s, t) w_n \left(\frac{|\partial^2 v_n/\partial s\partial t|}{q_n(s, t)} \right) ds dt \right) \right], \end{aligned} \tag{2.18}$$

where

$$S(s, t) = \sum_{i=1}^n \frac{\partial F}{\partial|u_i|} \frac{\partial|u_i|}{\partial s} \cdot \sum_{i=1}^n \frac{\partial G}{\partial|v_i|} \frac{\partial|v_i|}{\partial t} + \sum_{i=1}^n \frac{\partial F}{\partial|u_i|} \frac{\partial|u_i|}{\partial t} \cdot \sum_{i=1}^n \frac{\partial G}{\partial|v_i|} \frac{\partial|v_i|}{\partial s}. \tag{2.19}$$

Proof. From the hypotheses, we have

$$\begin{aligned} \int_a^b \int_c^d \left| \frac{\partial u_i^2}{\partial s\partial t} \right| ds dt &= \frac{\int_a^b \int_c^d p_i(s, t) |\partial u_i^2/\partial s\partial t| / p_i(s, t) ds dt}{\int_a^b \int_c^d p_i(s, t) ds dt}, \\ \int_a^b \int_c^d \left| \frac{\partial v_i^2}{\partial s\partial t} \right| ds dt &= \frac{\int_a^b \int_c^d q_i(s, t) |\partial v_i^2/\partial s\partial t| / q_i(s, t) ds dt}{\int_a^b \int_c^d q_i(s, t) ds dt}, \end{aligned} \tag{2.20}$$

for $i = 1, \dots, n$.

From (2.20), the hypotheses on h_i , w_i , $i = 1, \dots, n$, and in view of Jensen's inequality, we obtain

$$\begin{aligned} h_i \left(\int_a^b \int_c^d \left| \frac{\partial u_i^2}{\partial s \partial t} \right| ds dt \right) &\leq \int_a^b \int_c^d p_i(s, t) \cdot h_i \left(\frac{|\partial^2 u_i / \partial s \partial t|}{p_i(s, t)} \right) ds dt, \\ w_i \left(\int_a^b \int_c^d \left| \frac{\partial v_i^2}{\partial s \partial t} \right| ds dt \right) &\leq \int_a^b \int_c^d q_i(s, t) \cdot w_i \left(\frac{|\partial^2 v_i / \partial s \partial t|}{q_i(s, t)} \right) ds dt, \end{aligned} \quad (2.21)$$

for $i = 1, \dots, n$.

From (2.21), we observe that

$$\begin{aligned} \int_a^b \int_c^d \left| \frac{\partial u_i^2}{\partial s \partial t} \right| ds dt &\leq h_{(i-1)} \left(\int_a^b \int_c^d p_i(s, t) \cdot h_i \left(\frac{|\partial^2 u_i / \partial s \partial t|}{p_i(s, t)} \right) ds dt \right), \\ \int_a^b \int_c^d \left| \frac{\partial v_i^2}{\partial s \partial t} \right| ds dt &\leq w_{(i-1)} \left(\int_a^b \int_c^d q_i(s, t) \cdot w_i \left(\frac{|\partial^2 v_i / \partial s \partial t|}{q_i(s, t)} \right) ds dt \right), \end{aligned} \quad (2.22)$$

for $i = 1, \dots, n$.

From (2.22) and in view of inequality (2.1), we get inequality (2.18) and the proof is complete. \square

Remark 2.4. (i) Taking $G = 1$ in inequality (2.18), and in view of

$$\frac{\partial^2 G}{\partial |v_i|^2} \cdot \frac{\partial |v_i|}{\partial t} \cdot \frac{\partial |v_i|}{\partial s} + \frac{\partial G}{\partial |v_i|} \frac{\partial |v_i|}{\partial s \partial t} = 0, \quad (2.23)$$

for $i = 1, \dots, n$, and

$$S(s, t) = 0, \quad (2.24)$$

(2.18) becomes

$$\begin{aligned} &\int_a^b \int_c^d \left[\sum_{i=1}^n \left(\frac{\partial^2 F}{\partial |u_i|^2} \cdot \frac{\partial |u_i|}{\partial t} \cdot \frac{\partial |u_i|}{\partial s} + \frac{\partial F}{\partial |u_i|} \cdot \frac{\partial^2 |u_i|}{\partial s \partial t} \right) \right] ds dt \\ &\leq F \left[h_1^{-1} \left(\int_a^b \int_c^d p_1(s, t) h_1 \left(\frac{|\partial^2 u_1 / \partial s \partial t|}{p_1(s, t)} \right) ds dt \right), \dots, \right. \\ &\quad \left. h_n^{-1} \left(\int_a^b \int_c^d p_n(s, t) h_n \left(\frac{|\partial^2 u_n / \partial s \partial t|}{p_n(s, t)} \right) ds dt \right) \right], \end{aligned} \quad (2.25)$$

for $i = 1, \dots, n$.

Let $u_i(s, t)$, $h_i(s, t)$, and $p_i(s, t)$ change to $f_i(t)$, $h_i(t)$, and $p_i(t)$, respectively, where $i = 1, \dots, n$, then (2.25) reduces to the following inequality:

$$\int_a^b \left(\sum_{i=1}^n D_i F(|f_1(t)|, \dots, |f_n(t)| |f'_i(t)|) dt \right) \leq F \left(h_1^{-1} \left(\int_a^b p_1(t) h_1 \left(\frac{|f'_1(t)|}{p_1(t)} \right) dt \right), \dots, h_n^{-1} \left(\int_a^b p_n(t) h_n \left(\frac{|f'_n(t)|}{p_n(t)} \right) dt \right) \right), \tag{2.26}$$

where $D_i F$ is as in [18]. This is an inequality given by Pečarić in [18].

Taking $F(x_1, \dots, x_n) = \prod_{i=1}^n F_i(x_i)$, $i = 1, \dots, n$, (2.25) changes to a general form of the inequality which was given by Pachpatte [16]. Taking $n = 1$, (2.25) reduces to a general form of the inequality which was given by Godunova and Levin [12].

On the other hand, inequality (2.18) is also a general form of another inequality in Pečarić and Brnetić [20, Theorem 1].

(ii) Taking $G = F$ and $u_i(s, t) = v_i(s, t)$, $i = 1, \dots, n$, in inequality (2.18), we have

$$\int_a^b \int_c^d \left[F(|u_1(s, t)|, \dots, |u_n(s, t)|) \sum_{i=1}^n \left(\frac{\partial^2 G}{\partial |u_i|^2} \cdot \frac{\partial |u_i|}{\partial t} \cdot \frac{\partial |u_i|}{\partial s} + \frac{\partial G}{\partial |u_i|} \frac{\partial |u_i|}{\partial s \partial t} \right) + \sum_{i=1}^n \frac{\partial F}{\partial u_i} \frac{\partial u_i}{\partial s} \cdot \sum_{i=1}^n \frac{\partial F}{\partial u_i} \frac{\partial u_i}{\partial t} \right] ds dt \leq \frac{1}{2} \cdot F^2 \left[h_1^{-1} \left(\int_a^b \int_c^d p_1(s, t) h_1 \left(\frac{|\partial^2 u_1 / \partial s \partial t|}{p_1(s, t)} \right) ds dt \right), \dots, h_n^{-1} \left(\int_a^b \int_c^d p_n(s, t) h_n \left(\frac{|\partial^2 u_n / \partial s \partial t|}{p_n(s, t)} \right) ds dt \right) \right]. \tag{2.27}$$

Taking $n = 1$, (2.27) reduces to

$$\int_a^b \int_c^d \left[F(|u(s, t)|) \left(\frac{\partial^2 G}{\partial |u|^2} \cdot \frac{\partial |u|}{\partial t} \cdot \frac{\partial |u|}{\partial s} + \frac{\partial G}{\partial |u|} \frac{\partial |u|}{\partial s \partial t} \right) + \frac{\partial F}{\partial u} \frac{\partial u}{\partial s} \frac{\partial F}{\partial u} \frac{\partial u}{\partial t} \right] ds dt \leq \frac{1}{2} \cdot F^2 \left(h^{-1} \left(\int_a^b \int_c^d p(s, t) h \left(\frac{|\partial^2 u / \partial s \partial t|}{p(s, t)} \right) ds dt \right) \right). \tag{2.28}$$

This is a general form of the inequality which was given by Pachpatte [14].

(iii) Let $u_i(s, t)$, $v_i(s, t)$, $h_i(s, t)$, $w_i(s, t)$, $p_i(s, t)$, and $q_i(s, t)$ reduce to $u_i(t)$, $v_i(t)$, $h_i(t)$, $w_i(t)$, $p_i(t)$, and $q_i(t)$, respectively, and with suitable modifications (where $i = 1, \dots, n$),

then inequality (2.18) changes to the following inequality:

$$\begin{aligned}
 & \int_a^b \left[F(|u_1(t)|, \dots, |u_n(t)|) \sum_{i=1}^n G'_i(|v_1(t)|, \dots, |v_n(t)|) |v'_i(t)| \right. \\
 & \quad \left. + G(|v_1(t)|, \dots, |v_n(t)|) \sum_{i=1}^n F'_i(|u_1(t)|, \dots, |u_n(t)|) |u'_i(t)| \right] dt \\
 & \leq F\left(h_1^{-1}\left(\int_a^b p_1(t) h_1\left(\frac{|u'_1(t)|}{p_1(t)}\right) dt\right), \dots, h_n^{-1}\left(\int_a^b p_n(t) h_n\left(\frac{|u'_n(t)|}{p_n(t)}\right) dt\right)\right) \\
 & \quad \cdot G\left(w_1^{-1}\left(\int_a^b q_1(t) w_1\left(\frac{|v'_1(t)|}{q_1(t)}\right) dt\right), \dots, w_n^{-1}\left(\int_a^b q_n(t) w_n\left(\frac{|v'_n(t)|}{q_n(t)}\right) dt\right)\right).
 \end{aligned} \tag{2.29}$$

This is just an inequality given by Agarwal and Pang in [2].

THEOREM 2.5. *Let $u_i(s, t)$, $v_i(s, t)$, F , G , be as in Theorem 2.1. Let $\phi_i, \psi_i, i = 1, \dots, n$, be real-valued positive convex and increasing functions on $(0, \infty)^2$. Let $r_i(s, t) \geq 0$, $\partial^2 r_i / \partial s \partial t > 0$, $r_i(s, c) = r_i(a, t) = r_i(a, c) = 0$, $\partial^2 e_i / \partial s \partial t > 0$, $e_i(s, c) = e_i(a, t) = e_i(a, c) = 0$, $i = 1, \dots, n$. Let $\partial^2 F / \partial \overline{M}_i^2$, $\partial F / \partial \overline{M}_i$, $\partial^2 G / \partial \overline{N}_i^2$, $\partial G / \partial \overline{N}_i$, $i = 1, \dots, n$, be nonnegative continuous and nondecreasing functions on $[0, \infty)^n$. Let $\partial \overline{M}_i / \partial s$, $\partial \overline{M}_i / \partial t$, $\partial^2 \overline{M}_i / \partial s \partial t$, $\partial \overline{N}_i / \partial s$, $\partial \overline{N}_i / \partial t$, $\partial^2 \overline{N}_i / \partial s \partial t$, $i = 1, \dots, n$, be nonnegative continuous and nondecreasing functions on $[a, b] \times [c, d]$. Then the following inequality holds:*

$$\begin{aligned}
 & \int_a^b \int_c^d \left[F(\overline{M}_1(s, t), \dots, \overline{M}_n(s, t)) \cdot \sum_{i=1}^n \left(\frac{\partial^2 G}{\partial \overline{N}_i^2} \cdot \frac{\partial \overline{N}_i}{\partial t} \cdot \frac{\partial \overline{N}_i}{\partial s} + \frac{\partial G}{\partial \overline{N}_i} \cdot \frac{\partial^2 \overline{N}_i}{\partial s \partial t} \right) \right. \\
 & \quad \left. + G(\overline{N}_1(s, t), \dots, \overline{N}_n(s, t)) \cdot \sum_{i=1}^n \left(\frac{\partial^2 F}{\partial \overline{M}_i^2} \cdot \frac{\partial \overline{M}_i}{\partial t} \cdot \frac{\partial \overline{M}_i}{\partial s} + \frac{\partial F}{\partial \overline{M}_i} \cdot \frac{\partial^2 \overline{M}_i}{\partial s \partial t} \right) + \overline{S}(s, t) \right] ds dt \\
 & \leq F\left(\int_a^b \int_c^d \frac{\partial^2 r_1}{\partial s \partial t} \cdot \phi_1\left(\frac{|\partial^2 u_1 / \partial s \partial t|}{\partial^2 r_1 / \partial s \partial t}\right) ds dt, \dots, \int_a^b \int_c^d \frac{\partial^2 r_n}{\partial s \partial t} \cdot \phi_n\left(\frac{|\partial^2 u_n / \partial s \partial t|}{\partial^2 r_n / \partial s \partial t}\right) ds dt\right) \\
 & \quad \cdot G\left(\int_a^b \int_c^d \frac{\partial^2 e_1}{\partial s \partial t} \cdot \psi_1\left(\frac{|\partial^2 v_1 / \partial s \partial t|}{\partial^2 e_1 / \partial s \partial t}\right) ds dt, \dots, \int_a^b \int_c^d \frac{\partial^2 e_n}{\partial s \partial t} \cdot \psi_n\left(\frac{|\partial^2 v_n / \partial s \partial t|}{\partial^2 e_n / \partial s \partial t}\right) ds dt\right),
 \end{aligned} \tag{2.30}$$

where

$$\begin{aligned}
 \overline{M}_i(s, t) &= r_i(s, t) \cdot \phi_i\left(\frac{|u_i(s, t)|}{r_i(s, t)}\right), \\
 \overline{N}_i(s, t) &= e_i(s, t) \cdot \psi_i\left(\frac{|v_i(s, t)|}{e_i(s, t)}\right),
 \end{aligned} \tag{2.31}$$

for $i = 1, \dots, n$, and

$$\bar{S}(s, t) = \sum_{i=1}^n \frac{\partial F}{\partial \bar{M}_i} \frac{\partial \bar{M}_i}{\partial s} \cdot \sum_{i=1}^n \frac{\partial G}{\partial \bar{N}_i} \frac{\partial \bar{N}_i}{\partial t} + \sum_{i=1}^n \frac{\partial F}{\partial \bar{M}_i} \frac{\partial \bar{M}_i}{\partial t} \cdot \sum_{i=1}^n \frac{\partial G}{\partial \bar{N}_i} \frac{\partial \bar{N}_i}{\partial s}, \tag{2.32}$$

for $i = 1, \dots, n$.

Proof. From the hypotheses on $u_i(s, t)$, $v_i(s, t)$, $r_i(s, t)$, $e_i(s, t)$, $i = 1, \dots, n$, we have

$$\begin{aligned} |u_i(s, t)| &\leq \int_a^s \int_c^t \left| \frac{\partial^2 u_i}{\partial \sigma \partial \tau}(\sigma, \tau) \right| d\sigma d\tau, \\ |v_i(s, t)| &\leq \int_a^s \int_c^t \left| \frac{\partial^2 v_i}{\partial \sigma \partial \tau}(\sigma, \tau) \right| d\sigma d\tau, \\ r_i(s, t) &= \int_a^s \int_c^t \frac{\partial^2 r_i}{\partial \sigma \partial \tau}(\sigma, \tau) d\sigma d\tau, \\ e_i(s, t) &= \int_a^s \int_c^t \frac{\partial^2 e_i}{\partial \sigma \partial \tau}(\sigma, \tau) d\sigma d\tau, \end{aligned} \tag{2.33}$$

for $s \in [a, b]$, $t \in [c, d]$.

From (2.33) and using the hypotheses on ϕ_i , ψ_i , $i = 1, \dots, n$, and Jensen's inequality, we have

$$\begin{aligned} \bar{M}_i(s, t) &\leq \int_a^s \int_c^t \frac{\partial^2 r_i}{\partial \sigma \partial \tau} \cdot \phi_i \left(\frac{|\partial^2 u_i / \partial \sigma \partial \tau(\sigma, \tau)|}{(\partial^2 r_i / \partial \sigma \partial \tau)(\sigma, \tau)} \right) d\sigma d\tau, \\ \bar{N}_i(s, t) &\leq \int_a^s \int_c^t \frac{\partial^2 e_i}{\partial \sigma \partial \tau} \cdot \psi_i \left(\frac{|\partial^2 v_i / \partial \sigma \partial \tau(\sigma, \tau)|}{(\partial^2 e_i / \partial \sigma \partial \tau)(\sigma, \tau)} \right) d\sigma d\tau, \end{aligned} \tag{2.34}$$

for $s \in [a, b]$, $t \in [c, d]$.

From (2.34), using the hypotheses on all partial derivatives and in view of

$$\begin{aligned} M_i(s, t) &= \int_a^s \int_c^t \frac{\partial^2 r_i}{\partial \sigma \partial \tau} \cdot \phi_i \left(\frac{|\partial^2 u_i / \partial \sigma \partial \tau|}{\partial^2 r_i / \partial \sigma \partial \tau} \right) d\sigma d\tau, \\ N_i(s, t) &= \int_a^s \int_c^t \frac{\partial^2 e_i}{\partial \sigma \partial \tau} \cdot \psi_i \left(\frac{|\partial^2 v_i / \partial \sigma \partial \tau|}{\partial^2 e_i / \partial \sigma \partial \tau} \right) d\sigma d\tau, \end{aligned} \tag{2.35}$$

we have

$$\begin{aligned}
 & \int_a^b \int_c^d \left[F(\overline{M}_1(s,t), \dots, \overline{M}_n(s,t)) \cdot \sum_{i=1}^n \left(\frac{\partial^2 G}{\partial \overline{N}_i^2} \cdot \frac{\partial \overline{N}_i}{\partial t} \cdot \frac{\partial \overline{N}_i}{\partial s} + \frac{\partial G}{\partial \overline{N}_i} \cdot \frac{\partial^2 \overline{N}_i}{\partial s \partial t} \right) \right. \\
 & \quad \left. + G(\overline{N}_1(s,t), \dots, \overline{N}_n(s,t)) \cdot \sum_{i=1}^n \left(\frac{\partial^2 F}{\partial \overline{M}_i^2} \cdot \frac{\partial \overline{M}_i}{\partial t} \cdot \frac{\partial \overline{M}_i}{\partial s} + \frac{\partial F}{\partial \overline{M}_i} \cdot \frac{\partial^2 \overline{M}_i}{\partial s \partial t} \right) + \overline{S}(s,t) \right] ds dt \\
 & \leq \int_a^b \int_c^d \left[F(M_1(s,t), \dots, M_n(s,t)) \cdot \sum_{i=1}^n \left(\frac{\partial^2 G}{\partial N_i^2} \cdot \frac{\partial N_i}{\partial t} \cdot \frac{\partial N_i}{\partial s} + \frac{\partial G}{\partial N_i} \cdot \frac{\partial^2 N_i}{\partial s \partial t} \right) \right. \\
 & \quad \left. + G(N_1(s,t), \dots, N_n(s,t)) \cdot \sum_{i=1}^n \left(\frac{\partial^2 F}{\partial M_i^2} \cdot \frac{\partial M_i}{\partial t} \cdot \frac{\partial M_i}{\partial s} + \frac{\partial F}{\partial M_i} \cdot \frac{\partial^2 M_i}{\partial s \partial t} \right) \right. \\
 & \quad \left. + \sum_{i=1}^n \frac{\partial F}{\partial M_i} \frac{\partial M_i}{\partial s} \cdot \sum_{i=1}^n \frac{\partial G}{\partial N_i} \frac{\partial N_i}{\partial t} + \sum_{i=1}^n \frac{\partial F}{\partial M_i} \frac{\partial M_i}{\partial t} \cdot \sum_{i=1}^n \frac{\partial G}{\partial N_i} \frac{\partial N_i}{\partial s} \right] ds dt \\
 & = \int_a^b \int_c^d \frac{\partial^2}{\partial s \partial t} \left[F(M_1(s,t), \dots, M_n(s,t)) \cdot G(N_1(s,t), \dots, N_n(s,t)) \right] ds dt \\
 & = F(M_1(b,d), \dots, M_n(b,d)) \cdot G(N_1(b,d), \dots, N_n(b,d)) \\
 & = F \left(\int_a^b \int_c^d \frac{\partial^2 r_1}{\partial s \partial t} \cdot \phi_1 \left(\frac{|\partial^2 u_1 / \partial s \partial t|}{\partial^2 r_1 / \partial s \partial t} \right) ds dt, \dots, \int_a^b \int_c^d \frac{\partial^2 r_n}{\partial s \partial t} \cdot \phi_n \left(\frac{|\partial^2 u_n / \partial s \partial t|}{\partial^2 r_n / \partial s \partial t} \right) ds dt \right) \\
 & \quad \cdot G \left(\int_a^b \int_c^d \frac{\partial^2 e_1}{\partial s \partial t} \cdot \psi_1 \left(\frac{|\partial^2 v_1 / \partial s \partial t|}{\partial^2 e_1 / \partial s \partial t} \right) ds dt, \dots, \int_a^b \int_c^d \frac{\partial^2 e_n}{\partial s \partial t} \cdot \psi_n \left(\frac{|\partial^2 v_n / \partial s \partial t|}{\partial^2 e_n / \partial s \partial t} \right) ds dt \right). \tag{2.36}
 \end{aligned}$$

This completes the proof. □

Remark 2.6. (i) Taking $n = 1$, (2.30) changes to a general form of the inequality which was given by Pachpatte [17].

(ii) Taking $G = 1$, (2.30) changes to a general form of the inequality which was given by Pečarić and Brnetić [19].

(iii) Taking $n = 1$, $G = 1$, (2.30) changes to the following inequality:

$$\int_a^b \int_c^d \left(\frac{\partial^2 F}{\partial \overline{M}^2} \cdot \frac{\partial \overline{M}}{\partial t} \cdot \frac{\partial \overline{M}}{\partial s} + \frac{\partial F}{\partial \overline{M}} \cdot \frac{\partial^2 \overline{M}}{\partial s \partial t} \right) ds dt \leq F \left(\int_a^b \int_c^d \frac{\partial^2 r}{\partial s \partial t} \cdot \phi \left(\frac{|\partial^2 u / \partial s \partial t|}{\partial^2 r / \partial s \partial t} \right) ds dt \right), \tag{2.37}$$

which is a general form of the following inequality established by Rožanova [21]:

$$\int_a^b F' \left(r(t) \phi \left(\frac{|f(t)|}{r(t)} \right) \right) r'(t) \phi \left(\frac{|f'(t)|}{r'(t)} \right) dt \leq F \left(\int_a^b r'(t) \phi \left(\frac{|f'(x)|}{r'(t)} \right) dt \right). \tag{2.38}$$

(iv) Let $u_i(s, t)$, $v_i(s, t)$, $r_i(s, t)$, and $e_i(s, t)$ reduce to $u_i(t)$, $v_i(t)$, $r_i(t)$, and $e_i(t)$, respectively, and with suitable modifications (where $i = 1, \dots, n$), the inequality in Theorem 2.5 changes to the inequality in Agarwal and Pang [2, Theorem 3, page 305].

THEOREM 2.7. *Let $u_i(s, t)$, $v_i(s, t)$, F , G , be as in Theorem 2.1. Let p_i , q_i , h_i , w_i , $i = 1, \dots, n$, be as in Theorem 2.3. Let $\bar{S}(s, t)$, \bar{M}_i , \bar{N}_i , $\partial^2 F / \partial \bar{M}_i^2$, $\partial F / \partial \bar{M}_i$, $\partial^2 G / \partial \bar{N}_i^2$, $\partial G / \partial \bar{N}_i$, $i = 1, \dots, n$, $\partial \bar{M}_i / \partial s$, $\partial \bar{M}_i / \partial t$, $\partial^2 \bar{M}_i / \partial s \partial t$, $\partial \bar{N}_i / \partial s$, $\partial \bar{N}_i / \partial t$, $\partial^2 \bar{N}_i / \partial s \partial t$, $i = 1, \dots, n$, be as in Theorem 2.5. Then the following integral inequality holds:*

$$\begin{aligned} & \int_a^b \int_c^d \left[F(\bar{M}_1(s, t), \dots, \bar{M}_n(s, t)) \cdot \sum_{i=1}^n \left(\frac{\partial^2 G}{\partial \bar{N}_i^2} \cdot \frac{\partial \bar{N}_i}{\partial t} \cdot \frac{\partial \bar{N}_i}{\partial s} + \frac{\partial G}{\partial \bar{N}_i} \cdot \frac{\partial^2 \bar{N}_i}{\partial s \partial t} \right) \right. \\ & \quad \left. + G(\bar{N}_1(s, t), \dots, \bar{N}_n(s, t)) \cdot \sum_{i=1}^n \left(\frac{\partial^2 F}{\partial \bar{M}_i^2} \cdot \frac{\partial \bar{M}_i}{\partial t} \cdot \frac{\partial \bar{M}_i}{\partial s} + \frac{\partial F}{\partial \bar{M}_i} \cdot \frac{\partial^2 \bar{M}_i}{\partial s \partial t} \right) + \bar{S}(s, t) \right] ds dt \\ & \leq F \left[h_1^{-1} \left(\int_a^b \int_c^d p_1(s, t) h_1 \left(\frac{\partial^2 r_1}{\partial s \partial t} \cdot \phi_1 \left(\frac{|\partial^2 u_1 / \partial s \partial t|}{\partial^2 r_1} \partial s \partial t \right) p_1(s, t) \right) ds dt \right), \dots, \right. \\ & \quad \left. h_n^{-1} \left(\int_a^b \int_c^d p_n(s, t) h_n \left(\frac{\partial^2 r_n}{\partial s \partial t} \cdot \phi_n \left(\frac{|\partial^2 u_n / \partial s \partial t|}{\partial^2 r_n / \partial s \partial t} \right) p_n(s, t) \right) ds dt \right) \right] \\ & \quad \cdot G \left[w_1^{-1} \left(\int_a^b \int_c^d q_1(s, t) w_1 \left(\frac{\partial^2 e_1}{\partial s \partial t} \cdot \psi_1 \left(\frac{|\partial^2 v_1 / \partial s \partial t|}{\partial^2 e_1 / \partial s \partial t} \right) q_1(s, t) \right) ds dt \right), \dots, \right. \\ & \quad \left. w_n^{-1} \left(\int_a^b \int_c^d q_n(s, t) w_n \left(\frac{\partial^2 e_n}{\partial s \partial t} \cdot \psi_n \left(\frac{|\partial^2 v_n / \partial s \partial t|}{\partial^2 e_n / \partial s \partial t} \right) q_n(s, t) \right) ds dt \right) \right]. \end{aligned} \tag{2.39}$$

Proof. From the hypotheses of Theorem 2.7, we have

$$\begin{aligned} & \int_a^b \int_c^d \frac{\partial^2 r_i}{\partial s \partial t} \cdot \phi_i \left(\frac{|\partial^2 u_i / \partial s \partial t|}{\partial^2 r_i / \partial s \partial t} \right) ds dt \\ & = \frac{\int_a^b \int_c^d p_i(s, t) \left(\left((\partial^2 r_i / \partial s \partial t) \cdot \phi_i \right) \left(|\partial^2 u_i / \partial s \partial t| / (\partial^2 r_i / \partial s \partial t) \right) \right) / p_i(s, t) ds dt}{\int_a^b \int_c^d p_i(s, t) ds dt}, \end{aligned} \tag{2.40}$$

$$\begin{aligned} & \int_a^b \int_c^d \frac{\partial^2 e_i}{\partial s \partial t} \cdot \psi_i \left(\frac{|\partial^2 v_i / \partial s \partial t|}{\partial^2 e_i / \partial s \partial t} \right) ds dt \\ & = \frac{\int_a^b \int_c^d q_i(s, t) \left(\left((\partial^2 e_i / \partial s \partial t) \cdot \psi_i \right) \left(|\partial^2 v_i / \partial s \partial t| / (\partial^2 e_i / \partial s \partial t) \right) \right) / q_i(s, t) ds dt}{\int_a^b \int_c^d q_i(s, t) ds dt}, \end{aligned}$$

for $i = 1, \dots, n$.

From (2.40) and using the hypotheses on $h_i, w_i, i = 1, \dots, n$, and Jensen's inequality, we obtain

$$\begin{aligned}
 & h_i \left(\int_a^b \int_c^d \left(\frac{\partial^2 r_i}{\partial s \partial t} \cdot \phi_i \left(\frac{|\partial^2 u_i / \partial s \partial t|}{\partial^2 r_i / \partial s \partial t} \right) \right) ds dt \right) \\
 & \leq \int_a^b \int_c^d p_i(s, t) h_i \left(\frac{(\partial^2 r_i / \partial s \partial t) \cdot \phi_i (|\partial^2 u_i / \partial s \partial t| / \partial^2 r_i / \partial s \partial t)}{p_i(s, t)} \right) ds dt,
 \end{aligned}
 \tag{2.41}$$

$$\begin{aligned}
 & w_i \left(\int_a^b \int_c^d \left(\frac{\partial^2 e_i}{\partial s \partial t} \cdot \psi_i \left(\frac{|\partial^2 v_i / \partial s \partial t|}{\partial^2 e_i / \partial s \partial t} \right) \right) ds dt \right) \\
 & \leq \int_a^b \int_c^d q_i(s, t) w_i \left(\frac{(\partial^2 e_i / \partial s \partial t) \cdot \psi_i (|\partial^2 v_i / \partial s \partial t| / \partial^2 e_i / \partial s \partial t)}{q_i(s, t)} \right) ds dt,
 \end{aligned}
 \tag{2.42}$$

for $i = 1, \dots, n$.

Then

$$\begin{aligned}
 & \int_a^b \int_c^d \frac{\partial^2 r_i}{\partial s \partial t} \cdot \phi_i \left(\frac{|\partial^2 u_i / \partial s \partial t|}{\partial^2 r_i / \partial s \partial t} \right) ds dt \\
 & \leq h_i^{-1} \left(\int_a^b \int_c^d p_i(s, t) \cdot h_i \left(\frac{(\partial^2 r_i / \partial s \partial t) \cdot \phi_i (|\partial^2 u_i / \partial s \partial t| / \partial^2 r_i / \partial s \partial t)}{p_i(s, t)} \right) ds dt \right),
 \end{aligned}
 \tag{2.43}$$

$$\begin{aligned}
 & \int_a^b \int_c^d \frac{\partial^2 e_i}{\partial s \partial t} \cdot \psi_i \left(\frac{|\partial^2 v_i / \partial s \partial t|}{\partial^2 e_i / \partial s \partial t} \right) ds dt \\
 & \leq w_i^{-1} \left(\int_a^b \int_c^d q_i(s, t) \cdot w_i \left(\frac{(\partial^2 e_i / \partial s \partial t) \cdot \psi_i (|\partial^2 v_i / \partial s \partial t| / \partial^2 e_i / \partial s \partial t)}{q_i(s, t)} \right) ds dt \right).
 \end{aligned}
 \tag{2.44}$$

By applying (2.43) and (2.44) to the right-hand side of inequality (2.30), we get the desired inequality (2.39) and the proof is complete. \square

Remark 2.8. (i) Taking $n = 1$, (2.39) changes to a general form of the inequality which was given by Pachpatte [17].

(ii) Taking $G = 1$, (2.39) changes to a general form of the inequality which was given by Pečarić and Brnetić [19].

(iii) Let $u_i(s, t), v_i(s, t), h_i(s, t), w_i(s, t), r_i(s, t)$, and $e_i(s, t)$ reduce to $u_i(t), v_i(t), h_i(t), w_i(t), r_i(t)$, and $e_i(t)$, respectively, and with suitable modifications (where $i = 1, \dots, n$), then inequality (2.39) changes to the inequality in Agarwal and Pang [2, Theorem 4, page 308].

Acknowledgments

This research is partially supported by the Research Grants Council of the Hong Kong SAR, China (project no. HKU7017/05P), and supported by Zhejiang Provincial Natural Science Foundation of China (Y605065), National Natural Sciences Foundation of China

(10271071), Foundation of the Education Department of Zhejiang Province of China (20050392), and the Academic Mainstay of Middle-age and Youth Foundation of Shandong Province of China (200203).

References

- [1] Z. Opial, "Sur une inégalité," *Annales Polonici Mathematici*, vol. 8, pp. 29–32, 1960.
- [2] R. P. Agarwal and P. Y. H. Pang, *Opial Inequalities with Applications in Differential and Difference Equations*, vol. 320 of *Mathematics and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1995.
- [3] R. P. Agarwal and V. Lakshmikantham, *Uniqueness and Nonuniqueness Criteria for Ordinary Differential Equations*, vol. 6 of *Series in Real Analysis*, World Scientific, River Edge, NJ, USA, 1993.
- [4] D. Bařnov and P. Simeonov, *Integral Inequalities and Applications*, vol. 57 of *Mathematics and Its Applications (East European Series)*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1992.
- [5] J. D. Li, "Opial-type integral inequalities involving several higher order derivatives," *Journal of Mathematical Analysis and Applications*, vol. 167, no. 1, pp. 98–110, 1992.
- [6] D. S. Mitrinović, J. E. Pečarić, and A. M. Fink, *Inequalities Involving Functions and Their Integrals and Derivatives*, vol. 53 of *Mathematics and Its Applications (East European Series)*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1991.
- [7] W.-S. Cheung, "On Opial-type inequalities in two variables," *Aequationes Mathematicae*, vol. 38, no. 2-3, pp. 236–244, 1989.
- [8] W.-S. Cheung, "Some new Opial-type inequalities," *Mathematika*, vol. 37, no. 1, pp. 136–142, 1990.
- [9] W.-S. Cheung, "Some generalized Opial-type inequalities," *Journal of Mathematical Analysis and Applications*, vol. 162, no. 2, pp. 317–321, 1991.
- [10] W.-S. Cheung, "Opial-type inequalities with m functions in n variables," *Mathematika*, vol. 39, no. 2, pp. 319–326, 1992.
- [11] W.-S. Cheung, Z. Dandan, and J. E. Pečarić, "Opial-type inequalities for differential operators," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 66, no. 9, pp. 2028–2039, 2007.
- [12] E. K. Godunova and V. I. Levin, "An inequality of Maroni," *Matematicheskie Zametki*, vol. 2, pp. 221–224, 1967.
- [13] D. S. Mitrinović, *Analytic Inequalities*, vol. 1965 of *Die Grundlehren der mathematischen Wissenschaften*, Springer, New York, NY, USA, 1970.
- [14] B. G. Pachpatte, "On integral inequalities similar to Opial's inequality," *Demonstratio Mathematica*, vol. 22, no. 1, pp. 21–27, 1989.
- [15] B. G. Pachpatte, "On inequalities of the Opial type," *Demonstratio Mathematica*, vol. 25, pp. 35–45, 1992.
- [16] B. G. Pachpatte, "Some inequalities similar to Opial's inequality," *Demonstratio Mathematica*, vol. 26, no. 3-4, pp. 643–647, 1993.
- [17] B. G. Pachpatte, "A note on generalized Opial-type inequalities," *Tamkang Journal of Mathematics*, vol. 24, no. 2, pp. 229–235, 1993.
- [18] J. E. Pečarić, "An integral inequality," in *Analysis, Geometry and Groups: A Riemann Legacy Volume—Part II*, H. M. Srivastava and Th. M. Rassias, Eds., Hadronic Press Collect. Orig. Artic., pp. 471–478, Hadronic Press, Palm Harbor, Fla, USA, 1993.
- [19] J. E. Pečarić and I. Brnetić, "Note on generalization of Godunova-Levin-Opial inequality," *Demonstratio Mathematica*, vol. 30, no. 3, pp. 545–549, 1997.
- [20] J. E. Pečarić and I. Brnetić, "Note on the generalization of the Godunova-Levin-Opial inequality in several independent variables," *Journal of Mathematical Analysis and Applications*, vol. 215, no. 1, pp. 274–282, 1997.

- [21] G. I. Rozanova, "Integral inequalities with derivatives and with arbitrary convex functions," *Moskovskii Gosudarstvennyi Pedagogicheskii Institut imeni V. I. Lenina. Uchenye Zapiski*, vol. 460, pp. 58–65, 1972.
- [22] G. S. Yang, "Inequality of Opial-type in two variables," *Tamkang Journal of Mathematics*, vol. 13, no. 2, pp. 255–259, 1982.

Wing-Sum Cheung: Department of Mathematics, The University of Hong Kong, Pokfulam Road, Hong Kong
Email address: wscheung@hku.hk

Chang-Jian Zhao: Department of Information and Mathematics Sciences, College of Science, China Jiliang University, Hangzhou 310018, China
Email addresses: chjzhao@163.com; chjzhao315@yahoo.com.cn; chjzhao@cjlu.edu.cn