



<b>Title</b>	<b>Cytokines and junction restructuring during spermatogenesis - A lesson to learn from the testis</b>
<b>Author(s)</b>	<b>Xia, W; Mruk, DD; Lee, WM; Cheng, CY</b>
<b>Citation</b>	<b>Cytokine And Growth Factor Reviews, 2005, v. 16 n. 4-5, p. 469-493</b>
<b>Issued Date</b>	<b>2005</b>
<b>URL</b>	<b><a href="http://hdl.handle.net/10722/84892">http://hdl.handle.net/10722/84892</a></b>
<b>Rights</b>	<b>Cytokine &amp; Growth Factor Reviews. Copyright © Elsevier Ltd.</b>

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Cytokine &amp; Growth Factor Reviews xxx (2005) xxx–xxx

[www.elsevier.com/locate/cytogfr](http://www.elsevier.com/locate/cytogfr)

# Cytokines and junction restructuring during spermatogenesis—a lesson to learn from the testis

Weiliang Xia, Dolores D. Mruk, Will M. Lee, C. Yan Cheng\*

Population Council, Center for Biomedical Research, 1230 York Avenue, New York, NY 10021, USA

## Abstract

In the mammalian testis, preleptotene and leptotene spermatocytes residing in the basal compartment of the seminiferous epithelium must traverse the blood-testis barrier (BTB) during spermatogenesis, entering the adluminal compartment for further development. However, until recently the regulatory mechanisms that regulate BTB dynamics remained largely unknown. We provide a critical review regarding the significance of cytokines in regulating the ‘opening’ and ‘closing’ of the BTB. We also discuss how cytokines may be working in concert with adaptors that selectively govern the downstream signaling pathways. This process, in turn, regulates the dynamics of either Sertoli–Sertoli tight junction (TJ), Sertoli–germ cell adherens junction (AJ), or both junction types in the epithelium, thereby permitting TJ opening without compromising AJs, and vice versa. We also discuss how adaptors alter their protein–protein association with the integral membrane proteins at the cell–cell interface via changes in their phosphorylation status, thereby altering adhesion function at AJ. These findings illustrate that the testis is a novel *in vivo* model to study the biology of junction restructuring. Furthermore, a molecular model is presented regarding how cytokines selectively regulate TJ/AJ restructuring in the epithelium during spermatogenesis.

© 2005 Published by Elsevier Ltd.

**Keywords:** Spermatogenesis; Testis; Junction restructuring; Cytokines; TGF- $\beta$ 3; TNF $\alpha$ ; p38 MAPK; ERK; JNK; Blood-testis barrier; Adherens junction; Tight junction; Adaptors; Ectoplasmic specialization

## 1. Introduction

The production of mature spermatozoa (haploid,  $1n$ ) from spermatogonia (diploid,  $2n$ ) is essential for the perpetuation of all mammalian species. Such event, known as spermatogenesis in the male, takes place in the functional unit of the testis called the seminiferous tubule. Seminiferous tubules, in turn, coordinate with Leydig cells in the interstitium and the brain via the hypothalamic–pituitary–testicular axis to regulate spermatogenesis [1,2]. Although spermatogenesis varies in detail in different species (e.g., minks are seasonal breeders exhibiting seasonally or environmentally responsive phases in this process whereas spermatogenesis continues throughout the entire life span in humans and rodents), the cellular constituents and the basic physiology of the testes are rather similar [3]. We limit our discussion largely in rats, mice and/or men since most studies were conducted in these species.

Spermatogenesis can be divided into three distinct phases which provide an upward of  $150 \times 10^6$  spermatozoa per day per man [1,3]. The germline stem cells spermatogonia can either self-proliferate (phase 1) or differentiate into primary spermatocytes, which then undergo meiosis and differentiate into secondary spermatocytes and eventually haploid spermatids (phase 2). These cells, in turn, differentiate morphologically and functionally to spermatozoa via spermiogenesis (phase 3), which are released into the tubule lumen at spermiation [1,3]. This entire process of germ cell development in the seminiferous epithelium is dependent on temporal and spatial expression of unique sets of genes and proteins. In the rat testis, an epithelial cycle ( $\sim 12$ – $14$  days duration) can be divided into 14 stages which are classified according to the unique germ cell types that associate with Sertoli cells in the epithelium [3,4]. It takes  $\sim 58$  days for a single spermatogonium to fully differentiate and develop into 256 spermatozoa. As such, it takes  $\sim 4.5$  epithelial cycles for one spermatogonium to differentiate into 256 spermatids. For each stage, at least four germ cell types are present in the epithelium that are organized

\* Corresponding author. Tel.: +1 212 327 8738; fax: +1 212 327 8733.  
E-mail address: Y.Cheng@popcbr.rockefeller.edu (C.Y. Cheng).




Fig. 1. Spermatogenesis and cell junctions in the seminiferous epithelium of the mammalian testis (e.g. rats). (A) This is the cross-section of a seminiferous tubule from an adult rat testis showing the intimate relationship between Sertoli cells (N, Sertoli cell nucleus) and germ cells (e.g., pachytene spermatocyte PS, round spermatid RS). (B) Schematic drawing of developing germ cells and their intimate relationship with Sertoli cells during spermatogenesis in the seminiferous epithelium. Also shown is the relative location of different junction types in the epithelium between Sertoli cells as well as between Sertoli and germ cells. Sertoli and germ cells constitute the seminiferous epithelium that is adjacent to the tunica propria. Differentiating germ cells must migrate from the basal to the adluminal compartment, traversing the BTB, which has physically divided the epithelium into the basal and adluminal compartment. (C–E) Electron micrographs of cross sections of seminiferous epithelium illustrating the ultrastructural features of the blood-testis barrier at low (C) and high (D) magnification. The basal ES is characterized by the presence of actin filament bundles (white arrows) sandwiched between the cisternae of endoplasmic reticulum (er), and the

spatially into layers from the base to the lumen of the seminiferous tubule [3,4]. Furthermore, spermatogenesis cannot complete without the support of Sertoli cells, which are the only other cell type in the seminiferous epithelium behind the BTB besides germ cells (note: the BTB has physically divided the epithelium into the basal and adluminal compartment, see Fig. 1) [5–7]. Except for the spermatogonia, developing germ cells move progressively toward the lumen [8]. For instance, preleptotene and leptotene spermatocytes that lie at the periphery of the tubule and outside the BTB must traverse the BTB at late stages VIII and early IX of the epithelial cycle [8].

It is conceivable that enormous Sertoli–germ cell interactions take place in the seminiferous epithelium throughout spermatogenesis [1–4,6,9,10]. If one views spermatogenesis as a voyage of a germ cell that moves from the basal to the adluminal compartment while developing to a mature spermatozoon, this process involves numerous decision makings and executions. It also requires signalings in and out of germ cells to facilitate this event. Although it is not entirely clear regarding the sequence of these signals, there are at least two sources: external signals from outside the tubule (e.g., via Leydig cells, peritubular myoid cells, and both paracrine and hormonal factors including those from the pituitary gland), and internal crosstalks between germ and Sertoli cells (e.g., integrin-mediated signalings) [5,6,11,12]. The phenotypic consequence of these signalings is manifested, at least in part, via the constant remodeling at the Sertoli–Sertoli and Sertoli–germ cell interface where different cell junction types are present [1,2].

The identities of these signals and the details of the remodeling events have become increasingly clear in recent years [1,2]. For instance, there is accumulating evidence that illustrates the crucial roles of cytokines pertinent to spermatogenesis and junction restructuring [2]. In this review, we first give an update on the junction complexes that are found in the testis, highlighting how cytokines (e.g., TGF- $\beta$ 3, TNF $\alpha$ ) can affect junction dynamics and how these signals are being fine-tuned to allow their regulation of a particular junction type.

## 2. The seminiferous epithelium: Sertoli–germ cell junctions and spermatogenesis

### 2.1. Seminiferous epithelium

The seminiferous epithelium is composed of Sertoli and germ cells. The Sertoli cell is by and large a tall columnar

cell extending from the base to the apex of the seminiferous tubule [3]. It is physically reshaped by germ cells to possess many cytoplasmic processes because each Sertoli cell is ‘nursing’ about 30–50 germ cells at different stages of their development at any given time during the epithelial cycle [13,14]. In the rat, Sertoli cells cease to proliferate at about day 20 postnatal and the number of these nursing cells determines how many germ cells can be supported and produced via spermatogenesis in the testis [3], illustrating the crucial function of Sertoli cells. For instance, Sertoli cells provide structural support for germ cells and their translocation, create the BTB and define the polarity of the epithelium, secrete numerous biological factors and nutrients for germ cells, and conduct other vital functions pertinent to spermatogenesis (e.g., phagocytosis) [2,3,14].

### 2.2. Sertoli–Sertoli and Sertoli–germ cell junctions

The different junction types that are found in the seminiferous epithelium have recently been reviewed [1,2]. Similar to other epithelia or endothelia, virtually all major junction types are found in the testis. Besides the tight junctions that are restricted to the BTB, several anchoring junction types (four are found in most epithelia) are also detected in the testis: (a) adherens junction (including basal and apical ectoplasmic specialization [ES], basal and apical tubulobulbar complex [TBC]); (b) desmosome-like junctions; and (c) hemidesmosomes (for reviews, see [1,2,10,15]). ES is a testis-specific, actin-based adherens junction localized at two sites in the seminiferous epithelium: basal and apical compartment (see Fig. 1) [2,9,10,16,17]. Basal ES is limited to BTB and present side-by-side with TJ (Fig. 1). Apical ES is found between elongating/elongate spermatids and Sertoli cells. At least three protein complexes, namely, the cadherin/catenin, the nectin/afadin, and the integrin/laminin, are known to be ES components [2,17]. TBC is another modified AJ type found in the testis [2,10,18]. Apical TBC only appears a few days before spermiation in the epithelium at late stage VIII of the epithelial cycle when apical ES begins to disappear whereas basal TBC co-exists with TJ, basal ES, and desmosomal-like junctions at the BTB site. Desmosome-like junctions are present between Sertoli cells and spermatogonia, spermatocytes and round spermatids, being most prominent surrounding pachytene spermatocytes [10]. The BTB is not fully formed until 16–19 days postnatal in the rat testis [3]. Unlike barriers in other organs (e.g., the blood–brain barrier, the blood–retinal barrier) where TJs are localized to the apical region of the epithelium/endothelium, to be

Sertoli cell membrane (apposing arrowheads represent the apposing Sertoli cell membranes), which can be found on both sides of the apposing Sertoli cells. Tight junction (TJ) is found between the basal ES, the coexisting TJ and basal ES in turn constitute the BTB. Apical ES is shown in (E) which is typified by the presence of actin filament bundles (white arrowheads) sandwiches between the cisternae of er and Sertoli cell membrane (apposing white arrowheads represent the apposing Sertoli and germ cell membranes). However, this typical feature of ES, in contrast to the basal ES, is restricted only to the Sertoli cell side in apical ES. (F–G) Schematic drawings that illustrate the molecular architecture of the constituent proteins at the BTB (F) and apical ES (G), which include cytokines (e.g., TGF- $\beta$ 3 and TNF $\alpha$ ) released from Sertoli and/or germ cells can mediate Sertoli–germ cell crosstalk during spermatogenesis. The protein complexes known to exist at the apical ES site include cadherin/catenin, nectin/afadin, and  $\alpha$ 6 $\beta$ 1 integrin/laminin  $\gamma$ 3; whereas occludin/ZO-1, JAM/ZO-1, claudin/ZO-1, cadherin/catenin and nectin/afadin are found at the BTB site. Bar in E = 10  $\mu$ m, C = 3  $\mu$ m, D = 0.25  $\mu$ m and E = 0.3  $\mu$ m, respectively.

152 followed by AJ, and TJs are furthest away from the ECM; TJs  
 153 at the BTB lie closest to the basement membrane (a modified  
 154 form of ECM). Furthermore, BTB is a dynamic structure  
 155 which must ‘open’ and ‘close’ to permit preleptotene/  
 156 leptotene spermatocyte transmigration. BTB is a rather  
 157 complex barrier when compared to other barriers (e.g.,  
 158 gastric–mucosal barrier which is formed by epithelial cells,  
 159 blood–retinal barrier and blood–brain barrier which are  
 160 formed by endothelial cells) [19–21] (see Fig. 1). Recent  
 161 studies have also shown that apical ES is constituted and  
 162 regulated by proteins that are usually restricted to the focal  
 163 contact in cell–matrix interface in other epithelia [22]. This  
 164 hybrid cell–matrix–cell junction type may indeed be essential  
 165 for rapid junction remodeling to facilitate spermatids  
 166 orientation and movement at spermiation.

### 167 2.3. Constituent proteins of different junction types in 168 the testis

#### 169 2.3.1. Tight junction (TJ)

170 TJ is the only known example of occluding junction that  
 171 confers the barrier function of an epithelium or endothelium  
 172 by restricting the passages of molecules through the  
 173 intercellular spaces and creates a boundary that defines cell  
 174 polarity [23]. In the testis, TJ also creates an immunological  
 175 barrier that sequesters the post-meiotic germ cell antigens  
 176 from the immune system of the host animals. The currently  
 177 known TJ integral membrane proteins include JAMs  
 178 (junctional adhesion molecules), claudins and occludins,  
 179 which have recently been reviewed [1,2,23,24], as such, only  
 180 a brief update is provided in this section.

181 2.3.1.1. JAMs. JAMs are members of a distinct class of cell  
 182 adhesion molecules typified by the presence of two Ig-like  
 183 loops in the extracellular domain that are expressed in  
 184 leukocytes and are localized to tight junctions as integral  
 185 membrane proteins in epithelial and endothelial cells  
 186 [25,26]. Since the discovery of JAM-A in 1998 [27], other  
 187 members, including the more related JAM-B and JAM-C,  
 188 and the less related JAM4, coxsackie and adenovirus  
 189 receptor (CAR), and endothelial cell-selective adhesion  
 190 molecule (ESAM), have recently been added to the list  
 191 [25,26,28]. The presence of JAM-A, B and C in the testis  
 192 have now been confirmed [29,30]. JAM-A is present at the  
 193 BTB in the rat testis, co-localizing with ZO-1 [30].  
 194 Moreover, JAM-A expression is stage-specific, being  
 195 highest at IX–XIV, lowest at IV–VI [30]. This stage  
 196 specificity apparently is related to its possible involvement  
 197 in BTB dynamics, facilitating the passage of preleptotene/  
 198 leptotene spermatocytes across the BTB. Although *Jam-A*<sup>-/-</sup>  
 199 mice has been generated, it is not known if the BTB is  
 200 affected since a morphological examination of the testis has  
 201 yet to be reported [31]. A recent study on the *Jam-C*<sup>-/-</sup> mice  
 202 have shown that JAM-C is crucial to spermiogenesis since in  
 203 the viable mutants, mature spermatids are missing [29]. In  
 204 normal mice, JAM-C is localized to the developing round

205 and elongating spermatids [29]. Interestingly, JAM-B has  
 206 been localized both to the site of TJs at the basal  
 207 compartment and to the apical ES at the spermatid-Sertoli  
 208 cell interface in the seminiferous epithelium, outside the  
 209 BTB [29]. Besides their homophilic interactions amongst  
 210 JAM-A, B and C, JAM-C can interact with JAM-B  
 211 heterotypically [32]. Both JAM-B and JAM-C are localized  
 212 to the heads of spermatids at the apical ES, and this  
 213 heterophilic association may be important for the Sertoli  
 214 cell-spermatid adhesion function [29]. Based on currently  
 215 available data, two roles are suggested for JAMs: in the  
 216 immune system they are crucial to leukocyte transmigration;  
 217 and in polarized epithelial and endothelial cells, they  
 218 seem to take part in organizing TJ and cell polarity [26].  
 219 This latter physiological role has been extended to the testis  
 220 since the cell polarity complex [partitioning-defective (Par)  
 221 3/atypical protein kinase C (aPKC)/Cdc42] apparently is  
 222 recruited by JAM-C to facilitate round spermatid polariza-  
 223 tion and thus differentiation [26]. How JAMs assist  
 224 preleptotene/leptotene spermatocytes to traverse the BTB  
 225 similar to neutrophil transmigration across the endothelial  
 226 TJ-barrier remains to be investigated since germ cells per se,  
 227 unlike neutrophils or macrophages, are not actively migrating  
 228 cells. It is possible that JAMs are associated with other  
 229 motor proteins (e.g., myosin VIIa) and cytoskeletons (e.g.,  
 230 actin, tubulin) that facilitate germ cell movement using  
 231 the locomotive apparatus in Sertoli cells that provides the  
 232 necessary protrusive force to guide germ cell movement  
 233 (for review, see [2]).

234 JAMs are expressed in multiple epithelia, endothelia,  
 235 leukocytes and platelets [25,26]. The regulation of JAMs in  
 236 the testis is largely unknown. In the rat testis, when the  
 237 intratesticular T was suppressed by placing testosterone and  
 238 estrogen implants subdermally, spermatids (step 8 and  
 239 beyond) were depleted because of a disruption of the cell  
 240 adhesion at the ES [30,33–36]. However, the tight junctions  
 241 at the BTB remained intact which were associated with a  
 242 significant surge in the levels of JAM-A, occludin and ZO-1  
 243 in the epithelium [30]. Indeed, the JAM-A distribution at the  
 244 BTB site in the basal compartment of the seminiferous  
 245 epithelium was significantly induced and intensified,  
 246 becoming a thickened and prominent ring surrounding the  
 247 entire tubule [30]. It is apparent that a depletion of androgen  
 248 in the testis triggers a novel mechanism that leads to two  
 249 distinctive events: germ cell loss and a reinforced BTB [30].  
 250 Another model using Adjudin to induce germ cell sloughing  
 251 from the epithelium in adult rat testes has yield similar  
 252 results in which JAM-A expression was induced at the time  
 253 of germ cell depletion (unpublished observations). Although  
 254 the compounds that were used to trigger the changes in the  
 255 epithelium are different in these two models, namely  
 256 androgen suppression and Adjudin, the signaling events  
 257 (e.g., both treatments activate the integrin/focal adhesion  
 258 kinase signaling pathway) and the phenotypic outcome (e.g.,  
 259 germ cell loss from the epithelium and a reinforced BTB) are  
 260 similar [36,37]. This seemingly suggests that JAM-A is

261 regulated, at least in part, by a mechanism downstream of  
 262 lowered intratesticular T level that triggers germ cell  
 263 sloughing from the epithelium. It is not known if cytokines  
 264 are the upstream regulators of JAMs. An earlier report has  
 265 shown that TNF and IFN- $\gamma$  treatment of human umbilical  
 266 vein endothelial cells can reduce cell surface expression of  
 267 JAM-A, but these cytokines have no effects on the rate of  
 268 transmigration of neutrophils [38].

269 2.3.1.2. *Claudins*. The claudin superfamily of TJ integral  
 270 membrane proteins consists of at least 24 members with Mr  
 271 ranging between 20 and 27 kDa [39,40]. Claudins have a  
 272 unique expression profile in a tissue [39]. For instance,  
 273 claudin-1 and claudin-11 are expressed in the testis, mostly  
 274 restricted to Sertoli cells, and the brain, whereas more than 10  
 275 claudin members are expressed in the kidney [39]. In the  
 276 testis, the expression of claudin-3, -4, -5, -7, -8 has also been  
 277 reported [2,41]. Claudin-11, also known as oligodendrocyte-  
 278 specific protein (OSP), is the best studied claudin in the testis  
 279 [42–46]. *Cld11*<sup>-/-</sup> mice were sterile and were associated  
 280 with the absence of TJ strands in the seminiferous epithelium  
 281 and in the myelin sheath in the brain [46]. Claudin-11 is  
 282 known to be up-regulated by androgens [42,47] and down-  
 283 regulated by TGF- $\beta$ 3 [44] in Sertoli cells cultured in vitro.  
 284 Claudin-11 expression is high from postnatal days 10–16 in  
 285 the rat testis corresponding to the maturation of BTB [43].  
 286 Anti-androgen, such as flutamide, can also inhibit the  
 287 expression of claudin-11 in prepubertal rat testes [42].  
 288 Claudin-11 is also important for hearing function since  
 289 *Cld11*<sup>-/-</sup> mice lacking TJ in the basal cells of stria vascularis  
 290 in cochlea failed to compartmentalize the endolymph and  
 291 suppressed electrical potentials [48].

292 TJ strands in the intercellular junction are not a static but  
 293 dynamic structure. A recent study by real-time imaging to  
 294 examine the behavior of exogenously expressed claudin-1  
 295 in mouse L fibroblasts showed that the paired claudin  
 296 strands underwent constant and dynamic reorganization  
 297 while maintaining the structural integrity of the entire  
 298 TJ network [49]. Internalization of claudin-3 was also  
 299 observed via endocytosis in confluent epithelial cells, after  
 300 it was dissociated from other TJ components, such as JAM,  
 301 occludin and ZO-1 [50]. This dynamic nature of claudins,  
 302 plausibly applicable to other TJ constituent proteins, is  
 303 not entirely unexpected since TJ barriers must undergo  
 304 conformational changes to accommodate paracellular  
 305 transport of substances, such as during food adsorption  
 306 in the small intestine. For the BTB, it has to be ‘opened’  
 307 (or ‘dissolved’ ?) and then ‘closed’ (or ‘regenerated’ ?)  
 308 frequently to facilitate germ cell passage while maintaining  
 309 the barrier function during the epithelial cycle. It is likely  
 310 that such reorganization of claudin strands, possibly also of  
 311 occludin- and JAM-constituted TJ strands, are occurring at  
 312 the BTB. The uncoupling of TJ proteins may indeed be a  
 313 prerequisite for the dual roles played by the BTB during its  
 314 restructuring to permit germ cell passage while maintaining  
 315 the barrier function simultaneously.

2.3.1.3. *Occludin*. Occludin is the first TJ integral mem-  
 316 brane protein found in epithelia [51] and is the most studied  
 317 in this category. Occludin is known to be regulated by  
 318 cytokines in the testis (e.g., TGF- $\beta$ 2, TGF- $\beta$ 3) [44,52].  
 319 Other signaling events are recently shown to engage in its  
 320 regulation as well. In the androgen-suppressed rat testes to  
 321 induce germ cell loss from the epithelium, occludin  
 322 expression, similar to JAM-A, is significantly induced,  
 323 resulting in prominent staining at the BTB when Sertoli-  
 324 germ cell adhesion function was compromised [30]. This  
 325 also reinforces the notion that the regulation of TJ proteins is  
 326 essentially different from that of AJ proteins in the rat testis.  
 327 Occludin is also regulated, at least in part, by ubiquitination  
 328 [53]. Itch (an E3 ubiquitin ligase) and UBC4 (an ubiquitin-  
 329 conjugating enzyme) are reciprocally regulated versus  
 330 occludin during Sertoli cell TJ assembly or disassembly,  
 331 and ubiquitin-conjugated and Itch-conjugated occludin are  
 332 detected when the dibutyl-*c*AMP-induced degradation of  
 333 occludin is blocked by a proteasome inhibitor MG-132 [53].  
 334 Other cytokines, such as TNF $\alpha$ , IL-1 $\beta$ , IFN- $\gamma$ , can also  
 335 affect occludin expression and its distribution at TJs in  
 336 multiple epithelia and endothelia [54–56]. It is not known if  
 337 these cytokines can exert any effect on occludin expression  
 338 or its cellular distribution at the BTB. But TNF $\alpha$  has been  
 339 shown to perturb TJ-barrier function in Sertoli cell cultures  
 340 [57] and can cause germ cell exfoliation in the rat testis after  
 341 its systemic administration [58]. Recent studies have shown  
 342 that TNF and IFN- $\gamma$  can indeed regulate occludin  
 343 transcription by diminishing its promoter activity [55].  
 344

### 2.3.2. Anchoring junction 345

2.3.2.1. *The cadherin/catenin protein complex*. Cadherins  
 346 are transmembrane glycoproteins that mediate calcium-  
 347 dependent cell–cell adhesion in multiple epithelia including  
 348 the seminiferous epithelium in the testis [59,60]. The  
 349 cadherin superfamily consists of over 80 members that fall  
 350 into at least six subfamilies, which include (i) the type I  
 351 classical cadherins (e.g., E-cadherin, N-cadherin, P-cad-  
 352 herin) and its highly related; (ii) type II classical cadherins  
 353 (e.g., VE-cadherin); (iii) desmosomal cadherins (desmo-  
 354 collins and desmogleins); (iv) protocadherins; (v) seven-  
 355 pass transmembrane cadherins (Flamingo); and (vi) Fat-like  
 356 cadherins [59–62]. Type I classical cadherins are the best  
 357 studied cadherins in multiple tissues including the testis.  
 358 Besides the classical cadherins, the presence of other  
 359 subfamilies in the testis, such as protocadherins, Fat and  
 360 Flamingo, has also been detected by RT-PCR [63], but their  
 361 function in the seminiferous epithelium are less known.  
 362 During development, the expression of different cadherins is  
 363 highly dynamic [64] and this seems to be applicable to the  
 364 testis as well since the expression profile of cadherins varies  
 365 with the age and cell types in the rat testis where at least 24  
 366 cadherins are known to be present [63]. For instance, N-  
 367 cadherin is predominantly localized to the basal ES and the  
 368 periphery of the seminiferous tubules with restricted and  
 369 stage-specific localization at the apical ES [30,65–69],  
 370

whereas E-cadherin is relatively more abundant in germ cells [63,66]. A smaller amount of N-cadherin in the testis appears also to be a component of desmosomal-like junctions which is a hybrid junction type of desmosome and gap junctions [65,67]. Indeed, N-cadherin has been shown to link to both actin microfilament and microtubules in the testis [66]. A recent report has also illustrated that protocadherin  $\alpha 3$  is associated with spermatids at the acrosomal area, intercellular bridge as well as flagellum, distinct from the distribution of classical cadherins [70].

Classical cadherin-based protein complex comprising of the transmembrane protein cadherins and intracellular adaptor catenins is a well defined focal point of cell adhesion and signaling [59,71].  $\beta$ -Catenin and  $\gamma$ -catenin connects cadherins to  $\alpha$ -catenin and  $\alpha$ -actinin, which are two putative actin binding proteins [72]. Phosphorylation of  $\beta$ -catenin can in turn regulate the integrity of the cadherin/catenin complex [73]. In both Adjudin- and androgen suppression-induced germ cell loss models, the event of germ cell loss is facilitated by the dissociation of N-cadherin from  $\beta$ -catenin [30,35,68]. Indeed, increased tyrosine phosphorylation of  $\beta$ -catenin was detected at the time of germ cell depletion in these models [30]. Kinases and phosphatases are also known to regulate cadherin/catenin association [35,74,75]. For instance, myotubularin-related protein 2 (MTMR2), a lipid phosphoinositide phosphatase, was shown to interact with the kinase c-Src [35] and c-Src in turn associates with the N-cadherin/ $\beta$ -catenin complex [74]. This illustrates a novel regulatory mechanism may be in place in the testis regarding the cadherin/catenin-mediated cell adhesion function in which MTMR2 and c-Src regulate the phosphorylation status of the cadherin/catenin, which in turn determines its cell adhesive function. More recent studies have shown that the N-cadherin/ $\beta$ -catenin adhesion unit can also be regulated by the equilibrium between IQGAP-1 (IQ motif containing GTPase activating protein, an effector of Cdc42 GTPase) and Cdc42 in Sertoli-germ cell AJ [76]. For instance, using a  $\text{Ca}^{2+}$  switch model, it has been demonstrated that at low  $\text{Ca}^{2+}$  level, IQGAP-1 is released from Cdc42, and interacts with  $\beta$ -catenin instead, causing the dissociation of  $\beta$ -catenin from N-cadherin, and germ cell depletion from Sertoli cells [76].

E-Cadherin is also a tumor suppressor which is down-regulated while N-cadherin is up-regulated during epithelial tumor progression [64,77,78]. This 'cadherin switch' further illustrates the unique yet pivotal role of each cadherin in cell adhesion and cell motility. It is not clear if such dynamic switch-over between different cadherins occur during germ cell movement in the seminiferous epithelium. However, N-cadherin can become highly expressed in the testis of Adjudin treated rats during germ cell loss from the epithelium [66–68]. N-cadherin is also up-regulated in androgen suppressed rat testes during germ cell loss [30,35]. Yet such a surge in N-cadherin cannot rescue germ cell loss from the epithelium since a loss of association between N-

cadherin and  $\beta$ -catenin was detected at the time of germ cell sloughing in both models [30,35]. It seems that such an induction of cadherins reinforces the BTB integrity since N-cadherin is also a component protein of the BTB in the rat testis.

**2.3.2.2. The nectin/afadin/ponsin/ADIP complex.** The nectin/afadin/ponsin complex is another actin-based cell adhesion protein complex that plays a crucial role in the testis during spermatogenesis. It confers Sertoli-germ cell adhesion function particularly for elongating/elongate spermatids [1,2,68,79]. Four nectins (nectin-1, -2, -3, and -4) have been identified thus far, all of which are expressed in the testis with nectin-2 and nectin-3 being the highly expressed [2,80–82]. Nectin-3 is restricted exclusively to elongating/elongate spermatids which can heterotypically interacting with nectin-2 on the Sertoli cell side [68,83]. Spermatozoa from *nectin-2*<sup>-/-</sup> mice were morphologically aberrant and functionally impotent [83–85]. Since nectins are capable of activating Cdc42 via c-Src and a Cdc42 GEF (GDP/GTP exchange factor) [86], or activating Rac, thus recruiting the polarity complex Par3/aPKC/Par6 to the apical ES site [87], the absence of nectin-3 may also lead to malfunctioning of spermatid polarization, similar to *Jam-C*<sup>-/-</sup> mice [29]. Nectins are known to initiate cell-cell contacts by recruiting cadherin and JAM-A to establish functional AJ and TJ in epithelial cells [79,87–89]. It is likely that nectin-2/-3 and JAM-B/-C can also interact with each other since they are all localized to the elongating/elongate spermatids at the apical ES site, which should be investigated in future studies. In the Adjudin-induced germ cell loss model, it was found that the nectin-3/afadin interaction became severely weakened before any obvious reduction in their protein levels was detected [68], illustrating this cell adhesion unit must be compromised to facilitate spermatid loss (Table 1).

Besides afadin and ponsin, cytoplasmic adaptors that link nectin to the actin-based cytoskeleton [79], a new adaptor protein ADIP (afadin DIL domain-interacting protein) has recently been localized to AJ sites that interacts with both  $\alpha$ -actinin and afadin, providing additional cytoplasmic link between nectin- and cadherin-based cell adhesion units [90,91]. ADIP is highly expressed in the mouse testis [90]. Another possible linker that binds to both afadin and  $\alpha$ -actinin is LMO7 (LIM domain only 7), however, its presence in the rat testis failed to be confirmed by immunoblot analysis [92].

Nectin-like (Necl) molecules are similar to nectins, but do not bind to afadins [87,88]. This group of calcium-independent cell adhesion molecules consists of five members, capable of homo- or heterophilic interactions with nectins, and are important cell-cell adhesion molecules in various tissues [87,88]. At least Necl2 has been shown to be highly expressed in the rat testis [93]. It will be important to explore the significance of Necls in the testis, which is likely to involve in Sertoli-germ cell adhesion function.

Table 1  
Cytokine-mediated regulation of junction component proteins in epithelia including the testis

Junction component	Protein	Cytokine/hormone that modulates the steady-state mRNA/protein level (+/–) or protein distribution pattern (d) of the target junction protein	Selected references
TJ-integral membrane	JAM-A	TNF (d), IFN- $\gamma$ (d), T $\downarrow$ (+)	[30,38]
	Occludin	TGF- $\beta$ 3 (–), HGF (–/d), TNF (–/d), IFN- $\gamma$ (–/d), VEGF (–/d, inhibited by ANP), IL-1 $\beta$ (–/d), IL-4 (–), IL-13 (–), MCP-1 (–), T $\downarrow$ (+)	[44,52,54–56,223–225]
	Claudin	TGF- $\beta$ 3 (–), TNF (–), FSH/cAMP (–)	[43,44]
AJ-integral membrane	N-Cadherin	HGF (+), EGF (+), TGF- $\beta$ (+/d), T (+), T $\downarrow$ (+), IL-6 (–)	[30,66,78]
	E-Cadherin	TGF- $\beta$ (–/d), T (+)	[52,66,68]
	Nectin-3	TGF- $\beta$ 3(–/d)	[52,68]
	Integrin- $\beta$ 1	TGF- $\beta$ (+), T $\downarrow$ (+),	[30,226]
Adaptor	ZO-1	TGF- $\beta$ (d), IL-4 (–), IL-13 (–), T $\downarrow$ (+)	[30,225]
	Afadin	TGF- $\beta$ 3(–/d)	[68]
	$\beta$ -Catenin	TGF- $\beta$ (d), T $\downarrow$ (+)	[30,139,227]
	$\alpha$ -Catenin	TGF- $\beta$ (d), T $\downarrow$ (+)	[30,139,227]

T $\downarrow$ , suppression of intratesticular testosterone level with the use of testosterone (T) and estradiol implants; +, stimulation; –, inhibition. Protein distribution pattern was assessed by either immunofluorescent microscopy or immunohistochemistry using testicular cells cultured in vitro or seminiferous epithelium in vivo.

482 2.3.2.3. *The integrin/laminin complex.* The integrin/lami- 516  
 483 nin protein complex has recently been identified at the apical 517  
 484 ES which confers Sertoli–germ cell adhesion and provides a 518  
 485 new platform regarding how these two cell types interact 519  
 486 with each other and coordinate spermatogenesis [37,94].  
 487 Integrin-based protein complexes are usually found at the  
 488 cell–matrix junctions, such as hemidesmosomes or focal  
 489 adhesion, which further connects to the intermediate  
 490 filament or actin bundles, with integrin also capable of  
 491 serving as a cell receptor for the ECM [95,96]. Interestingly,  
 492 the junctions between Sertoli and germ cells are not simple  
 493 cell–cell junction types; rather, they are a hybrid of both  
 494 cell–cell and cell–matrix junction types, probably to  
 495 facilitate rapid junction turnover and germ cell migration  
 496 during spermatogenesis [22]. Several recent reviews on the  
 497 role of integrins and ECM in the testis are available, thus this  
 498 information is not discussed herein [22,97].

### 499 3. Cytokines are key regulators of junction dynamics 520 500 in the testis

501 Cytokines are regulatory peptides (usually  $\leq 30$  kDa in 521  
 502 size) produced virtually by every nucleated cells in 522  
 503 mammals and have pleiotropic actions on cell physiology 523  
 504 as an autocrine or paracrine factor [98]. In the testis, Sertoli 524  
 505 and germ cells produce a number of cytokines, including 525  
 506 members of the TGF- $\beta$  superfamily (e.g., TGF- $\beta$ s, activins, 526  
 507 inhibins), platelet-derived growth factor (PDGF), interleu- 527  
 508 kins (e.g., IL-1, IL-6, IL-11), tumor necrosis factor (e.g., 528  
 509 TNF $\alpha$ , Fas ligand), interferons (e.g., IFN- $\alpha$ , IFN- $\gamma$ ), 529  
 510 fibroblast growth factor (FGF), nerve growth factor 530  
 511 (NGF), and stem cell factor (or steel factor) (for reviews, 531  
 512 see [1,2,11,99,100]) (see also Table 2). These cytokines likely 532  
 513 mediate crosstalk between Sertoli and germ cells to facilitate 533  
 514 germ cell movement across the seminiferous epithelium and 534  
 515 other cellular events in the epithelium during the epithelial 535

cycle such as germ cell differentiation. Herein, we critically 516  
 evaluate two best studied cytokines, namely TNF $\alpha$  and TGF- 517  
 $\beta$ 3, regarding their significance in spermatogenesis in the 518  
 testis and briefly summarize the action of other cytokines. 519

#### 520 3.1. TNF

521 TNF, also known as TNF $\alpha$  or cachectin, is synthesized as 521  
 a 26 kDa type II transmembrane prepeptide (pro-TNF), 522  
 which is subsequently activated by proteolytic cleavage to 523  
 release the C-terminal 17 kDa mature protein by the TNF- 524  
 converting enzyme (TACE). The mature protein is formed 525  
 by aggregates creating a homotrimer that can bind to two 526  
 types of receptors: TNFR1 and TNFR2 [101,102]. The 527  
 major source of TNF $\alpha$  in mammalian body is immune cells 528  
 such as macrophage and monocytes, but TNF $\alpha$  is also 529  
 produced by other non-immune cells including astrocytes, 530  
 keratinocytes, Sertoli cells and germ cells [57,101]. TNF 531  
 signaling is mediated mainly through TNFR1, which has 532  
 distinct domains that facilitate the recruitment of other 533  
 intracellular adaptors to activate signaling pathways. The net 534  
 result of such activation can modulate apoptosis, inflamma- 535  
 tion and cell proliferation [101,103]. These adaptors include 536  
 TNFR1-associated death domain protein (TRADD) which 537  
 can recruit Fas-associated death domain protein (FADD), 538  
 TNF receptor associated factor-2 (TRAF-2), or receptor- 539  
 interacting protein (RIP), to induce the caspase-mediated 540  
 apoptosis, activate transcription factors (e.g., c-jun, c-fos, 541  
 ATF-2) via MAPK (ERK, JNK and p38), or activate nuclear 542  
 factor kappa B (NF $\kappa$ B) through inhibitor of NF $\kappa$ B kinase 543  
 (IKK), respectively [101–103]. A TNFR1 scaffolding 544  
 protein called TGFR-associated ubiquitous scaffolding 545  
 and signaling protein (TRUSS) has recently been cloned 546  
 and characterized [104]. The expression of TRUSS is 547  
 enriched in heart, liver and testes, it is also known to interact 548  
 with TRADD, TRAF-2 and IKK [104]. In addition to these 549  
 complex signaling networks that can be activated down- 550



Table 2  
Cytokines and their functions in the testis

Group	Cytokine	KO mice	Cellular expression	Function in the testis	References
TGF- $\beta$	TGF- $\beta$ 1	Perinatal/neonatal lethal	Sertoli, Leydig, germ, myoid cells	Testicular development	[109,112,119,123]
	TGF- $\beta$ 2	Perinatal lethal	Sertoli, Leydig and germ cells	Testicular development	
	TGF- $\beta$ 3	Perinatal lethal	Sertoli and germ cells	Junction dynamics	
	Activin/inhibin $\beta$ A	Perinatal lethal	Sertoli and peritubular cells	Regulate FSH production, testicular development	[109,228]
	Activin/inhibin $\beta$ B	Viable/reproductive abnormality (female)	Sertoli cells, germ cells		
	BMP-4	Embryonic lethal (–/–)/ lowered fecundity (+/–)	Pachytene spermatocytes, Sertoli cells (early postnatal)	Maintain spermatogenesis; spermatogonia differentiation	[109,229–231]
TNF	TNF $\alpha$	Viable/fertile	Sertoli, germ cells	Repress steroidogenesis, disrupt TJ, inhibit GC apoptosis	[57,106–108,218,232]
	FasL	Viable	Spermatocytes/spermatids; Sertoli cell (?)	Induce apoptosis, preserve immune-privilege	[233–235]
	TRAIL	Viable/fertile	Germ, Leydig cells	GC apoptosis	[236,237]
Growth factors	EGF	Viable/fertile	Sertoli, germ cells	Maintain spermatogenesis; stimulate steroidogenesis;	[238–240]
	FGF4	Embryonic lethal	Sertoli cells	Enhance spermatogenesis	[241–243]
	HGF	Embryonic lethal	Spermatozoa, myoid cells	Initiate sperm motility, induce testicular cord formation	[244–246]
	MIF	Viable/fertile	Leydig cells	Leydig-Sertoli cell paracrine mediator/inhibit inhibin production	[247,248]
	SCF	Perinatal lethal (–/–), sterile (+/–)	Sertoli cells (c-Kit receptor on differentiating spermatogonia)	Spermatogenesis; Sertoli cell-spermatogonia adhesion (membrane bound form)	[11,249,250]
	VEGF (A)	Embryonic lethal (+/–)	Sertoli cells (receptor on germ cells), Leydig cells	Spermatogonial proliferation, spermiogenesis	[251–253]
Interleukin/interferon	IL-1 $\alpha$ /-1 $\beta$	Viable/lowered fecundity	Sertoli cells, spermatocytes, spermatids	Inhibit steroidogenesis, Regulate Sertoli secretion	[254,255]
	IFN- $\gamma$	Viable/fertile	Spermatogonia, interstitium	Inhibit steroidogenesis, stimulate FasL expression	[256,257]

Abbreviation: bone morphogenetic protein (BMP), epidermal growth factor (EGF), fibroblast growth factor (FGF), hepatocyte growth factor (HGF, or scatter factor, SF), interferon (IFN), interleukin (IL), macrophage migration inhibitory factor (MIF), Stem cell factor (SCF, or Steel factor, SLF), transforming growth factor  $\beta$  (TGF- $\beta$ ), tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ), tumor necrosis factor- $\alpha$ -related apoptosis-inducing ligand (TRAIL), vascular endothelial growth factor (VEGF).

stream of TNF, at least 19 ligands and more than 20 receptors have been identified in the TNF superfamily [103], which can mediate an array of physiological processes and diseases [103]. It has been known that TNF can disrupt TJ integrity in multiple epithelial cells. As aforementioned, TNF down-regulates occludin expression through its promoter activity [55] or reduces JAM-A distribution on the vascular endothelial cell surface [38]. TNF level is also elevated in Crohn's disease, a chronic granulomatous inflammatory disease that affects the gastrointestinal tract, which is manifested by impaired intestinal barrier function with leaky TJs [105]. With the exception of TNF and Fas ligand (FasL), the roles of other members of TNF superfamily in the testis remain elusive.

In the testis, TNF $\alpha$  has been shown to play a role in regulating germ cell apoptosis, junction remodeling and Leydig cell steroidogenesis [57,106,107]. For instance, it is known that TNF $\alpha$  represses the expression of steroidogenic-enzyme genes in Leydig cells through an activation of NF $\kappa$ B, which can in turn inhibit the transactivation of orphan nuclear receptors [106]. Intratesticular injection of TNF in normal and hypophysectomized rats has also demonstrated its suppressive effect on testosterone production in vivo [108]. Chronic infusion of TNF caused germ cell (in particular spermatocytes and spermatids) depletion from the epithelium, a loss of testis weight and a plunge in testosterone level [58]. It remains unknown regarding the mechanism(s) by which TNF $\alpha$  utilized to induce these changes, but this could involve a suppression of Leydig cells steroidogenesis, or an inhibition of Sertoli cell TJ protein production at the BTB, or via its direct effect on germ cells. Other recent studies have shown that TNF $\alpha$  can perturb the TJ-permeability barrier in cultured Sertoli cells dose-dependently and reversibly since the disrupted TJ-barrier can be resealed upon the removal of the cytokine [57]. This inhibitory effect of TNF $\alpha$  on Sertoli cell TJ function is likely mediated via an induced production of collagen  $\alpha$ 3(IV), matrix metalloprotease (MMP)-9 and tissue inhibitor of metalloprotease (TIMP)-1 which collectively affect the homeostasis of ECM, thereby altering the association of the Sertoli cell epithelium with the basement membrane and perturbing the TJ-barrier [57]. Also, TNF $\alpha$  can activate the integrin/integrin linked kinase (ILK)/glycogen synthase kinase (GSK)  $\beta$ -3/p130 Cas/JNK signaling pathway which also contribute to changes in the TJ-protein expression and/or distribution at the BTB [22,57,97].

### 3.2. TGF- $\beta$

The TGF- $\beta$  superfamily comprises of TGF- $\beta$ s, activins, inhibins, bone morphogenetic proteins (BMPs), growth differentiation factors (GDFs), Müllerian-inhibiting substance (MIS) and others, totaling more than 35 members [109]. TGF- $\beta$  superfamily proteins are crucial in the regulation of a variety of biological processes, including cell proliferation, differentiation, apoptosis, and tissue

remodeling [110]. Some members, like activins and inhibins, were initially identified in the male gonad for their ability to regulate the pituitary follicle stimulating hormone (FSH) production [11]. MIS is known for its role in sexual differentiation causing the regression of the Müllerian ducts in the male [111]. The functions of TGF- $\beta$  superfamily proteins in reproduction have been recently reviewed [109,112] hence we only focus on regulation of junction restructuring by TGF- $\beta$ s herein, which is elaborated in Section 4. Table 2 summarizes other cytokines that are known regulators of junction dynamics.

### 3.3. Cytokines working in concert with other ECM proteins to regulate junction dynamics

Recent reviews have summarized how cytokines regulate the homeostasis of proteases and their inhibitors, and ECM proteins to coordinate spermatogenesis [2,22,97]. It is not at all surprising that these molecules are working in concert since their production, activation, and termination are all interdependent and connected. Their homeostasis and regulation are essential to almost all biological processes. In the testis, for instance, when the BTB is disrupted by cadmium, TGF- $\beta$ 3/p38 MAPK signaling is activated to down-regulate the steady-state of TJ and AJ protein levels that leads to the breakdown of both junctions and germ cell exfoliation [44,52,113]. Proteases (e.g., cathepsin L) and protease inhibitors (e.g.,  $\alpha$ <sub>2</sub>-macroglobulin) are induced to coordinate the junction restructuring event [52]. Using a p38 MAPK inhibitor SB202190, the damage to the BTB and the plunge of TJ and AJ proteins induced by CdCl<sub>2</sub> can be delayed but it cannot prevent the overexpression of protease inhibitor  $\alpha$ <sub>2</sub>-MG [52]. Further study revealed that  $\alpha$ <sub>2</sub>-MG production is regulated by JNK signaling pathway in the testis, independent of the p38 MAPK pathway [114]. This yin and yang relation of protease and protease inhibitor regulation that utilizes distinct signaling pathways, and their connection with cytokines (e.g., TGF- $\beta$ 3) have illustrated that the testis is equipped with some delicate regulatory mechanisms to orchestrate junction restructuring at spermatogenesis.

## 4. TGF- $\beta$ 3 as a junction regulator—versatility realized through selectivity

### 4.1. Signaling conduits and versatile players in biological processes

TGF- $\beta$ s ( $\beta$ 1,  $\beta$ 2 and  $\beta$ 3) are key regulators in a plethora of biological processes (for reviews, see [110,115–120]). These cytokines, when activated by releasing from the latency-associated proteins (LAPs), can bind to their receptors—first to the type II receptor, T $\beta$ R<sub>II</sub>, which then recruits the type I receptor, T $\beta$ R<sub>I</sub> (or ALT5, activin-like kinase)—although TGF- $\beta$ 2 requires binding of the two

receptors more or less at the same time and the assistance from the type III receptor, betaglycan. The binding of the cytokine to type I and type II receptors initiates a series of phosphorylation mediated activation—autophosphorylation of T $\beta$ RII and T $\beta$ RI phosphorylation by T $\beta$ RII—and triggers consequent intracellular signaling events (the canonical Smad-mediated signalings and Smad-independent pathways). Despite their structural similarities and shared signaling mechanisms, the three TGF- $\beta$ s are spatiotemporally expressed and play non-redundant roles, particularly under in vivo conditions. This in part is attributed to their unique promoter sequences [121]. For instance, in the mouse testis, expression of TGF- $\beta$ 1 and TGF- $\beta$ 2 are much higher in embryonic and early postnatal stages, and TGF- $\beta$ 3 becomes the highest expressed among the three isoforms in adulthood [122]. Similarly, in postnatal day 5 to day 60 rats, TGF- $\beta$ 1 and TGF- $\beta$ 2 expression are predominant in immature testes, which decrease at the onset of puberty; whereas TGF- $\beta$ 3 expression is most abundant at the pubertal stage, coinciding with the initiation of spermatogenesis [123]. These thus illustrate TGF- $\beta$ s have unique roles in distinct phases of testicular development: TGF- $\beta$ 1 and TGF- $\beta$ 2 are important for the development while TGF- $\beta$ 3 takes the center stage during spermatogenesis. Herein, we summarize the TGF- $\beta$ -mediated signaling conduits, focusing on their regulation of junction remodeling.

#### 4.1.1. Smad-mediated signaling

The smad-mediated TGF- $\beta$  signaling pathways have been extensively characterized and recently reviewed [110,116,124,125]. Among the 8 Smad proteins (Smad1–8), receptor-regulated R-Smad (Smad2 and Smad3), common-partner Co-Smad (Smad4) and inhibitory I-Smad (Smad 7) are involved in TGF- $\beta$ /T $\beta$ RII/T $\beta$ RI signaling. However, many of these Smad proteins have not been subjected to rigorous investigation in the testis. In the testis, the expression of Smad2 and Smad3 are developmentally regulated and stage-specific: being more prominent in prepubertal than in sexually mature rats, and at the lowest levels at stages VII–VIII of the epithelial cycle in adult rats [126]. Expression of Smad3, 4, 6 and 7 are also detected in embryonic mouse testes [127]. It is not surprising that Smad proteins are highly expressed in younger animals since TGF- $\beta$  superfamily members are essential for development. TGF- $\beta$ 1 and TGF- $\beta$ 2 may be more important in the testis at the early stages through Smad-mediated signaling pathways. Yet the regulation and maintenance of spermatogenesis by TGF- $\beta$ 3 in adult testes is likely mediated via Smad-independent signalings, such as TJ and BTB dynamics [68]. For instance, TGF- $\beta$ 3 activates ERK without activation of Smad2 and Smad3 in the Adjudin-induced germ cell loss model [68].

#### 4.1.2. MAPK-mediated signaling

There are accumulating evidence in the literature regarding Smad-independent TGF- $\beta$  signalings that regulate

diverse biological function, which has recently been reviewed [115,116,118,128]. Amongst these, the best studied is the MAPK signalings [129–131]. For instance, TGF- $\beta$  is capable of activating all three MAPK pathways [115,118,125]. In the testis, all three pathways have been implicated in the regulation of junction dynamics pertinent to spermatogenesis. First, JNK pathway is involved in TNF- $\alpha$ -induced TJ restructuring and  $\alpha$ <sub>2</sub>-MG regulation [57,114]. Second, ERK pathway can be activated via either integrin or TGF- $\beta$ 3, which can in turn regulate AJ dynamics [36,37,68]. Third, p38 MAPK is responsible for TGF- $\beta$ 3-activated TJ and AJ restructuring [52,113,132]. Nonetheless, little is known about the expression and distribution of MAPKs and their upstream kinases in the testis. ERK1/2 and p-ERK1/2 have been localized to the elongate spermatids at the apical ES/TBC site in the epithelium at stages VII–VIII [68,133], illustrating its role in spermiation. ERK1/2 is also detected at the basal compartment of the epithelium [133]. Indeed, when induced by Adjudin, p-ERK1/2 is activated at the site of apical ES in depleting elongate/elongating spermatids in tubules other than stages VII–VIII, probably facilitating germ cell exfoliation [68].

The complexity of TGF- $\beta$ -mediated signaling pathways is manifested by the presence of multiple intracellular interacting points. Recent studies have identified different interacting proteins with TGF- $\beta$  receptors, illustrating these proteins may play a role in selecting the downstream signaling events. For example, occludin is known to associate with T $\beta$ RI and as such, TGF- $\beta$  can efficiently regulate TJ disruption during epithelial-mesenchymal transition (EMT) [134]. Indeed, the proximity of TGF- $\beta$  receptors with TJ proteins has created an efficient regulatory mechanism where TGF- $\beta$ -induced TJ dissolution is mediated through the cell polarity complex. Upon activation by TGF- $\beta$ , T $\beta$ RII is recruited to the T $\beta$ RI/occludin/Par6 complex, thereby phosphorylating Par6, this in turn stimulates Par6 which binds to Smurf1 (an E3 ubiquitin ligase), and causing degradation of RhoA that leads to TJ disassembly [135,136]. Although it has not yet been confirmed for T $\beta$ RII, proteins that associate with type II receptor of BMP have recently been identified, which include MAPK, PKC, and cytoskeleton tubulin  $\beta$ 5 [137]. These proteins associate not only with the kinase domain of the receptor but also its C-terminus [137], illustrating receptors of the TGF- $\beta$  family proteins can affect junction dynamics via protein-protein interactions with junction protein complexes.

#### 4.1.3. TGF- $\beta$ s regulate junction restructuring

TGF- $\beta$ s regulate junction dynamics in various cell types. For instance, TGF- $\beta$ 1 can perturb the permeability of the blood–retinal barrier via a stimulation of MMP-9 production [138]. TGF- $\beta$ 1 also perturbs the TJ-permeability barrier in pulmonary endothelial monolayers by inducing AJ proteins to move away from the cell–cell contact site, possibly via a myosin light chain kinase mediated mechanism [139]. TGF- $\beta$ 1 and Ras can also work synergistically to promote cell

764 invasiveness in intestinal epithelial cells by down-regulating  
 765 E-cadherin expression and subcellular redistribution of  $\beta$ -  
 766 catenin [140]. In addition, TGF- $\beta$ 1 can induce AJ disruption  
 767 in renal proximal tubular epithelial cells, which cannot be  
 768 reproduced by transient overexpression of Smad2/4 or  
 769 Smad3/4 [141], illustrating this is an Smad-independent  
 770 signaling event. On the other hand, a blockage of TGF- $\beta$   
 771 signaling by treatment of a TGF- $\beta$  receptor kinase inhibitor  
 772 up-regulates TJ protein production (e.g., claudin-5) in  
 773 embryonic stem cell-derived endothelial cells [142].  
 774 Interestingly, in almost all of these epithelial/endothelial  
 775 cells, a disruption of either TJ or AJ can affect the integrity  
 776 of the other junction type following an induction by TGF- $\beta$ s.  
 777 Yet the functional inter-relationship of AJ and TJ in the  
 778 seminiferous epithelium is significantly different from all  
 779 other epithelia and endothelia. For instance, TGF- $\beta$ 3 (and  
 780 also TGF- $\beta$ 2 in vitro) can disrupt the Sertoli–Sertoli TJ-  
 781 barrier by down-regulating TJ proteins (e.g., occludin) via  
 782 p38 MAPK signaling pathway and this effect is indeed  
 783 confirmed using an in vivo model to study the BTB  
 784 dynamics [44,113,132] (see Fig. 2). Analogous to other  
 785 epithelia and endothelia, a breakdown of TJ can indeed  
 786 affect the integrity of AJ, resulting in a loss of Sertoli–germ  
 787 cell adhesion [52]. However, a disruption of AJ between  
 788 Sertoli–germ and Sertoli–Sertoli cells seems to reinforce the  
 789 TJ at the BTB instead, let alone its disruption, in the  
 790 Adjudin- and intratesticular testosterone suppression-  
 791 induced germ cell loss models [2,30]. Recent studies have  
 792 shown that TGF- $\beta$ 3 can exert its effects on AJ integrity via  
 793 a signaling pathway different from the one that regulates TJ  
 794 dynamics in the testis [68], so that Sertoli–germ cell AJ can  
 795 undergo restructuring without perturbing the BTB integrity  
 796 (Fig. 2). This unique relation of AJ and TJ in the  
 797 seminiferous epithelium may be a physiological require-  
 798 ment for the testis to facilitate germ cell migration (i.e., AJ  
 799 restructuring) while maintaining TJ integrity. This concept  
 800 will be revisited and discussed in detail in Section 5.

#### 801 4.2. Signaling regulation and selectivity

##### 802 4.2.1. Multilayers of signal modulation

803 Regulation of TGF- $\beta$ -mediated signalings occurs at  
 804 multiple levels: ligand production and activation, ligand–  
 805 receptor coupling, intracellular signal pathway selection,  
 806 nucleocytoplasmic shuttling of transcription factors, an  
 807 interaction of multiple transcription factors that finally  
 808 determines the activation or repression of gene expression,  
 809 and signal termination [110,143–146]. Less is known  
 810 regarding how the expression of TGF- $\beta$ s is regulated. The  
 811 promoter sequences of human TGF- $\beta$ s have been char-  
 812 acterized. For instance, TGF- $\beta$ 1 is mostly regulated by AP-1  
 813 site lacking TATA box, whereas TGF- $\beta$ 2 and TGF- $\beta$ 3 are  
 814 regulated by AP-2 site and cAMP-responsive elements,  
 815 containing TATA box [121], and the most potent activator of  
 816 TGF- $\beta$ 1 expression known thus far is the cytokine itself  
 817 [147]. It has been shown that JNK suppresses the autocrine

818 expression of TGF- $\beta$ 1 in fibroblasts [148]. A recent in vivo  
 819 study in the testis has shown that JNK signaling is required  
 820 for the production of  $\alpha_2$ -MG in the seminiferous epithelium,  
 821 which tethers TGF- $\beta$ 3 and antagonizes the cytokine [114].  
 822 These results thus illustrate the TGF- $\beta$  action is regulated at  
 823 multiple levels and can induce diversified biological  
 824 responses. Upon secretion, TGF- $\beta$ s are tightly but non-  
 825 covalently bound to LAPs, which are further tethered to  
 826 latent transforming growth factor- $\beta$  binding proteins  
 827 (LTBPs) via covalent bonds [149]. LTBP can covalently  
 828 bind to ECM, enabling cytokines to be retained in the matrix  
 829 and creates a reservoir [149]. This biologically inactive  
 830 cytokine pool can be activated by low pH, protease (e.g.,  
 831 plasmin, MMP-2 and MMP-9), thrombospondin-1 (TSP-1),  
 832 integrin- $\alpha$ v $\beta$ 6 or - $\alpha$ v $\beta$ 8 [145,146,149,150]. At least one  
 833 LTBP called LTBP-1L (long form) is highly expressed in  
 834 testes [150]. MMP-2 and MMP-9 are also found in the testis  
 835 [97]. Other antagonists of TGF- $\beta$ s include  $\alpha_2$ -MG and  
 836 decorin, which can ‘lock’ the ligand and prevent its binding  
 837 with receptors, and endoglin, which binds to T $\beta$ RII-  
 838 associated TGF- $\beta$ 1 or - $\beta$ 3 and attenuates T $\beta$ RI mediated  
 839 signaling [110,151,152]. After TGF- $\beta$  binds to its receptors,  
 840 signaling is triggered but can be directed to a distinctive  
 841 pathway, and can sometimes activate multiple pathways.  
 842 Because of such diversified signaling capacity, a mechanism  
 843 must be in place to choose the needed downstream signaling  
 844 pathway. It is likely that adaptor proteins play the decision-  
 845 making role. For instance, activation of Smad2/3 is  
 846 facilitated by the adaptor SARA. Yet the detail of this  
 847 selection still remains elusive. To transmit the signaling to  
 848 the corresponding genes for their transcriptional induction,  
 849 activated transcription factors (e.g., Smad2 and Smad3)  
 850 must enter the nucleus. As such, there is constant  
 851 nucleocytoplasmic shuttling of the R-Smads between active  
 852 (phosphorylated) and inactive (dephosphorylated) status to  
 853 keep sensing the signals at real-time [110,144]. The cell-  
 854 specific and non-specific transcription factors/coactivator/  
 855 co-repressors can determine the final gene expression  
 856 outcome in a particular cell type at the end of TGF- $\beta$   
 857 activation [110,116]. Receptor internalization and degrada-  
 858 tion, Smad shuttling and ubiquitination, and expression  
 859 feedback can all contribute to the signal termination [118].

##### 860 4.2.2. Adaptors as molecular switches for TGF- $\beta$ 861 signaling in the testis

862 It is of interest to note that in the testis, the TGF- $\beta$ 3-  
 863 activated signaling can have distinctive effects on the  
 864 junction restructuring. When p38 MAPK is activated by  
 865 TGF- $\beta$ 3, the BTB in the seminiferous epithelium is  
 866 disrupted concomitant with Sertoli–germ cell AJ disas-  
 867 sembly [52] (Fig. 2). In contrast, when ERK1/2 is activated  
 868 by TGF- $\beta$ 3, only AJs are affected without affecting the BTB  
 869 integrity [68] (Fig. 2). Indeed, a blockade of the TGF- $\beta$ 3-  
 870 mediated signaling by using an antagonist (e.g., T $\beta$ RII/Fc  
 871 conjugate) can prevent the activation of ERK1/2 and  
 872 significantly delay the Adjudin-induced germ cell loss from




Fig. 2. A schematic illustration of how cytokines (e.g., TGF- $\beta$ 3, TNF $\alpha$ ) can regulate junction dynamics in the testis via their effects on the steady-state levels of proteins (e.g., TJ- and AJ-proteins, protease inhibitors such as  $\alpha_2$ -MG) at the BTB and apical ES. This model was prepared based on recent studies from this laboratory using different animal models as reviewed herein. In brief, cytokines released from either Sertoli or germ cells can activate at least three different signaling pathways upon their binding to receptors. For instance, TGF- $\beta$ 3 can activate p38 MAPK signaling pathway to down-regulate both TJ and AJ proteins, resulting in the disruption of the BTB and Sertoli-germ cell adhesion function, eventually leading to germ cell loss from the epithelium (green sector), which was identified in studies using the cadmium model [52,113,132]. This also illustrates that when TGF- $\beta$ 3 utilizes the p38 MAPK pathway for its signaling function, it can perturb both the BTB and apical ES integrity. When rats were treated with Adjudin, or testosterone/estradiol implants to reduce intratesticular androgen level, the testis responds to these treatments with an induction of TGF- $\beta$ 3 that can activate only the ERK signaling pathway to compromise Sertoli-germ cell adhesion function by lowering the steady-state protein levels at the apical ES or weakening protein-protein interactions at this site via changes in the phosphorylation status of adaptors (e.g.,  $\beta$ -catenin); this in turn leads to the loss of germ cells from the epithelium, and this event does not affect the BTB

873 the epithelium [68]. By blocking these two MAPK signaling  
 874 pathways using kinase inhibitors can also rescue the  
 875 epithelium from the disruptive effects of CdCl<sub>2</sub> and Adjudin  
 876 on the BTB and Sertoli–germ cell AJ, respectively  
 877 [52,68,113]. As such, TGF-β<sub>3</sub> serves as a key regulator  
 878 that decides whether BTB is affected or not. It is tempting to  
 879 speculate that this TGF-β-activated MAPK signaling  
 880 cascade requires the recruitment of adaptors to the site  
 881 which can in turn shuttle to the correct signaling pathway  
 882 downstream. Recent studies have shown that TGF-β-  
 883 induced p38 MAPK activation is mediated through a protein  
 884 scaffold complex XIAP (X-linked inhibitor of apoptosis)/  
 885 TAB1 (TAK binding protein)/TAK1 (TGF-β-activated  
 886 kinase), in which adaptor XIAP may link adaptor TAB1  
 887 and MAPKKK TAK1 to the TβRI. TβRI can activate  
 888 TAK1, which further activates either MKK3/6 or MKK4 that  
 889 in turn activates p38 MAPK or JNK, respectively [118].  
 890 Besides, MAPKK-independent autophosphorylation of p38  
 891 is also possible, which is TAB1 dependent [153]. On the  
 892 other hand, TGF-β can activate Ras, which further activates  
 893 ERK signaling pathway and regulates various cellular  
 894 processes including junction dynamics [68]. TGF-β-induced  
 895 ERK activation also requires the adaptor CD2-associated  
 896 protein (CD2AP). When this adaptor is not involved, p38  
 897 MAPK is preferentially activated instead. In its presence,  
 898 TGF-β activates both the PI 3-kinase/Akt and the Ras/ERK  
 899 pathways [154]. Interestingly, CD2AP is not involved in PI  
 900 3-kinase/Akt activation by EGF and insulin, nor in the  
 901 activation of Smad2 by TGF-β, suggesting it plays a role in  
 902 TGF-β-activated, Smad-independent signaling [155]. The  
 903 association between TGF-β receptors and CD2AP is further  
 904 supported by the evidence that both are present in lipid rafts  
 905 [156,157]. Thus CD2AP can serve as a molecular switch to  
 906 determine the downstream signaling direction of TGF-β.  
 907 CD2AP belongs to a family of ubiquitously expressed  
 908 adaptors containing three Src-homology 3 (SH3) domains, a  
 909 proline-rich region and a coiled-coil domain [158] and is  
 910 expressed in human testes [159]. The SH3 domain mediates  
 911 interaction with the p85 subunit of PI 3-kinase and the  
 912 proline-rich region mediates association with p130 Cas and  
 913 Src family kinases (for a review, see [158]). p130 Cas and  
 914 Src are components of a signaling machinery connecting  
 915 FAK, paxillin, ERK and myosin light chain kinase (MLCK),  
 916 which, in turn regulate cell adhesion during cell migration  
 917 process [160]. Interestingly, virtually all of these proteins  
 918 have recently been found in the testis and they are likely  
 919 involved in junction dynamics during spermatogenesis (for

reviews, see [2,22,97]). It is likely that CD2AP is a crucial  
 adaptor of TGF-β mediated and integrin/FAK mediated  
 signaling events in the testis. Although much of the  
 information on CD2AP derives from studies in the kidney,  
 the testis may employ this molecular switch to select the  
 downstream signaling pathways to be activated by TGF-β<sub>3</sub>,  
 affecting either AJ alone or TJ and basal ES at the BTB. This  
 should be vigorously validated in future studies.

## 5. What lessons we learn from the testis as a model to study junction restructuring?

As we have discussed above, the testis is an intriguing  
 organ where extensive junction restructuring occurs in the  
 seminiferous epithelium at each stage of the epithelial cycle.  
 Recent studies aiming to delineate the mechanisms that  
 regulate the junction restructuring events in the testis have  
 yielded some crucial information, which is likely applicable  
 to general cellular physiology as a whole. Herein, we  
 summarize several *in vivo* models that have been established  
 and used in recent studies (see Table 3). We only highlight  
 some of the latest development using these models and  
 readers are encouraged to refer to several recent reviews  
 [1,2,9,75,161,162].

### 5.1. Adjudin model

Formerly called AF-2364 [1-(2,4-dichlorobenzyl)-1H-indazole-3-carbohydrazide], Adjudin is a molecule that  
 selectively induces adherens junction disruption. It is a well  
 studied potential male contraceptive derived from indazole-3-  
 carboxylic acid [163,164]. It is also one of the best studied  
 compounds that induce germ cell sloughing in the testis.  
 Adjudin apparently exerts its effects on the Sertoli–germ cell  
 adhesion unit to induce a loss of AJ function by triggering a  
 couple of signaling events, including a surge of the ES-  
 associated signaling molecule testin, an induction of integrin-  
 and cadherin-initiated pathways, as well as TGF-β<sub>3</sub> activa-  
 tion [37,66–68,74,163,165–167]. Moreover, the ES-based  
 AJs are compromised due to a loss of protein-protein  
 association in the N-cadherin/β-catenin and nectin/afadin  
 protein complexes, which is likely the result of a coordinated  
 regulation by protein and lipid kinases and phosphatases,  
 proteases and protease inhibitors [35,67,68,74,94]. These  
 signalings are triggered within a few hours after adult male  
 rats are treated with a single or multiple doses of Adjudin at

integrity (blue sector) [2,68]. This thus suggests that TGF-β<sub>3</sub> can limit its action at the apical ES without compromising the BTB when the ERK signaling pathway is being utilized. Using the cadmium model, it is presently known that JNK is activated during the cadmium-induced BTB damage. This induces the production of α<sub>2</sub>-MG, which either bind to the free cytokines, limiting their biological action and/or blocking protease activity to limit the BTB damage. Since it is known that by blocking the production of α<sub>2</sub>-MG, it can worsen the damaging effect of cadmium on testicular junctions (red sector), illustrating this JNK-α<sub>2</sub>-MG pathway is crucial to maintain the normal physiology in the seminiferous epithelium [57,114]. This pathway is likely utilized by TNFα to regulate the steady-state protein level of α<sub>2</sub>-MG. The coordinated action of these three interacting signaling pathways that are intriguingly regulated by cytokines (e.g., TGF-β<sub>3</sub> and TNFα) is crucial to maintain the integrity of the seminiferous epithelium during spermatogenesis, permitting selective disruption of either TJ, AJ, or TJ and AJ. As reviewed herein, it is likely that adaptors play a crucial role upstream to select which signaling pathway should be activated by these cytokines, which in turn determines if either BTB, apical ES, or both BTB and apical ES should be compromised during spermatogenesis.

Table 3  
Chemicals that target the testis and can potentially serve as models to study junction dynamics

Affected junction types	Chemical	Classification	Target junction types in the testis and manifestations	References
TJ/AJ	Cadmium	Heavy metal	BTB/ES disruption; germ cell loss/apoptosis, irreversible	[52,114,178,179,200]
TJ	Cisplatin	Chemotherapeutic drug	BTB disruption; azoospermia, irreversible	[177,258,259]
TJ	Glycerol	1,2,3-Propanetriol	BTB disruption; germ cell loss, irreversible	[260,261]
TJ	Occludin peptide	22-a.a. from 2nd extracellular loop of occludin	BTB disruption; germ cell loss, reversible	[262]
AJ/TJ	Gossypol	Extract from cotton seed oil	ES/AJ/prevent BTB formation in neonatal animal; germ cell loss; irreversible in neonatal animal	[177,263–265]
AJ	Adjudin (AF-2364)	Indazole-3-carboxylic acid analog	ES/AJ disruption; germ cell loss, reversible	[163,164]
AJ	AF-2785		ES/AJ disruption; germ cell loss, reversible	[163,164]
AJ	Lonidamine (AF-1890)		ES/AJ disruption; germ cell loss; irreversible in selected subjects	[266]
AJ	Testosterone/estrogen implants	Steroid hormone	ES/AJ disruption; germ cell loss, reversible	[30,33,35,36,267,268]
AJ	Vinclozolin	Fungicide/antiandrogen	ES/AJ (?); germ cell apoptosis	[180,191,269,270]
AJ	Phthalate	Widely used as a plasticizer and in cosmetics/antiandrogen	Basal and apical ES disruption; seminiferous tubule atrophy, germ cell loss	[180,181,183,186]
AJ	Bisphenol A	Plastics/estrogenic	Apical ES disruption; abnormal spermatids, acrosomal defects	[177,182,198,271–273]

40–50 mg/kg b.w. either via i.p. or by gavage. Thereafter, morphological changes (i.e., germ cell depletion) are typically seen by 6–8 h [168]. The effect of Adjudin is limited to AJs, since the BTB remains intact in Adjudin treated rats. Furthermore, spermatogonia cell population apparently is unaffected [2]. Its antifertility effects are reversible, since the voided tubules treated with Adjudin can become repopulated with germ cells, making them almost indistinguishable from normal testes [164]. Studies from Adjudin treated rat testes have revealed some regulatory mechanisms that affect Sertoli–germ cell adhesion function pertinent to spermatogenesis. For instance, the integrin/FAK signaling is activated during Adjudin-induced germ cell loss from the epithelium [37]. This information has recently been validated and expanded using an androgen suppression-induced germ cell loss model in which rats received androgen and estrogen implants to suppress the intratesticular androgen level thereby perturbing Sertoli–germ cell apical ES function [36,37]. More important, TGF-β3 is also induced in androgen-suppressed rat testes, similar to the Adjudin model ([68] and unpublished observations), illustrating the involvement of cytokines in cell adhesion function. It is possible that the migration of germ cells across the seminiferous epithelium during spermatogenesis is controlled by several independent signaling pathways. When an agent activates these signalings, though the initial responses are different for different agents, the net outcome (i.e., alteration in Sertoli–germ cell adhesion function and the subsequent germ cell sloughing) is similar. Indeed, the signaling events in the rat testis identified using the Adjudin model have shown that this organ is utilizing the junction restructuring events usually restricted cell-ECM interface to regulate cell adhesion, migration, tissue remodeling and

development, and tumor cell metastasis [115,169,170], illustrating the cell–cell anchoring junction in the testis is indeed a hybrid cell–cell and cell–matrix junction type [22].

5.2. Cadmium model

Cd is a heavy metal and an environmental pollutant that is widely used in industry. It poses significant threat to human health and is classified as an endocrine disruptor [171–173]. It adversely affects a number of organs including the testis, kidney, lung, liver, pancreas and placenta [171,173]. The molecular mechanisms of action of cadmium toxicity are rather diverse which include: (i) binding to estrogen receptors, mimicking estrogen in the uterus and mammary gland [174]; (ii) disrupting the cadherin-based cell–cell adhesion [172]; (iii) inhibiting the DNA mismatch repair [175]; and (iv) disrupting endothelial and blood-testis barriers [52]. The testis is very sensitive to Cd exposure and the Cd-induced testicular effects (e.g., necrosis) is common across all animal species [176]. The antifertility effect of Cd has been known for decades. A recent study has identified a metal transporter of Cd (ZIP8, ZRT-, IRT-like protein 8) that is highly expressed in Sertoli cells [176], which likely explains, at least in part, why this cell type is sensitive to Cd-induced damages in the testis. Indeed, the junctional proteins are the early targets of a panel of toxicants, including Cd, in Sertoli cells cultured in vitro [177]. At a relative low dosage of Cd (e.g., 0.1–1 μM), it can reversibly perturb the Sertoli cell TJ-barrier in vitro when testosterone and FSH are present in the media [178]. Intraperitoneal administration of cadmium (1–3 mg/kg b.w.) to adult rats can irreversibly damage the BTB, which has been used as a model to study

1024 BTB dynamics in the testis [52,113,114,179]. Apparently,  
 1025 when absorbed by Sertoli cells, Cd targets the microfila-  
 1026 ment, causing a disorganization of actin bundles [179].  
 1027 Furthermore, Cd induces the dissolution of TJ proteins (e.g.,  
 1028 occludin) from the seminiferous epithelium, and down-  
 1029 regulates AJ-proteins (e.g., cadherin, nectin) to induce a  
 1030 secondary disruption of basal ES, leading to germ cell  
 1031 sloughing [52]. Using this model, it has been shown that the  
 1032 TGF- $\beta$ 3/MEKK/p38 MAPK mediated signaling pathway is  
 1033 a putative mechanism that regulates TJ dynamics at the BTB  
 1034 in vivo, and a disruption of this pathway using specific  
 1035 inhibitors can indeed significantly delay the Cd damage to  
 1036 the BTB as well as the subsequent germ cell loss from the  
 1037 epithelium [44,113,132].

### 1038 5.3. Possible in vivo models to study junction dynamics 1039 in the testis

1040 Recent studies have illustrated a number of chemicals  
 1041 that can affect testicular junctions, which may be developed  
 1042 into useful in vivo models. Many of these molecules are  
 1043 endocrine disruptors, which include phthalate, bisphenol A,  
 1044 vinclozolin and others [180–183]. Because of their wide-  
 1045 spread distribution in the environment and potential health  
 1046 hazards (e.g., reproductive organs), these compounds have  
 1047 attracted great attention of research, particularly on their  
 1048 effects to the reproductive organs (e.g., testes). They affect  
 1049 the endocrine system either by acting as antiandrogens  
 1050 (e.g., phthalate, vinclozolin) or estrogens (e.g., bisphenol A)  
 1051 [180–183].

1052 Phthalate and vinclozolin are compounds that can  
 1053 antagonize androgens. However, they exert these effects  
 1054 via different mechanisms: phthalate affects androgen  
 1055 synthesis [183] whereas the metabolites of vinclozolin are  
 1056 antagonists of androgen receptors [184,185]. Phthalate is  
 1057 found in cosmetic products (e.g., nail polishes, perfumes,  
 1058 hair sprays) and is widely used as a plasticizer, which can be  
 1059 non-covalently bound to the matrix and thereby slowly  
 1060 releases to the environment, and can be inhaled or adsorbed  
 1061 dermally [181,183]. Its toxic effects in male neonatal  
 1062 animals include hypospadias, reduced anogenital distance,  
 1063 vaginal pouch, some of which (e.g., hypospadias) are  
 1064 detected in humans [180,181,183,186]. When adults rats  
 1065 were treated with a single dose of di-*n*-pentyl phthalate  
 1066 (DPP) (2.2 g/kg b.w.), Sertoli cell junctions displayed  
 1067 abnormalities with disrupted basal ES, and apical ES was  
 1068 either absent or badly disorganized [187]. Interestingly, the  
 1069 disrupted basal ES between apposing Sertoli cells were  
 1070 reformed by 48 h after DPP treatment [187], illustrating this  
 1071 is a potentially useful model to study basal ES dynamics if  
 1072 adequately characterized. Furthermore, in DPP-treated  
 1073 prepubertal rats, extensive vacuolation occurs in Sertoli  
 1074 cells, to be followed by sloughing of germinal cells [188].  
 1075 Apparently the observed effects of phthalate on germ cell  
 1076 loss are mediated via disruption of Sertoli-Sertoli and  
 1077 Sertoli-germ cell adhesion function.

Vinclozolin [3-(3,5-dichlorophenyl)-5-methyl-5-viny-  
 1078 loxazolidine-2,4-dione] is a fungicide that is widely used  
 1079 in farming industry. When adsorbed by humans or rodents,  
 1080 vinclozolin is metabolized to M1 (2-[[[3,5-dichlorophenyl]-  
 1081 carbamoyl]oxy]-2-methyl-3-butenoic acid) and M2 (3',5'-  
 1082 dichloro-2-hydroxy-2-methylbut-3-enamide), which can  
 1083 bind to androgen receptors, antagonizing androgen function  
 1084 in vivo [184,185]. Besides its disruptive effects on  
 1085 reproductive organs (e.g., hypospadias, reduced anogenital  
 1086 distance) in male rats when exposed to vinclozolin in utero, it  
 1087 is neurotoxic and is an endocrine disruptor [189]. In the testis,  
 1088 vinclozolin can induce Leydig cell hypertrophy, reduce testis  
 1089 weight as a result of germ cell loss, and subsequently impair  
 1090 sperm production [190–192]. The direct structural damage of  
 1091 junctions at the Sertoli-Sertoli and Sertoli-germ cell interface  
 1092 following vinclozolin treatment remains to be examined.  
 1093

Bisphenol A is a xenoestrogen although its bioactivity is  
 1094 1000–1500-fold lower than 17 $\beta$ -estradiol. An ultrastructural  
 1095 examination of adult rat and mouse testes after treatment  
 1096 with bisphenol A has revealed that the apical ES was absent  
 1097 or badly damaged versus control animals; but, interestingly,  
 1098 the basal ES and BTB were not affected [182]. This  
 1099 information has also strengthened the notion that AJ  
 1100 disruption in the seminiferous epithelium can be restricted  
 1101 to the ES site without perturbing the TJ-barrier function at  
 1102 the BTB [2,30] in contrast to other epithelia where a  
 1103 disruption of AJ can lead to a secondary damage of the TJ-  
 1104 barrier function and vice versa [193–197]. The disruptive  
 1105 effects of bisphenol A and estrogens (17 $\beta$ -estradiol and  $\beta$ -  
 1106 estradiol-3-benzoate) on apical ES (but not basal ES and  
 1107 BTB, which remained intact) was also detected in  
 1108 maturing rats and mice that had been exposed to bisphenol  
 1109 A at neonatal [198]. However, these effects were not found  
 1110 when rats were fully mature [198]. In short, one of the target  
 1111 structures of this endocrine disruptor is the apical ES.  
 1112

### 1113 5.4. Why is the testis a vulnerable target of 1114 environmental toxicants? A lesson to learn from the 1115 testis

1116 Studies on different environmental toxicants have  
 1117 unequivocally demonstrated that the testis is extremely  
 1118 vulnerable to these toxicants (for a review, see [199]). When  
 1119 exposed to these chemicals, Sertoli-Sertoli and Sertoli-germ  
 1120 cell junctions are the early targets and their subsequent  
 1121 dissolution is likely the result of down-regulation of junction  
 1122 proteins or changes in protein-protein association of the  
 1123 junction protein complexes [177]. Indeed, recent studies  
 1124 have shown that when adult rats were exposed to cadmium  
 1125 chloride, the BTB damage had occurred at least 24 h before  
 1126 the TJ-barrier of the microvessel in the interstitium  
 1127 [114,200], indicating the BTB is more sensitive than the  
 1128 endothelial TJ-barrier in microvessels to cadmium toxicity.  
 1129 Subsequent analyses by immunohistochemistry and fluor-  
 1130 escent microscopy using these rats have conclusively  
 1131 demonstrated a significant loss of TJ- and AJ-integral



proteins from the BTB site, consistent with results of immunoblot analyses [52,114]. Furthermore, a loss of protein-protein interactions of the AJ integral membrane proteins and their adaptors namely cadherin-catenin and nectin-afadin was also detected when rats were exposed to Adjudin [68], a chemical known to induce germ cell loss from the seminiferous epithelium without disrupting the TJ-barrier at the BTB (for reviews, see [1,2,9]). Taking collectively, these data have clearly illustrated the vulnerability of the testis to environmental toxicants (e.g., cadmium) and that the proteins at the TJ and AJ sites are some of the primary targets of these toxicants. While the precise mechanism underlying such vulnerability is not fully understood, recent studies have shed new lights on this issue, which also highlights a unique opportunity to use these toxicant-induced BTB or AJ damage to the testis as novel models to study BTB dynamics, AJ restructuring pertinent to spermatogenesis, and their regulation. Furthermore, these studies can plausibly provide new insights in developing preventive measures to antagonize these toxicants.

First, Sertoli cells are secretory cells that actively provide virtually all the necessary nutrients for germ cell development behind the BTB including metal transporters, such as transferrin, ceruoplasmin, and metallothioneins (MTs). MTs are small Mr proteins having high affinities for heavy metal ions including cadmium, zinc, copper and mercury. MTs are produced in virtually all mammalian tissues in response to metal ions exposure, which can detoxify heavy metals, such as cadmium (for reviews, see [201,202–205]). MTs are also important to maintain the homeostasis of essential trace elements, such as zinc and copper and are scavengers of free radicals [202,204] and protect cells from the cytotoxic effects of cadmium [206]. In the rat testis, MTs, such as MT1 and MT2, have been identified and isolated [207,208]. Recent studies have found a novel testis-specific MT-like protein called tesmin which is specifically expressed by spermatogenic cells [209]. MT1 and MT2 are products of Sertoli and germ cells, which are significantly induced after cadmium exposure [210]. Yet the production of MTs by Sertoli and germ cells are significantly lower when compared to hepatocytes in vitro in response to cadmium exposure [210]. Indeed, the quiescence of MT expression in the ventral prostate and the testis is the possible cause of their susceptibility to cadmium cytotoxicity and carcinogenicity [204,211–213].

Second, recent studies on the effects of cadmium on different cell lines, including MDCK, LLC-PK1, and Caco-2 cells, have shown that its primary target is E-cadherin (for a review, see [214]). For instance, cells that were exposed to cadmium were found to have their E-cadherin moving away from the cell–cell interface and became diffusely localized in the cytoplasm. It was postulated that cadmium may be competing to the binding of calcium to the E-cadherin, thereby perturbing the AJ function [214–217]. If this is the case, cadmium (and possibly other environmental toxicants) must first gain access to AJ to disrupt E-cadherin. In all other

epithelia found in mammals, AJ is physically located behind the TJ since the TJ-barrier is located to the apical portion of the cell epithelium, and behind TJ lies desmosomes, which collectively known as the junctional complex. Behind the junctional complex are the gap junctions to be followed by the cell–matrix adhesion complex. As such, the TJ would seal most of the environmental toxicants off the epithelium in virtually all organs. Yet in the testis, TJ coexists with AJ and desmosome-like junctions at the BTB, which collectively lies adjacent to the basement membrane (a modified form of ECM, for a review, see [12]), closest to the interstitium. Thus, toxicants (e.g., cadmium) diffuses from the microvessels will have immediate access to the E-cadherin in the AJ (which is the cellular target of cadmium) at the BTB because there is no TJ-barrier that seals off cadmium. This, in turn, disrupts AJ, inducing germ cell loss from the epithelium as manifested by germ cell sloughing in many of these animal models using environmental toxicants. It is of interest to note that recent studies have begun to shed light on the physiological significance of such coexisting TJ and AJ at the BTB in relation to spermatogenesis. For instance, it is well understood that spermatogenesis is associated with extensive restructuring of Sertoli–Sertoli and Sertoli–germ cell interface because of the constant reshaping of germ cell shapes as a result of differentiation and germ cell movement from the basal to the adluminal compartment. If such AJ restructuring leads to TJ-barrier disruption as it is the case in other epithelia [193,195,197], the BTB integrity cannot be maintained, and haploid germ cell antigens cannot be sequestered from the host immune system; and such a disruption, even transiently, of the immunological barrier is detrimental to spermatogenesis. Thus, the fact that the BTB is constituted by co-existing AJ and TJ is to ensure such transient disruption of TJ during AJ restructuring in the seminiferous epithelium does not occur. Recent studies have shown that a signal that induces AJ disruption [e.g., via treatment of rats with Adjudin to induce extensive AJ restructuring that leads to germ cell loss from the epithelium, or a decline in endogenous intratesticular T level using androgen/estradiol transdermal implants] can lead to a surge in the production of both AJ (e.g., cadherins, catenins) and TJ (e.g., occludin, ZO-1) proteins [30,35,68] (Fig. 2). The increased TJ proteins are being used to reinforce the TJ-barrier integrity at the BTB at the time of extensive AJ restructuring. While the levels of AJ proteins are also induced, germ cells can still be dissociated from Sertoli cells because the AJ-integral membrane protein-AJ adaptor (e.g., the N-cadherin- $\beta$ -catenin protein complex) association is found to be weakened via an increase in tyrosine phosphorylation of  $\beta$ -catenin [30,35,68]. These findings are significant because it depicts the presence of a novel mechanism utilized by the testis to ensure TJ-barrier integrity while permitting AJ restructuring within a microenvironment such as the seminiferous epithelium. Fig. 2 is a schematic drawing that illustrates this novel mechanism of increasing AJ and TJ proteins, which is likely

regulated by cytokines [e.g., TGF- $\beta$ 3, TNF- $\alpha$  released from either Sertoli or germ cells [57,113,123,218]] via the ERK signaling pathway that maintains the BTB integrity while permitting AJ restructuring, facilitating germ cell movement across the seminiferous epithelium. However, this same mechanism that is physiologically necessary to facilitate germ cell movement while maintaining BTB integrity during spermatogenesis also makes the BTB extremely vulnerable to environmental toxicants because of the unusual exposure of AJ structural proteins (e.g., E-cadherin) to the toxicants (e.g., cadmium).

## 6. Conclusion and future perspectives

In all animal species, cell migration and junction remodeling are naturally occurring processes. For instance, there are three types of signals that control different aspects of *Drosophila* border cell migration: a global steroid-hormone signal to determine the timing, a highly localized cytokine signal to induce migration, and a growth factor to guide cells to their destination (for a review, see [219]). In the testis, FSH released from the pituitary and testosterone from Leydig cells may serve as the global regulatory factors. Cytokines that function as either paracrine or autocrine can in turn regulate localized signaling and processes. Several theories of germ cell movement during spermatogenesis have been proposed, and recently, we have put forth a junction restructuring theory in which cytokines, protease/protease inhibitors, cytoskeleton regulators and junctional complex proteins are all coordinated to facilitate germ cell movement in the epithelium [2]. Stem cell research has also offered new insights on Sertoli–germ cell interactions and may facilitate the research regarding the local regulatory function of cytokines in this event. When rat spermatogonia are transplanted into recipient mouse testes, the rat stem cells develop according to their only timing (~7 week instead of ~5 week), irrespective to the surrounding mouse spermatogenesis milieu [220]. It seems that this internal preprogrammed rhythm autonomously determines the fate of rat spermatogonia differentiation, and creates a suitable localized environment through dialogues with mouse Sertoli cells, probably via cytokines for crosstalk. This intrinsic preprogrammed timing may be controlled or executed by homeobox genes. A homeobox gene cluster *Rhox* (reproductive homeobox on the X chromosome) has recently been identified in mice [221]. The 12 *Rhox* genes are expressed mostly in reproductive organs (placenta, ovary, testis and epididymis), arranged into three subclusters and manifested temporal and quantitative colinearity in expression patterns [221]. In the testis, the majority of *Rhox* genes are primarily expressed in Sertoli cells and androgen responsive [221]. During the first wave of spermatogenesis, the timing of *Rhox* genes expression corresponds to the specific phases of germ cell differentiation. Hence these transcription factors may direct the expression of an array of proteins required for

germ cell development, and may also define the corresponding timing of epithelial cycle and length of spermatogenesis in rodents [221].

Several approaches can be used in future studies to aid the understanding of Sertoli–germ cell crosstalk and junction restructuring. First, development of testis-specific knockout mice against crucial proteins pertinent to junction restructuring and spermatogenesis to identify the function of these proteins in the testis. For many cytokines (e.g., TGF- $\beta$ s), their deletion can lead to lethality of the null mice. As such, their roles in spermatogenesis at adulthood cannot be examined. In the rat, the first wave of spermiation occurs only by 30–40 days of age. Recently conditional knockout technique has allowed investigators to elucidate protein function in a tissue- and time-specific manner in testis using specific Sertoli cell KOs, such as androgen receptor [222]. The generation of testis-specific KOs (e.g., TGF- $\beta$ 3) will help define the roles of these cytokines in junction dynamics at spermatogenesis. Second, germline stem cell transplantation with traceable markers to follow germ cell differentiation as well as junction remodeling during spermatogenesis can assist the study of cell–cell interactions pertinent to germ cell movement. For instance, when spermatogonia are transplanted into the recipient testis, they can migrate to the basal niche and initiate spermatogenesis in the preprogrammed cycle independent of the host environment. This migration must traverse the BTB, differing from gonocyte migration in the tubule when BTB has not yet formed. Third, using microarray technique to identify the expression profiles of various cytokines, proteases and protease inhibitors, junctional proteins, adaptors and transcription factors in staged tubules and in testes obtained from selected *in vivo* models. This approach can also pinpoint the leading and supporting biological factors pertinent to spermatogenesis. It is hopeful that using these approaches, a better understanding of spermatogenesis can emerge, which should be helpful for various applications such as treating male infertility or for contraception.

## Acknowledgements

Supported in part by grants from the National Institutes of Health (NICHD, 5U01 HD045908 to CYC; 5U54 HD029990, Project 3 to CYC), and the CONRAD Program (CICCR, C1G 01-72 to CYC, C1G 01-74 to DDM).

## References

- [1] Cheng CY, Mruk DD. Cell junction dynamics in the testis: Sertoli–germ cell interactions and male contraceptive development. *Physiol Rev* 2002;82:825–74.
- [2] Mruk DD, Cheng CY. Sertoli–Sertoli and Sertoli–germ cell interactions and their significance in germ cell movement in the seminiferous epithelium during spermatogenesis. *Endocr Rev* 2004;25:747–806.

- 1348 [3] de Kretser DM, Kerr JB. The cytology of the testis. In: Knobil E,  
1349 Neill JD, editors. The physiology of reproduction. New York: Raven  
1350 Press; 1994. p. 1177–300.
- 1351 [4] Parvinen M. Regulation of the seminiferous epithelium. *Endocr Rev*  
1352 1982;3:404–17.
- 1353 [5] Griswold M. Interactions between germ cells and Sertoli cells in the  
1354 testis. *Biol Reprod* 1995;52:211–6.
- 1355 [6] Kierszenbaum AL. Mammalian spermatogenesis in vivo and in vitro:  
1356 a partnership of spermatogenic and somatic cell lineages. *Endocr Rev*  
1357 1994;15:116–34.
- 1358 [7] Griswold MD. The central role of Sertoli cells in spermatogenesis.  
1359 *Semin Cell Dev Biol* 1998;9:411–6.
- 1360 [8] Russell LD. Movement of spermatocytes from the basal to the adluminal  
1361 compartment of the rat testis. *Am J Anat* 1977;148:313–28.
- 1362 [9] Mruk DD, Cheng CY. Cell–cell interactions at the ectoplasmic  
1363 specialization in the testis. *Trends Endocr Metab* 2004;15:439–47.
- 1364 [10] Russell L. Sertoli–germ cell interactions: a review. *Gamete Res*  
1365 1980;3:179–202.
- 1366 [11] Gnassi L, Fabbri A, Spera G. Gonadal peptides as mediators of  
1367 development and functional control of the testis: an integrated system  
1368 with hormones and local environment. *Endocr Rev* 1997;18:541–609.
- 1369 [12] Dym M. Basement membrane regulation of Sertoli cells. *Endocr Rev*  
1370 1994;15:102–15.
- 1371 [13] Weber JE, Russell LD, Wong V, Peterson RN. Three-dimensional  
1372 reconstruction of a rat stage V Sertoli cell. II. Morphometry of  
1373 Sertoli–Sertoli and Sertoli–germ-cell relationships. *Am J Anat*  
1374 1983;167:163–79.
- 1375 [14] Hess RA, Franca LR. Structure of the Sertoli cell. In: Skinner MK,  
1376 Griswold MD, editors. *Sertoli cell biology*. San Diego: Elsevier  
1377 Academic Press; 2005. p. 19–40.
- 1378 [15] Alberts B, Johnson A, Lewis J, Raff M, Roberts K, Watter P.  
1379 *Molecular Biology of the Cell*. New York: Garland Science, 2002.
- 1380 [16] Toyama Y, Maekawa M, Yuasa S. Ectoplasmic specializations in the  
1381 Sertoli cell: news vistas based on genetic defects and testicular  
1382 toxicology. *Anat Sci Int* 2003;78:1–16.
- 1383 [17] Lee NPY, Cheng CY. Ectoplasmic specialization, a testis-specific  
1384 cell–cell actin-based adherens junction type: is this a potential target  
1385 for male contraceptive development? *Hum Reprod Update*  
1386 2004;10:349–69.
- 1387 [18] Guttman JA, Takai Y, Vogl AW. Evidence that tubulobulbar com-  
1388 plexes in the seminiferous epithelium are involved with internaliza-  
1389 tion of adhesion junctions. *Biol Reprod* 2004;71:548–59.
- 1390 [19] Cunha-Vaz JG. The blood–retinal barriers system. Basic concepts  
1391 and clinical evaluation. *Exp Eye Res* 2004;78:715–21.
- 1392 [20] Rubin LL, Staddon JM. The cell biology of the blood–brain barrier.  
1393 *Annu Rev Neurosci* 1999;22:11–28.
- 1394 [21] Dejana E. Endothelial cell–cell junctions: happy together. *Nat Rev*  
1395 *Mol Cell Biol* 2004;5:261–70.
- 1396 [22] Siu MKY, Cheng CY. Dynamic cross-talk between cells and the  
1397 extracellular matrix in the testis. *BioEssays* 2004;26:978–92.
- 1398 [23] Tsukita S, Furuse M, Itoh M. Multifunctional strands in tight junc-  
1399 tions. *Nat Rev Mol Cell Biol* 2001;2:285–93.
- 1400 [24] Schneeberger EE, Lynch RD. The tight junction: a multifunctional  
1401 complex. *Am J Physiol Cell Physiol* 2004;286:C1213–28.
- 1402 [25] Bazzoni G. The JAM family of junctional adhesion molecules. *Curr*  
1403 *Opin Cell Biol* 2003;15:525–30.
- 1404 [26] Ebnet K, Suzuki A, Ohno S, Vestweber D. Junctional adhesion  
1405 molecules (JAMs): more molecules with dual functions? *J Cell*  
1406 *Sci* 2004;117:19–29.
- 1407 [27] Martin-Padura I, Lostaglio S, Schneemann M, Williams L, Romano  
1408 M, Fruscella P, et al. Junctional adhesion molecule, a novel member  
1409 of the immunoglobulin superfamily that distributes at intercellular  
1410 junctions and modulates monocyte transmigration. *J Cell Biol*  
1411 1998;142:117–27.
- 1412 [28] Hirabayashi S, Tajima M, Yao I, Nishimura W, Mori H, Hata Y.  
1413 JAM4, a junctional cell adhesion molecule interacting with a tight  
1414 junction protein, MAGI-1. *Mol Cell Biol* 2003;23:4267–82.
- [29] Gliki G, Ebnet K, Aurrand-Lions M, Lmhot BA, Adams RH. 1415  
Spermatid differentiation requires the assembly of a cell polarity 1416  
complex downstream of junctional adhesion molecule-C. *Nature* 1417  
2004;431:320–4. 1418
- [30] Xia W, Wong CH, Lee NPY, Lee WM, Cheng CY. Disruption of 1419  
Sertoli–germ cell adhesion function in the seminiferous epithelium of 1420  
the rat testis can be limited to adherens junctions without affecting 1421  
the blood–testis barrier integrity: an in vivo study using an androgen 1422  
suppression model. *J Cell Physiol*, in press. 1423
- [31] Cera MR, Del Prete A, Vecchi A, Corada M, Martin-Padura I, 1424  
Motoike T, et al. Increased DC trafficking to lymph nodes and 1425  
contact hypersensitivity in junctional adhesion molecule-A-deficient 1426  
mice. *J Clin Invest* 2004;114:729–38. 1427
- [32] Liang TW, Chiu HH, Gurney A, Sidle A, Tumas DB, Schow P, et al. 1428  
Vascular endothelial-junctional adhesion molecule (VE-JAM)/JAM 1429  
2 interacts with T, NK, and dendritic cells through JAM 3. *J Immunol* 1430  
2002;168:1618–26. 1431
- [33] McLachlan RI, Wreford NG, Meachem SJ, Kretser DMD, Robertson 1432  
DM. Effects of testosterone on spermatogenic cell populations in the 1433  
adult rat. *Biol Reprod* 1994;51:945–55. 1434
- [34] O'Donnell L, McLachlan RI, Wreford NG, de Kretser DM, Robert- 1435  
son DM. Testosterone withdrawal promotes stage-specific detach- 1436  
ment of round spermatids from the at seminiferous epithelium. *Biol* 1437  
*Reprod* 1996;55:895–901. 1438
- [35] Zhang J, Wong CH, Xia W, Mruk DD, Lee NPY, Lee WM, et al. 1439  
Regulation of Sertoli–germ cell adherens junction dynamics via 1440  
changes in protein–protein interactions of the N-cadherin- $\beta$ -catenin 1441  
protein complex which are possibly mediated by c-Src and MTMR2: 1442  
an in vivo study using an androgen suppression model. *Endocrinol-* 1443  
*ogy* 2005;146:1268–84. 1444
- [36] Wong CH, Xia W, Lee NPY, Mruk DD, Lee WM, Cheng CY. 1445  
Regulation of ectoplasmic specialization dynamics in the semini- 1446  
ferous epithelium by focal adhesion-associated proteins in 1447  
testosterone-suppressed rat testes. *Endocrinology* 2005;146: 1448  
1192–204. 1449
- [37] Siu MKY, Mruk DD, Lee WM, Cheng CY. Adhering junction 1450  
dynamics in the testis are regulated by an interplay of  $\beta$ 1-integrin 1451  
and the focal adhesion complex-associated proteins. *Endocrinology* 1452  
2003;144:2141–63. 1453
- [38] Shaw SK, Perkins BN, Lim YC, Liu Y, Nusrat A, Schnell FJ, et al. 1454  
Reduced expression of junctional adhesion molecule and platelet/ 1455  
endothelial cell adhesion molecule-1 (CD31) at human vascular 1456  
endothelial junctions by cytokines tumor necrosis factor- $\alpha$  plus 1457  
interferon-gamma does not reduce leukocyte transmigration under 1458  
flow. *Am J Pathol* 2001;159:2281–91. 1459
- [39] Turksen K, Troy T-C. Barriers built on claudins. *J Cell Sci* 2004; 1460  
117:2435–47. 1461
- [40] Tsukita S, Furuse M. The structure and function of claudins, cell 1462  
adhesion molecules at tight junctions. *Ann N Y Acad Sci* 1463  
2000;915:129–35. 1464
- [41] Heiskala M, Peterson PA, Yang Y. The roles of claudin superfamily 1465  
proteins in paracellular transport. *Traffic* 2001;2:92–8. 1466
- [42] Florin A, Maire M, Bozec A, Hellani A, Chater S, Bars R, 1467  
et al. Androgens and postmeiotic germ cells regulate claudin- 1468  
11 expression in rat Sertoli cells. *Endocrinology* 2005;146:1532– 1469  
40. 1470
- [43] Hellani A, Ji J, Mauduit C, Deschildre C, Tabone E, Benahmed M. 1471  
Developmental and hormonal regulation of the expression of oligo- 1472  
dendrocyte-specific protein/claudin 11 in mouse testis. *Endocrinol-* 1473  
*ogy* 2000;141:3012–9. 1474
- [44] Lui WY, Lee WM, Cheng CY. Transforming growth factor- $\beta$ 3 1475  
perturbs the inter-Sertoli tight junction permeability barrier in vitro 1476  
possibly mediated via its effects on occludin, zonula occludens-1, 1477  
and claudin-11. *Endocrinology* 2001;142:1865–77. 1478
- [45] Morita K, Sasaki H, Fujimoto K, Furuse M, Tsukita S. Claudin- 1479  
11/OSP-based tight junctions of myelin sheaths in brain and Sertoli 1480  
cells in testis. *J Cell Biol* 1999;145:579–88. 1481

- 1482 [46] Gow A, Southwood CM, Li JS, Pariali M, Riordan GP, Brodie SE, et al. CNS myelin and Sertoli cell tight junction strands are absent in *Osp/claudin-11* null mice. *Cell* 1999;99:649–59. 1549
- 1483 [47] Gye MC. Changes in the expression of claudins and transepithelial 1550
- 1484 electrical resistance of mouse Sertoli cells by Leydig cell coculture. 1551
- 1485 *Int J Androl* 2003;26:271–8. 1552
- 1486 [48] Kitajiri S-I, Miyamoto T, Mineharu A, Sonoda N, Furuse K, Hata M, 1553
- 1487 et al. Compartmentalization established by claudin-11-based tight 1554
- 1488 junctions in stria vascularis is required for hearing through generation 1555
- 1489 of endocochlear potential. *J Cell Sci* 2004;117:5087–96. 1556
- 1490 [49] Sasaki H, Matsui C, Furuse K, Mimori-Kiyosue Y, Furuse M, Tsukita 1557
- 1491 S. Dynamic behavior of paired claudin strands within apposing 1558
- 1492 plasma membranes. *Proc Natl Acad Sci USA* 2003;100:3971–6. 1559
- 1493 [50] Matsuda M, Kubo A, Furuse M, Tsukita S. A peculiar internalization 1560
- 1494 of claudins, tight junction-specific adhesion molecules, during the 1561
- 1495 intercellular movement of epithelial cells. *J Cell Sci* 2004;117:1247– 1562
- 1496 57. 1563
- 1497 [51] Furuse M, Hirase T, Itoh M, Nagafuchi A, Yonemura S, Tsukita S. 1564
- 1498 Occludin: a novel integral membrane protein localizing at tight 1565
- 1499 junctions. *J Cell Biol* 1993;123:1777–88. 1566
- 1500 [52] Wong CH, Mruk DD, Lui WY, Cheng CY. Regulation of blood-testis 1567
- 1501 barrier dynamics: an in vivo study. *J Cell Sci* 2004;117:783–98. 1568
- 1502 [53] Lui WY, Lee WM. cAMP perturbs inter-Sertoli tight junction 1569
- 1503 permeability barrier in vitro via its effect on proteasome-sensitive 1570
- 1504 ubiquitination of occludin. *J Cell Physiol* 2005;203:564–72. 1571
- 1505 [54] Coyne CB, Vanhook MK, Gambling TM, Carson JL, Boucher RC, 1572
- 1506 Johnson LG. Regulation of airway tight junctions by proinflammatory 1573
- 1507 cytokines. *Mol Biol Cell* 2002;13:3218–34. 1574
- 1508 [55] Mankertz J, Tavalali S, Schmitz H, Mankertz A, Riecken EO, Fromm 1575
- 1509 M, et al. Expression from the human occludin promoter is affected by 1576
- 1510 tumor necrosis factor alpha and interferon gamma. *J Cell Sci* 1577
- 1511 2000;113:2085–90. 1578
- 1512 [56] Yamamoto T, Kojima T, Murata M, Takano K, Go M, Chiba H, et al. 1579
- 1513 IL-1beta regulates expression of Cx32, occludin, and claudin-2 of rat 1580
- 1514 hepatocytes via distinct signal transduction pathways. *Exp Cell Res* 1581
- 1515 2004;299:427–41. 1582
- 1516 [57] Siu MKY, Lee WM, Cheng CY. The interplay of collagen IV, tumor 1583
- 1517 necrosis factor- $\alpha$ , gelatinase B (matrix metalloproteinase-9), and tissue 1584
- 1518 inhibitor of metalloproteinases-1 in the basal lamina regulates Sertoli 1585
- 1519 cell-tight junction dynamics in the rat testis. *Endocrinology* 2003; 1586
- 1520 144:371–87. 1587
- 1521 [58] Mealy K, Robinson B, Millette CF, Majzoub J, Wilmore DW. The 1588
- 1522 testicular effects of tumor necrosis factor. *Ann Surg* 1990;211:470–5. 1589
- 1523 [59] Wheelock MJ, Johnson KR. Cadherins as modulators of cellular 1590
- 1524 phenotype. *Annu Rev Cell Dev Biol* 2003;19:207–35. 1591
- 1525 [60] Yagi T, Takeichi M. Cadherin superfamily genes: functions, genomic 1592
- 1526 organization, and neurologic diversity. *Genes Dev* 2000;14:1169–80. 1593
- 1527 [61] Gooding JM, Yap KL, Ikura M. The cadherin-catenin complex as a 1594
- 1528 focal point of cell adhesion and signalling: new insights from three- 1595
- 1529 dimensional structures. *BioEssays* 2004;26:497–511. 1596
- 1530 [62] Angst BD, Marozzi C, Magee AI. The cadherin superfamily: 1597
- 1531 diversity in form and function. *J Cell Sci* 2001;114:629–41. 1598
- 1532 [63] Johnson KJ, Patel SR, Boekelheide K. Multiple cadherin superfamily 1599
- 1533 members with unique expression profiles are produced in rat testis. 1600
- 1534 *Endocrinology* 2000;141:675–83. 1601
- 1535 [64] Peinado H, Portillo F, Cano A. Transcriptional regulation of cadherins 1602
- 1536 during development and carcinogenesis. *Int J Dev Biol* 1603
- 1537 2004;48:365–75. 1604
- 1538 [65] Johnson KJ, Boekelheide K. Dynamic testicular adhesion junctions 1605
- 1539 are immunologically unique. II. Localization of classic cadherins in 1606
- 1540 rat testis. *Biol Reprod* 2002;66:992–1000. 1607
- 1541 [66] Lee NPY, Mruk D, Lee WM, Cheng CY. Is the cadherin/catenin 1608
- 1542 complex a functional unit of cell–cell actin-based adherens junctions 1609
- 1543 in the rat testis? *Biol Reprod* 2003;68:489–508. 1610
- 1544 [67] Lee NPY, Mruk DD, Conway AM, Cheng CY. Zyxin, axin, and 1611
- 1545 Wiskott-Aldrich syndrome protein are adaptors that link the cadherin/catenin protein complex to the cytoskeleton at adherens junctions in the seminiferous epithelium of the rat testis. *J Androl* 2004;25:200–15. 1612
- 1546 [68] Xia W, Cheng CY. TGF- $\beta$ 3 regulates anchoring junction dynamics 1613
- 1547 in the seminiferous epithelium of the rat testis via the Ras/ 1614
- 1548 ERK signaling pathway: an in vivo study. *Dev Biol* 2005;280: 1615
- 321–43. 1616
- [69] Wine RN, Chapin RE. Adhesion and signaling proteins spatiotemporally associated with spermiation in the rat. *J Androl* 1999;20:198–213. 1617
- [70] Johnson KJ, Zecevic A, Kwon EJ. Protocadherin  $\alpha$ 3 acts at sites distinct from classic cadherins in rat testis and sperm. *Biol Reprod* 2004;70:303–12. 1618
- [71] Wheelock MJ, Johnson KR. Cadherin-mediated cellular signaling. *Curr Opin Cell Biol* 2003;15:509–14. 1619
- [72] Perez-Moreno M, Jamora C, Fuchs E. Sticky business: orchestrating cellular signals at adherens junctions. *Cell* 2003;112:535–48. 1620
- [73] Daniel JM, Reynolds AB. Tyrosine phosphorylation and cadherin/catenin function. *BioEssays* 1997;19:883–91. 1621
- [74] Lee NPY, Cheng CY. Protein kinases and adherens junction dynamics in the seminiferous epithelium of the rat testis. *J Cell Physiol* 2005;202:344–60. 1622
- [75] Zhang J, Mruk DD, Cheng CY. Myotubularin phosphoinositide phosphatases, proteins phosphatases, and kinases: their roles in junction dynamics and spermatogenesis. *J Cell Physiol*, in press. 1623
- [76] Lui WY, Mruk D, Cheng CY. Interactions among IQGAP1, Cdc42, and the cadherin/catenin protein complex regulate Sertoli–germ cell adherens junction dynamics in the testis. *J Cell Physiol* 2005;202:49–66. 1624
- [77] Hazan RB, Qiao R, Keren R, Badano I, Suyama K. Cadherin switch in tumor progression. *Ann N Y Acad Sci* 2004;1014:155–63. 1625
- [78] Derycke LDM, Bracke ME. N-cadherin in the spotlight of cell–cell adhesion, differentiation, embryogenesis, invasion and signaling. *Int J Dev Biol* 2004;48:463–76. 1626
- [79] Takai Y, Nakanishi H. Nectin and afadin: novel organizers of intercellular junctions. *J Cell Sci* 2003;116:17–27. 1627
- [80] Satoh-Horikawa K, Nakanishi H, Takahashi K, Miyahara M, Nishimura M, Tachibana K, et al. Nectin-3, a new member of immunoglobulin-like cell adhesion molecules that shows homophilic and heterophilic cell–cell adhesion activities. *J Biol Chem* 2000;275:10291–9. 1628
- [81] Reymond N, Fabre S, Lecocq E, Adelaide J, Dubreuil P, Lopez M. Nectin4/PRR4, a new afadin-associated member of the nectin family that trans-interacts with nectin1/PRR1 through V domain interaction. *J Biol Chem* 2001;276:43205–15. 1629
- [82] Takahashi K, Nakanishi H, Miyahara M, Mandai K, Satoh K, Satoh A, et al. Nectin/PRR: an immunoglobulin-like cell Adhesion molecule recruited to cadherin-based adherens junctions through interaction with afadin, a PDZ domain-containing protein. *J Cell Biol* 1999;145:539–49. 1630
- [83] Ozaki-Kuroda K, Nakanishi H, Ohta H, Tanaka H, Kurihara H, Mueller S, et al. Nectin couples cell–cell adhesion and the actin scaffold at heterotypic testicular junctions. *Curr Biol* 2002;12:1145–50. 1631
- [84] Bouchard MJ, Dong Jr Y, McDermott Jr BM, Lam D-H, Brown KR, Shelanski M, et al. Defects in nuclear and cytoskeletal morphology and mitochondrial localization in spermatozoa of mice lacking nectin-2, a component of cell–cell adherens junctions. *Mol Cell Biol* 2000;20:2865–73. 1632
- [85] Mueller S, Rosenquist TA, Takai Y, Bronson RA, Wimmer E. Loss of nectin-2 at Sertoli–spermatid junctions leads to male infertility and correlates with severe spermatozoan head and midpiece malformation, impaired binding to the zona pellucida, and oocyte penetration. *Biol Reprod* 2003;69:1330–40. 1633
- [86] Fukuhara T, Shimizu K, Kawakatsu T, Fukuyama T, Minami Y, Honda T, et al. Activation of Cdc42 by trans interactions of the cell adhesion molecules nectins through c-Src and Cdc42-GEF FRG. *J Cell Biol* 2004;166:393–405. 1634

- 1616 [87] Takai Y, Irie K, Shimizu K, Sakisaka T, Ikeda W. Nectins and nectin-like molecules: roles in cell adhesion, migration, and polarization. *Cancer Sci* 2003;94:655–67. 1684
- 1617 1685
- 1618 [88] Sakisaka T, Takai Y. Biology and pathology of nectins and nectin-like molecules. *Curr Opin Cell Biol* 2004;16:513–21. 1686
- 1619 1687
- 1620 [89] Irie K, Shimizu K, Sakisaka T, Ikeda W, Takai Y. Roles and modes of action of nectins in cell–cell adhesion. *Semin Cell Dev Biol* 2004;15:643–56. 1688
- 1621 1689
- 1622 [90] Asada M, Irie K, Morimoto K, Yamada A, Ikeda W, Takeuchi M, et al. ADIP, a novel afadin- and  $\alpha$ -actinin-binding protein localized at cell–cell adherens junctions. *J Biol Chem* 2003;278:4103–11. 1690
- 1623 1691
- 1624 [91] Tachibana K, Nakanishi H, Mandai K, Ozaki K, Ikeda W, Yamamoto Y, et al. Two cell adhesion molecules, nectin and cadherin, interact through their cytoplasmic domain-associated proteins. *J Cell Biol* 2000;150:1161–76. 1692
- 1625 1693
- 1626 [92] Ooshio T, Irie K, Morimoto K, Fukuhara A, Imai T, Takai Y. Involvement of LMO7 in the association of two cell–cell adhesion molecules, nectin and E-cadherin, through afadin and  $\alpha$ -actinin in epithelial cells. *J Biol Chem* 2004;279:31365–73. 1694
- 1627 1695
- 1628 [93] Shingai T, Ikeda W, Kakunaga S, Morimoto K, Takekuni K, Itoh S, et al. Implications of nectin-like molecule-2/IGSF4/RA175/SgIGSF/TSLC1/SynCAM1 in cell–cell adhesion and transmembrane protein localization in epithelial cells. *J Biol Chem* 2003;278:35421–7. 1696
- 1629 1697
- 1630 [94] Siu MKY, Cheng CY. Interactions of proteases, protease inhibitors, and the  $\beta 1$  integrin/laminin  $\gamma 3$  protein complex in the regulation of ectoplasmic specialization dynamics in the rat testis. *Biol Reprod* 2004;70:945–64. 1698
- 1631 1699
- 1632 [95] Hynes RO. Integrins: bidirectional, allosteric signaling machines. *Cell* 2002;110:673–87. 1700
- 1633 1701
- 1634 [96] Jin H, Varner J. Integrins: roles in cancer development and as treatment targets. *Br J Cancer* 2004;90:561–5. 1702
- 1635 1703
- 1636 [97] Siu MKY, Cheng CY. Extracellular matrix: recent advances on its role in junction dynamics in the seminiferous epithelium during spermatogenesis. *Biol Reprod* 2004;375–91. 1704
- 1637 1705
- 1638 [98] Vilcek J. The cytokines: an overview. In: Thomson AW, Lotze MT, editors. *The cytokine handbook*. Academic Press; 2003. p. 3–18. 1706
- 1639 1707
- 1640 [99] Mruk DD, Cheng CY. Sertoli cell proteins in testicular paracrine. In: Jegou B, Pineau C, Saez J, editors. *Testis, epididymis and technologies in the year 2000*. Berlin: Springer-Verlag; 2000. p. 197–228. 1708
- 1641 1709
- 1642 [100] Bardin CW, Cheng CY, Mustow NA, Gunsalus GL. The Sertoli cell. In: Knobil E, Neill JD, editors. *The physiology of reproduction*. New York: Raven Press; 1994. p. 1291–332. 1710
- 1643 1711
- 1644 [101] Wang H, Czura CJ, Tracey KJ. Tumor necrosis factor. In: Thomson AW, Lotze MT, editors. *The cytokine handbook*. Academic Press; 2003. p. 837–60. 1712
- 1645 1713
- 1646 [102] Mocellin S, Rossi CR, Pilati P, Nitti D. Tumor necrosis factor, cancer and anticancer therapy. *Cytokine Growth Factor Rev* 2005;16:35–53. 1714
- 1647 1715
- 1648 [103] Aggarwal BB. Signalling pathways of the TNF superfamily: a double-edged sword. *Nat Rev Immunol* 2003;3:745–56. 1716
- 1649 1717
- 1650 [104] Soond SM, Terry JL, Colbert JD, Riches DWH. TRUSS, a novel tumor necrosis factor receptor 1 scaffolding protein that mediates activation of the transcription factor NF- $\kappa$ B. *Mol Cell Biol* 2003;23:8334–44. 1718
- 1651 1719
- 1652 [105] Gibson PR. Increased gut permeability in Crohn's disease: is TNF the link? *Gut* 2004;53:1724–5. 1720
- 1653 1721
- 1654 [106] Hong CY, Park JH, Ahn RS, Im SY, Choi H-S, Soh J, et al. Molecular mechanism of suppression of testicular steroidogenesis by proinflammatory cytokine tumor necrosis factor  $\alpha$ . *Mol Cell Biol* 2004;24:2593–604. 1722
- 1655 1723
- 1656 [107] Pentikainen V, Erkkila K, Suomalainen L, Ojala M, Pentikainen MO, Parvinen M, et al. TNF $\alpha$  down-regulates the Fas ligand and inhibits germ cell apoptosis in the human testis. *J Clin Endocrinol Metab* 2001;86:4480–8. 1724
- 1657 1725
- 1658 [108] Morales V, Santana P, Diaz R, Tabraue C, Gallardo G, Blanco FL, et al. Intratesticular delivery of tumor necrosis factor- $\alpha$  and ceramide directly abrogates steroidogenic acute regulatory protein expression and Leydig cell steroidogenesis in adult rats. *Endocrinology* 2003; 144:4763–72. 1726
- 1659 1727
- 1660 [109] Chang H, Brown CW, Matzuk MM. Genetic analysis of the mammalian transforming growth factor- $\beta$  superfamily. *Endocr Rev* 2002;23:787–823. 1728
- 1661 1729
- 1662 [110] Shi Y, Massague J. Mechanisms of TGF- $\beta$  signaling from cell membrane to the nucleus. *Cell* 2003;113:685–700. 1730
- 1663 1731
- 1664 [111] Behringer RR, Finegold MJ, Cate RL. Mullerian-inhibiting substance function during mammalian sexual development. *Cell* 1994;79:415–25. 1732
- 1665 1733
- 1666 [112] Ingman WV, Robertson SA. Defining the action of transforming growth factor  $\beta$  in reproduction. *BioEssays* 2002;24:904–14. 1734
- 1667 1735
- 1668 [113] Lui WY, Wong CH, Mruk DD, Cheng CY. TGF- $\beta 3$  regulates the blood-testis barrier dynamics via the p38 mitogen activated protein (MAP) kinase pathway: an in vivo study. *Endocrinology* 2003;144: 1139–42. 1736
- 1669 1737
- 1670 [114] Wong CH, Mruk DD, Siu MKY, Cheng CY. Blood-testis barrier dynamics are regulated by  $\alpha 2$ -macroglobulin via the c-Jun N-terminal protein kinase pathway. *Endocrinology* 2005;146:1893–908. 1738
- 1671 1739
- 1672 [115] Wakefield LM, Roberts AB. TGF- $\beta$  signaling: positive and negative effects on tumorigenesis. *Curr Opin Genet Dev* 2002;12:22–9. 1740
- 1673 1741
- 1674 [116] Massague J. How cells read TGF- $\beta$  signals. *Nat Rev Mol Cell Biol* 2000;1:169–78. 1742
- 1675 1743
- 1676 [117] Leask A, Abraham DJ. TGF- $\beta$  signaling and the fibrotic response. *FASEB J* 2004;18:816–27. 1744
- 1677 1745
- 1678 [118] Derynck R, Zhang YE. Smad-dependent and Smad-independent pathways in TGF- $\beta$  family signaling. *Nature* 2003;425:577–84. 1746
- 1679 1747
- 1680 [119] Lui WY, Lee WM, Cheng CY. TGF- $\beta$ s: their role in testicular function and Sertoli cell tight junction dynamics. *Int J Androl* 2003;26:1–14. 1748
- 1681 1749
- 1682 [120] Howe PH. Transforming growth factor  $\beta$ . In: Thomson AW, Lotze MT, editors. *The cytokine handbook*. Academic Press; 2003. p. 1119–52. 1750
- 1683 1751
- 1684 [121] Roberts AB, Kim SJ, Noma T, Glick AB, Lafyatis R, Lechleider R, et al. Multiple forms of TGF- $\beta$ : distinct promoters and differential expression. *Ciba Found Symp* 1991;157:7–15. 1752
- 1685 1753
- 1686 [122] Miller DA, Pelton RW, Derynck R, Moses HL. Transforming growth factor- $\beta$ -a family of growth regulatory peptides. *Ann N Y Acad Sci* 1990;593:209–17. 1754
- 1687 1755
- 1688 [123] Mullaney BP, Skinner MK. Transforming growth factor- $\beta$  ( $\beta 1$ ,  $\beta 2$ , and  $\beta 3$ ) gene expression and action during pubertal development of the seminiferous tubule: potential role at the onset of spermatogenesis. *Mol Endocrinol* 1993;7:67–76. 1756
- 1689 1757
- 1690 [124] Massague J. TGF- $\beta$  signal transduction. *Annu Rev Biochem* 1998;67:753–91. 1758
- 1691 1759
- 1692 [125] Massague J, Chen Y-G. Controlling TGF- $\beta$  signaling. *Genes Dev* 2000;14:627–44. 1760
- 1693 1761
- 1694 [126] Xu J, Beyer AR, Walker WH, McGee EA. Developmental and stage-specific expression of Smad2 and Smad3 in rat testis. *J Androl* 2003;24:192–200. 1762
- 1695 1763
- 1696 [127] Luukko K, Ylikorkala A, Makela TP. Developmentally regulated expression of Smad3, Smad4, Smad6, and Smad7 involved in TGF- $\beta$  signaling. *Mech Dev* 2001;101:209–12. 1764
- 1697 1765
- 1698 [128] Yingling JM, Blanchard KL, Sawyer JS. Development of TGF- $\beta$  signalling inhibitors for cancer therapy. *Nat Rev Drug Disc* 2004;3:1011–22. 1766
- 1699 1767
- 1700 [129] Widmann C, Gibson S, Jarpe MB, Johnson GL. Mitogen-activated protein kinases: conservation of a three-kinase module from yeast to human. *Physiol Rev* 1999;79:143–80. 1768
- 1701 1769
- 1702 [130] Peason G, Robinson F, Bibson TB, Xu B-E, Karandikar M, Berman K, et al. Mitogen-activated protein (MAP) kinase pathways: regulation and physiological functions. *Endocr Rev* 2001;22:153–83. 1770
- 1703 1771
- 1704 [131] Kyriakis JM, Avruch J. Mammalian mitogen-activated protein kinase signal transduction pathways activated by stress and inflammation. *Physiol Rev* 2001;81:807–69. 1772
- 1705 1773
- 1706 [132] Lui WY, Lee WM, Cheng CY. Transforming growth factor- $\beta 3$  regulates the dynamics of Sertoli cell tight junctions via the p38

- mitogen-activated protein kinase pathway. *Biol Reprod* 2003;68:1597–612.
- [133] Chapin RE, Wine RN, Harris MW, Borchers CH, Haseman JK. Structure and control of a cell–cell adhesion complex associated with spermiation in rat seminiferous epithelium. *J Androl* 2001;22:1030–52.
- [134] Barrios-Rodiles M, Brown KR, Ozdamar B, Bose R, Liu Z, Donovan RS, et al. High-throughput mapping of a dynamic signaling network in mammalian cells. *Science* 2005;307:1621–5.
- [135] Ozdamar B, Bose R, Barrios-Rodiles M, Wang H-R, Zhang Y, Wrana JL. Regulation of the polarity protein Par6 by TGF $\beta$  receptors controls epithelial cell plasticity. *Science* 2005;307:1603–9.
- [136] Wang H-R, Zhang Y, Ozdamar B, Ogunjimi AA, Alexandrova E, Thomsen GH, et al. Regulation of cell polarity and protrusion formation by targeting RhoA for degradation. *Science* 2003;302:1775–9.
- [137] Hassel S, Eichner A, Yakymovych M, Hellman U, Knaus P, Souchelnyskiy S. Proteins associated with type II bone morphogenetic protein receptor (BMPRII) and identified by two-dimensional gel electrophoresis and mass spectrometry. *Proteomics* 2004;4:1346–58.
- [138] Behzadian MA, Wang X-L, Windsor LJ, Ghaly N, Caldwell RB. TGF- $\beta$  increases retinal endothelial cell permeability by increasing MMP-9: possible role of glial cells in endothelial barrier function. *Invest Ophthalmol Vis Sci* 2001;42:853–9.
- [139] Hurst IV V, Goldberg PL, Minnear FL, Heimark RL, Vincent PA. Rearrangement of adherens junctions by transforming growth factor- $\beta$ 1: role of contraction. *Am J Physiol Lung Cell Mol Physiol* 1999;276:L582–95.
- [140] Fujimoto K, Sheng H, Shao J, Beauchamp RD. Transforming growth factor- $\beta$ 1 promotes invasiveness after cellular transformation with activated Ras in intestinal epithelial cells. *Exp Cell Res* 2001;266:239–49.
- [141] Tian YC, Fraser D, Attisano L, Phillips AO. TGF- $\beta$ 1-mediated alterations of renal proximal tubular epithelial cell phenotype. *Am J Physiol Renal Physiol* 2003;285:F130–42.
- [142] Watabe T, Nishihara A, Mishima K, Yamashita J, Shimizu K, Miyazawa K, et al. TGF- $\beta$  receptor kinase inhibitor enhances growth and integrity of embryonic stem cell-derived endothelial cells. *J Cell Biol* 2003;163:1303–11.
- [143] Sheppard D. Roles of  $\alpha$ v integrins in vascular biology and pulmonary pathology. *Curr Opin Cell Biol* 2004;16:552–7.
- [144] Xu L, Massague J. Nucleocytoplasmic shuttling of signal transducers. *Nat Rev Mol Cell Biol* 2004;5:1–11.
- [145] Keski-Oja J, Koli K, von Melchner H. TGF- $\beta$  activation by traction? *Trends Cell Biol* 2004;14:657–9.
- [146] Gumienny TL, Padgett RW. The other side of TGF- $\beta$  superfamily signal regulation: thinking outside the cell. *Trends Endocr Metab* 2002;13:295–9.
- [147] Pardoux C, Derynck R. JNK regulates expression and autocrine signaling of TGF- $\beta$ 1. *Mol Cell* 2004;15:170–1.
- [148] Ventura J-J, Kennedy NJ, Flavell RA, Davis RJ. JNK regulates autocrine expression of TGF- $\beta$ 1. *Mol Cell* 2004;15:269–78.
- [149] Annes JP, Munger JS, Rifkin DB. Making sense of latent TGF $\beta$  activation. *J Cell Sci* 2003;116:217–24.
- [150] Rifkin DB. Latent transforming growth factor- $\beta$  (TGF- $\beta$ ) binding proteins: orchestrators of TGF- $\beta$  availability. *J Biol Chem* 2005;280:7409–12.
- [151] Piek E, Heldin C-H, ten Dijke P. Specificity, diversity, and regulation in TGF- $\beta$  superfamily signaling. *FASEB J* 1999;13:2105–24.
- [152] Lebrin F, Deckers M, Bertolino P, ten Dijke P. TGF- $\beta$  receptor function in the endothelium. *Cardiovasc Res* 2005;65:599–608.
- [153] Ge B, Gram H, Di Padova F, Huang B, New L, Ulevitch RJ, et al. MAPKK-independent activation of p38 $\alpha$  mediated by TAB1-dependent autophosphorylation of p38 $\alpha$ . *Science* 2002;295:1291–4.
- [154] Schiffer M, Mundel P, Shaw AS, Bottinger EP. A novel role for the adaptor molecule CD2-associated protein in transforming growth factor- $\beta$ -induced apoptosis. *J Biol Chem* 2004;279:37004–12.
- [155] Schiffer M, Mundel P, Shaw AS, Bottinger EP. A novel role for the adaptor molecule CD2-associated protein in transforming growth factor- $\beta$ -induced apoptosis. *J Biol Chem* 2004;279:37004–12.
- [156] Razani B, Zhang XL, Bitzer M, von Gersdorff G, Bottinger EP, Lisanti MP. Caveolin-1 regulates transforming growth factor (TGF)- $\beta$ /Smad signaling through an interaction with the TGF- $\beta$  type I receptor. *J Biol Chem* 2001;276:6727–38.
- [157] Schwarz K, Simons M, Reiser J, Saleem MA, Faul C, Kriz W, et al. Podocin, a raft-associated component of the glomerular slit diaphragm, interacts with CD2AP and nephrin. *J Clin Invest* 2001;108:1621–9.
- [158] Dikic I. CIN85/CMS family of adaptor molecules. *FEBS Lett* 2002;529:110–5.
- [159] Kirsch KH, Georgescu M-M, Ishimaru S, Hanafusa H. CMS: An adapter molecule involved in cytoskeletal rearrangements. *Proc Natl Acad Sci USA* 1999;96:6211–6.
- [160] Webb DJ, Donais K, Whitmore LA, Thomas SM, Turner CE, Parsons JT, et al. FAK-Src signalling through paxillin, ERK and MLCK regulates adhesion disassembly. *Nat Cell Biol* 2004;6:154–61.
- [161] Lui WY, Mruk D, Lee WM, Cheng CY. Sertoli cell tight junction dynamics: their regulation during spermatogenesis. *Biol Reprod* 2003;68:1087–97.
- [162] Lui WY, Mruk DD, Lee WM, Cheng CY. Adherens junction dynamics in the testis and spermatogenesis. *J Androl* 2003;24:1–14.
- [163] Grima J, Silvestrini B, Cheng CY. Reversible inhibition of spermatogenesis in rats using a new male contraceptive, 1-(2,4-dichlorobenzyl)-indazole-3-carbohydrazide. *Biol Reprod* 2001;64:1500–8.
- [164] Cheng CY, Silvestrini B, Grima J, Mo M-Y, Zhu L-J, Jahansson E, et al. Two new male contraceptives exert their effects by depleting germ cells prematurely from the testis. *Biol Reprod* 2001;65:449–61.
- [165] Grima J, Wong CCS, Zhu L-J, Zong S-D, Cheng CY. Testin secreted by Sertoli cells is associated with the cell surface, and its expression correlates with the disruption of Sertoli–germ cell junctions but not the inter-Sertoli tight junction. *J Biol Chem* 1998;273:21040–53.
- [166] Grima J, Cheng CY. Testin induction: the role of cyclic 3',5'-adenosine monophosphate/protein kinase A signaling in the regulation of basal and lomidamine-induced testin expression by rat Sertoli cells. *Biol Reprod* 2000;63:1648–60.
- [167] Lui WY, Lee WM, Cheng CY. Sertoli–germ cell adherens junction dynamics in the testis are regulated by RhoG GTPase via the ROCK/LIMK signaling pathway. *Biol Reprod* 2003;68:2189–206.
- [168] Chen Y-M, Lee NPY, Mruk DD, Lee WM, Cheng CY. Fer kinase/FerT and adherens junction dynamics in the testis: an in vitro and in vivo study. *Biol Reprod* 2003;69:656–72.
- [169] Werner S, Grose R. Regulation of wound healing by growth factors and cytokines. *Physiol Rev* 2003;83:835–70.
- [170] Sahai E. Mechanisms of cancer cell invasion. *Curr Opin Genet Dev* 2005;15:87–96.
- [171] Zalups RK, Ahmad S. Molecular handling of cadmium in transporting epithelia. *Toxicol Appl Pharmac* 2003;186:163–88.
- [172] Waisberg M, Joseph P, Hale B, Beyersmann D. Molecular and cellular mechanisms of cadmium carcinogenesis. *Toxicology* 2003;192:95–117.
- [173] Henson MC, Chedrese PJ. Endocrine disruption by cadmium, a common environmental toxicant with paradoxical effects on reproduction. *Expt Biol Med* 2004;229:383–92.
- [174] Johnson MD, Kenney N, Stoica A, Hlaskivi-Clarke L, Singh B, Chepko G, et al. Cadmium mimics the in vivo effects of estrogen in the uterus and mammary gland. *Nat Med* 2003;9:1081–4.
- [175] Jin YH, Clark AB, Slebos RJC, Al-Refai H, Taylor JA, Kunkel TA, et al. Cadmium is a mutagen that acts by inhibiting mismatch repair. *Nat Genet* 2003;34:326–9.
- [176] Dalton TP, He L, Wang B, Miller ML, Jin L, Stringer KF, et al. Identification of mouse SLC39A8 as the transporter responsible for cadmium-induced toxicity in the testis. *Proc Natl Acad Sci USA* 2005;102:3401–6.

- 1884 [177] Fiorini C, Tilloy-Ellul A, Schevalier S, Charuel C, Pointis G. Sertoli  
1885 cell junctional proteins as early targets for different classes of  
1886 reproductive toxicants. *Reprod Toxicol* 2004;18:413–21.
- 1887 [178] Chung NPY, Cheng CY. Is cadmium chloride-induced inter-Sertoli  
1888 tight junction permeability barrier disruption a suitable in vitro model  
1889 to study the events of junctional disassembly during spermatogenesis  
1890 in the rat testis? *Endocrinology* 2001;142:1878–88.
- 1891 [179] Hew KW, Heath GL, Jiwa AH, Welsh MJ. Cadmium in vivo causes  
1892 disruption of tight junction-associated microfilaments in rat Sertoli  
1893 cells. *Biol Reprod* 1993;49:840–9.
- 1894 [180] Gray LE, Ostby J, Furr J, Wolf CJ, Lambright C, Parks L, et al.  
1895 Effects of environmental antiandrogens on reproductive development  
1896 in experimental animals. *Hum Reprod Update* 2001;7:248–64.
- 1897 [181] Boekelheide K, Johnson KJ, Richburg JH. Sertoli cell toxicants. In:  
1898 Skinner MK, Griswold MD, editors. *Sertoli cell biology*. San Diego:  
1899 Elsevier Academic Press; 2005. p. 345–82.
- 1900 [182] Toyama Y, Suzuki-Toyota F, Maekawa M, Ito C, Toshimori K.  
1901 Adverse effects of bisphenol A to spermiogenesis in mice and rats.  
1902 *Arch Histol Cytol* 2004;67:373–81.
- 1903 [183] Fisher JS. Environmental anti-androgens and male reproductive  
1904 health: focus on phthalates and testicular dysgenesis syndrome.  
1905 *Reproduction* 2004;127:305–15.
- 1906 [184] Kelce WR, Monosson E, Gamcsik MP, Laws SC, Gray LEJ. Environ-  
1907 mental hormone disruptors: evidence that vinclozolin develop-  
1908 mental toxicity is mediated by antiandrogenic metabolites. *Toxicol  
1909 Appl Pharmacol* 1994;126.
- 1910 [185] Wong C-I, Kelce WR, Sar M, Wilson EM. Androgen receptor  
1911 antagonist versus agonist activities of the fungicide vinclozolin  
1912 relative to hydroxyflutamide. *J Biol Chem* 1995;270:19998–20003.
- 1913 [186] Foster PMD, Mylchreest E, Gaido KW, Sar M. Effects of phthalate  
1914 esters on the developing reproductive tract of male rats. *Hum Reprod  
1915 Update* 2001;7:231–5.
- 1916 [187] Creasy DM, Beech LM, Gray TJ, Butler WH. The ultrastructural  
1917 effects of di-*n*-pentyl phthalate on the testis of the mature rats. *Exp  
1918 Mol Pathol* 1987;46:357–71.
- 1919 [188] Creasy DM, Foster JR, Foster PM. The morphological development  
1920 of di-*N*-pentyl phthalate induced testicular atrophy in the rat. *J Pathol  
1921* 1983;139:309–21.
- 1922 [189] Hotchkiss AK, Ostby JS, Vandenberg JG, Gray J, Earl L. An  
1923 environmental antiandrogen, vinclozolin, alters the organization of  
1924 play behavior. *Physiol Behav* 2003;79:151–6.
- 1925 [190] Gray LEJ, Ostby J, Monosson E, Kelce WR. Environmental anti-  
1926 androgens: low doses of the fungicide vinclozolin alter sexual  
1927 differentiation of the male rat. *Toxicol Ind Health* 1999;15:48–64.
- 1928 [191] Yu WJ, Lee BJ, Nam SY, Ahn B, Hong JT, Do JC, et al. Reproductive  
1929 disorders in pubertal and adult phase of the male rats exposed to  
1930 vinclozolin during puberty. *J Vet Med Sci* 2004;66:847–53.
- 1931 [192] Kubota K, Ohsako S, Kurosawa S, Takeda K, Qing W, Sakaue M, et  
1932 al. Effects of vinclozolin administration on sperm production and  
1933 testosterone biosynthetic pathway in adult male rat. *J Reprod Dev  
1934* 2003;49:403–12.
- 1935 [193] Venkiteswaran K, Xiao K, Summers S, Calkins CC, Vincent PA,  
1936 Pumiglia K, et al. Regulation of endothelial barrier function and  
1937 growth by VE-cadherin, plakoglobin, and  $\beta$ -catenin. *Am J Physiol  
1938 Cell Physiol* 2002;283:C811–21.
- 1939 [194] Troxell ML, Chen Y-T, Cobb N, Nelson WJ, Marrs JA. Cadherin  
1940 function in junctional complex rearrangement and posttranslational  
1941 control of cadherin expression. *Am J Physiol Cell Physiol* 1999;  
1942 276:C404–18.
- 1943 [195] Man Y, Hart VJ, Ring CJA, Sanjar S, West MR. Loss of epithelial  
1944 integrity resulting from E-cadherin dysfunction predisposes airway  
1945 epithelial cells to adenoviral infection. *Am J Respir Cell Mol Biol*  
1946 2000;23:610–7.
- 1947 [196] Guo X, Rao JN, Liu L, Zou TT, Turner DJ, Bass BL, et al. Regulation  
1948 of adherens junctions and epithelial paracellular permeability: a  
1949 novel function for polyamines. *Am J Physiol Cell Physiol* 2003;  
1950 285:C1174–87.
- [197] Gassler N, Rohr C, Schneider A, Kartenbeck J, Bach A, Obermuller N, et al. Inflammatory bowel disease is associated with changes of enterocytic junctions. *Am J Physiol Gastrointest Liver Physiol* 2001;281:G216–28.
- [198] Toyama Y, Yuasa S. Effects of neonatal administration of 17 $\beta$ -estradiol,  $\beta$ -estradiol 3-benzoate, or bisphenol A on mouse and rat spermatogenesis. *Reprod Toxicol* 2004;19:181–8.
- [199] Li LH, Heindel JJ. Sertoli cell toxicants. In: Korach KS, editor. *Reproductive and developmental toxicology*. New York: Dekker Marcel; 1998. p. 655–91.
- [200] Setchell BP, Waites GMH. Changes in the permeability of testicular capillaries and of the “blood-testis barrier” after injection of cadmium chloride in the rat. *J Endocrinol* 1970;47:81–6.
- [201] Kagi J, Schaffer A. *Biochemistry of metallothionein*. *Biochemistry* 1988;27:8509–15.
- [202] Nath R, Kambadur R, Gulati S, Paliwal V, Sharma M. Molecular aspects, physiological function, and clinical significance of metallothioneins. *Crit Rev Food Nutr* 1988;27:41–85.
- [203] Nordberg G. Cadmium metabolism and toxicity. *Environ Physiol Biochem* 1972;2:7–36.
- [204] Vallee B. The function of metallothionein. *Neurochem Int* 1995;27:23–33.
- [205] Waalkes M, Goering P. Metallothionein and other cadmium-binding proteins: Recent developments. *Chem Res Toxicol* 1990;3:281–8.
- [206] Sciavolino P, Lee T, Vilcek J. Overexpression of metallothionein confers resistance to the cytotoxic effect of TNF with cadmium in MCF-7 breast carcinoma cells. *Lymphokine Cytokine Res* 1992;11:265–70.
- [207] Dufresne J, Cyr D. Effects of short-term methylmercury exposure on metallothionein mRNA levels in the testis and epididymis of the rat. *J Androl* 1999;20:769–78.
- [208] Suzuki J, Kodama N, Molotkov A, Aoko E, Tohyama C. Isolation and identification of metallothionein isoforms (MT-1 and MP-2) in the rat testis. *Biochem J* 1998;334:695–701.
- [209] Sugihara T, Wadhwa R, Kaul S, Mitsui Y. A novel testis-specific metallothionein-like protein, tesmin, is an early marker of male germ cell differentiation. *Genomics* 1999;57:130–6.
- [210] Ren X, Zhou Y, Zhang J, Feng W, Jiao B. Metallothionein gene expression under different time in testicular Sertoli and spermatogenic cells of rats treated with cadmium. *Reprod Toxicol* 2003;17:219–27.
- [211] Coogan T, Shiraiishi N, Waalkes M. Minimal basal activity and lack of metal-induced activation of the metallothionein gene correlates with lobe-specific sensitivity to the carcinogenic effects of cadmium in the rat prostate. *Toxicol Appl Pharmacol* 1995;132:164–73.
- [212] Coogan T, Shiraiishi N, Waalkes M. Metallothionein gene expression in the reproductive tissues of the Wistar rat: Effects of treatment with metals and glucocorticoids. *Toxic Subst Mech* 1997;16:357–70.
- [213] Lee K, Lau K, Ho S. Effects of cadmium on metallothionein-I and metallothionein-II mRNA expression in rat ventral, lateral, and dorsal prostatic lobes: Quantification by competitive RT-PCR. *Toxicol Appl Pharmacol* 1999;154:20–7.
- [214] Prozialeck WC. Evidence that E-cadherin may be a target for cadmium toxicity in epithelial cells. *Toxicol Appl Pharmacol* 2000;164:231–49.
- [215] Prozialeck WC, Lamar PC. Interaction of cadmium (Cd<sup>2+</sup>) with a 13-residue polypeptide analog of a putative calcium-binding motif of E-cadherin. *Biochim Biophys Acta* 1999;1451:93–100.
- [216] Prozialeck WC, Niewenhuis RJ. Cadmium (Cd<sup>2+</sup>) disrupts intercellular junctions and actin filaments in LLC-PK1 cells. *Toxicol Appl Pharmacol* 1991;107:81–97.
- [217] Plow EF, Haas TA, Zhang L, Loftus J, Smith JW. Ligand binding to integrins. *J Biol Chem* 2000;275:21785–8.
- [218] De SK, Chen HL, Pace JL, Hunt JS, Terranova PF, Enders GC. Expression of tumor necrosis factor- $\alpha$  in mouse spermatogenic cells. *Endocrinology* 1993;133:389–96.
- [219] Montell DJ. Border-cell migration: the race is on. *Nat Rev Mol Cell Biol* 2003;4:13–24.

- 2018 [220] Matzuk MM. Germ-line immortality. *Proc Natl Acad Sci USA* 2004;101:16395–6.
- 2019 [221] MacLean IIA, Chen MA, Wayne CM, Bruce SR, Rao M, Meistrich ML, et al. Rhox: a new homeobox gene cluster. *Cell* 2005;120:369–82.
- 2020 [222] De Gendt K, Swinnen JV, Saunders PTK, Schoonjans L, Dewerchin M, Devos A, et al. A Sertoli cell-selective knockout of the androgen receptor causes spermatogenic arrest in meiosis. *Proc Natl Acad Sci USA* 2004;101:1327–32.
- 2021 [223] Stamatovic SM, Keep RF, Kunkel SL, Andjelkovic AV. Potential role of MCP-1 in endothelial cell tight junction ‘opening’: signaling via Rho and Rho kinase. *J Cell Sci* 2003;116:4615–28.
- 2022 [224] Pedram A, Razandi M, Levin ER. Deciphering vascular endothelial cell growth factor/vascular permeability factor signaling to vascular permeability. Inhibition by atrial natriuretic peptide. *J Biol Chem* 2002;277:44385–98.
- 2023 [225] Ahdieh M, Vandenbos T, Youakim A. Lung epithelial barrier function and wound healing are decreased by IL-4 and IL-13 and enhanced by IFN- $\gamma$ . *Am J Physiol Cell Physiol* 2001;281:C2029–38.
- 2024 [226] Kagami S, Kuhara T, Yasutomo K, Okada K, Loster K, Reutter W, et al. Transforming growth factor- $\beta$  (TGF- $\beta$ ) stimulates the expression of  $\beta$ 1 integrins and adhesion by rat mesangial cells. *Exp Cell Res* 1996;229:1–6.
- 2025 [227] Tian YC, Fraser D, Attisano L, Phillips AO. TGF- $\beta$ 1-mediated alterations of renal proximal tubular epithelial cell phenotype. *Am J Physiol Renal Physiol* 2003;285:F130–42.
- 2026 [228] de Kretser DM, Buzzard JJ, Okuma Y, O’Connor AE, Hayashi T, Lin S-Y, et al. The role of activin, follistatin and inhibin in testicular physiology. *Mol Cell Endocr* 2004;225:57–64.
- 2027 [229] Shimasaki S, Moore RK, Otsuka F, Erickson GF. The bone morphogenetic protein system in mammalian reproduction. *Endocr Rev* 2004;25:72–101.
- 2028 [230] Pellegrini M, Grimaldi P, Rossi P, Geremia R, Dolci S. Developmental expression of BMP4/ALK3/SMAD5 signaling pathway in the mouse testis: a potential role of BMP4 in spermatogonia differentiation. *J Cell Sci* 2003;116:3363–72.
- 2029 [231] Hua J, Chen Y-X, Wang D, Qi X, Li T-G, Hao J, et al. Developmental expression and function of Bmp4 in spermatogenesis and in maintaining epididymal integrity. *Dev Biol* 2004;276:158–71.
- 2030 [232] Pasparakis M, Alexopoulou L, Episkopou V, Kollias G. Immune and inflammatory responses in TNF $\alpha$ -deficient mice: a critical requirement for TNF $\alpha$  in the formation of primary B cell follicles, follicular dendritic cell networks and germinal centers, and in the maturation of the humoral immune response. *J Exp Med* 1996;184:1397–411.
- 2031 [233] D’Alessio A, Riccioli A, Lauretti P, Padula F, Muciaccia B, De Cesaris P, et al. Testicular FasL is expressed by sperm cells. *Proc Natl Acad Sci USA* 2001;98:3316–21.
- 2032 [234] Lee J, Richburg JH, Younkin SC, Boekelheide K. The Fas system is a key regulator of germ cell apoptosis in the testis. *Endocrinology* 1997;138:2081–8.
- 2033 [235] Adachi M, Suematsu S, Kondo T, Ogasawara J, Tanaka T, Yoshida N, et al. Targeted mutation in the Fas gene causes hyperplasia in peripheral lymphoid organs and liver. *1995;11:294–300*.
- 2034 [236] Grataroli R, Vindrieux D, Gougeon A, Benahmed M. Expression of tumor necrosis factor- $\alpha$ -related apoptosis-inducing ligand and its receptors in rat testis during development. *Biol Reprod* 2002;66:1707–15.
- 2035 [237] Cretney E, Takeda K, Yagita H, Glaccum M, Peschon JJ, Smyth MJ. Increased susceptibility to tumor initiation and metastasis in TNF-related apoptosis-inducing ligand-deficient mice. *J Immunol* 2002;168:1356–61.
- 2036 [238] Luetke N, Qiu T, Fenton S, Troyer K, Riedel R, Chang A, et al. Targeted inactivation of the EGF and amphiregulin genes reveals distinct roles for EGF receptor ligands in mouse mammary gland development. *Development* 1999;126:2739–50.
- 2037 [239] Radhakrishnan B, Oke B, Papadopoulos V, DiAugustine R, Suarez-Quian C. Characterization of epidermal growth factor in mouse testis. *Endocrinology* 1992;131:3091–9.
- 2038 [240] Wahab-Wahlgren A, Martinelle N, Holst M, Jahnukainen K, Parvinen M, Soder O. EGF stimulates rat spermatogonial DNA synthesis in seminiferous tubule segments in vitro. *Mol Cell Endocr* 2003;201:39–46.
- 2039 [241] Bottcher RT, Niehrs C. Fibroblast growth factor signaling during early vertebrate development. *Endocr Rev* 2005;26:63–77.
- 2040 [242] Yamamoto H, Ochiya T, Tamamushi S, Toriyama-Baba H, Takahama Y, Hirai K, et al. HST-1/FGF-4 gene activation induces spermatogenesis and prevents adriamycin-induced testicular toxicity. *Oncogene* 2002;21:899–908.
- 2041 [243] Yamamoto H, Ochiya T, Takahama Y, Ishii Y, Osumi N, Sakamoto H, et al. Detection of spatial localization of Hst-1/Fgf-4 gene expression in brain and testis from adult mice. *Oncogene* 2000;19:3805–10.
- 2042 [244] Ricci G, Catizone A, Galdieri M. Pleiotropic activity of hepatocyte growth factor during embryonic mouse testis development. *Mech Dev* 2002;118:19–28.
- 2043 [245] Catizone A, Ricci G, Galdieri M. Expression and functional role of hepatocyte growth factor receptor (C-MET) during postnatal rat testis development. *Endocrinology* 2001;142:1828–34.
- 2044 [246] Uehara Y, Minowa O, Mori C, Shiota K, Kuno J, Noda T, et al. Placental defect and embryonic lethality in mice lacking hepatocyte growth factor/scatter factor. *Nature* 1995;373:702–5.
- 2045 [247] Meinhardt A, Bacher M, Wennemuth G, Eickhoff R, Hedger M. Macrophage migration inhibitory factor (MIF) as a paracrine mediator in the interaction of testicular somatic cells. *2000;32:46–48*.
- 2046 [248] Honma N, Koseki H, Akasaka T, Nakayama T, Taniguchi M, Serizawa I, et al. Deficiency of the macrophage migration inhibitory factor gene has no significant effect on endotoxaemia. *Immunology* 2000;100:84–90.
- 2047 [249] Marziali G, Lazzaro D, Sorrentino V. Binding of germ cells to mutant S1<sup>d</sup> Sertoli cells is defective and is rescued by expression of the transmembrane form of the c-kit ligand. *Dev Biol* 1993;157:182–90.
- 2048 [250] Bedell MA, Zama AM. Genetic analysis of Kit ligand functions during mouse spermatogenesis. *J Androl* 2004;25:188–99.
- 2049 [251] Nalbandian A, Dettin L, Dym M, Ravindranath N. Expression of vascular endothelial growth factor receptors during male germ cell differentiation in the mouse. *Biol Reprod* 2003;69:985–94.
- 2050 [252] Rudolfsson SH, Wikstrom P, Jonsson A, Collin O, Bergh A. Hormonal regulation and functional role of vascular endothelial growth factor A in the rat testis. *Biol Reprod* 2004;70:340–7.
- 2051 [253] Ferrara N, Carver-Moore K, Chen H, Dowd M, Lu L, O’Shea KS, et al. Heterozygous embryonic lethality induced by targeted inactivation of the VEGF gene. *Nature* 1996;380:439–42.
- 2052 [254] Horai R, Asano M, Sudo K, Kanuka H, Suzuki M, Nishihara M, et al. Production of mice deficient in genes for interleukin (IL)-1 $\alpha$ , IL-1 $\beta$ , IL-1 $\alpha/\beta$ , and IL-1 receptor antagonist shows that IL-1 $\beta$  is crucial in turpentine-induced fever development and glucocorticoid secretion. *J Exp Med* 1998;187:1463–75.
- 2053 [255] Hedger MP, Meinhardt A. Cytokines and the immune-testicular axis. *J Reprod Immunol* 2003;58:1–26.
- 2054 [256] Dalton D. IFN- $\gamma$  and IFN- $\gamma$  receptor knockout mice. In: Fantuzzi G, editor. *Cytokine knockouts*. Totowa, NJ: Humana Press; 2003. p. 347–59.
- 2055 [257] Dejuq N, Lienard M-O, Guillaume E, Dorval I, Jegou B. Expression of interferons- $\alpha$  and - $\gamma$  in testicular interstitial tissue and spermatogonia of the rat. *Endocrinology* 1998;139:3081–7.
- 2056 [258] Pogach LM, Lee Y, Gould S, Giglio W, Meyenhofer M, Huang HF. Characterization of cis-platinum-induced Sertoli cell dysfunction in rodents. *Toxicol Appl Pharmacol* 1989;98:350–61.
- 2057 [259] Peterson PM, Giwerzman A, Skakkebaek NE, Rorth M. Gonadal function in men with testicular cancer. *Semin Oncol* 1998;25:224–33.
- 2058 [260] Wiebe JP, Kowalik A, Gallardi RL, Egeler O, Clubb BH. Glycerol disrupts tight junction-associated actin microfilaments, occludin, and microtubules in Sertoli cells. *J Androl* 2000;21:625–35.
- 2059 [261] Wiebe JP, Barr KJ, Buckingham KD, Geddes PD, Kudo PA. Prospects of a male contraceptive based on selective antispermatogenic action of 1,2,3-trihydroxypropane (THP; glycerol). In: Zatzchni GI,



- 2152 Goldsmith A, Spiler AJM, Sciarra JJ, editors. Male contraceptio- 2174  
 2153 n:advances and future prospects. Philadelphia: Harper and Row; 2175  
 2154 1986. p. 252–70.
- [262] Chung NPY, Mruk D, Mo M-Y, Lee WM, Cheng CY. A 22-amino 2176  
 2155 acid synthetic peptide corresponding to the second extracellular loop 2177  
 2156 of rat occludin perturbs the blood-testis barrier and disrupts sperma- 2178  
 2157 togenesis reversibly in vivo. *Biol Reprod* 2001;65:1340–51. 2179
- [263] Waites GM, Wang C, Griffin PD. Gossypol: reasons for its failure to 2180  
 2159 be accepted as a safe, reversible male antifertility drug. *Int J Androl* 2181  
 2160 1998;21:8–12. 2182
- [264] Qian S, Wang Z. Gossypol: a potential antifertility agent for males. 2183  
 2162 *Annu Rev Pharmacol Toxicol* 1984;24:329–60. 2184
- [265] Pelletier RM, Friend DS. Sertoli cell junctional complexes in gos- 2185  
 2164 sypol-treated neonatal and adult guinea pigs. *J Androl* 1986;7: 2186  
 2165 127–39. 2187
- [266] Silvestrini B, Palazzo G, De Gregorio M. Lonidamine and related 2188  
 2167 compounds. *Prog Med Chem* 1984;21:111–35. 2189
- [267] O'Donnell L, Stanton PG, Bartles JR, Robertson DM. Sertoli cell 2190  
 2169 ectoplasmic specializations in the seminiferous epithelium of the 2191  
 2170 testosterone-suppressed adult rat. *Biol Reprod* 2000;63:99–108. 2192
- [268] McLachlan RI, O'Donnell L, Meachem SJ, Stanton PG, Kretser 2193  
 2172 DMD, Pratis K, et al. Hormonal regulation of spermatogenesis in 2194  
 2173 primates and man: insights for development of the male hormonal 2195  
 contraceptive. *J Androl* 2002;23:149–62. 2196
- [269] Kelce WR, Lambright CR, Gray J, Earl L, Roberts KP. Vinclozolin 2176  
 and p,p'-DDE alter androgen-dependent gene expression: in vivo 2177  
 confirmation of an androgen receptor-mediated mechanism. *Toxicol 2178  
 Appl Pharmacol* 1997;142:192–200. 2179
- [270] Uzumcu M, Suzuki H, Skinner MK. Effect of the anti-androgenic 2180  
 endocrine disruptor vinclozolin on embryonic testis cord formation 2181  
 and postnatal testis development and function. *Reprod Toxicol* 2182  
 2004;18:765–74. 2183
- [271] Ashby J, Tinwell H, Lefevre PA, Joiner R, Haseman J. The effect on 2184  
 sperm production in adult Sprague-Dawley rats exposed by gavage to 2185  
 bisphenol A between postnatal days 91–97. *Toxicol Sci* 2003;74: 2186  
 129–38. 2187
- [272] Williams K, McKinnell C, Saunders PTK, Walker M, Fisher JS, Turner 2188  
 KJ, et al. Neonatal exposure to potent and environmental oestrogens 2189  
 and abnormalities of the male reproductive system in the rat: evidence 2190  
 for importance of the androgen-oestrogen balance and assessment of 2191  
 the relevance to man. *Hum Reprod Update* 2001;7:236–47. 2192
- [273] Wang Y, Thuillier R, Culty M. Prenatal estrogen exposure differen- 2193  
 tially affects estrogen receptor-associated proteins in rat testis gono- 2194  
 cytes. *Biol Reprod* 2004;71:1652–64. 2195  
 2196

UNCORRECTED PROOF