



Title	Life course approach in assessment of ECC in 2-year-old children
Author(s)	Zhou, Y; Lin, HC; Lo, ECM
Citation	The 2nd Meeting of the International Association for Dental Research (IADR) Pan Asian Pacific Federation (PAPF) and the 1st Meeting of the International Association for Dental Research (IADR) Asia/Pacific Region (APR), Wuhan, China, 22-24 September 2009. In Journal of Dental Research, 2009, v. 88, special issue B, abstract no. 5
Issued Date	2009
URL	http://hdl.handle.net/10722/94572
Rights	Creative Commons: Attribution 3.0 Hong Kong License



[Start](#) | [Browse by Day](#) | [Author Index](#) | [Keyword Index](#)

5 Life Course Approach in Assessment of ECC in 2-Year-Old Children

Location: Function Room Xiang Fan (3rd Floor) (Shangri-La Hotel)

Y. ZHOU, Sun Yat-sen University, Guangzhou, China, H.-C. LIN, Sun Yat-sen University, Guang Zhou, China, and E.C.M. LO, University of Hong Kong, Hong Kong, Hong Kong

Objectives: To explore the relationships between early childhood caries (ECC) and relevant biological, behavioral, and socio-economic experiences in the life course of the children.

Methods: A case-control study of 2 years old children was undertaken with an oral health examination and questionnaire investigation. The questionnaire collected socioeconomic and behavioral information at two time points of the participant's life (at birth and 2-year-old). Clinical examinations were performed using standardized criteria. Caries experience was recorded at the level of cavitation and the level of oral hygiene was assessed using the Visible Plaque Index (VPI). Microbiology examination collected information on colony-forming units (CFU) of streptococcus mutans. The outcome was whether the child had early childhood caries (ECC). Unconditional bivariate and multiple logistic regression analysis were performed.

Results: The study included 378 children. At age 2 years, 109 children had ECC (case) while 269 had not (control). The mean dmft of the cases was 3.62 ± 3.12 and 100% of the score was made up of untreated decayed teeth. Mother's occupation, unemployed at the child's birth v.s employers/Professional [OR=3.982; 95%CI=1.146-13.822], Mother's schooling <12 years at child's birth v.s =12 years [OR=1.758; 95%CI=1.098-2.815], Breastfeeding duration=1years v.s no breastfeeding [OR=2.918; 95%CI=1.166-7.301], VPI >60% and 41%-60% v.s 0-20% [OR=5.231, 95%CI=1.840-14.871; OR=2.514, 95%CI=1.097-5.765], Streptococcus mutans= 10^5 CFU/ml v.s 0 CFU/ml [OR=3.867; 95%CI=1.608-9.301] were independent risk factors for ECC through their life course.

Conclusion: The results of this study support the hypothesis linking social, biological and behavioral exposures and development of ECC during the life course of 2-year-old children.

See more of: [Oral Session A \(Cariology Research & Instruments and Equipment\)](#)
See more of: [Scientific Groups](#)

[<< Previous Abstract](#) | [Next Abstract >>](#)