



Title	Effects of increased cholesterol level on BK channels
Author(s)	Wu, W; Lau, CP; Tse, HF; Li, GR
Citation	The 16th Medical Research Conference (MRC), The University of Hong Kong, Hong Kong, China, 22 January 2011. In Hong Kong Medical Journal, 2011, v. 17, suppl. 1, p. 61, abstract no. 101
Issued Date	2011
URL	http://hdl.handle.net/10722/133386
Rights	Creative Commons: Attribution 3.0 Hong Kong License

Effects of increased cholesterol level on BK channels

W Wu, CP Lau, HT Fat, GR Li

Department of Medicine, The University of Hong Kong, Hong Kong

Background: The large conductance Ca^{2+} -activated K^+ (BK, or slo) channels are ubiquitously expressed in different tissues and play an important role in regulating various physiological processes such as cell excitability, hormone secretion, vascular activity, etc. The present study was designed to investigate how/whether BK channels are regulated by increased cholesterol level.

Methods: Whole-cell BK current and BK single channel current were recorded in whole-cell patch clamp mode and cell-attached single channel recording, respectively, in HEK 293 cells stably expressing Maxi-K with beta1-subunit.

Results: Whole-cell BK current was significantly suppressed in cells expressing both with α and $\beta 1$ subunits with cholesterol-enrichment by cholesterol-saturated methyl-beta-cyclodextrin (M β CD), whereas cholesterol depletion by M β CD had no effect on the current amplitude. Low-density lipoprotein (LDL) also decreased BK current. Single channel recording showed that cholesterol enrichment significantly reduced the open probability of BK channels. However, in cells expressing only α -subunit of BK channels (without $\beta 1$ -subunit), cholesterol-saturated M β CD had no significant effect on the current amplitude of BK channels.

Conclusion: Our results demonstrate the important evidence that BK channels exhibit $\beta 1$ -subunit-dependent response to cholesterol. The enriched-cholesterol and LDL reduce the activity of BK channels in cells co-expressed with both α and $\beta 1$ subunits, which may at least in part accounts for the occurrence of hypertension in patients with high plasma cholesterol level, since both of α and $\beta 1$ subunit transcripts are abundant in vascular smooth muscle.

Association of *JAG1* gene and its variant sequence with bone mineral density and osteoporotic fractures: a genome-wide association study and follow-up replication studies

SM Xiao^{1,2}, S Cherny³, G Li¹, Y Gao¹, G Tso¹, KS Lau¹, K Luk⁴, PC Sham³, AWC Kung^{1,2}

¹Department of Medicine, The University of Hong Kong, Hong Kong

²Research Centre of Heart, Brain, Hormone & Healthy Aging, Faculty of Medicine, The University of Hong Kong, Hong Kong

³Department of Psychiatry, The University of Hong Kong, Hong Kong

⁴Department of Orthopaedics and Traumatology, The University of Hong Kong, Hong Kong

Introduction: Bone mineral density (BMD), a diagnostic parameter for osteoporosis and a clinical predictor of fracture, is a polygenic trait with high heritability. The identified variants so far demonstrate only a small effect size on BMD variation. Many additional associated loci remain to be identified. The identification of additional genes underlying BMD variation would therefore provide insight into the pathogenetic mechanisms of osteoporosis and offer strategies for therapeutic development.

Methods: We performed a genome-wide association study (GWAS) on 800 unrelated Southern Chinese women with extreme BMD and follow-up replication studies in six independent European descent and Asian populations including 18 098 subjects. The potential biological function of the identified variant was further validated via electrophoretic mobility shift assay (EMSA) and gene expression study in human bone-derived cells (HBDCs) and peripheral blood mononuclear cells (PBMCs).

Results: In the meta-analysis, rs2273061 of the Jagged 1 (*JAG1*) gene was associated with high BMD ($P=5.27 \times 10^{-8}$ for lumbar spine [LS] and $P=4.15 \times 10^{-5}$ for femoral neck [FN], $n=18\ 898$). This SNP was also associated with the low risk of osteoporotic fracture ($P=0.009$, $OR=0.7$, 95% CI 0.57-0.93, $n=1881$). Furthermore, we performed an EMSA which demonstrated the binding of c-Myc to the 'G' but not 'A' allele of rs2273061. An mRNA expression study in both HBDCs and PBMCs confirmed the high BMD-related allele 'G' of rs2273061 was associated with higher *JAG1* expression.

Conclusion: Our results support the *JAG1* gene as a novel candidate for BMD regulation and it is a potential key factor for fracture pathogenesis.

Acknowledgement: This project was supported by Hong Kong Research Grant Council; The KC Wong Education Foundation; The Bone Health Fund of HKU Foundation; Matching Grant, CRCG Grant and The Osteoporosis Research Fund of The University of Hong Kong.