



Title	Differential actions of glycodelin-A on Th-1 and Th-2 cells: A paracrine mechanism that could produce the Th-2 dominant environment during pregnancy
Author(s)	Lee, CL; Chiu, PCN; Lam, KKW; Siu, SO; Chu, IK; Koistinen, R; Koistinen, H; Seppl, M; Lee, KF; Yeung, WSB
Citation	Human Reproduction, 2011, v. 26 n. 3, p. 517-526
Issued Date	2011
URL	http://hdl.handle.net/10722/135025
Rights	Creative Commons: Attribution 3.0 Hong Kong License

1 **Differential actions of glycodelin-A on Th-1 and Th-2 cells: A paracrine mechanism that could**
2 **produce the Th-2 dominant environment during pregnancy**

3

4 Cheuk-Lun Lee^{1,2,3,#}, Philip C.N. Chiu^{1,3,#,*}, Kevin K.W. Lam¹, Siu-On Siu², Ivan K. Chu², Riitta
5 Koistinen⁴, Hannu Koistinen⁴, Markku Seppälä⁴, Kai-Fai Lee^{1,3}, William S.B. Yeung^{1,3}

6 ¹Department of Obstetrics and Gynaecology, ²Department of Chemistry, ³Centre for Reproduction,
7 Development and Growth, University of Hong Kong, Pokfulam Road, Hong Kong, China.

8 ⁴Department of Clinical Chemistry, Helsinki University Central Hospital and University of Helsinki,
9 00029 HUS Helsinki, Finland.

10 # contributed equally to the report

11

12 Running Title: Differential actions of GdA on Th-1/Th-2 cells

13 Key terms: Glycodelin, T-helper cells, Th-1/Th-2, cell death, Fas, extracellular signal-regulated
14 kinases

15

16 * To whom correspondence should be addressed: Dr. P.C.N. Chiu, Tel: 852-28199388. Fax:

17 852-28161947, email: pchiucn@hkucc.hku.hk

18

19 **Abstract**

20 **BACKGROUND:** The maternal-fetal interface has unique immunological response towards the
21 implanting placenta. It is generally accepted that a T-helper type-2 (Th-2) cytokine prevailing
22 environment is important in pregnancy. The proportion of Th-2 cells in the peripheral blood and
23 deciduas is significantly higher in pregnant women than in nonpregnant women in the first trimester.
24 Glycodelin-A (GdA) is a major endocrine-regulated decidual glycoprotein thought to be related to
25 feto-maternal defense. Yet the relationship between its immunoregulatory activities and the shift
26 towards Th-2 cytokine profile during pregnancy is unclear. **METHODS:** GdA was immunoaffinity
27 purified from human amniotic fluid. T-helper, T-helper type-1 (Th-1) and Th-2 cells were isolated
28 from peripheral blood. Viability of these cells was studied by XTT assay. Immunophenotyping of
29 CD4/CD294, cell death and GdA-binding were determined by flow cytometry. The mRNA
30 expression, surface expression and secretion of Fas/Fas ligand (FasL) were determined by qPCR,
31 flow cytometry and ELISA, respectively. The activities of caspase-3, -8 and -9 were measured. The
32 phosphorylation of extracellular signal-regulated kinases (ERK), p38 and c-Jun N-terminal kinase
33 were determined by Western blotting. **RESULTS:** Although GdA bound to both Th-1 and Th-2
34 cells, it had differential actions on the two cell types. GdA induced cell death of the Th-1 cells but
35 not the Th-2 cells. The cell death was mediated through activation of caspase-3, -8 and -9 activities.
36 GdA up-regulated the expression of Fas and inhibited the ERK activation in the Th-1 cells, which
37 might enhance the vulnerability of the cells to cell death caused by trophoblast-derived FasL.

38 **CONCLUSION:** The data suggest that GdA could be an endometrial factor that contributes to
39 enhancing the Th-1/Th-2 shift during pregnancy.

40 **Introduction**

41 Placenta is genetically a fetal semiallograft in the maternal body, and mechanisms have
42 evolved to suppress the maternal immune response in the uterine tissue during pregnancy
43 (Trowsdale and Betz, 2006). One of these mechanisms is change in the decidual leukocyte
44 population (Luppi, 2003). The altered population of immune cells at the maternal-fetal interface not
45 only allows the mother to tolerate the fetus but also to interact with the trophoblasts, thereby
46 creating an environment that is favorable for fetal development (Luppi, 2003). Contrary to their
47 abundance in the peripheral blood, T-cells represent a minor population of immune cells in early
48 decidua (Loke *et al.*, 1995), partly due to apoptosis of the leukocytes. The trophoblast cells express
49 Fas ligand (FasL), which induces apoptosis of the Fas-expressing leukocytes (Runic *et al.*, 1996;
50 Green and Ferguson, 2001). Other proposed mechanisms for the causing T-cells a minority
51 population in the deciduas include inhibition of T-helper cells proliferation by indoleamine
52 2,3-dioxygenase from antigen-presenting cells (Mellor *et al.*, 2002), ligation of the inhibitory
53 programmed death ligand 1 on uterine T cells (Guleria *et al.*, 2005), and selective enrichment of
54 decidual natural killer cell (Bulmer and Lash, 2005).

55 Despite the reduction of T-cell population in the decidua, considerable amount of T-cells are
56 present around the extravillous trophoblasts, decidual stroma, endometrial gland and decidual
57 vessels (Vassiliadou and Bulmer, 1998; Michimata *et al.*, 2002). T-helper cells are classified into
58 T-helper type 1 (Th-1) and Th-2 according to the cytokines they secrete (Mosmann *et al.*, 1986). It

59 is generally thought that successful pregnancy is a Th-2 type cytokine predominant phenomenon.
60 The percentage of peripheral blood (Saito *et al.*, 1999b) and decidual (Michimata *et al.*, 2002) Th-2
61 cells is significantly higher in pregnant women than in nonpregnant women in the first trimester. The
62 shift from the production of inflammatory Th-1 cytokines towards Th-2 type cytokines promotes
63 immune protection of the trophoblasts (Dealtry *et al.*, 2000; Michimata *et al.*, 2002;
64 Straszewski-Chavez *et al.*, 2005). Pregnancy loss is associated with increased Th-1/Th-2 cytokine
65 ratio (Daher *et al.*, 2004). However, several Th-1 cytokines such as IFN- γ and TNF- α have shown to
66 be important in uterine vascular remodelling and implantation (Chaouat, 2007), suggesting that the
67 Th1/Th2 paradigm for pregnancy may be too simplistic.

68 Glycodelin-A (GdA) is an immunosuppressive glycoprotein abundantly expressed in the
69 decidualized endometrium (Seppala *et al.*, 2002; Seppala *et al.*, 2007). It induces apoptosis of
70 lymphocytes (Lee *et al.*, 2009) and monocytes (Tee *et al.*, 2008), skewing of T-cell response towards
71 Th-2 phenotype (Mishan-Eisenberg *et al.*, 2004), and modulates the activities of natural killer cells
72 (Lee *et al.*, 2010), B-cells (Yaniv *et al.*, 2003), and dendritic cells (Scholz *et al.*, 2008). Recently,
73 GdA was demonstrated to suppress the cytolytic activity of CD8⁺ T-cell (Soni and Karande, 2010).
74 The immunosuppressive activities of GdA are believed to be related to fetomaternal defense (Clark
75 *et al.*, 1996). The serum and decidual concentration of GdA peaks around week 10 of pregnancy,
76 consistent with a role in survival of the fetoplacental unit (Seppala *et al.*, 2002). Decreased maternal
77 serum glycodelin is associated with early spontaneous abortion (Salim *et al.*, 2007) and recurrent

78 miscarriage (Dalton *et al.*, 1998).

79 The mechanisms that generate the Th-2 cytokine-rich environment during pregnancy are not
80 fully known. We hypothesized that GdA has differential actions on Th-1 and Th-2 cells,
81 contributing to the Th1/Th2 shift. Therefore, the objectives of this report were to study the actions of
82 GdA on Th-1/Th-2 cell ratio, and to compare the actions of GdA on the two cell types in terms of
83 cell death, Fas/FasL expression and intracellular signaling.

84

85 **Materials and Methods**

86 **Purification of glycodelin from human amniotic fluid, seminal plasma and cumulus matrix**

87 The Institutional Review Board of the University of Hong Kong/Hospital Authority Hong
88 Kong West Cluster approved the protocol of this study. Glycodelin isoforms including glycodelin-A,
89 glycodelin-S (GdS) and glycodelin-C (GdC) were purified from amniotic fluid, seminal plasma and
90 cumulus matrix, respectively, by affinity chromatography using monoclonal anti-glycodelin
91 antibody (Clone F43-7F9) as described (Riittinen *et al.*, 1989; Lee *et al.*, 2009). In brief, the
92 collected samples were diluted with tris-buffered saline (TBS, pH 7.4) and 0.1% Triton X-100 in a
93 ratio of 1:3-1:5 was added. They were loaded onto anti-glycodelin column, which was then washed
94 successively by TBS, 1M NaCl with 1% isopropanol, 10 mM ammonium acetate with 0.1%
95 isopropanol, pH 5 and TBS. Glycodelin was eluted by 20 mM CaCl₂ with 0.1% trifluoroacetic acid.
96 The eluted GdS and GdC were further purified with anion-exchange Mono-Q (GE Healthcare)

97 column by AKTA purifier 10 (GE Healthcare). Deglycosylated glycodefin was prepared by
98 denaturation of GdA in 0.1% β -mecaptoethanol before incubation with 0.5 mU N-Glycosidase F at
99 37°C for 24 hours (Lee *et al.*, 2009). The concentrations of glycodefin were determined by a protein
100 assay kit (Bio-Rad, Hercules, USA).

101

102 **Isolation of human peripheral T-helper cells and enrichment of Th-2 cells**

103 Human non-pregnant female peripheral blood was obtained from the Hong Kong Red Cross.
104 Ficoll-Paque (GE Healthcare, Uppsala, Sweden) density gradient centrifugation was used to isolate
105 the PBMCs. The contaminated red blood cells and the adherent cells were removed by the red blood
106 cell lysing buffer (0.084% NaHCO_3 , 0.83% NH_4Cl and 0.003% ethylenediaminetetra-acetic acid)
107 and by adhesion to plastic culture flask, respectively. T-helper cells ($\text{CD3}^+\text{CD4}^+$) were isolated by
108 negative immuno-magnetic separation using CD4^+ T-cell isolation kit II (Miltenyi Biotec Inc.,
109 Bergisch Gladbach, Germany). The purity of the $\text{CD3}^+\text{CD4}^+$ cells increased to 90-95% after
110 processing.

111 Th-2 cells were then positively selected by the anti-CD294 (Chemoattractant receptor of Th-2
112 cells, CRTH-2) MicroBead Kit (Miltenyi Biotec Inc.). The purity of the $\text{CD4}^+\text{CD294}^+$ Th-2 cells
113 was >85% after processing (Supporting information Figure S1). The cell population that did not
114 bind to the anti-CD294 antibody column ($\text{CD4}^+\text{CD294}^-$) was considered as an enriched Th-1 cell
115 preparation. To verify that the enriched Th1 and Th2 cells were producing polarised cytokines we

116 have analyzed cytokines in the culture supernatant by ELISA. The isolated Th-2 cells secreted
117 significantly ($P < 0.05$) less IL-2, IL-8 and IFN- γ (Th-1 cytokines) and more IL-10 (Th-2 cytokine)
118 than that of the enriched Th-1 cells as determined by ELISA (Supporting information Table ST1).
119 The cells were resuspended in 10% fetal bovine serum supplemented RPMI 1640 medium (Sigma,
120 St. Louis, MO).

121

122 **Immunophenotyping of T-helper cells**

123 Cells (5×10^5) were treated with different concentrations (0.01-1 $\mu\text{g/mL}$) of GdA in 500 μL of
124 culture medium for 48 hours before the immunophenotyping. In brief, treated cells were
125 successively washed twice with PBS and once with 1% BSA containing 0.1% sodium azide in PBS.
126 The cells were then incubated with anti-CD4-FITC (T-helper cell marker); anti-CD294-PE (Th-2
127 cell marker); anti-Fas-FITC; anti-FasL-PE and PE/FITC-conjugated mouse isotypic control (BD
128 Biosciences, San Jose, CA) in 1% BSA and 0.1% sodium azide in PBS. The cells were analyzed by
129 a flow cytometer using 525 nm and 575 nm band pass filters and the results were evaluated by the
130 WinMDI 2.8 (The Scripps Research Institute Cytometry Software,
131 <http://facs.Scripps.edu/software.html>). The non-viable cells were removed by gating with forward
132 scatter/side-scatter.

133

134 **Cell viability assay**

135 Cells (3×10^4) were incubated with 0.001-1 $\mu\text{g}/\text{mL}$ of GdA, GdS, GdC or deglycosylated
136 glycodelin in 100 μL of culture medium for 36 hours before cell viability determination. Cell
137 viability was determined by the XTT assay (Roche Diagnostics Co., Basel, Switzerland). In brief,
138 freshly prepared XTT labeling mixture (50 μL) was added to the cell culture 12 hours before the end
139 of the experiment. The absorbance was measured at 450 nm with λ correction at 595 nm. The cell
140 viability was expressed as Suppression Index = (Absorbance of treated cells - Absorbance of
141 blank)/(Absorbance of control - Absorbance of blank) \times 100%.

142

143 **Cell death analysis**

144 Cells (5×10^5) were treated with 0.01-1 $\mu\text{g}/\text{mL}$ of GdA in 500 μL of culture medium for 48
145 hours. Apoptotic and necrotic cell deaths were determined by flow cytometry using Yo-Pro®-1 and
146 propidium iodide dye (Invitrogen, Carlsbad, CA). The treated cells were washed twice with PBS,
147 incubated with Yo-Pro®-1 (1 μL) and propidium iodide (1 μL) in 1 mL PBS for 15 minutes, and
148 analyzed immediately by flow cytometer using the 525 nm and 610 nm band pass filters. The data
149 were analyzed by WinMDI 2.8.

150

151 **Determination of Fas/FasL mRNA and secreted sFas/sFasL**

152 Cells (5×10^5) were treated with GdA (0.01-1 $\mu\text{g}/\text{mL}$) in 500 μL of culture medium for 48 hours.
153 The QuickPrep RNA extraction kit (Stratagene, La Jolla, CA) was then used to extract total RNA

154 from the cells. RNA was reverse transcribed using the TaqMan reverse transcription reagent kit
155 (Applied Biosystems, Foster City, CA) and multiscript reverse transcriptase. qPCR was performed
156 using the TaqMan PCR core reagent kit (N8080228, ABI Biosystems). In brief, cDNA sample (1 μ L)
157 was mixed with 2x TaqMan universal PCR master mix, Fas or FasL target primers and probe (ABI
158 Biosystems) and 18S internal control primers and probe (ABI Biosystems) in a 96-well reaction
159 plate (ABI Biosystems). The reactions were performed in triplicates. PCR was performed at 50°C
160 for 2 minutes and 95°C for 10 minutes, followed by 40 cycles at 95°C for 15 seconds, and 60°C for 1
161 minute in an ABI 7500 system (ABI Biosystems). The Ct values of the Fas and FasL experiments
162 was <35 and that of 18S was <20, respectively. The relative quantification value (RQ) was
163 calculated by $2^{-\Delta\Delta CT}$ method (Schmittgen and Livak, 2008). The data was present as relative
164 expression = (RQ of glycodelin-treated cells - RQ of negative control)/(RQ of control cells - RQ of
165 negative control).

166 The levels of soluble Fas (sFas) and FasL (sFasL) in the culture supernatant of the treated cells
167 were measured by ELISA according to the manufacturer's protocol (Bender Medsystem®,
168 Burlingame, CA). Briefly, the microwell coated with sFas or FasL monoclonal antibodies were
169 washed twice with 300 μ L of wash buffer. One hundred microlitres of culture medium or standard
170 was then added to the well, washed and incubated with biotin-conjugated detector antibodies for 2
171 hours at 37°C. The unbound material was washed away and streptavidin-HRP (100 μ L per well) was
172 added. Color development was performed using 100 μ L of 3,3',5,5'-tetramethylbenzidine as

173 chromogen. The reaction was stopped by the addition of 2 M sulphuric acid (1000 μ L/well) and the
174 absorbance was measured immediately at 450 nm with λ correction at 595 nm in an ELISA plate
175 reader (Infinite F200, Tecan, Männedorf, Switzerland).

176

177 **Caspase-3, -8 and -9 activity assays**

178 Cells (1×10^6) were treated with different concentrations (0.01-1 μ g/mL) of GdA in 500 μ L of
179 culture medium for 48 hours before the caspases activity assay. Caspase activities were determined
180 with the use of synthetic substrates of caspase-3 (Z-DEVD-R110, Invitrogen), caspase-8 (IETD-pNA,
181 Invitrogen) and caspase-9 (Ac-LEHD-pNA, Millipore). Cells were washed with PBS and lysed with
182 50 μ L of cell lysis buffer provided by the assay kit at 4°C for 1 hour. The cell lysate (50 μ L) was then
183 mixed with different caspase substrates in 50 μ L of reaction buffer and incubated in dark according to
184 the manufacturer's protocols. The fluorescence intensity or absorbance was measured immediately
185 after incubation. Caspase activities were expressed as relative activity (%) = (Absorbance of
186 glycodelin-treated cells - Absorbance of blank)/(Absorbance of control cells - Absorbance of blank) \times
187 100%.

188

189 **Western blot analysis of MAPK/ERK activation**

190 The action of GdA on the MAPK/ERK activation in T-helper cells (1×10^6) was determined by
191 western blot analyses after treatment of GdA (1 μ g/mL) for 6 hours followed by the

192 phytohaemagglutinin (5 µg/mL) stimulation for 30 minutes. Cells were lysed with CytoBluster™
193 protein extraction reagent (Novagen®, Darmstadt, Germany) at 4°C for 2 hours in the presence of a
194 cocktail of protease inhibitors (Calbiochem, San Diego, CA). The protein lysates were resolved by
195 SDS-PAGE and transferred to a PVDF membrane for Western blot analysis of the components of the
196 MAPK pathways using anti-ERK, anti-phospho-ERK, anti-p38 (BD Biosciences), anti-phospho-p38,
197 anti-JNK and anti-phospho-JNK antibodies (Cell signaling, MA). Density of the protein bands were
198 measured by Quantity One software (Bio-Rad) and the density values are present as relative quantities
199 = (Density of glycodeclin treated cells)/(Density of control).

200

201 **GdA binding assay**

202 GdA was labeled with Alexa Flour®488 according to the manufacturer's protocol (Invitrogen).
203 Cells (5×10^5) were incubated with the labeled GdA (1 µg/mL) for 2 hours. Unbound GdA was
204 removed by washing with PBS twice. The samples were then re-suspended in 500 µL of PBS and
205 analyzed by flow cytometry. The fluorescence signals were measured using 525 nm band pass filter
206 and the data were analyzed by WinMDI 2.8 software.

207

208 **Data Analysis**

209 All values were expressed as mean ± SEM. For all experiments, the non-parametric ANOVA
210 on Rank test for comparisons was used to identify differences between groups. If the data were

211 normally distributed, parametric Student's t-test or non-parametric Mann Whitney U test were used
212 where appropriate as the post-test. The data were analyzed by SigmaStat 2.03 (Jandel Scientific, San
213 Rafael, CA) with a P-value less than 0.05 was considered as significant.

214

215 **Results**

216 **GdA increased the proportion of Th-2 cells in the T-helper cell population**

217 T-helper cells ($CD3^+CD4^+$) were isolated by negative immuno-magnetic separation and treated
218 with different concentrations of GdA. The non-viable cells were removed by gating with forward
219 scatter/side-scatter. In the viable population, the treatment significantly increased the proportion of
220 $CD4^+CD294^+$ cells (Th-2 cells) after 48 hours (Figure 1). Treatment with 1 $\mu\text{g/mL}$ of GdA increased
221 the percentage of Th-2 cells from $1.05\pm 0.07\%$ (Control) to $1.34\pm 0.08\%$ ($P<0.05$). As a result, the ratio
222 of viable $CD4^+CD294^-$ (Th-1) cells to $CD4^+CD294^+$ (Th-2) cells decreased from 94.33 to 72.56
223 ($P<0.05$). The change in Th1/Th2 ratio could be due to a decrease in Th-1 cells or an increase in Th-2
224 cells. Therefore, we studied the action of GdA on the two isolated sub-populations.

225

226 **GdA induced cell death of the Th-1 cells but not the Th-2 cells**

227 GdA at concentrations $\geq 0.1 \mu\text{g/mL}$ significantly decreased ($P<0.05$) the viability of the isolated
228 T-helper cells as demonstrated by the XTT assay (Table 1). The suppression index of the cells
229 treated with 1 $\mu\text{g/mL}$ of GdA was $53.67\pm 8.94\%$ ($P<0.05$), which was significantly lower than that of

230 the untreated control (100%).

231 Consistently, GdA dose-dependently induced both apoptosis and necrosis of T-helper cells
232 (Figure 2). The viability of the cells treated with 1 $\mu\text{g}/\text{mL}$ of GdA decreased from $91.33\pm 1.05\%$
233 (Control) to $69.34\pm 4.81\%$ ($P<0.05$), whereas the proportion of apoptotic and necrotic cells increased
234 from 3.56 ± 0.16 (Control) to 15.74% ($P<0.05$) and from $5.08\pm 0.67\%$ (Control) to $14.69\pm 2.09\%$
235 ($P<0.05$), respectively. Differentially glycosylated glycodelin isoforms, glycodelin-S and
236 glycodelin-C, and deglycosylated glycodelin had no effects on the viability and cell death of the
237 T-helper cells.

238 Positive selection process for $\text{CD}294^+$ was used to isolate Th-2 cells from the total T-helper cells.
239 The selection process decreased the viability of the Th-2 cells ($\sim 80\%$), which was lower than that of
240 the Th-1 cells ($\sim 90\%$). Similar reduction in Th-1 cell viability was observed after incubating the cells
241 with CD45 microbeads for positive selection (Supporting information Figure S2). GdA did not affect
242 cell death of the isolated Th-2 cells. This was in striking contrast to that of the Th-1 sub-population,
243 the viability of which was significantly ($P<0.05$) reduced from $90.13 \pm 0.90\%$ to $68.82 \pm 4.58\%$ as a
244 result of increases in apoptotic and necrotic cell death after GdA ($1 \mu\text{g}/\text{mL}$) treatment (Figure 2, Table
245 2).

246

247 **GdA upregulated the Fas expression in isolated T-helper cells and Th-1 cell**

248 The effects of GdA on Fas/FasL expression were analyzed by quantitative polymerase chain

249 reaction, flow cytometry and ELISA (Table 2). GdA dose-dependently increased the mRNA and
250 surface expression of Fas. At 1 $\mu\text{g}/\text{mL}$ of GdA, the Fas mRNA expression was significantly ($P<0.05$)
251 upregulated by 2.34 ± 0.62 fold in isolated T-helper cells, and the cell surface Fas expression was also
252 significantly ($P<0.05$) increased from $34.10\pm 1.41\%$ (Control) to $45.26\pm 2.17\%$. In contrast, GdA did
253 not affect the FasL expression and sFas secretion in the isolated T-helper cells. The level of sFasL in
254 the culture medium was low and barely detectable.

255 The effect of GdA on Fas expression was also determined in the enriched Th-1 and Th-2 cells
256 by flow cytometry (Table 2). GdA treatment (1 $\mu\text{g}/\text{mL}$) significantly enhanced ($P<0.05$) the Fas
257 expression of Th-1 cells from $33.17\pm 3.80\%$ to $54.78\pm 3.97\%$, but had no effect on the Th-2 cells.

258

259 **GdA enhances the caspase activity in isolated T-helper cells and Th-1 cells**

260 The caspase-3, -8 and -9 activities of T-helper cells were investigated using specific substrates
261 (Table 3). As compared to the controls without treatment, 1 $\mu\text{g}/\text{mL}$ of GdA significantly ($P<0.05$)
262 increased the caspase-3, -8 and -9 activity to $151.25\pm 24.66\%$, $120.10\pm 11.85\%$ and $130.34\pm 10.75\%$,
263 respectively of the control values. GdA had no significant effect on the expression level of
264 non-activated pro-caspase-3, -8 and -9 as determined by Western blotting analysis.

265 In the enriched Th-1 cell sub-population, incubation with 1 $\mu\text{g}/\text{mL}$ GdA significantly increased
266 ($P<0.05$) the caspase-3 and caspase-9 activities to $163.89\pm 8.18\%$ and $133.72\pm 10.24\%$ of the control
267 values. Although the caspase-8 activity was increased to $119.11\pm 6.81\%$, the difference did not reach

268 statistical significance. GdA had no effect on caspase-3, -8 and -9 expression in the Th-2 cells.

269

270 **GdA suppresses ERK activation in isolated T-helper cells and Th-1 cells**

271 The expression and activation by PHA of ERK, p38 and JNK were determined by Western blot
272 analysis. GdA dose-dependently reduced the phosphorylated-ERK level, but not that of
273 phosphorylated-JNK nor phosphorylated-p38 in the T-helper cells (Figure 3A). GdA at a concentration
274 of 1 $\mu\text{g}/\text{mL}$ significantly reduced the levels of the 42 kDa and the 44 kDa phosphorylated-ERK to
275 0.48 ± 0.11 and 0.61 ± 0.09 ($P<0.05$) respectively, as compared to the control without treatment. GdA
276 treatment did not affect the expression or phosphorylation of p38 or JNK.

277 GdA suppressed ($P<0.05$) the phosphorylated-ERK level in the PHA-stimulated Th-1 cells but
278 not the Th-2 cells (Figure 3B). Treatment with 1 $\mu\text{g}/\text{mL}$ of GdA decreased ($P<0.05$) the expression
279 of phosphorylated-p42 and p44 ERKs in the Th-1 cells to 0.58 ± 0.09 and 0.57 ± 0.10 , respectively, but
280 had no effect on the non-phosphorylated ERK.

281

282 **GdA had similar binding on Th-1 and Th-2 cells**

283 The binding of fluorescent labeled GdA on Th-1 and Th-2 cells was determined by flow
284 cytometry (Supporting information Figure S3). The percentage of Th-1 cells with bound GdA
285 ($91.17\pm 0.90\%$) was not significantly different from that of Th-2 cells ($93.77\pm 0.84\%$), indicating that
286 the binding of GdA to these cells was similar.

287

288 **Discussion**

289 The maintenance of pregnancy requires a Th-2 cytokine dominant environment (Dealtry *et al.*,
290 2000; Michimata *et al.*, 2002). Dysregulation of Th-1 and Th-2 cells is associated with implantation
291 failure and recurrent pregnancy loss (Daher *et al.*, 2004). The ratio of Th-1/Th-2 cells decrease
292 drastically in the peripheral blood and early decidua of pregnant women when compared with the
293 nonpregnant one (Saito *et al.*, 1999a; Saito *et al.*, 1999b; Michimata *et al.*, 2002). Such a change has
294 been proposed to be resulted from selective modulation of differentiation, chemoattraction, and
295 proliferation of Th-2 cells and death of Th-1 cells. GdA stimulates the Th-2 type cytokine shift in
296 T-cells (Mishan-Eisenberg *et al.*, 2004). However, the specific mechanisms of action are still unclear.
297 This study provides evidence for a novel mechanism of GdA in shifting the Th-1/Th-2 balance, i.e.
298 differential actions on Th-1 and Th-2 cells in terms of selective induction of cell death, concomitant
299 with increased expression of Fas in the Th-1 cells.

300 Cytotoxic action of GdA on T-cells has been reported (Sundarraj *et al.*, 2008; Lee *et al.*, 2009).
301 In this study, both the T-helper and enriched Th-1 cells responded similarly to GdA-mediated cell
302 death. It is because the major population in the peripheral blood T-helper cells was the Th-1 cells,
303 and Th-2 cells only constitute a minor proportion ranging from 0.4-6.5% (Nagata *et al.*, 1999). The
304 cytotoxic action of GdA on Th-1 cells is glycan-dependent, consistent with the reported contribution
305 of the glycosylation of GdA to its binding to (Ish-Shalom *et al.*, 2006) and induction of cell death in

306 lymphocytes (Lee *et al.*, 2009). On the other hand, GdA has no effect on the isolated Th-2
307 population.

308 The percentage of peripheral blood Th-2 cells is significantly higher in pregnant women than
309 in nonpregnant women in the first trimester of pregnancy (Saito *et al.*, 1999a). The dosage (0.1-1
310 µg/ml) of GdA exerting its biological activities in this study is within the concentration range of
311 GdA (0.2-1.2 µg/ml) in the peripheral blood of women in their first trimester of pregnancy (Seppala
312 *et al.*, 2002), suggesting that the observations could be physiologically relevant. Due to the high
313 abundance of GdA in the first-trimester decidual tissue, it is possible that GdA may possess similar
314 actions on decidual T-cells. However, experimental evidence on this possibility is still lacking.

315 GdA increased apoptotic and necrotic cell death of the Th-1 subpopulation. Previous studies
316 had reported differential effects of apoptotic and necrotic cells in modulating the activities of other
317 immune cells. For example, the presence of apoptotic, but not the necrotic T-cells, up-regulates
318 IL-10 production from macrophages (Chung *et al.*, 2007). The development of dendritic cells is also
319 affected by apoptotic T-cells (Newton *et al.*, 2003). Therefore, the changes in apoptotic and necrotic
320 T-cell population induced by GdA may further modulate the immune response of pregnant women.

321 The results of this study showed that GdA-induced cell death in both T-helper cells and Th-1
322 cells are associated with the increase in caspase-3, -8, -9 activities. This suggests that both
323 mitochondrial-independent and mitochondrial-dependent pathways are involved. Previous studies
324 showed that both native and recombinant GdA induced apoptosis in T-cells through the

325 mitochondrial-dependent pathway as indicated by influx of mitochondrial membrane calcium ion
326 and involvement of caspase-9 and Bcl-2 activities (Sundarraaj *et al.*, 2008). There are no reports on
327 the involvement of mitochondrion-independent pathway in GdA-mediated cell death of lymphocytes,
328 whereas both recombinant glycodelin and native GdA have been reported to induce cell death of
329 monocytes via caspase-8 pathway (Tee *et al.*, 2008).

330 A novel observation in this study is the differential induction of GdA on Fas expression in
331 T-cells; it was enhanced by GdA in the Th-1, but not the Th-2 cells. Fas/FasL pathway regulates
332 clonal deletion of T-cells at the fetomaternal interface (Coumans *et al.*, 1999; Jerzak and Bischof,
333 2002), as well as in some other immunologically privileged sites, such as the anterior chamber of the
334 eye and the testis (Green and Ferguson, 2001). The trophoblasts express FasL (Runic *et al.*, 1996),
335 while the T-helper cells produce both Fas and FasL (Ramsdell *et al.*, 1994). Binding of FasL to Fas
336 receptor induces trimerization of the Fas receptor, which activates the Fas-associated death domain
337 and the caspase cascade leading to apoptosis. Most importantly, the Th-1 cells express more Fas
338 receptors than the Th-2 cells do, and they are more susceptible to Fas/FasL-induced cell death
339 (Roberts *et al.*, 2003). Thus, the up-regulation by GdA of Fas receptor in the Th-1 cells may further
340 increase vulnerability of these cells to death-induction by the FasL derived from the trophoblasts or
341 lymphocytes.

342 In this study, we further demonstrate the suppressive effect of GdA on ERK activation in the
343 Th-1 cells. ERK activation is important in T-cell activation, homeostasis and cytokine secretion

344 (Dong *et al.*, 2002). It has been correlated with Th-1 response (Borovsky *et al.*, 2002), and
345 regulation of the expression of Th-1 cytokines including IL-1 (Wang *et al.*, 2004) and IFN- γ
346 (Mainiero *et al.*, 1998) in different cell types. The ERK activation in T-cells also inhibited the cells
347 to Fas-mediated apoptosis (Holmstrom *et al.*, 2000). Therefore, apart from inducing Fas expression,
348 GdA treatment may sensitize Th-1 cells to Fas receptor-mediated apoptosis by suppressing ERK
349 activation in the cells, as has been shown in primary peripheral T-cells after inhibition of ERK
350 signaling (Holmstrom *et al.*, 2000). It is of interest that GdA also suppresses ERK activation in
351 spermatozoa (Yeung *et al.*, 2009) and trophoblast cells (Lam KKW and Chiu PCN, unpublished
352 data), suggesting that ERK may have a central role in GdA signaling in different biological systems.
353 Consistently, the lack of cytotoxic activity of GdA on Th-2 cells is associated with absence of ERK
354 suppression in these cells.

355 The observation of similar binding of GdA onto Th-1 and Th-2 cells suggests that GdA binding
356 alone does not contribute to the differential response of Th-1 and Th-2 cells to GdA treatment.
357 However, the observation does not exclude the possible presence of different GdA receptors with the
358 same affinity in the two cell types. CD45 has been proposed as a possible GdA receptor in T-cells
359 (Rachmilewitz *et al.*, 2003). Differential expression of the CD45RC isoform in CD4⁺ T-cells
360 sub-populations is associated with differences in cytokine production upon stimulation; the
361 CD45RC^{high} sub-population produces mainly type-1 cytokines including IL-2, while the CD45RC^{low}
362 sub-population produces IL-17, IL-10 and Th-2 cytokines (Ordonez *et al.*, 2009). Alternatively, the

363 two cell types may have the same GdA receptor but different intracellular signaling leading to the
364 observed differential action of GdA. The current data cannot distinguish between these possibilities.

365 We concluded that GdA may involved in shifting the Th-1/Th-2 ratio in the peripheral blood
366 and decidua by selectively reducing the Th-1 cell population, both directly through induction of cell
367 death of Th-1 cells and indirectly through enhancing the expression of Fas and suppression of ERK
368 activation in the Th-1 cells, thereby enhancing their vulnerability to cell death induced by
369 trophoblast-derived FasL. Overall, the present data uncover the mechanisms in part by which GdA
370 contributes to immunoprotection of the fetoplacental unit during human pregnancy.

371

372 **Acknowledgments**

373 This study is supported in part by the Helsinki University Central Hospital Research Fund.
374 C.L.L. and P.C.N.C. performed the experiments and analysed the data, contributed to discussion,
375 and wrote the manuscript. K.K.W.L., R.K., H.K. and M.S. performed the experiments, contributed
376 to discussion and revised the manuscript. S.O.S, I.K.C., K.F.L. and W.S.B.Y. contributed to
377 experimental design, discussion and reviewed/edited the manuscript.

378

379 **References**

- 380 Borovsky Z, Mishan-Eisenberg G, Yaniv E and Rachmilewitz J. Serial triggering of T cell receptors
381 results in incremental accumulation of signaling intermediates. *J Biol Chem*
382 2002;**24**;21529-21536.
- 383 Bulmer JN and Lash GE. Human uterine natural killer cells: a reappraisal. *Mol Immunol*
384 2005;**4**;511-521.
- 385 Chaouat G. The Th1/Th2 paradigm: still important in pregnancy? *Semin Immunopathol*
386 2007;**2**;95-113.
- 387 Chung EY, Liu J, Homma Y, Zhang Y, Brendolan A, Saggese M, Han J, Silverstein R, Selleri L and
388 Ma X. Interleukin-10 expression in macrophages during phagocytosis of apoptotic cells is
389 mediated by homeodomain proteins Pbx1 and Prep-1. *Immunity* 2007;**6**;952-964.
- 390 Clark GF, Oehninger S, Patankar MS, Koistinen R, Dell A, Morris HR, Koistinen H and Seppala M.
391 A role for glycoconjugates in human development: the human foeto-embryonic defence system
392 hypothesis. *Hum Reprod* 1996;**3**;467-473.
- 393 Coumans B, Thellin O, Zorzi W, Melot F, Bougoussa M, Melen L, Zorzi D, Hennen G, Igout A and
394 Heinen E. Lymphoid cell apoptosis induced by trophoblastic cells: a model of active
395 foeto-placental tolerance. *J Immunol Methods* 1999;**1-2**;185-196.
- 396 Daher S, de Arruda G, Denardi K, Blotta MH, Mamoni RL, Reck AP, Camano L and Mattar R.
397 Cytokines in recurrent pregnancy loss. *J Reprod Immunol* 2004;**1-2**;151-157.

398 Dalton CF, Laird SM, Estdale SE, Saravelos HG and Li TC. Endometrial protein PP14 and CA-125
399 in recurrent miscarriage patients; correlation with pregnancy outcome. *Hum Reprod*
400 1998;**11**;3197-3202.

401 Dealtry GB, O'Farrell MK and Fernandez N. The Th2 cytokine environment of the placenta. *Int*
402 *Arch Allergy Immunol* 2000;**2**;107-119.

403 Dong C, Davis RJ and Flavell RA. MAP kinases in the immune response. *Annu Rev Immunol*
404 2002;**55**-72.

405 Green DR and Ferguson TA. The role of Fas ligand in immune privilege. *Nat Rev Mol Cell Biol*
406 2001;**12**;917-924.

407 Guleria I, Khosroshahi A, Ansari MJ, Habicht A, Azuma M, Yagita H, Noelle RJ, Coyle A, Mellor
408 AL, Khoury SJ, *et al.* A critical role for the programmed death ligand 1 in fetomaternal
409 tolerance. *J Exp Med* 2005;**2**;231-237.

410 Holmstrom TH, Schmitz I, Soderstrom TS, Poukkula M, Johnson VL, Chow SC, Krammer PH and
411 Eriksson JE. MAPK/ERK signaling in activated T cells inhibits CD95/Fas-mediated apoptosis
412 downstream of DISC assembly. *Embo J* 2000;**20**;5418-5428.

413 Ish-Shalom E, Gargir A, Andre S, Borovsky Z, Ochanuna Z, Gabius HJ, Tykocinski ML and
414 Rachmilewitz J. alpha2,6-Sialylation promotes binding of placental protein 14 via its
415 Ca²⁺-dependent lectin activity: insights into differential effects on CD45RO and CD45RA T
416 cells. *Glycobiology* 2006;**3**;173-183.

417 Jerzak M and Bischof P. Apoptosis in the first trimester human placenta: the role in maintaining
418 immune privilege at the maternal-foetal interface and in the trophoblast remodelling. *Eur J*
419 *Obstet Gynecol Reprod Biol* 2002;**2**;138-142.

420 Lee CL, Chiu PC, Lam KK, Chan RW, Chu IK, Koistinen R, Koistinen H, Seppala M, Lee KF and
421 Yeung WS. Glycodelin-A modulates cytokine production of peripheral blood natural killer cells.
422 *Fertil Steril* 2010;**2**;769-771.

423 Lee CL, Pang PC, Yeung WS, Tissot B, Panico M, Lao TT, Chu IK, Lee KF, Chung MK, Lam KK,
424 *et al.* Effects of differential glycosylation of glycodelins on lymphocyte survival. *J Biol Chem*
425 2009;**22**;15084-15096.

426 Loke YW, King A and Burrows TD. Decidua in human implantation. *Hum Reprod* 1995:14-21.

427 Luppi P. How immune mechanisms are affected by pregnancy. *Vaccine* 2003;**24**;3352-3357.

428 Mainiero F, Gismondi A, Soriani A, Cippitelli M, Palmieri G, Jacobelli J, Piccoli M, Frati L and
429 Santoni A. Integrin-mediated ras-extracellular regulated kinase (ERK) signaling regulates
430 interferon gamma production in human natural killer cells. *J Exp Med* 1998;**7**;1267-1275.

431 Mellor AL, Chandler P, Lee GK, Johnson T, Keskin DB, Lee J and Munn DH. Indoleamine
432 2,3-dioxygenase, immunosuppression and pregnancy. *J Reprod Immunol* 2002;**1-2**;143-150.

433 Michimata T, Tsuda H, Sakai M, Fujimura M, Nagata K, Nakamura M and Saito S. Accumulation of
434 CRTH2-positive T-helper 2 and T-cytotoxic 2 cells at implantation sites of human decidua in a
435 prostaglandin D(2)-mediated manner. *Mol Hum Reprod* 2002;**2**;181-187.

436 Mishan-Eisenberg G, Borovsky Z, Weber MC, Gazit R, Tykocinski ML and Rachmilewitz J.
437 Differential regulation of Th1/Th2 cytokine responses by placental protein 14. *J Immunol*
438 2004;**9**;5524-5530.

439 Mosmann TR, Cherwinski H, Bond MW, Giedlin MA and Coffman RL. Two types of murine helper
440 T cell clone. I. Definition according to profiles of lymphokine activities and secreted proteins. *J*
441 *Immunol* 1986;**7**;2348-2357.

442 Nagata K, Tanaka K, Ogawa K, Kemmotsu K, Imai T, Yoshie O, Abe H, Tada K, Nakamura M,
443 Sugamura K, *et al.* Selective expression of a novel surface molecule by human Th2 cells in
444 vivo. *J Immunol* 1999;**3**;1278-1286.

445 Newton PJ, Weller IV, Katz DR and Chain BM. Autologous apoptotic T cells interact with dendritic
446 cells, but do not affect their surface phenotype or their ability to induce recall immune
447 responses. *Clin Exp Immunol* 2003;**1**;50-58.

448 Ordonez L, Bernard I, L'Faqihi-Olive FE, Tervaert JW, Damoiseaux J and Saoudi A. CD45RC
449 isoform expression identifies functionally distinct T cell subsets differentially distributed
450 between healthy individuals and AAV patients. *PLoS One* 2009;**4**;e5287.

451 Rachmilewitz J, Borovsky Z, Riely GJ, Miller R and Tykocinski ML. Negative regulation of T cell
452 activation by placental protein 14 is mediated by the tyrosine phosphatase receptor CD45. *J*
453 *Biol Chem* 2003;**16**;14059-14065.

454 Ramsdell F, Seaman MS, Miller RE, Picha KS, Kennedy MK and Lynch DH. Differential ability of

455 Th1 and Th2 T cells to express Fas ligand and to undergo activation-induced cell death. *Int*
456 *Immunol* 1994;**10**;1545-1553.

457 Riittinen L, Julkunen M, Seppala M, Koistinen R and Huhtala ML. Purification and characterization
458 of endometrial protein PP14 from mid-trimester amniotic fluid. *Clin Chim Acta* 1989;**1**;19-29.

459 Roberts AI, Devadas S, Zhang X, Zhang L, Keegan A, Greeneltch K, Solomon J, Wei L, Das J, Sun
460 E, *et al.* The role of activation-induced cell death in the differentiation of T-helper-cell subsets.
461 *Immunol Res* 2003;**3**;285-293.

462 Runic R, Lockwood CJ, Ma Y, Dipasquale B and Guller S. Expression of Fas ligand by human
463 cytotrophoblasts: implications in placentation and fetal survival. *J Clin Endocrinol Metab*
464 1996;**8**;3119-3122.

465 Saito S, Sakai M, Sasaki Y, Tanebe K, Tsuda H and Michimata T. Quantitative analysis of peripheral
466 blood Th0, Th1, Th2 and the Th1:Th2 cell ratio during normal human pregnancy and
467 preeclampsia. *Clin Exp Immunol* 1999a;**3**;550-555.

468 Saito S, Tsukaguchi N, Hasegawa T, Michimata T, Tsuda H and Narita N. Distribution of Th1, Th2,
469 and Th0 and the Th1/Th2 cell ratios in human peripheral and endometrial T cells. *Am J Reprod*
470 *Immunol* 1999b;**4**;240-245.

471 Salim R, Miel J, Savvas M, Lee C and Jurkovic D. A comparative study of glycodeclin
472 concentrations in uterine flushings in women with subseptate uteri, history of unexplained
473 recurrent miscarriage and healthy controls. *Eur J Obstet Gynecol Reprod Biol* 2007;**1**;76-80.

474 Schmittgen TD and Livak KJ. Analyzing real-time PCR data by the comparative C(T) method. *Nat*
475 *Protoc* 2008;**6**;1101-1108.

476 Scholz C, Toth B, Brunnhuber R, Rampf E, Weissenbacher T, Santoso L, Friese K and Jeschke U.
477 Glycodelin A induces a tolerogenic phenotype in monocyte-derived dendritic cells in vitro. *Am*
478 *J Reprod Immunol* 2008;**6**;501-512.

479 Seppala M, Koistinen H, Koistinen R, Chiu PC and Yeung WS. Glycosylation related actions of
480 glycodelin: gamete, cumulus cell, immune cell and clinical associations. *Hum Reprod Update*
481 2007;**3**;275-287.

482 Seppala M, Taylor RN, Koistinen H, Koistinen R and Milgrom E. Glycodelin: a major lipocalin
483 protein of the reproductive axis with diverse actions in cell recognition and differentiation.
484 *Endocr Rev* 2002;**4**;401-430.

485 Soni C and Karande AA. Glycodelin A suppresses the cytolytic activity of CD8+ T lymphocytes.
486 *Mol Immunol* 2010;**15**;2458-2466.

487 Straszewski-Chavez SL, Abrahams VM and Mor G. The role of apoptosis in the regulation of
488 trophoblast survival and differentiation during pregnancy. *Endocr Rev* 2005;**7**;877-897.

489 Sundarraj S, Mukhopadhyay D and Karande AA. Glycodelin A triggers mitochondrial stress and
490 apoptosis in T cells by a mechanism distinct and independent of TCR signaling. *Mol Immunol*
491 2008;**8**;2391-2400.

492 Tee MK, Vigne JL, Yu J and Taylor RN. Natural and recombinant human glycodelin activate a

493 proapoptotic gene cascade in monocyte cells. *J Leukoc Biol* 2008;4;843-852.

494 Trowsdale J and Betz AG. Mother's little helpers: mechanisms of maternal-fetal tolerance. *Nat*
495 *Immunol* 2006;3;241-246.

496 Vassiliadou N and Bulmer JN. Expression of CD69 activation marker by endometrial granulated
497 lymphocytes throughout the menstrual cycle and in early pregnancy. *Immunology*
498 1998;3;368-375.

499 Wang ZQ, Wu DC, Huang FP and Yang GY. Inhibition of MEK/ERK 1/2 pathway reduces
500 pro-inflammatory cytokine interleukin-1 expression in focal cerebral ischemia. *Brain Res*
501 2004;1;55-66.

502 Yaniv E, Borovsky Z, Mishan-Eisenberg G and Rachmilewitz J. Placental protein 14 regulates
503 selective B cell responses. *Cell Immunol* 2003;2;156-163.

504 Yeung WS, Lee KF, Koistinen R, Koistinen H, Seppala M and Chiu PC. Effects of glycodefins on
505 functional competence of spermatozoa. *J Reprod Immunol* 2009;1-2;26-30.

506

507

508 **Figure legends**

509 **Figure 1 Effect of GdA on T-helper type-1 and type-2 cells population.** T-helper cells (5×10^5)
510 were incubated with 0.01, 0.1 and 1 $\mu\text{g}/\text{mL}$ of GdA for 48 hours. Cells were stained with CD4-FITC
511 and CD294-PE for immunophenotyping of Th-1 and Th-2 cells. Data are mean \pm SEM, N=9, *
512 $P < 0.05$, P values are shown for significant differences as compared to control.

513

514 **Figure 2 Effect of glycodeilins on cell death of T-helper, Th-1 and Th-2 cells.** T-helper, Th-1 and
515 Th-2 cells (5×10^5) were incubated with 0.01, 0.1 and 1 $\mu\text{g}/\text{mL}$ of GdA, -S, -C and deglycosylated
516 glycodeilin (De-Gd) for 48 hours. Viable, necrotic and apoptotic cells were quantified by bivariate
517 Yo-Pro®-1/PI flow cytometry. Cells without stain were viable. Cells labeled with Yo-Pro®-1 only
518 were apoptotic cells. Cells labeled with Yo-Pro®-1 and PI were necrotic cells. Data are mean \pm
519 SEM, N=5 (T-helper cells), N=9 (Th-1 and Th-2 cells), * $P < 0.05$, P values are shown for significant
520 differences as compared to control.

521

522 **Figure 3 Effect of GdA on ERK activation in T-helper, Th-1 and Th-2 cells. (A) T-helper, (B)**
523 Th-1 and Th-2 cells (1×10^6) were incubated with 1 $\mu\text{g}/\text{mL}$ of GdA for 6 hours followed by PHA
524 stimulation for 30 minutes. Protein expressions were determined by Western blotting. Representative
525 blots are shown. The ERK/pERK protein bands were measured by densitometry. Data are mean \pm
526 SEM, N=4. * $P < 0.05$, P values are shown for significant differences as compared to control. The

527 density values are present as relative quantities = (Density of glycodelin treated cells)/(Density of
528 control).

529 **Table 1. Effect of Glycodelins on viability of T-helper cells in XTT assay.**

Glycodelin ($\mu\text{g/mL}$)	Suppression Index (S.I.)			
	GdA	GdS	GdC	De-Gd
0.001	88.98 ± 6.23	100.07 ± 1.54	102.22 ± 1.08	106.60 ± 2.42
0.01	70.16 ± 11.38	99.39 ± 1.60	102.01 ± 1.75	103.18 ± 2.64
0.1	$54.62 \pm 9.26^*$	101.56 ± 2.30	103.28 ± 2.03	100.57 ± 2.66
1	$53.67 \pm 8.94^*$	105.92 ± 3.94	104.94 ± 3.30	105.61 ± 6.16

530 T-helper cells (3×10^4) were incubated with 0.001, 0.01, 0.1 and $1 \mu\text{g/mL}$ of GdA, -S, -C and
531 deglycosylated glycodelin (De-Gd) for 48 hours. XTT labeling mixture was added 12 hr before
532 measurement. Data are mean \pm SEM, N=4, * $P < 0.05$, P values are shown for significant differences
533 as compared to control. Suppression index (%) = (Absorbance of Gd - Absorbance of
534 blank)/(Absorbance of control - Absorbance of blank) \times 100%

535 **Table 2. Effect of GdA on Fas/FasL mRNA expression of T-helper, Th-1 and Th-2 cells.**

GdA ($\mu\text{g/mL}$)	Surface expression (%)			mRNA expression (Relative expression)	Soluble component secretion (pg/mL)
	T-helper cells	Th-1 cells	Th-2 cells	T-helper cells	T-helper cells
<u>Fas</u>					
0	34.10 \pm 1.41	33.17 \pm 3.80	16.98 \pm 1.54	1	50.98 \pm 2.72
0.01	41.70 \pm 5.07	-	-	1.36 \pm 0.44	52.93 \pm 7.32
0.1	41.89 \pm 5.67	-	-	1.45 \pm 0.27	63.04 \pm 6.19
1	45.26 \pm 2.17 *	54.78 \pm 3.97 *	19.43 \pm 1.99	2.34 \pm 0.62 *	54.28 \pm 14.58
<u>FasL</u>					
0	22.50 \pm 3.61	-	-	1	undetectable
0.01	25.51 \pm 4.48	-	-	0.82 \pm 0.15	undetectable
0.1	24.46 \pm 4.98	-	-	0.87 \pm 0.12	undetectable
1	24.33 \pm 4.16	-	-	1.80 \pm 0.86	undetectable

536 T-helper, Th-1 and Th-2 cells (5×10^5) were incubated with 0.01, 0.1 and 1 $\mu\text{g/mL}$ of GdA for 48 hours. Fas and FasL surface expression were determined by
537 flow cytometry. Fas and FasL mRNA expression was quantified by qPCR. sFas and sFasL secretion to the culture medium were determined by ELISA. Data
538 are mean \pm SEM, N=5, * $P < 0.05$, P values are shown for significant differences as compared to control. The level of sFasL was undetectably low. Relative
539 expression = (RQ of glycodelin treated cells - RQ of negative control)/(RQ of control treated cells - RQ of negative control)

540 **Table 3. Caspase-3, -8 and -9 activities of GdA treated T-helper cells.**


GdA (µg/mL)	T-helper cell	Relative activity (%)							
		Caspase-3		Caspase-8			Caspase-9		
		Th-1 cell	Th-2 cell	T-helper cell	Th-1 cell	Th-2 cell	T-helper cell	Th-1 cell	Th-2 cell
0.01	112.25 ± 4.60*	-	-	115.03 ± 12.19	-	-	111.34 ± 11.22	-	-
0.1	116.83 ± 6.70*	-	-	111.45 ± 10.22	-	-	123.22 ± 8.22	-	-
1	151.25 ± 24.66*	163.89 ± 8.18*	110.33 ± 4.69	120.10 ± 11.85*	119.11 ± 6.81	107.40 ± 3.51	130.34 ± 10.75*	133.72 ± 10.24*	95.96 ± 14.91

541 T-helper, Th-1 and Th-2 cells (1×10^6) were incubated with 0.01, 0.1 and 1 µg/mL of GdA for 48 hours. Activities of caspase-3, -8 and -9 were determined by
542 caspase activity assay (N=6). Data are mean ± SEM, * P<0.05, P values are shown for significant differences as compared to control. Caspase activities were
543 expressed as relative activity (%) = (Absorbance of glycodelin-treated cells - Absorbance of blank) / (Absorbance of control cells - Absorbance of blank) ×
544 100%.

545 **Figure 1.**


546

Control




CD4+CD294⁻ (Th-1): 95.53 ± 0.27%
CD4+CD294⁺ (Th-2): 1.05 ± 0.07%
Th-1/Th-2: 94.33 ± 6.33

GdA (0.01 μg/mL)




CD4+CD294⁻ (Th-1): 95.26 ± 0.39%
CD4+CD294⁺ (Th-2): 1.15 ± 0.08%
Th-1/Th-2: 86.42 ± 6.56

GdA (0.1 μg/mL)



CD4+CD294⁻ (Th-1): 95.02 ± 0.42%
CD4+CD294⁺ (Th-2): 1.13 ± 0.07%
Th-1/Th-2: 87.08 ± 5.99

GdA (1 μg/mL)




CD4+CD294⁻ (Th-1): 94.98 ± 0.34%
CD4+CD294⁺ (Th-2): 1.34 ± 0.08% *
Th-1/Th-2: 72.56 ± 4.16 *

547

548

549 **Figure 2.**
550



551
552

		Viable	Apoptosis	Necrosis	
T-helper cell	Control	91.33 ± 1.05	3.56 ± 0.16	5.08 ± 0.67	
	GdA	0.01 µg/mL	89.45 ± 0.24	3.98 ± 0.28	6.57 ± 0.48
		0.1 µg/mL	71.82 ± 0.33 *	15.59 ± 0.36 *	12.60 ± 0.16 *
		1 µg/mL	69.34 ± 4.81 *	15.74 ± 3.82 *	14.69 ± 2.09 *
	GdS	1 µg/mL	92.40 ± 0.96	3.83 ± 0.50	3.65 ± 0.58
	GdC	1 µg/mL	92.52 ± 0.80	3.51 ± 0.51	3.75 ± 0.54
De-Gd	1 µg/mL	92.09 ± 0.91	3.42 ± 0.36	4.33 ± 0.73	
Th-1 Cell	Control	90.13 ± 0.90	3.19 ± 0.57	6.68 ± 0.90	
	GdA	1 µg/mL	68.82 ± 4.58 *	14.44 ± 1.89 *	16.73 ± 3.87 *
Th-2 Cell	Control	80.38 ± 1.29	7.86 ± 1.22	11.63 ± 1.55	
	GdA	1 µg/mL	76.64 ± 2.72	9.55 ± 1.58	13.89 ± 2.01

553 Figure 3.

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591


592

593

594

595

596

A**B**