



<b>Title</b>	<b>The antioxidative effect of propofol on angiotensin II-induced apoptosis in neonatal rat cardiomyocytes</b>
<b>Author(s)</b>	<b>Wong, K; Leung, YM; Lu, DY; Cheng, TH</b>
<b>Citation</b>	<b>The 15th World Congress on Heart Disease, International Academy of Cardiology Annual Scientific Sessions 2010, Vancouver, B.C., Canada, 24-27 July 2010.</b>
<b>Issued Date</b>	<b>2010</b>
<b>URL</b>	<b><a href="http://hdl.handle.net/10722/140858">http://hdl.handle.net/10722/140858</a></b>
<b>Rights</b>	<b>Creative Commons: Attribution 3.0 Hong Kong License</b>

**THE ANTI-OXIDATIVE EFFECT OF PROPOFOL ON ANGIOTENSIN-II INDUCED APOPTOSIS IN NEONATAL RAT CARDIOMYOCYTES**

**K.L. Wong**<sup>1</sup>, C.W. Cheung<sup>2</sup>, Y.M. Leung<sup>3</sup>, D.Y. Lu<sup>4</sup>, T.H. Cheng<sup>5</sup>

<sup>1</sup>Department of Anesthesiology, and Institute of Clinical Medical Sciences, and Research Group of Cardiovascular Biology, China Medical University and Hospital, Taichung, Taiwan, <sup>2</sup>Department of Anesthesiology, Li Ka Shing Faculty of Medicine, University of Hong Kong, Hong Kong, China, <sup>3</sup>Graduate Institute of Neural and Cognitive Sciences, College of Life Sciences, China Medical University, Taichung, Taiwan, <sup>4</sup>Graduate Institute of Neural and Cognitive Sciences, College of Life Sciences, China Medical University, Taichung, Taiwan, <sup>5</sup>Department of Biological Science and Technology, College of Life Sciences, China Medical University, Taichung, Taiwan

Propofol (2,6-diisopropylphenol), an intravenous sedative Vhypnotic agent popular for sedation, has been found to be effective in protecting against pathological states characterized by an increase in basal rate of reactive oxygen species (ROS) production in hearts, but the cardioprotective mechanism is not well established. Angiotensin-II (Ang-II), an important neurohormonal factor during heart failure, can induce cardiomyocyte apoptosis which has an important role in the transition from compensatory cardiac remodeling to heart failure. In the present study, we evaluated the effects of propofol on Ang-II-mediated cardiomyocyte apoptosis. Cultured cardiomyocytes from neonatal rats were stimulated with Ang-II. Apoptosis was evaluated by measuring caspase 3 activity and by TdT-mediated dUTP nick-end labeling (TUNEL) method. It was found that incubation with Ang-II (0.1 micromolar) for 48 h increased cardiomyocyte apoptosis. Administration of propofol (3-10 micromolar) significantly decreased this Ang-II-induced apoptosis. To further investigate the underlying mechanisms, the quantity of cleaved caspase-3, cytosol cytochrome c release, BcL-xL expression, and ROS generation were examined. These results suggest that propofol abates cardiomyocytes from Ang II-induced apoptosis possibly via reduced the quantity of cleaved caspase-3, and cytosol cytochrome c, and increased BcL-xL expression, and inhibiting the increased ROS generation. In addition, propofol was found to increase the Akt phosphorylation in cardiomyocytes. The siRNA transfection for Akt significantly reduced propofol-induced Akt phosphorylation and propofol's protective effect. Our data provide the first evidence that propofol prevents Ang-II-induced apoptosis, suggesting that propofol may provide a new therapeutic target for the prevention of the cardiac remodeling process.