



Title	Sex determining region Y-box 2 (SOX2) is a potential cell-lineage gene highly expressed in the pathogenesis of squamous cell carcinomas of the lung
Author(s)	Yuan, P; Kadara, H; Behrens, C; Tang, X; Woods, D; Solis, LM; Huang, J; Spinola, M; Dong, W; Yin, G; Fujimoto, J; Kim, E; Xie, Y; Girard, L; Moran, C; Hong, WK; Minna, JD; Wistuba, II
Citation	Plos One, 2010, v. 5 n. 2
Issued Date	2010
URL	http://hdl.handle.net/10722/146595
Rights	

Sex Determining Region Y-Box 2 (SOX2) Is a Potential Cell-Lineage Gene Highly Expressed in the Pathogenesis of Squamous Cell Carcinomas of the Lung

Ping Yuan^{1,9}, Humam Kadara^{2,9}, Carmen Behrens², Ximing Tang², Denise Woods¹, Luisa M. Solis¹, Jiaoti Huang⁴, Monica Spinola⁵, Wenli Dong³, Guosheng Yin³, Junya Fujimoto², Edward Kim², Yang Xie⁶, Luc Girard⁵, Cesar Moran¹, Waun Ki Hong², John D. Minna^{5,7,8}, Ignacio I. Wistuba^{1,2*}

1 Department of Pathology, The University of Texas MD Anderson Cancer Center, Houston, Texas, United States of America, **2** Department of Thoracic/Head and Neck Medical Oncology, The University of Texas MD Anderson Cancer Center, Houston, Texas, United States of America, **3** Department of Biostatistics, The University of Texas MD Anderson Cancer Center, Houston, Texas, United States of America, **4** Department of Pathology and Laboratory Medicine, David Geffen School of Medicine, The University of California Los Angeles, Los Angeles, California, United States of America, **5** The Hamon Center for Therapeutic Oncology, The University of Texas Southwestern Medical Center, Dallas, Texas, United States of America, **6** Department of Biostatistics, The University of Texas Southwestern Medical Center, Dallas, Texas, United States of America, **7** Department of Internal Medicine, The University of Texas Southwestern Medical Center, Dallas, Texas, United States of America, **8** Department of Pharmacology, The University of Texas Southwestern Medical Center, Dallas, Texas, United States of America

Abstract

Background: Non-small cell lung cancer (NSCLC) represents the majority (85%) of lung cancers and is comprised mainly of adenocarcinomas and squamous cell carcinomas (SCCs). The sequential pathogenesis of lung adenocarcinomas and SCCs occurs through dissimilar phases as the former tumors typically arise in the lung periphery whereas the latter normally arise near the central airway.

Methodology/Principal Findings: We assessed the expression of SOX2, an embryonic stem cell transcriptional factor that also plays important roles in the proliferation of basal tracheal cells and whose expression is restricted to the main and central airways and bronchioles of the developing and adult mouse lung, in NSCLC by various methodologies. Here, we found that SOX2 mRNA levels, from various published datasets, were significantly elevated in lung SCCs compared to adenocarcinomas (all $p < 0.001$). Moreover, a previously characterized *OCT4/SOX2/NANOG* signature effectively separated lung SCCs from adenocarcinomas in two independent publicly available datasets which correlated with increased SOX2 mRNA in SCCs. Immunohistochemical analysis of various histological lung tissue specimens demonstrated marked nuclear SOX2 protein expression in all normal bronchial epithelia, alveolar bronchiolization structures and premalignant lesions in SCC development (hyperplasia, dysplasia and carcinoma *in situ*) and absence of expression in all normal alveoli and atypical adenomatous hyperplasias. Moreover, SOX2 protein expression was greatly higher in lung SCCs compared to adenocarcinomas following analyses in two independent large TMA sets (TMA set I, $n = 287$; TMA set II, $n = 511$ both $p < 0.001$). Furthermore, amplification of SOX2 DNA was detected in 20% of lung SCCs tested ($n = 40$) and in none of the adenocarcinomas ($n = 17$).

Conclusions/Significance: Our findings highlight a cell-lineage gene expression pattern for the stem cell transcriptional factor SOX2 in the pathogenesis of lung SCCs and suggest a differential activation of stem cell-related pathways between squamous cell carcinomas and adenocarcinomas of the lung.

Citation: Yuan P, Kadara H, Behrens C, Tang X, Woods D, et al. (2010) Sex Determining Region Y-Box 2 (SOX2) Is a Potential Cell-Lineage Gene Highly Expressed in the Pathogenesis of Squamous Cell Carcinomas of the Lung. PLoS ONE 5(2): e9112. doi:10.1371/journal.pone.0009112

Editor: Andy T. Y. Lau, University of Minnesota, United States of America

Received: December 4, 2009; **Accepted:** January 17, 2010; **Published:** February 9, 2010

Copyright: © 2010 Yuan et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was supported in part by grants from the Department of Defense (W81XWH-04-1-0142 and W81XWH-07-1-03060 to W.K.H., J.D.M., I.I.W.) and by the Specialized Program of Research Excellence in Lung Cancer Grant P50CA70907 (J.D.M., I.I.W.) and Cancer Center Support Grant CA-16672 from the National Cancer Institute. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: iiwistuba@mdanderson.org

⁹ These authors contributed equally to this work.

Introduction

Lung cancer continues to be the leading cause of cancer-related deaths in the United States and worldwide with over one million deaths each year [1]. The majority of lung cancers (85%) are NSCLCs that include SCCs and adenocarcinomas [2]. The pathogenesis of NSCLC involves the accumulation of genetic and epigenetic alterations in a long multi-step process due in part to

chronic exposure to carcinogens such as tobacco smoke. Few early changes that occur during NSCLC pathogenesis have been identified. For example, mutations in *KRAS* [3] and in the epidermal growth factor receptor (*EGFR*) [4] typically occur early in the development of lung adenocarcinomas, whereas amplification of *EGFR* and *PI3KCA* [2] and epigenetic inactivation of the p16 tumor suppressor [5] are more frequent in SCC pathogenesis relative to adenocarcinomas. Moreover, the developmental

transcription factor Thyroid transcriptional factor 1 (*TTF1*) has been shown to be a lineage-survival oncogene over-expressed and amplified in lung adenocarcinoma development [6]. Despite recent progress in the delineation of cellular pathways aberrantly modulated in NSCLC, our understanding of the molecular changes occurring early in NSCLC pathogenesis is still lacking.

Given that the embryonic stem cell *SOX2* transcriptional factor [7] plays important roles in tracheal epithelial cells [8] and is only expressed in the main airways and non-branching bronchioles in the developing and adult mouse lung [9], we hypothesized that it may be a cell-lineage gene highly and specifically expressed in SCCs that originate from central and upper airway and bronchial epithelial cells relative to adenocarcinomas that typically arise from the lung periphery [10]. In this study, we found that *SOX2* mRNA was largely higher in lung SCCs relative to adenocarcinomas from various microarray datasets. Moreover, when we analyzed an *OCT4/SOX2/NANOG* stem cell gene signature previously characterized by Boyer *et al.* [11] in independent publicly available NSCLC datasets, the signature was found to effectively discriminate both major NSCLC subtypes. In addition, we found marked *SOX2* protein expression only in the pathogenesis of SCC which was greatly higher in SCCs relative

to lung adenocarcinomas following analyses of two independent and large tissue microarray (TMA) sets. Lastly, amplification of *SOX2* DNA assessed by quantitative PCR (qPCR) was evident in 20% of lung SCCs studied (n=40) and was absent in all adenocarcinoma cases tested (n=17).

Results

Elevated Expression of *SOX2* mRNA in Lung SCCs Relative to Adenocarcinomas in Various Microarray Datasets

Analysis of *SOX2* mRNA expression in NSCLC samples of various publicly available datasets revealed the significant elevated expression of this stem cell-related transcriptional factor in lung SCCs compared to adenocarcinomas (all $p < 0.001$) (Figure 1A). This differential expression pattern was also evident when microarray analysis was performed on FFPE tissue sections of NSCLC ($p < 0.001$) (Figure S1).

A Previously Characterized *OCT4/SOX2/NANOG* Signature Effectively Discriminates Lung SCCs from Adenocarcinomas

Since *SOX2* mRNA levels were largely higher in SCCs than adenocarcinomas of the lung, we questioned whether a *SOX2*-




Figure 1. Increased expression of *SOX2* mRNA in lung SCCs relative to adenocarcinomas. **A.** Normalized centered *SOX2* mRNA expression levels downloaded from four published microarray data sets as described in the Methods. The lead author of each published microarray cohort data is indicated in each panel. The number of analyzed samples is indicated below each column bar. p -values represent statistical significance assessed by independent two-sided t -tests. **B.** Hierarchical cluster analysis with average linkage of the expression of the previously characterized *OCT4/SOX2/NANOG* signature [11] using present probe sets features in the array platforms of the datasets by Bhattacherjee *et al.* [12] (**B**), and Bild *et al.* [13] (**C**). Data are represented in a matrix format in which individual rows represent single gene features and columns represent experiments. High or low gene expression levels are indicated by red or green color, respectively as indicated by the \log_2 transformed scale bars. **D.** Principal component analysis of the signature in the Bild *et al.* dataset using metric centered correlation. doi:10.1371/journal.pone.0009112.g001

related signature can discriminate between the two subtypes of NSCLC. Using only gene features represented in the published dataset platforms, the previously characterized *OCT4/SOX2/NANOG* signature by Boyer *et al.* [11] was analyzed alone using human gene expression data from the studies by Bhattacharjee *et al* and Bild *et al* [12,13]. In both datasets, the *OCT4/SOX2/NANOG* signature effectively separated lung tumors based on histology where most SCCs clustered alone (Figures 1B and 1C). In hierarchical cluster analysis of the Bild dataset, 43 out of 53 SCCs clustered together (Figure 1C) ($p = 4.3 \times 10^{-10}$ of the χ -test). The sensitivity and specificity for the classification of SCCs and lung adenocarcinomas were 0.81 and 0.78, respectively (Table S1). Moreover, the separation of SCCs from lung adenocarcinomas by the signature was also evident by principal component analysis in three-dimensional space (Figure 1D). In addition, *SOX2* expression in the cluster analyses was higher in lung SCCs relative to adenocarcinomas. Moreover, the expression of *SOX2* mRNA correlated with that of genes such as fibroblast growth factor receptor 2 (*FGFR2*) and parathyroid hormone-like hormone (*pTHLH*) in cluster analyses of both independent datasets (Figures 1B and 1C). In an attempt to validate these findings, we correlated the protein expression of *SOX2* with that of *FGFR2* by TMA immunohistochemistry analysis and found that nuclear *SOX2* protein levels correlated significantly with cytoplasmic *FGFR2* levels in lung SCC histological tissue specimens (Figure S2) but not in adenocarcinomas (data not shown).

Differential Expression of SOX2 Protein between Lung Adenocarcinoma and SCC Development Phases As Revealed by Immunohistochemical Analyses

We next attempted to analyze *SOX2* expression at the protein level in histological tissue specimens representing different stages in the pathogenesis of lung adenocarcinomas and SCCs. Strikingly, *SOX2* protein expression was absent in all normal alveoli ($n = 52$) and AAH ($n = 37$) (Figure 2 and Table 1). In sharp contrast, *SOX2* positive expression was mainly nuclear and evident in all normal bronchial epithelia ($n = 52$), all four alveolar bronchiolization structures and in all ($n = 32$) but one cases of dysplasia and carcinoma *in situ* representing the sequence of SCC pathogenesis (Figure 2 and Table 1).

Elevated Expression of SOX2 DNA and Protein in Lung SCCs Relative to Adenocarcinomas

We next sought to analyze *SOX2* protein expression in NSCLC tissue microarrays. The characteristics of the NSCLC patients from which the tissue specimens were obtained are summarized in Table S2. Nuclear *SOX2* levels were largely and statistically significantly higher in lung SCCs ($n = 109$, set I; $n = 177$ set II) relative to the levels in adenocarcinomas (set I, $n = 178$; set II, $n = 334$) (both $p < 0.001$) (Figure 3A). Notably, the median level of *SOX2* protein was 8.2 and 39.7 higher in lung SCCs relative to adenocarcinomas when analyzed in TMA sets I and II, respectively (set I, adenocarcinoma, 30, SCC, 245; set II, adenocarcinoma 5.8,




Figure 2. SOX2 protein is highly expressed in the pathogenesis of lung SCC but not of adenocarcinoma. **A.** Representative photomicrographs displaying the immunohistochemical expression of SOX2 protein in histological tissue sections of normal bronchial epithelia, preneoplastic lesions representing SCC development (Hyperplasia, Dysplasia and carcinoma *in situ*), and in well (Well SCC) and poorly (Poor SCC) differentiated SCCs. **B.** Representative photomicrographs of SOX2 expression in lung parenchyma (LP), atypical adenomatous hyperplasia (AAH), alveolar bronchiolization structures (AB), and in acinar adenocarcinoma and bronchioalveolar carcinoma (BAC).

doi:10.1371/journal.pone.0009112.g002

Table 1. SOX2 expression in normal and preneoplastic lung lesions.

Tissue type	N	Immunohistochemistry Expression	
		Positive	Negative
Normal bronchial epithelia	52	52	0
Normal alveoli	52	0	52
Atypical adenomatous hyperplasia (AAH)	37	0	37
Alveolar bronchiolization	4	4	0
Squamous dysplasia and carcinoma <i>in situ</i>	32	31	1

doi:10.1371/journal.pone.0009112.t001

SCC, 230). In addition and irrespective of tumor histology, SOX2 protein expression was statistically significantly higher in ever compared to never smokers (set I, $p=0.002$; set II, $p<0.001$) (Figure 3B) and was also significantly increased in current or former smokers compared to never smoker NSCLC patients (set I, $p=0.008$; set II, $p<0.001$) (Figure 3C). Furthermore, in analysis

of both TMA sets, SOX2 protein expression was comparable between former and current smoker NSCLC patients.

We also assessed for the levels of SOX2 DNA copy number in 57 NSCLC tumors. We found 20% SCCs had gene copy gain (8/40; 6/29 with high SOX2 protein expression and 2/11 with low SOX2 expression) (Table S3). In contrast, none of the 17 adenocarcinomas tested displayed SOX2 gene copy gain with most cases (83%) exhibiting SOX2 DNA RQs <1 .

Discussion

In this study, we sought to investigate the expression patterns of SOX2 in NSCLC pathogenesis based on its role and function in the developing and adult mouse lung and trachea [8,9]. We found that SOX2 mRNA levels are significantly higher in lung SCCs relative to adenocarcinomas or large-cell lung carcinomas from various published microarray datasets. Moreover, a previously characterized *OCT4/SOX2/NANOG* embryonic stem cell expression signature [11] effectively separated SCCs from lung adenocarcinomas when analyzed in publicly available NSCLC microarray datasets. In addition, we found that SOX2 protein was completely absent in lung adenocarcinoma pathogenesis and highly expressed in SCC development. Lastly, we demonstrated that SOX2 protein expression was largely significantly elevated in lung SCCs relative




Figure 3. Increased SOX2 expression in lung SCCs compared to adenocarcinomas and its association with smoking patterns. Box-plot depicting statistical analysis by the Wilcoxon-rank test of differences in nuclear SOX2 protein score in both TMA sets I and II between lung adenocarcinomas (ADC) and SCCs (A), ever and never NSCLC smokers (B) and between never, former and current smokers (C). doi:10.1371/journal.pone.0009112.g003

to adenocarcinomas following analyses of two independent TMA sets and copy gain of the gene was evident in 20% of lung SCCs studied and absent in adenocarcinomas. These findings entail the possible clinical application of *SOX2* expression as a diagnostic biomarker to discriminate lung SCCs from adenocarcinomas, in an analogous fashion to the potential application of *TTF1* as a biomarker for lung adenocarcinomas but not SCCs [2,6].

It is noteworthy that the *OCT4/SOX2/NANOG* signature is comprised of genes under direct transcriptional control by the *SOX2* transactivating factor [11]. Given the effective separation of lung SCCs and adenocarcinomas by this signature, it is possible that other genes directly or indirectly related to *SOX2* signaling are also dissimilar in expression between both NSCLC subtypes. In accordance and based on analysis of the expression signature in two independent microarray datasets, we validated and demonstrated the correlative expression of both *SOX2* and *FGFR2* protein in lung SCC tissue histological specimens. Interestingly, *FGFR2* has been shown to be critical for the formation of the lung bud, which typically is associated with *SOX2* expression and *TTF1* repression, and mice lacking this gene do not form lungs [9,14,15]. It is plausible to suggest that the reduction in *SOX2* expression in adenocarcinomas of the lung may be evolutionarily conserved.

While completing our study, Bass *et al* reported an amplification of *SOX2* in esophageal (15%) and lung SCCs (23%) [16] assessed by Affymetrix single nucleotide polymorphism (SNP) arrays. Similarly, we found copy gain of the *SOX2* gene in 20% of lung SCCs as revealed by our quantitative PCR analysis. Notably, we also analyzed a set of lung adenocarcinomas and found no significant *SOX2* copy number gain further demonstrating a cell-lineage expression pattern for *SOX2* specifically in lung SCCs. In contrast to the study by Bass *et al*, we also assessed *SOX2* protein levels in lung SCCs and adenocarcinomas using two independent and large tissue microarrays and correlated the expression of *SOX2* protein expression and copy number gain in 57 NSCLC samples. It is note worthy that several cases displayed discordant *SOX2* protein and copy number gain. It is reasonable to suggest that *SOX2* expression may be regulated at different levels; at the DNA, mRNA and protein and that it may be important to assess for its expression at all three levels.

In conclusion, we demonstrated a vast increase in expression of the pluripotent stem cell related transcriptional factor, *SOX2*, in squamous cell carcinomas relative to adenocarcinomas of the lung. Moreover, our findings raise the possibility of the activation of *SOX2*-dependent stem cell-related pathways in squamous cell carcinomas of the lung.

Materials and Methods

Analysis of *SOX2* mRNA in Published Microarray Datasets and Integration of the *OCT4/SOX2/NANOG* Signature

The integrated cancer microarray database and data-mining platform, OncoPrint [17], was utilized to analyze the expression of *SOX2* in publicly available microarray datasets of human lung carcinomas available on-line. *SOX2* mRNA (probe sets 213721_at and 228038_at) was also compared by microarray analysis between formalin-fixed paraffin embedded (FFPE) lung SCCs and adenocarcinomas (unpublished observations). Genes characterizing an *OCT4/SOX2/NANOG* expression signature [11] were compiled and queried using NetAffxTM from Affymetrix (<http://www.affymetrix.com/analysis/index.affx>) to search for corresponding probe set annotations in the HG-U95A and HG-U133 plus 2.0 platforms used in the lung cancer microarray studies by Bhattacharjee *et al.* [12] and Bild *et al.* [13], respectively. The

OCT4/SOX2/NANOG signature gene features were then analyzed alone in published human gene expression data obtained from the reports by Bhattacharjee *et al.* and Bild *et al.* Raw microarray data files from the two published datasets were imported and analyzed using the BRB-ArrayTools v.3.7.0 developed by Dr. Richard Simon and BRB-ArrayTools Development Team [18]. Gene expression data were normalized by Robust multi-array analysis (RMA) in R language environment [19] and median-centered across all samples in each data set before hierarchical clustering analysis. Clustering by average linkage was performed with Cluster 2.11, and results were visualized with TreeView programs (Michael Eisen Laboratory, Lawrence Berkeley National Laboratory and University of California, Berkeley; <http://rana.lbl.gov/EisenSoftware.htm>). Principal component analysis following gene centering was performed using the BRB-ArrayTools software.

Immunohistochemistry Analysis, Human Lung Tissues and Tissue Microarray Sets I and II

Detailed description of the normal and preneoplastic histological tissue specimens analyzed and two TMA sets is available in Methods S1. Cytoplasmic *FGFR2* protein expression was available based on previous analysis [20]. Details of the immunohistochemical analysis for *SOX2* protein expression are also available in Methods S1.

DNA Extraction and Quantitative PCR (qPCR)

Tumor tissues (40 SCCs, 17 adenocarcinomas) were dissected from FFPE Hematoxylin-stained tissue sections using manual microdissection to ensure that tumor cell proportions are greater than 70% for subsequent DNA extraction. Tumor DNA was extracted using the PicoPure DNA extraction Kit (Arcturus, Mt View, CA) according to the manufacturer's instructions. Five μ l of DNA was added to a 20 μ l final volume reaction mixture consisting of 10 μ l Power SYBR[®] Green PCR Master Mix (Applied Biosystems, Foster City, CA) and 0.5 μ mol/l of each of forward and reverse primers which span 102 dinucleotides (613 to 714) of the *SOX2* gene (ID: NM 003106) as follows: 5'- GAACCCCAA-GATGCACAACCTC and 5'-CGCTTAGCCTCGTCGATGAC. β -actin was used as an endogenous reference gene (TaqMan[®] Control Human Genomic DNA, Applied Biosystems) and was amplified as a standard control for calibration. All samples and standard DNA reactions were carried out in triplicates. qPCR was performed using an ABI 7300 Real Time PCR System Sequence (Applied Biosystems) at 50°C for 2 min, 95°C for 10 min, followed by 40 cycles at 95°C for 15 s and 60°C for 1 min. The quantity of the target genes were normalized using the level of the β -actin gene, and expressed as relative quantities (RQ) compared with the value of the Human Genomic DNA. RQ equal or larger than 2 was considered as gene copy gain.

Statistical Analyses

The data were summarized using standard descriptive statistics. The rank-based non-parametric Wilcoxon rank-sum test and the Kruskal-Wallis test were used to assess the statistical significance of the differences in nuclear *SOX2* staining intensity score between lung SCCs and adenocarcinomas and based on tobacco history (ever vs never smokers) and type of smokers (never vs former vs current). All tests were two sided. p-values <0.05 were considered statistically significant.

Supporting Information

Methods S1 Supplementary methods.

Found at: doi:10.1371/journal.pone.0009112.s001 (0.04 MB DOC)

Figure S1 Increased expression of SOX2 mRNA in lung SCCs relative to adenocarcinomas in FFPE NSCLC specimens. SOX2 levels were analyzed from microarray analysis of FFPE NSCLC specimens using the Affymetrix HG-U133A platform. P-values were obtained by the Student's t-test.

Found at: doi:10.1371/journal.pone.0009112.s002 (0.22 MB TIF)

Figure S2 Correlation of expression of SOX2 and FGFR2 protein in lung SCC tissue specimens. SOX2 and FGFR2 protein levels were assessed by immunohistochemistry as described in Methods S1. Assessment of significance in correlation between SOX2 nuclear and FGFR2 cytoplasmic protein levels was performed using the Spearman Rank correlation test.

Found at: doi:10.1371/journal.pone.0009112.s003 (0.31 MB TIF)

Table S1 Separation of 111 NSCLCs from the study of Bild et al. by the OCT4/SOX2/NANOG signature. *P-value was obtained by Fisher's exact test. Sensitivity (probability for an SCC sample to be correctly predicted as SCC)=0.811. Specificity (probability for an ADC sample to be correctly predicted as an ADC)=0.776.

References

- Jemal A, Siegel R, Ward E, Hao Y, Xu J, et al. (2009) Cancer statistics, 2009. *CA Cancer J Clin* 59: 225–249.
- Herbst RS, Heymach JV, Lippman SM (2008) Lung cancer. *N Engl J Med* 359: 1367–1380.
- Westra WH, Baas IO, Hruban RH, Askin FB, Wilson K, et al. (1996) K-ras oncogene activation in atypical alveolar hyperplasias of the human lung. *Cancer Res* 56: 2224–2228.
- Tang X, Varella-Garcia M, Xavier AC, Massarelli E, Ozburn N, et al. (2008) Epidermal growth factor receptor abnormalities in the pathogenesis and progression of lung adenocarcinomas. *Cancer Prev Res (Phila Pa)* 1: 192–200.
- Belinsky SA, Nikula KJ, Palmisano WA, Michels R, Saccomanno G, et al. (1998) Aberrant methylation of p16(INK4a) is an early event in lung cancer and a potential biomarker for early diagnosis. *Proc Natl Acad Sci U S A* 95: 11891–11896.
- Kwei KA, Kim YH, Girard L, Kao J, Pacyna-Gengelbach M, et al. (2008) Genomic profiling identifies TTF1 as a lineage-specific oncogene amplified in lung cancer. *Oncogene* 27: 3635–3640.
- Boiani M, Scholer HR (2005) Regulatory networks in embryo-derived pluripotent stem cells. *Nat Rev Mol Cell Biol* 6: 872–884.
- Que J, Luo X, Schwartz RJ, Hogan BL (2009) Multiple roles for Sox2 in the developing and adult mouse trachea. *Development* 136: 1899–1907.
- Gontan C, de Munck A, Vermeij M, Grosveld F, Tibboel D, et al. (2008) Sox2 is important for two crucial processes in lung development: branching morphogenesis and epithelial cell differentiation. *Dev Biol* 317: 296–309.
- Wistuba II, Gazdar AF (2006) Lung cancer preneoplasia. *Annu Rev Pathol* 1: 331–348.
- Boyer LA, Lee TI, Cole MF, Johnstone SE, Levine SS, et al. (2005) Core transcriptional regulatory circuitry in human embryonic stem cells. *Cell* 122: 947–956.
- Bhattacharjee A, Richards WG, Staunton J, Li C, Monti S, et al. (2001) Classification of human lung carcinomas by mRNA expression profiling reveals distinct adenocarcinoma subclasses. *Proc Natl Acad Sci U S A* 98: 13790–13795.
- Bild AH, Yao G, Chang JT, Wang Q, Potti A, et al. (2006) Oncogenic pathway signatures in human cancers as a guide to targeted therapies. *Nature* 439: 353–357.
- Bellucci S, Grindley J, Emoto H, Itoh N, Hogan BL (1997) Fibroblast growth factor 10 (FGF10) and branching morphogenesis in the embryonic mouse lung. *Development* 124: 4867–4878.
- Min H, Danilenko DM, Scully SA, Bolon B, Ring BD, et al. (1998) Fgf-10 is required for both limb and lung development and exhibits striking functional similarity to Drosophila branchless. *Genes Dev* 12: 3156–3161.
- Bass AJ, Watanabe H, Mermel CH, Yu S, Perner S, et al. (2009) SOX2 is an amplified lineage-survival oncogene in lung and esophageal squamous cell carcinomas. *Nat Genet* 41: 1238–1242.
- Rhodes DR, Yu J, Shanker K, Deshpande N, Varambally R, et al. (2004) ONCOMINE: a cancer microarray database and integrated data-mining platform. *Neoplasia* 6: 1–6.
- Simon R, Lam A, Li MC, Ngan M, Menendez S, et al. (2007) Analysis of Gene Expression Data Using BRB-Array Tools. *Cancer Inform* 3: 11–17.
- Irizarry RA, Bolstad BM, Collin F, Cope LM, Hobbs B, et al. (2003) Summaries of Affymetrix GeneChip probe level data. *Nucleic Acids Res* 31: e15.
- Behrens C, Lin HY, Lee JJ, Raso MG, Hong WK, et al. (2008) Immunohistochemical expression of basic fibroblast growth factor and fibroblast growth factor receptors 1 and 2 in the pathogenesis of lung cancer. *Clin Cancer Res* 14: 6014–6022.

Found at: doi:10.1371/journal.pone.0009112.s004 (0.03 MB DOC)

Table S2 Patient characteristics in tissue microarray sets I and II.

Found at: doi:10.1371/journal.pone.0009112.s005 (0.05 MB DOC)

Table S3 Analysis of SOX2 gene copy gain in lung SCCs and adenocarcinomas and its correlation with its corresponding protein levels. CN, SOX2 gene copy number assessed as relative quantities (RQs) to β -actin.

Found at: doi:10.1371/journal.pone.0009112.s006 (0.03 MB DOC)

Author Contributions

Conceived and designed the experiments: PY HK JDM IIW. Performed the experiments: PY HK CB XT DW LMS MS JF. Analyzed the data: PY HK CB XT DW LMS JH MS WD GY EK YX LG CM JDM IIW. Contributed reagents/materials/analysis tools: JH EK LG CM WKH JDM IIW. Wrote the paper: PY HK CB XT WD WKH JDM IIW.