



Title	Dissemination of pHK01-like incompatibility group IncFII plasmids encoding CTX-M-14 in Escherichia coli from human and animal sources
Author(s)	Ho, PL; Lo, WU; Yeung, MK; Li, Z; Chan, J; Chow, KH; Yam, WC; Tong, AHY; Bao, JYJ; Lin, CH; Lok, S; Chiu, SS
Citation	Veterinary Microbiology, 2012, v. 158 n. 1-2, p. 172-179
Issued Date	2012
URL	http://hdl.handle.net/10722/152771
Rights	NOTICE: this is the author's version of a work that was accepted for publication in Veterinary Microbiology. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Veterinary Microbiology, 2012, v. 158 n. 1-2, p. 172-179. DOI: 10.1016/j.vetmic.2012.02.004

1 Original research paper (VetMic-D-11-6397 revised)

2 **Dissemination of pHK01-like incompatibility group IncFII plasmids encoding CTX-M-**
3 **14 in *Escherichia coli* from human and animal sources, 2002-2010**

4

5 P. L. Ho^{a,b*}, W. U Lo^a, M. K. Yeung^a, Zhen Li^a, Jane Chan^a, K. H. Chow^a, W. C. Yam^a,
6 Amy H.Y. Tong^c, Jessie Y. J. Bao^c, C. H. Lin^c, Si Lok^c, Susan S. Chiu^d

7

8 ^a Department of Microbiology, University of Hong Kong, Pokfulam Road, Pokfulam, Hong
9 Kong, People's Republic of China,

10 ^b Carol Yu Center for Infection, University of Hong Kong, Pokfulam Road, Pokfulam, Hong
11 Kong, People's Republic of China,

12 ^c Genome Research Centre, University of Hong Kong, Pokfulam Road, Pokfulam, Hong
13 Kong, People's Republic of China,

14 ^d Department of Paediatrics and Adolescent Medicine, University of Hong Kong, Pokfulam
15 Road, Pokfulam, Hong Kong, People's Republic of China

16

17 *Corresponding author. Fax: 852-2855-1241; Tel: 852-2255-4892;

18 E-mail: plho@hkucc.hku.hk (P.L. Ho)

19 Word counts: abstract (250 words), body text (3627 words)

20

21

Abstract

22 Few studies have compared CTX-M encoding plasmids identified in different ecological
23 sources. This study aimed to analyze and compare the molecular epidemiology of plasmids
24 encoding CTX-M-14 among strains from humans and animals. The CTX-M-14 encoding
25 plasmids in 160 *E. coli* isolates from animal faecal (14 pigs, 16 chicken, 12 cats, 8 cattle, 5
26 dogs and 3 rodents), human faecal (45 adults and 20 children) and human urine (37 adults)
27 sources in 2002-2010 were characterized by molecular methods. The replicon types of the
28 CTX-M-14 encoding plasmids were IncFII ($n=61$), I1-I γ ($n=24$), other F types ($n=23$), B/O
29 ($n=10$), K ($n=6$), N ($n=3$), A/C ($n=1$), HI1 ($n=1$), HI2 ($n=1$) and nontypeable ($n=30$). The
30 genetic environment, *ISEcp1* - *bla*_{CTX-M-14} - *IS903* was found in 89.7% (52/58), 87.7% (57/65)
31 and 86.5% (32/37) of the animal faecal, human faecal and human urine isolates, respectively.
32 Subtyping of the 61 IncFII incompatibility group plasmids by replicon sequence typing,
33 plasmid PCR-restriction fragment length polymorphism and marker genes (*vac*, *malB*,
34 *eitA/eitC* and *parB/A*) profiles showed that 31% (18/58), 30.6% (20/65) and 37.8% (14/37) of
35 the plasmids originating from animal faecal, human faecal and human urine isolates,
36 respectively, were pHK01-like. These 52 pHK01-like plasmids originated from diverse
37 human (20 faecal isolates from 2002, 2007-2008, 14 urinary isolates from 2004) and animal
38 (all faecal, 1 cattle, 1 chicken, 5 pigs, 9 cats, 1 dogs, 1 rodents from 2008-2010) sources. In
39 conclusion, this study highlights the importance of the IncFII group, pHK01-like plasmids in
40 the dissemination of CTX-M-14 among isolates from diverse sources.

41

42 **Key words:** pets, pigs, chicken, plasmid, extended-spectrum beta-lactamases

43

44 **Introduction**

45 Asia is one of the epicentres of antimicrobial resistance (Hawkey, 2008; Ho et al., 2011c; Ho
46 et al., 2011a). In this part of the world, the CTX-M type extended-spectrum β -lactamases
47 (ESBLs) have spread extensively among bacteria in human, animals and the environment
48 (Hawkey, 2008; Ho et al., 2011a). Many studies have reported the overwhelming dominance
49 of the CTX-M-14 allele among clinical isolates of *Escherichia coli* and *Klebsiella* spp in
50 mainland China, Hong Kong and South Korea (Hawkey, 2008; Ho et al., 2007; Naseer and
51 Sundsfjord, 2011). In general, bacterial strains producing CTX-M-14 are genetically diverse
52 (Ho et al., 2007; Lo et al., 2010; Valverde et al., 2009). Hence, the dissemination of CTX-M-
53 14 has been attributed to conjugative plasmids and other mobile genetic elements rather than
54 clonal expansion (Naseer and Sundsfjord, 2011).

55 In Spain, the spread of CTX-M-14 from 2000 to 2005 was largely due to pRYC105-
56 like plasmids of the IncK incompatibility group disseminated among diverse *E. coli* lineages
57 (Valverde et al., 2009). It has further been shown that pRYC105 is related to the pCT plasmid
58 that has been found in CTX-M-14 producing strains from the United Kingdom, mainland
59 China and Australia (Cottell et al., 2011). Recently, we sequenced the IncFII incompatibility
60 group plasmid, pHK01 encoding CTX-M-14 and showed that it has disseminated widely
61 among *E. coli* isolates causing community-acquired urinary tract infections in women (Ho et
62 al., 2011b). Plasmids closely related to pHK01 have also been identified among *Klebsiella*
63 strains from mainland China (Yi et al., 2010). Besides plasmids, the acquisition and
64 horizontal transmission of *bla*_{CTX-M} genes have been associated with insertion sequences (ISs),
65 putative transposons and class 1 integrons (Canton and Coque, 2006). These elements may
66 have mobilized the *bla*_{CTX-M} gene from its progenitor and may drive the expression of the β -
67 lactamase (Canton and Coque, 2006). In addition, these elements might accumulate
68 additional resistance genes to the *bla*_{CTX-M} flanking regions (Canton and Coque, 2006). It has

69 also been suggested that *ISEcp*, *bla*_{CTX-M} and *IS903* form a putative transposon and this block
70 of genes could disseminate by transposition (Poirel et al., 2005).

71 Although the roles of plasmids and other genetic elements are recognized to be
72 important in the dissemination of the CTX-M genes, few studies have compared the
73 distribution of these mobile genetic elements among strains from humans and animals and
74 their inter-relationship (Garcia-Fernandez et al., 2008). In this study, we investigated the
75 molecular epidemiology of plasmids encoding CTX-M-14 and their *bla*_{CTX-M-14} genetic
76 environment for a collection of *E. coli* isolates collected from diverse human and animal
77 sources between 2002 and 2010.

78

79 **Methods**

80 **Bacterial strains**

81 One hundred and sixty CTX-M-14 producing *E. coli* isolates were studied. The strains were
82 chosen to provide representation from different time periods and animal and human sources.
83 All viable CTX-M-14 producing *E. coli* isolates recovered from four regional antimicrobial
84 resistance surveillances during 2002-2010 were eligible for inclusion (Ho et al., 2007; Ho et
85 al., 2008; Ho et al., 2011a; Lo et al., 2010). These included 37 isolates from female
86 outpatients with urinary tract infections, 65 human faecal isolates (45 adults and 20 children)
87 and 58 faecal isolates from animals (14 pigs, 16 chicken, 12 cats, 8 cattle, 5 dogs and 3
88 rodents) (Ho et al., 2007; Ho et al., 2008; Ho et al., 2011b; Ho et al., 2011a). The human
89 urine isolates were recovered in 2004 from female outpatients with urinary tract infections
90 (age range, 27 –80 years) and all CTX-M-14 isolates in the collection were included (Ho et
91 al., 2007). The 65 human faecal isolates include all of 62 CTX-M-14 isolates identified in a
92 study that examined faecal carriage in hospitalized children and their household contacts in
93 October 2007-September 2008 (Lo et al., 2010) and three (out of seven CTX-M-14 isolates

94 found) isolates from a study that assessed carriage in non-hospitalized individuals in 2002
95 (Ho et al., 2008). The remaining four isolates obtained in the 2002 faecal study were lost
96 during storage and therefore not included. All the animal isolates were obtained during
97 September 2008-August 2010 as part of an ongoing antimicrobial resistance by trained staff
98 in two government departments (Ho et al., 2011a). For cattle and pigs, rectal swabs were
99 obtained from fresh carcasses at a centralized slaughterhouse in Hong Kong (Ho et al.,
100 2011a). Chicken samples were obtained while the animals were temporarily held for
101 inspection before sale at wet markets in Hong Kong. All the cattle were imported from
102 mainland China. The pigs and chicken included animals produced at local farms and those
103 imported from mainland China. Samples from the stray dogs, stray cats and rodents were
104 collected by trained staff at governmental animal management centres. These animals were
105 captured from urban areas from all over Hong Kong. The animals were sampled in batches:
106 chicken (20 animals per batch), cattle (10 animals per batch), pigs (2 to 7 animals per batch),
107 stray cats (1-10 animals batch), stray dogs (1-10 animals per batch) and urban rodents (2 to
108 23 animals per batch). In total, 2106 animals from 298 batches were tested. In 179 batches,
109 at least one animal was found to carry ESBL-producing *E. coli*. A subset of 132 isolates was
110 chosen randomly according to the collection period and batch number for CTX-M PCR and
111 sequencing. The isolates were selected randomly according to the collection period and batch
112 number. This subset included isolates recovered from 84 batches and covered the entire
113 specimen collection periods. Sequencing showed that 58 of the 132 isolates had CTX-M-14.
114 All the 58 CXT-M-14 producing isolates were included (Ho et al., 2011a).

115

116 **Susceptibility testing, conjugation and replicon typing**

117 Antimicrobial susceptibility to amoxicillin-clavulanic acid, amikacin, cefotaxime,
118 ceftazidime, ciprofloxacin, chloramphenicol, cotrimoxazole, gentamicin, nalidixic acid,

119 nitrofurantoin, tetracycline was tested by the disc diffusion method (Clinical and Laboratory
120 Standards Institute, 2011). ESBL production was determined by the double-disc synergy test
121 using cefotaxime, amoxicillin-clavulanic acid and ceftazidime at inter-disc distances (centre
122 to centre) of 20 and 25 mm (Ho et al., 2000). Conjugation was carried out in filters with *E.*
123 *coli* J53Az^r as the recipient. Donor and recipient cells were mixed at 1:10 ratio.
124 Transconjugants were selected on trypticase soy agar plates containing sodium azide (150
125 mg/L; Sigma Chemical Co.) for counterselection and cefotaxime (1 mg/L; Sigma Chemical
126 Co.) to select for plasmid-encoded resistance. For detecting plasmids, bacterial cells were
127 embedded in agarose plugs and disrupted by alkaline lysis. Subsequently, the plasmids were
128 converted to the linear forms by incubation with *Aspergillus oryzae* S1 nuclease (Sigma
129 Chemical Co., St Louis, MO, USA) and were sized by pulsed-field gel electrophoresis (PFGE)
130 as previously described (Ho et al., 2010; Ho et al., 2011b). The replicon types for *E. coli*
131 transconjugants with CTX-M encoding plasmids were determined by a scheme previously
132 described (Carattoli et al., 2005). Eighteen pairs of primers were tested in five multiplex and
133 three simplex PCR assays. The method allowed recognition of the following plasmid
134 incompatibility groups (Inc): FIA, FIB, FIC, FIIA, HI1, HI2, I1-I γ , L/M, N, P, W, T, A/C, K,
135 B/O, X, Y, F. An additional primer pair was used for identification IncFII replicons (Osborn
136 et al., 2000). The FII plasmids were further subtyped by the replicon sequence typing (RST)
137 scheme (Villa et al., 2010). This involved PCR and sequencing of DNA fragments containing
138 the *copA* region of the FII replicon, the iterons-*repE* region of the FIA replicon, the *repB*
139 gene of the FIB replicon and the *copA* region of the FIC replicon. Alleles were assigned to
140 each sequence and the replicon types were expressed according to the FAB formula (Villa et
141 al., 2010). Where a transconjugant could not be obtained, the replicon type for the plasmid
142 encoding CTX-M-14 was determined in the parent strains. In all the isolates, the replicon
143 location in the plasmids was confirmed by hybridization with probes specific for *bla*_{CTX-M} and

144 *rep* amplified by PCR from different samples. For non-transconjugants, S1-PFGE was used
145 to separate all the plasmids and the plasmid carrying the *bla*_{CTX-M} gene was identified by
146 hybridization probe. PCR was used to determine all the plasmid replicons carried by the non-
147 transconjugants. Afterwards, the non-transconjugants were tested by hybridization using
148 probes for all positive replicon types. The *rep* probes that hybridized to the *bla*_{CTX-M} carrying
149 plasmid were used to define the replicon type. The four main phylogenetic groups (A, B1, B2,
150 and D) of the *E. coli* isolates were determined by a multiplex PCR assay (Clermont et al.,
151 2000).

152

153 **PCR-RFLP analysis of IncFII plasmids**

154 The IncFII plasmids were analysed further by a PCR-RFLP scheme previously described (Ho
155 et al., 2011b). The PCR-RFLP scheme involved PCR amplifications of seven loci or regions
156 (designated A, B1, B2, and C to F) with sizes ranging from 1.7 kb to 7.8 kb. Following PCR
157 amplifications, the amplicons were digested with 5 U each of the following restriction
158 enzymes (New England BioLabs) according to the manufacturer's instructions: *Rsa*I (locus
159 A); *Sma*I (locus B1); *Sfc*I (locus B2); *Hha*I (locus C); *Bsm*I (locus D); *Bst*EII (locus E); and
160 *Drd*I (locus F). For each locus, the PCR-RFLP patterns were assigned numbers. Patterns with
161 one or more band differences were assigned different numbers. The profiles for the seven loci
162 were used to designate the pRFLP type. Primers for amplification of the A, B2, C to F loci
163 were those described previously (Ho et al., 2011b). Locus B1 was included in this study to
164 map the region upstream of *bla*_{CTX-M-14}. The primer pair used to amplify locus B1 was:
165 *repA1F* (forward), 5'- CGCTCCTTCTGCGCATTGTAA -3' and *CTX-M-9F* (backward), 5'-
166 CAAAGAGAGTGCAACGGATG -3'(Woodford et al., 2006). The content of the amplified
167 regions in pHK01 were (Ho et al., 2011b): locus A (*finO*, *orf99-102*, *repA2*, *repA3* and
168 *repA1*), locus B1 (*repA1*, *yacABC*, *yadA*, *malB*, *IS903*, *bla*_{CTX-M-14}), locus B2, (*bla*_{CTX-M-14},

169 ISECp1, *eitD*, *eitC*, *eitB*, *eitA*, *orf18-19*, *parB*), locus C (***parB***, ***parA***), locus D (*orf37*, *orf38-*
170 *40*, *ssb*, *orf42-43*, *psiB*, *psiA*, *orf45*), locus E (*traM*, *J*, *Y*, *A*, *L*, *E*, *K*, *B*, *P*, *trbD*), and locus F
171 (*traI*, *X*, *orf97*, *finO*). The primers were designed within the underlined gene or
172 upstream/downstream of the gene in **boldface**. Plasmid with PCR-RFLP patterns identical to
173 the reference plasmid (pHK01) in three or more loci were considered to be pHK01-like. PCR
174 for the pHK01-associated *malB*, *yac*, *eitA*, *eitC* and *parA/B* genes was performed using
175 previously described primers (Ho et al., 2011b).

176

177 **Analysis of *bla*_{CTX-M-14} genetic environment**

178 The genetic environment may be defined as the sequences flanking the open reading frame of
179 the *bla*_{CTX-M-14} gene. This includes insertion elements that may be involved in the
180 mobilization and expression of the *bla*_{CTX-M-14} gene. Based on published studies and
181 sequences deposited in the GenBank database, eight different types of *bla*_{CTX-M-14} genetic
182 environment have been reported (Table 1). For purpose of description, they were denoted
183 types I to VIII. Monoplex PCRs using different primer pair combinations were used to map
184 the genetic environment to one of the recognized types (Table 1).

185

186 **Plasmid sequencing**

187 The Illumina Genome Analyzer Iix was used for sequencing of plasmids at approximately
188 500-fold coverage, as described previously (Ho et al., 2011b). In brief, plasmids were
189 extracted from the transconjugants by using the Qiagen Large Construct kit (Qiagen, Hong
190 Kong). Purified plasmid DNA was fragmented by nebulization. The fragments were
191 amplified and a library was constructed as described previously (Ho et al., 2011c). Based on
192 the qPCR quantified concentration of the barcoded plasmid library, it was diluted to generate
193 approximately 500,000 clusters and seeded with other samples in the same Solexa sample

194 lane. Sequencing run of 76-base pair-end reads was performed according to the
195 manufacturer's recommendations. The Illumina Off-Line Basecaller (version v1.6), WebACT
196 and Geneious Pro (Version 5.0.1, Biomatters Limited, Auckland, New Zealand) softwares
197 were used for bioinformatics analysis (Ho et al., 2011c).

198

199 **Results**

200 In conjugation experiments, the plasmids harbouring CTX-M 14 could be transferred in
201 77.5% (124/160) of the isolates at frequencies of 10^{-6} to 10^{-1} per donor cells. In addition to
202 cefotaxime resistance, the plasmids in 19 transconjugants encoded co-resistance to non- β -
203 lactam antibiotics with profiles involving chloramphenicol, gentamicin, cotrimoxazole and/or
204 tetracycline. These plasmids have origins from 17 faecal (chicken 8, pigs 3, dog 1, humans 5)
205 and two human urine isolates (Supplementary Table S1).

206 Ten different replicons (IncFII, IncFIA, IncFIB, IncI1-I γ , IncB/O, IncK, IncN, IncA/C,
207 IncHI1, IncHI2), either alone or in combinations were found among the plasmids encoding
208 CTX-M-14 (Table 2). The two most common replicon types were IncFII (38.1%, 61/160) and
209 IncI1-I γ (15%, 24/160). The frequencies of IncFII replicon were similar among isolates from
210 the three sources: animal faecal (36.2%), human faecal (38.5%) and human urine (40.5%, chi
211 square for trend, $P = 0.7$). Replicon IncI1-I γ was more common among animal faecal isolates
212 (20.7%) than in human faecal isolates (10.8%) and human urine isolates (13.5%), but the
213 difference was not statistically significant ($P = 0.3$). The replicons of the CTX-M-14
214 encoding plasmids for 19 transconjugants with co-resistance to non- β -lactam antibiotics were
215 as follows: IncFII (n=1), IncFII, FIB (n=2), IncFIB (n=2), IncFII, I1 γ (n=2), IncFII, FIA, FIB
216 (n=1), Inc I1 γ (n=5), IncB/O (n=2), A/C (n=1), HI2 (n=1) and nontypeable (n=2)

217 Overall, 37.5% (60/160) isolates were phylogenetic group D, 22.5% (36/160) were
218 group B2, 20.6% (33/160) were group A and 19.4% (31/160) were group B1. Analysis of

219 plasmid replicon distribution showed that the frequency of IncFII replicon in the four
220 phylogenetic groups were similar. The IncFII frequencies for the virulent groups B2 and D
221 were 47.2% (17/36) and 40% (24/60), respectively; that for the commensal groups A and B1
222 were 39.4% (13/33) and 22.6% (7/31), respectively (virulent vs. commensal groups, $P = 0.1$).

223

224 PCR mapping and sequencing of representative products revealed six of the eight
225 reported types of *bla*_{CTX-M-14} genetic environment (Table 3 and Supplementary Table S2).
226 One hundred forty-one (88.1%) isolates were found to have a genetic environment consisting
227 of an *ISEcp1* element upstream and an *IS903* element downstream of the *bla*_{CTX-M-14} gene
228 (type II). The proportions of isolates from animal faecal, human faecal and human urine
229 sources with type II CTX-M-14 genetic environment were 89.7%, 87.7% and 86.5%,
230 respectively. In addition to *ISEcp1* and *IS903*, four other insertion sequences (*IS10*, *IS26*,
231 *IS6100* and *ISCR1*) and class 1 integron were found in the other types of genetic environment
232 (III to V, VII and VIII) in different combinations. Type VII and VIII genetic environments
233 were found to have genes encoding resistance to aminoglycosides (aminoglycoside
234 acetyltransferases, *aacA4* and *aadA2*), chloramphenicol (*cml*), sulphonamides (*sulI*) and/or
235 trimethoprim (dihydrofolate reductase, *dfrA12*).

236 Table 4 summarized the plasmid subtyping result for the 61 CTX-M-14 encoding
237 IncFII plasmids. RST revealed that the IncFII plasmids had four allelic variants: F2 (n=42),
238 F35 (n=17), F43 (n=1) and F51 (n=1). According to the PCR-RFLP patterns for the seven
239 loci (Figure 1), 32 unique plasmid RFLP (pRFLP) profiles were obtained for the 61 plasmids
240 encoding *bla*_{CTX-M-14} (Supplementary Table S3). These included 16 plasmids (group 1) with
241 all seven PCR-RFLP patterns identical to the reference plasmid (pHK01), 36 plasmids (group
242 2) with identity over three to six loci, and nine plasmids (group 3) without any identical PCR-
243 RFLP pattern or identical patterns in one or two loci only. Hence, 85.2% (52/61) of the

244 IncFII plasmids (group 1 and 2) were considered to be pHK01-like. These 52 pHK01-like
245 plasmids had their origins from human faecal ($n=20$), human urine ($n=14$) and animal faecal
246 ($n=18$, including 1 cattle, 1 chicken, 5 pigs, 9 cats, 1 dog and 1 rodent) sources. There were
247 15 unique pRFLP profiles for plasmids with the F2 and F35 replicon allele each. The
248 presence of the pHK01-associated *malB*, *yac*, *eitA*, *eitC* and *parA/B* genes was assessed by
249 monoplex PCRs using the purified plasmids as template. The result showed that 90.4%
250 (47/52) of the pHK01-like plasmids were positive for all targets. In contrast, these genes were
251 variably found among group 3 plasmids (Supplementary Table S3). The proportions of
252 animal faecal, human faecal and human urine isolates with pHK01-like plasmids were 31%
253 (18/58), 30.6% (20/65) and 37.8% (14/37), respectively.

254 The CTX-M-14 encoding plasmids, pHK09 (*E. coli* strain C017e) and pHK17a (*E.*
255 *coli* strain P0014ST) originating from the faecal samples of a child aged one year in 2007 and
256 a pig in 2008, respectively were sequenced (Ho et al., 2011a; Lo et al., 2010). The sizes of
257 pHK09 (GenBank accession JN087528) and pHK17a (GenBank accession JF779678) were
258 70382 bp and 70060 bp, respectively. The sequences of the two plasmids were highly similar
259 (>95%) to pHK01 (Hong Kong, 2004, GenBank accession HM355591, pRFLP 1-1-1-1-1-1-1)
260 and the pHK01-like plasmid, pKF3-70 (China, 2006, GenBank accession FJ494913, *in silico*
261 pRFLP 1-1-1-1-1-2-4). Figure 1 showed an alignment of the four plasmids (pHK01, pHK09,
262 pHK17a and pKF3-70) according to the functional regions. Except for some sequence
263 variations and possible insertions/deletions, the four plasmids were virtually identical. With
264 reference to pHK01, sequence variations were found in the *repA4* gene (94.5% similarity,
265 pHK17a), the *ssb* gene (98.9% similarity, pHK17a) and the *ISEcp1* element (99.8% similarity,
266 pKF3-70). The transfer region from *traM* to *traX* in pHK09 was more similar to pHK01
267 (99.3%) than to pHK17a (91.6%) or pKF3-70 (91.5%). The regions (*traM* to *traX*) in
268 pHK17a and pKF3-70 were almost identical to each other (99.6% sequence identity).

269 Sequence variations in the transfer region correlated with the observed difference in the
270 restriction digestion pattern over locus E and F in pHK17a and pKF3-70. In pHK17a,
271 sequence variations within *orf37* (encoding an adenine-specific DNA methylase gene)
272 correlated with negative PCR result for locus D. The *yfiA* gene (hypothetical protein), which
273 was found between *traR* and *traC* in pHK01 and pHK09, was absent in pHK17a and pKF3-
274 70. In contrast, an *artA* gene (hypothetical protein), absent in pHK01 and pHK09, was
275 inserted between *traF* and *traQ* in pHK17a and pKF3-70.

276

277 **Discussion**

278 This study demonstrated the widespread dissemination of pHK01-like plasmids among
279 isolates originating from multiple human and animal sources. The finding showed that 85%
280 (52/61) of the IncFII plasmids and 32.5% (52/160) of all the plasmids encoding CTX-M-14
281 were pHK01-like. Our previous work showed that pHK01-like plasmids play an important
282 role in the community emergence of CTX-M-14 among urinary *E. coli* in 2004 (Ho et al.,
283 2007; Ho et al., 2011b). The present work extends the observation by showing that pHK01-
284 like plasmids were found among human and animal faecal isolates collected in different time
285 periods. Of note, pHK01-like plasmids appeared to have approximately equal representation
286 among *E. coli* isolates of the four phylogenetic groups. This may partially explain its
287 widespread prevalence in bacteria of both animal and human origin. In Hong Kong, studies
288 have revealed that identical gentamicin resistance genes, integron and cassette arrays were
289 shared by isolates from animal faecal, human faecal and human urinary sources at similar
290 prevalence (Ho et al., 2009; Ho et al., 2010).

291 Since CTX-M-14 is the only antibiotic resistance gene in most pHK01-like plasmids,
292 persistence and spread of the plasmids cannot be attributed to co-selection by non- β -lactam
293 antibiotics. This single resistance feature is shared by the group of epidemic pCT-like

294 plasmids (IncK) encoding CTX-M-14 that have been reported in bacteria from human and
295 animals (Cottell et al., 2011). Furthermore, most pHK01-like plasmids had the 1-1 restriction
296 pattern at locus B1 and B2, and were PCR positive for *yac*, *malB*, and *eitA/eitC*. The finding
297 correlated with a conserved array of genes in the variable region (*yac-malB-IS903-bla_{CTX-M-14}-ISEcp1-eitABCD*). It is possible that the genes in the variable region may be advantageous
298 to the host bacteria. The toxin-antitoxin plasmid stability system (*yac*) is recognized to play
299 roles in the stable maintenance of large plasmids while the other transport proteins (*malB* and
300 *eitABCD* operon) could provide the host bacteria with some metabolic or virulence
301 advantages (Cheng et al., 2009; Zhao et al., 2009). Interestingly, the *yac* system and *eitABCD*
302 operon combination were also found in two major avian pathogenic plasmids, pAPEC-O2-
303 ColV (GenBank accession AY545598) and pAPEC-O1-ColBM (GenBank accession
304 DQ381420) that are widespread among poultry isolates (Johnson et al., 2006).

306 This study subtyped the IncFII plasmids by several methods. The RST scheme
307 discerned related from unrelated plasmids according to the *copA* sequence, while the pRFLP
308 scheme analysed the plasmid scaffold (Ho et al., 2011b; Villa et al., 2010). Our experience
309 showed that the RST scheme was easy to apply and the discriminatory power was good.
310 However, the RST allele did not consistently predict the plasmid scaffolds. Despite the fact
311 that the two FII alleles (F2 and F35) cluster into two distinct replicon subgroups (Villa et al.,
312 2010), plasmids of the two replicon alleles could share highly similar pRFLP profiles.
313 Complete sequencing of two other pHK01-like plasmids confirmed that variations in the
314 pRFLP profiles correlate with DNA insertion, deletion, point mutations and possible
315 homologous recombination. The finding suggests that multiple genetic processes are involved
316 in the evolution of the pHK01-like plasmids.

317 Among isolates of all sources, the predominant genetic environment of *bla_{CTX-M-14}*
318 consisted of an upstream *ISEcp1* and a downstream *IS903* (type II). This type of genetic

319 environment was similarly prevalent among the IncFII and non-IncFII plasmids. This study
320 also revealed the presence of several other mobile genetic elements (*IS10*, *IS26*, *IS6100*,
321 *ISCR1* and class 1 integron) in association with *bla*_{CTX-M-14} and/or *ISEcp1/IS903* in six
322 different combinations. Thus, many genetic elements might be involved in the mobilization
323 of *bla*_{CTX-M-14} into different plasmids (Barlow et al., 2008). In the present study, almost all
324 recognized genetic environment types were found. This could possibly be due to the diverse
325 origins of animals.

326 This study has limitations. Since pigs and chicken from different farms had been
327 mixed at the time of sampling, the geographic origins of these samples could not be traced
328 individually. Therefore, the observed variations in CTX-M-14 genetic environment and
329 divergence in plasmid sequences could be partly related to mixed origins of the samples from
330 local farms and importation from mainland China. Secondly, only two plasmids were
331 completely sequenced. As a result, the full spectrum of sequence divergence in the plasmids
332 could not be comprehensively resolved. Nonetheless, multiple pRFLP profiles involving
333 variations in different loci were found for isolates that originated from the same animal and
334 human sources. The finding highlights plasticity among related plasmids in different modules
335 of the plasmid backbones and the *bla*_{CTX-M-14} genetic environment. In the future, additional
336 plasmids with farm and geographical origins clearly defined should be investigated to clarify
337 the epidemiologic pattern of different plasmid sequences. Preferably, a larger number of
338 pHK01-like plasmids from different time periods should be completely sequenced to
339 delineate how plasmid sequence divergence correlates with the epidemiologic information
340 and how that changes over time.

341

342 **Conclusion**

343 This study showed that the pHK01-like plasmids encoding CTX-M-14 were shared among *E.*
344 *coli* isolates collected in different time periods from multiple human and animal sources.
345 Public health authorities should adopt measures to reduce the direct and indirect transfer of
346 resistant bacteria or resistance determinants within animal populations, from animals to
347 human and vice versa (Codex Alimentarius Commission, 2005).

348

349

350 **Acknowledgements**

351 This work was supported by grants from the Research Fund for the Control of Infectious
352 Diseases (RFCID) of the Health, Welfare and Food Bureau of the Government of the
353 HKSAR, and the Consultancy Service for Enhancing Laboratory Surveillance of Emerging
354 Infectious Disease for the HKSAR Department of Health.

355

356 **Transparency Declaration**

357 Conflicts of interest: Nothing to declare.

358

359 **Appendix A**

360 Supplementary data

361 **Table 1**
 362 Oligonucleotide primers for PCR mapping of CTX-M-14 genetic environment
 363

Primer name ^{a, b}	Direction	Nucleotide sequence (5' to 3')	Position	Accession No.	Amplicon size (bp)	Source
<i>ISEcpIU2</i>	F	AATACTACCTTGCTTTCTGA	7650-7669	HM355591	1464	(Saladin et al., 2002)
M9L	B	CCCTTCGGCGATGATTCTC	6206-6224			(Saladin et al., 2002)
<i>ISEcpIU1</i>	F	AAAAATGATTGAAAGGTGGT	7210-7229	HM355591	366	(Saladin et al., 2002)
CTX-M-9B	B	ATTGGAAAGCGTTCATCACC	6864-6883			(Woodford et al., 2006)
<i>ISCRIF</i>	F	GCCACCAACCCGACCAGAC	4001-4019	EF450247	2019	This study
CTX-M-9B	B	ATTGGAAAGCGTTCATCACC	6000-6019			(Woodford et al., 2006)
<i>tnpIS26-R</i>	F	AACTCTGCTTACCAGGCG	1413-1430	GQ385314	1507	This study
CTX-M-9B	B	ATTGGAAAGCGTTCATCACC	2900-2919			(Woodford et al., 2006)
<i>ISEcp_uw</i>	F	AACATCAAACGAATCGACCG	799-818	EU136400	1468	This study
<i>M9L_dw</i>	B	CACCTGCGTATTATCTGCGG	2247-2266			This study
Int-F	F	GCCACTGCGCCGTTACCACC	322-341	EF450247		(Ho et al., 2010)
Int1-285B	B	GCACAGCACCTTGCCGTAGAA	66-86		276	This study
CTX-M-9B	B	ATTGGAAAGCGTTCATCACC	6000-6019		5698	(Woodford et al., 2006)

M9U	F	ATGGTGACAAAGAGAGTGCA	7056-7075	HM355591	1265	(Saladin et al., 2002)
IS903D-5811R	B	TAACCGACTTTGCCCGCCTG	5811-5830			This study

364 ^aBased on published studies and sequences deposited in the GenBank database, eight different types of *bla*_{CTX-M-14} genetic environment have
365 been reported. Type I (*ISEcp1* - *bla*_{CTX-M-14}) consisted of an upstream *ISEcp1* but the downstream *IS903* was not detected by PCR (Kim et al.,
366 2011). However, the presence of a truncated *IS903* has not been confirmed by plasmid sequencing. The GenBank accession numbers for the
367 other types were HM355591 (type II, *ISEcp1* - *bla*_{CTX-M-14} - *IS903*), EU136400 (type III, Δ *ISEcp1* - *IS10* - *bla*_{CTX-M-14} - *IS903*), GQ385314 (type
368 IV, *IS26* - Δ *ISEcp1* - *bla*_{CTX-M-14} - *IS903*), AB180674 (type V, *IS26* - Δ *IS10* - *bla*_{CTX-M-14} - *IS903*), EU056266 (type VI, Class 1 integron- *orf5* -
369 *IS6100* -*ISCR1* - Δ *ISEcp1* - *bla*_{CTX-M-14} - *IS903*), FQ482074 (type VII, Class 1 integron- *orf1* -*ISEcp1* - *bla*_{CTX-M-14} - *IS903*) and EF450247 (type
370 VIII, Class 1 integron- *ISCR1*-*bla*_{CTX-M-14} - *IS903*). The class 1 integron include type VI and VIII contain *int11-dfrA12-orfF-aadA2-qacEA1-sul1*
371 (type VI and VIII) or *Aint11-aacA4-cml-qacEA1-sul1* (type VII).

372 ^bMonoplex PCRs using different primer pair combination was used to map the genetic environment to one of the eight recognized types. All the
373 assigned types yielded amplicons of the expected sizes. Representative amplicons were sequenced for confirmation.

374

375

376

377 **Table 2**

378 Replicon types for plasmids encoding CTX-M-14 analysed in this study

Replicon type ^a	No. of isolates	Plasmid sizes, kb	No. (%) according to source		
			Animal faecal	Human faecal	Human UTI
FII	61	55-100	21 (36.2)	25 (38.5)	15 (40.5)
FII, FIB	7	100-150	3	3	1
FIB	6	55-190	1	3	2
FII, I1-I γ	3	70-100	1	2	-
FIA	3	55-100	-	1	2
FII, FIA	2	100-130	-	2	-
FIA, FIB	1	80	1	-	-
FII, FIA, FIB	1	150	-	1	-
I1-I γ	24	40-120	12 (20.7)	7 (10.8)	5 (13.5)
B/O	10	60-140	5	5	-
K	6	80-100	-	5	1
Others ^b	36	50-250	14	11	11
Total	160	-	58	65	37

379 ^a The replicon types were determined by probe hybridization in the transconjugants (n=84) or
 380 the parent strains (n=76).

381 ^bThese included N (n=3), A/C (n=1), HI1 (n=1), HI2 (n=1) and nontypeable (n=30).

382

383

384

385 **Table 3**

386 CTX-M-14 genetic environment of *E. coli* isolates from humans ($n=102$) and animals ($n=58$),

387 Hong Kong.

Type ^a	Genetic environment of <i>bla</i> _{CTX-M-14}	No (%)		
		Animal- faecal	Human- faecal	Human- urine
II	<i>ISEcp1</i> - <i>bla</i> _{CTX-M-14} - IS903	52 (89.7)	57 (87.7)	32 (86.5)
III	Δ <i>ISEcp1</i> - IS10 - <i>bla</i> _{CTX-M-14} - IS903 ^b	-	4	1
IV	IS26 - Δ <i>ISEcp1</i> - <i>bla</i> _{CTX-M-14} - IS903	3	1	-
V	IS26 - Δ IS10 - Δ <i>ISEcp1</i> - <i>bla</i> _{CTX-M-14} - IS903 ^c	1	-	1
VII	Class 1 integron- <i>orf1</i> - <i>ISEcp1</i> - <i>bla</i> _{CTX-M-14} - IS903	2	1	3
VIII	Class 1 integron- <i>ISCR1</i> - <i>bla</i> _{CTX-M-14} - IS903	-	2	-
Subtotal		58	65	37

388 ^a The Genbank accession numbers were as follows: HM355591 (type II), EU136400 (type III),

389 GQ385314 (type IV), AB180674 (type V), EU056266 (type VI, Class 1 integron- *orf5* -

390 IS6100 - *ISCR1* - Δ *ISEcp1* - *bla*_{CTX-M-14} - IS903), FQ482074 (type VII) and EF450247 (type

391 VIII). The class 1 integron includes type VI and VIII containing *intl1-dfrA12-orfF-aadA2-*

392 *qacEΔ1-sul1* (type VI and VIII) or Δ *intl1-aacA4-cml-qacEΔ1-sul1* (type VII).

393 ^b The Δ *ISEcp1* was interrupted by an IS10 element.

394 ^c The Δ *ISEcp1* element was very short and limited to a putative *ISEcp1* promoter.

395

396

397 **Table 4**

398 Characteristics of the IncFII plasmids encoding *bla*_{CTX-M-14} in 61 *Escherichia coli* isolates
 399 from humans and animals.

	No of plasmids ^a			
	Group 1	Group 2	Group 3	All
Number of plasmids	16	36	9	61
FAB formula				
F2:A-:B-	16	26	-	42
F35:A-:B-	-	10	7	17
F43:A-:B-	-	-	1	1
F51:A-:B-	-	-	1	1
PCR profile				
<i>malB-yac-eitA-eitC-parAB</i>	16	31	2	49
Others ^b	-	1	5	6
None	-	4	2	6
Source				
Human UTI	5	9	1	15
Human faecal	4	16	5	25
Chicken	1	-	-	1
Cattle	1	-	-	1
Pig	1	4	1	6
Cats	3	6	1	10
Dogs	0	1	-	1
Rodents	1	-	1	2

400 ^aThe plasmids were categorised according to their similarity to the reference plasmid, pHK01:
401 group 1, all seven PCR-RFLP patterns were identical to reference; group 2, three to six
402 restriction patterns were identical; and group 3, less than three restriction patterns were
403 identical.


404 ^bIncluding the following profiles: *yac-eitA-eitC-parAB* (*n*=2), *yac-eitC-parAB* (*n*=1), *mal-*
405 *eitA-eitC* (*n*=1), *parAB* (*n*=2).

406

407

FIGURE 1. Schematic representation of pHK09 and pHK17a and comparison with two other CTX-M-14 encoding plasmids, pHK01 and pKF3-70.

The sequence of pHK01 was used as the reference for all comparisons. The five functional regions were indicated by arrows on top while the seven sets of long-range PCR targets (locus A to locus F) were indicated by arrows in the bottom. The genes and ORF of interest were indicated in boxes and annotated. Potential deletions were indicated by dotted line. The *rep* genes in the replication and *tra* genes in the transfer region were indicated by white boxes with number and capital letters, respectively. The percentage identity over each indicated block in the transfer region were given above the boxes. The genes or ORFs with sequence variations were indicated in **boldface**. *Orf37* is named according to pHK01. Its counterpart is *orf35* in pHK09 and pHK17 and *orf81* in pKF3-70. The boxes according to shading patterns were: dotted lines (partition-associated genes, *parA/B*, *ssb*, *parB-like*, *psiA/B*), dots (toxin-antitoxin genes, *yacABC*), oblique lines (transportation-related proteins, *malB* and *eitABCD*), gray (insertion sequences, *IS903* and *ISEcp1*) and black (resistance gene, *bla_{CTX-M-14}*). The plasmids' origins and their pRFLP profiles were labelled on the two sides..



References

1. Barlow M, Reik RA, Jacobs SD, Medina M, Meyer MP, McGowan JE, Jr., Tenover FC (2008) High rate of mobilization for blaCTX-Ms. *Emerg Infect Dis* 14: 423-428.
2. Canton R and Coque TM (2006) The CTX-M beta-lactamase pandemic. *Curr Opin Microbiol* 9: 466-475.
3. Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ (2005) Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods* 63: 219-228.
4. Cheng VC, To KK, Li IW, Tang BS, Chan JF, Kwan S, Mak R, Tai J, Ching P, Ho PL, Seto WH (2009) Antimicrobial stewardship program directed at broad-spectrum intravenous antibiotics prescription in a tertiary hospital. *Eur J Clin Microbiol Infect Dis* 28: 1447-1456.
5. Clermont O, Bonacorsi S, Bingen E (2000) Rapid and simple determination of the *Escherichia coli* phylogenetic group. *Appl Environ Microbiol* 66: 4555-4558.
6. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing: twenty-first informational supplement M100-S21. 2011. CLSI, Wayne, PA, USA.

Ref Type: Generic

7. Codex Alimentarius Commission. Code of practice to minimize and contain antimicrobial resistance. 2005. http://www.codexalimentarius.net/web/standard_list.jsp.

Ref Type: Generic

8. Cottell JL, Webber MA, Coldham NG, Taylor DL, Cerdeno-Tarraga AM, Hauser H, Thomson NR, Woodward MJ, Piddock LJ (2011) Complete sequence and molecular epidemiology of IncK epidemic plasmid encoding blaCTX-M-14. *Emerg Infect Dis* 17: 645-652.
9. Garcia-Fernandez A, Chiaretto G, Bertini A, Villa L, Fortini D, Ricci A, Carattoli A (2008) Multilocus sequence typing of IncII plasmids carrying extended-spectrum beta-lactamases in *Escherichia coli* and *Salmonella* of human and animal origin. *J Antimicrob Chemother* 61: 1229-1233.
10. Hawkey PM (2008) Prevalence and clonality of extended-spectrum beta-lactamases in Asia. *Clin Microbiol Infect* 14 Suppl 1: 159-165.
11. Ho PL, Chow KH, Lai EL, Lo WU, Yeung MK, Chan J, Chan PY, Yuen KY (2011a) Extensive dissemination of CTX-M-producing *Escherichia coli* with multidrug resistance to 'critically important' antibiotics among food animals in Hong Kong, 2008-10. *J Antimicrob Chemother* 66: 765-768.
12. Ho PL, Lo WU, Wong RC, Yeung MK, Chow KH, Que TL, Tong AH, Bao JY, Lok S, Wong SS (2011b) Complete sequencing of the FII plasmid pHK01, encoding CTX-M-14, and molecular analysis of its variants among *Escherichia coli* from Hong Kong. *J Antimicrob Chemother* 66: 752-756.
13. Ho PL, Lo WU, Yeung MK, Lin CH, Chow KH, Ang I, Tong AH, Bao JY, Lok S, Lo JY (2011c) Complete sequencing of pNDM-HK encoding NDM-1 carbapenemase from a multidrug-resistant *Escherichia coli* strain isolated in Hong Kong. *PLoS One* 6: e17989.

14. Ho PL, Poon WW, Loke SL, Leung MS, Chow KH, Wong RC, Yip KS, Lai EL, Tsang KW (2007) Community emergence of CTX-M type extended-spectrum beta-lactamases among urinary *Escherichia coli* from women. *J Antimicrob Chemother* 60: 140-144.
15. Ho PL, Tsang DN, Que TL, Ho M, Yuen KY (2000) Comparison of screening methods for detection of extended-spectrum beta-lactamases and their prevalence among *Escherichia coli* and *Klebsiella* species in Hong Kong. *APMIS* 108: 237-240.
16. Ho PL, Wong RC, Chow KH, Que TL (2009) Distribution of integron-associated trimethoprim-sulfamethoxazole resistance determinants among *Escherichia coli* from humans and food-producing animals. *Lett Appl Microbiol* 49: 627-634.
17. Ho PL, Wong RC, Chow KH, Yip K, Wong SY, Que TL (2008) CTX-M type beta-lactamases among fecal *Escherichia coli* and *Klebsiella pneumoniae* isolates in non-hospitalized children and adults. *J Microbiol Immuno Infect* 41: 428-432.
18. Ho PL, Wong RC, Lo SW, Chow KH, Wong SS, Que TL (2010) Genetic identity of aminoglycoside-resistance genes in *Escherichia coli* isolates from human and animal sources. *J Med Microbiol* 59: 702-707.
19. Johnson TJ, Johnson SJ, Nolan LK (2006) Complete DNA sequence of a ColBM plasmid from avian pathogenic *Escherichia coli* suggests that it evolved from closely related ColV virulence plasmids. *J Bacteriol* 188: 5975-5983.
20. Kim J, Bae IK, Jeong SH, Chang CL, Lee CH, Lee K (2011) Characterization of IncF plasmids carrying the blaCTX-M-14 gene in clinical isolates of *Escherichia coli* from Korea. *J Antimicrob Chemother* 66: 1263-1268.

21. Lo WU, Ho PL, Chow KH, Lai EL, Yeung F, Chiu SS (2010) Fecal carriage of CTXM type extended-spectrum beta-lactamase-producing organisms by children and their household contacts. *J Infect* 60: 286-292.
22. Naseer U and Sundsfjord A (2011) The CTX-M conundrum: dissemination of plasmids and *Escherichia coli* clones. *Microb Drug Resist* 17: 83-97.
23. Osborn AM, Silva Tatley FM, Steyn LM, Pickup RW, Saunders JR (2000) Mosaic plasmids and mosaic replicons: evolutionary lessons from the analysis of genetic diversity in IncFII-related replicons. *Microbiology* 146: 2267-2275.
24. Poirel L, Lartigue MF, Decousser JW, Nordmann P (2005) ISEcp1B-mediated transposition of blaCTX-M in *Escherichia coli*. *Antimicrob Agents Chemother* 49: 447-450.
25. Saladin M, Cao VT, Lambert T, Donay JL, Herrmann JL, Ould-Hocine Z, Verdet C, Delisle F, Philippon A, Arlet G (2002) Diversity of CTX-M beta-lactamases and their promoter regions from *Enterobacteriaceae* isolated in three Parisian hospitals. *FEMS Microbiol Lett* 209: 161-168.
26. Valverde A, Canton R, Garcillan-Barcia MP, Novais A, Galan JC, Alvarado A, de la CF, Baquero F, Coque TM (2009) Spread of blaCTX-M-14 is driven mainly by IncK plasmids disseminated among *Escherichia coli* phylogroups A, B1, and D in Spain. *Antimicrob Agents Chemother* 53: 5204-5212.
27. Villa L, Garcia-Fernandez A, Fortini D, Carattoli A (2010) Replicon sequence typing of IncF plasmids carrying virulence and resistance determinants. *J Antimicrob Chemother* 65: 2518-2529.

28. Woodford N, Fagan EJ, Ellington MJ (2006) Multiplex PCR for rapid detection of genes encoding CTX-M extended-spectrum (beta)-lactamases. *J Antimicrob Chemother* 57: 154-155.
29. Yi H, Xi Y, Liu J, Wang J, Wu J, Xu T, Chen W, Chen B, Lin M, Wang H, Zhou M, Li J, Xu Z, Jin S, Bao Q (2010) Sequence analysis of pKF3-70 in *Klebsiella pneumoniae*: probable origin from R100-like plasmid of *Escherichia coli*. *PLoS One* 5: e8601.
30. Zhao L, Gao S, Huan H, Xu X, Zhu X, Yang W, Gao Q, Liu X (2009) Comparison of virulence factors and expression of specific genes between uropathogenic *Escherichia coli* and avian pathogenic *E. coli* in a murine urinary tract infection model and a chicken challenge model. *Microbiology* 155: 1634-1644.