



Title	Exact Meromorphic Stationary Solutions of the Real Cubic Swift-Hohenberg Equation
Author(s)	Conte, BR; Ng, TW; Wong, KK
Citation	Studies In Applied Mathematics, 2012
Issued Date	2012
URL	http://hdl.handle.net/10722/156285
Rights	Creative Commons: Attribution 3.0 Hong Kong License

Exact meromorphic stationary solutions of the real cubic Swift-Hohenberg equation

ROBERT CONTE^{1,2*}, TUEN-WAI NG^{1†} AND KWOK-KIN WONG^{1‡}

23 November 2011, revised 14 January 2012

Abstract. We show that all meromorphic solutions of the stationary reduction of the real cubic Swift-Hohenberg equation are elliptic or degenerate elliptic. We then obtain them all explicitly by the subequation method, and one of them appears to be a new elliptic solution.

1 Introduction

The real cubic Swift-Hohenberg (RCSH) equation

$$\frac{\partial u}{\partial t} = \varepsilon u - \left(1 + \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right)^2 u - u^3, \quad \varepsilon \in \mathbb{R} \quad (1)$$

originally proposed in [34], is a model for Rayleigh-Bénard convection in hydrodynamics. Since then, the equation and its generalizations have also been used in various areas, in particular those where the field u needs to be a complex amplitude, such as laser [26] and nonlinear optics [27]. Much attention of the researches about

*Partially supported by PROCORE - France/Hong Kong joint research grants F-HK29/05T, F-HK39/11T and RGC grant HKU 703807P.

†Partially supported by PROCORE - France/Hong Kong joint research grants F-HK29/05T, F-HK39/11T and RGC grant HKU 703807P.

‡Partially supported by RGC grant HKU 703807P and a post-graduate studentship at HKU.

2000 *Mathematics Subject Classification*: Primary 35Q53.

Key words and phrases. Real cubic Swift-Hohenberg equation, exact meromorphic solutions, elliptic solutions, Nevanlinna theory, subequation method.

1. Department of Mathematics, The University of Hong Kong, Pokfulam Road.

2. LRC MESO, Centre de mathématiques et de leurs applications (UMR 8536) et CEA-DAM, École normale supérieure de Cachan, 61, avenue du Président Wilson, F-94235 Cachan Cedex, France.

E-mail: Robert.Conte@cea.fr, ntw@maths.hku.hk, wkkm@graduate.hku.hk

RCSH has been put into its relation to the theory of pattern formation. The phenomena of pattern formation occur in a large variety of physical systems. We refer the readers to the extensive review [13] and the monographs [12] and [21].

The Swift-Hohenberg equation (1) admits a stationary reduction by taking $u(x, y) = \sqrt{\varepsilon - 1}U(Z)$, with the rescaling $Z = (\varepsilon - 1)^{1/4}(x + y)/2$:

$$U'''' + aU'' + U^3 - U = 0, \quad ' := \frac{d}{dZ}. \quad (2)$$

Here $a = 2/\sqrt{\varepsilon - 1}$. The ODE (2) is sometimes mentioned as “extended Fisher-Kolmogorov equation” [1, 24] for $a < 0$.

The equation (2) has attracted intensive studies concerning, for example, the existence of various types of solutions and qualitative properties of the solution curves [2, 3, 21, 24, 31, 30, 32]. For instance, when U is a smooth function defined on the real line such that $U - 1 \in H^2(\mathbb{R})$, Santra and Wei [32] proved the existence of a homoclinic solution U for each $a \in [0, a_0]$, where $a_0 = 1.228\dots$ ¹

We shall see in section 4 that a meromorphic homoclinic solution can only exist when $a = 10/\sqrt{11} = 3.015\dots$, which is outside the above specified range, and we shall write down this exact meromorphic homoclinic solution in terms of the tanh function (i.e. solution (30)). Indeed, in this paper we shall find all the meromorphic solutions of (2) explicitly.

The method to derive all meromorphic solutions of (2) is twofold. We first use an analysis based on Nevanlinna theory proposed by Eremenko [15] and slightly modified in [10] to prove that any meromorphic solution is necessarily elliptic or degenerate of elliptic. In a second step, we use the subequation method first proposed in [29] to characterize each elliptic or degenerate elliptic solution by some first order ODE, which are easily integrated by classical methods.

The overall advantage of the method is that, once the algorithm is completed, all possible meromorphic solutions are exhausted. The detailed procedure of the method can be found in [10].

In section 2, we show that all meromorphic solutions of (2) belong to class W (like Weierstrass), which consists of elliptic functions and their successive degeneracies, i.e.: elliptic functions, rational functions of one exponential $\exp(kz)$, $k \in \mathbb{C}$ and rational functions of z . We then apply the subequation method to find explicitly all these class W solutions in section 3. Finally, in section 4, we shall compare the meromorphic solutions obtained in section 3 with some known results concerning (2).

Let us finally mention a recent paper [6] which uses similar ideas. Given a specific nonlinear second order ODE, these authors use Wiman-Valiron theory combined with local series analysis to explicitly determine all entire solutions.

¹ The value a_0 here is different from that in [32]. In [32, p. 2040], $a_0 = \beta_0^2$ where $\beta_0 = \sqrt{\sqrt{2}/k_0}$, and $4k_0^2 - 2k_0 - 3 = 0$. The value a_0 stated in [32] is probably a typographical error.

2 Class W solutions

Only five polynomial terms $(UU'', U''', U^2, U', 1)$ can be added to (2) while retaining the double pole behaviour displayed below, Eq. (13). For a reason to be explained soon, we only retain two of them $(U^2, 1)$ and consider the more general form of the equation (2) in which the three fixed points $U = -1, 0, 1$ are not necessarily equispaced,

$$u'''' + 60bu'' - \frac{120}{c^2} (u^3 - 3s_1u^2 + 3s_2u - s_3) = 0, \quad ' := \frac{d}{dz}. \quad (3)$$

The complex constant c , which could be scaled out, will be useful for parity considerations, for example to determine the second Laurent series (13) by simply changing c to $-c$ in the first one.

Its three fixed points a_j are defined by

$$u^3 - 3s_1u^2 + 3s_2u - s_3 = (u - a_1)(u - a_2)(u - a_3) \quad (4)$$

When the three fixed points a_j of (3) are equispaced,

$$3s_1s_2 - 2s_1^3 - s_3 = 0, \quad (2a_1 - a_2 - a_3)(2a_2 - a_3 - a_1)(2a_3 - a_1 - a_2) = 0, \quad (5)$$

equation (3) is indeed equivalent to (2) under the rescaling,

$$U = \lambda(u - s_1), \quad z = \mu Z, \\ \lambda^2 = -\frac{1}{3(s_2 - s_1^2)} = \frac{6}{(a_2 - a_3)^2 + (a_3 - a_1)^2 + (a_1 - a_2)^2}, \quad \mu^2 = \frac{a}{60b}. \quad (6)$$

Equation (3) admits the integrating factor u' , thus yielding the first integral

$$K = u'u'''' - \frac{1}{2}u''^2 + 30bu'^2 - \frac{30}{c^2} (u^4 - 4s_1u^3 + 6s_2u^2 - 4s_3u + 3s_1^4 - 6s_1^2s_2 + 4s_1s_3), \quad (7)$$

where K is the integration constant, and the added constant terms will make later expressions simpler.

Remark 1. The reason for excluding terms (uu'', u''', u') in (3) is to allow the first integral (7) to exist.

Let u be a meromorphic solution (defined on \mathbb{C}) of (7). To find all such u , we shall first prove the following result.

Theorem 1. *If the ODE (7) has a particular meromorphic solution u , then u belongs to the class W .*

The proof requires the use of the Nevanlinna theory, whose main features required here are introduced below. Some good references of Nevanlinna theory are [18] and [25].

The method for proving 1 comes from a paper of Eremenko [14] which shows that all meromorphic solutions of odd order Briot-Bouquet differential equations with at least one pole must belong to class W . The even order case was recently proven to be also true [16]. Eremenko's method applied to (7) or other autonomous nonlinear algebraic ODEs consists of two main steps:

1. Show that there are *finitely many* Laurent series with a pole at any point z_0 which satisfy (7).
2. Show that any *transcendental* meromorphic solution u of (7) must have infinitely many poles.

Step 1 usually involves computing the Fuchs indices of (7) and Step 2 requires applying Nevanlinna theory or Wiman-Valiron theory. Once we have established the two steps, we can then show easily that the solution u must be periodic and hence belong to class W .

For the convenience of readers, we shall include a proof of 1 which is very similar to the one given in [15] or [10].

We first introduce some notations commonly used in Nevanlinna theory (see [18], also see [25] for a quick introduction). Assume f to be a non-constant meromorphic function on the open disc $D(r)$ where r can be ∞ .

Denote the number of poles of f on the closed disc $\overline{D}(r)$ by $n(r, f)$, counting multiplicity. The function $n(r, f)$ is usually called the **unintegrated counting function**. Note that $n(r, \frac{1}{f})$ would be the number of zeros of f on $\overline{D}(r)$ and $n(r, \frac{1}{f-a})$ would be the number of times f takes a .

Define the **integrated counting function** $N(r, f)$ as

$$N(r, f) = n(0, f) \log r + \int_0^r [n(t, f) - n(0, f)] \frac{dt}{t}, \quad (8)$$

and the **proximity function** $m(r, f)$ as

$$m(r, f) = \int_0^{2\pi} \log^+ f(re^{i\theta}) \frac{d\theta}{2\pi}, \quad (9)$$

where $\log^+ x = \max\{0, \log x\}$.

Finally, we define the Nevanlinna's characteristic function $T(r, f)$ as

$$T(r, f) = m(r, f) + N(r, f). \quad (10)$$

A basic fact concerning the functions T, N, m is the following:

Theorem 2 (Nevanlinna's first fundamental theorem). *Let f be a meromorphic function and $a \in \mathbb{C}$. Then*

$$T(r, \frac{1}{f-a}) = m(r, f) + N(r, f) + O(1)$$

as $r \rightarrow \infty$.

We also need the following well known results (detailed discussions can be found in [25]):

Lemma 1. *A meromorphic function f is rational if and only if $T(r, f) = O(\log r)$.*

Lemma 2 (Clunie’s lemma). *Let f be a transcendental meromorphic solution of*

$$f^n P(z, f, f', \dots) = Q(z, f, f', \dots), \quad (11)$$

where n is a nonzero positive integer, P and Q are polynomials in f and its derivatives with meromorphic coefficients $\{a_\lambda | \lambda \in I\}$, such that for all $\lambda \in I$, $m(r, a_\lambda) = S(r, f)$, where I is some known index set. If the total degree² of Q as a polynomial in f and its derivatives is less than or equal to n , then

$$m(r, P(z, f, f', \dots)) = S(r, f). \quad (12)$$

Here $S(r, f)$ is called the “small” function and $m(r, P(z, f, f', \dots)) = S(r, f)$ means that the function on left hand side has growth $o(T(r, f))$ as $r \rightarrow \infty$ outside a possible exceptional set of finite linear measure.

Proof of Theorem 1. Let’s begin with Step 1. Suppose u is a meromorphic solution of (7). If u has a movable pole at $z = z_0$, then it must be a double pole. Let

$$u(z) = \alpha_0(z - z_0)^{-2} + \alpha_1(z - z_0)^{-1} + \alpha_2 + \dots \quad (13)$$

be the Laurent series of u in a neighborhood of z_0 . Inserting (13) into (7) and balancing the leading terms, we get $\alpha_0 = \pm c$. Using for example the procedure given in [7], the Fuchs indices of the ODE (7) are found to be independent of the sign of α_0 and equal to -1 , $(7 \pm \sqrt{-71})/2$. This implies that all other Laurent series coefficients in (13) are uniquely determined by the leading coefficients α_0 and are independent of z_0 [7], thus we only have two distinct Laurent series. This shows that there exist at most two meromorphic functions with a pole at $z = z_0$ satisfying (7).

If u is rational, then u belongs to W and we are done. Now assume u to be transcendental. By putting (7) into the form:

$$\frac{30}{c^2}u^4 = u'u''' - \frac{1}{2}u''^2 + 30bu'^2 - \frac{30}{c^2}(-4s_1u^3 + 6s_2u^2 - 4s_3u) + \text{constant}, \quad (14)$$

we conclude from Clunie’s lemma ($f = u, n = 3, P = u$) that $m(r, u) = S(r, u)$ and hence $(1 - o(1))T(r, u) = N(r, u)$. If u has only finitely many poles, then $N(r, u) = O(\log r)$ and hence $T(r, u) = O(\log r)$. By Lemma 1, u must be rational, which is a contradiction. Therefore u must have infinitely many poles and this completes Step 2.

We are now ready to prove that u belongs to class W . We first claim that, if u is transcendental, it must be periodic. Suppose u has a pole at $z = z_0$. By the previous analysis, there exist at most two meromorphic solutions of (7) with poles at $z = z_0$. Let $z_j, j = 1, 2, \dots$, be the infinitely many poles of $u(z)$; then for all $j = 1, 2, \dots$, $w_j(z) \equiv u(z + z_j - z_0)$ are also solutions of (7) with a pole at z_0 .

Since there exist at most two Laurent series around the pole z_0 , some of the w_j ’s must be the same, otherwise we get a contradiction to the maximum number

² Defined as the global degree in all derivatives $f^{(j)}, j \geq 0$.

of possible Laurent series. But this implies that for some $j \neq i$, $u(z - z_0 + z_j) \equiv u(z - z_0 + z_i)$ and hence $u(z) \equiv u(z + z_i - z_j)$ in a neighborhood of z_0 . Since u is meromorphic, we can conclude that u is periodic in \mathbb{C} with period $z_i - z_j$.

By a suitable rescaling, we may assume that $2\pi i$ is a primitive period of u . Let $D = \{z : 0 \leq \Im z < 2\pi\}$. If u has more than two poles in D , we can also consider the solutions defined by $w_j(z) = u(z + z_j - z_0)$ similar to the last paragraph. Then since the number Laurent series is at most two, for some $j \neq i$, $u(z - z_0 + z_j) \equiv u(z - z_0 + z_i)$ and u must be periodic in D with period $z_i - z_j$. Note that $z_i - z_j \neq 2k\pi i$ for any $k \in \mathbb{Z}$ because $z_i, z_j \in D$. Thus u is doubly periodic and therefore elliptic.

If u has one or two poles in D , then since u is a periodic function with period $2\pi i$, we have $N(r, u) = O(r)$ as $r \rightarrow \infty$. Since $(1 - o(1))T(r, u) = N(r, u)$, we have $T(r, u) = O(r)$. By the Nevanlinna's first fundamental theorem, for any $\zeta \in \mathbb{C}$, $N(r, 1/(u - \zeta)) = O(r)$ as $r \rightarrow \infty$. Since u is periodic with period $2\pi i$ and $N(r, 1/(u - \zeta)) = O(r)$, it must take each ζ only finitely many times in D . This implies that the function $R(z) = u(\ln z)$ is a single-valued analytic function defined in the punctured complex plane $\mathbb{C} - \{0\}$ which takes each ζ only finitely many times. So 0 is a removable singularity of R . This implies that R is rational. Since $u(z) = R(e^z)$, we conclude that u must belong to the class W and 1 is proven.

Remark 2. From the proof, we conclude that, if the solution u is elliptic, the number of poles of u in the fundamental parallelogram must be at most two.

Remark 3. The ODE (3) admits infinitely many Laurent series parametrized by the arbitrary constant K , therefore 1 cannot apply to it.

3 Explicit meromorphic solutions

On the one hand, Theorem 1 shows that any meromorphic solution of (7) must be in the class W . On the other hand, there exists a method (the subequation method proposed in [29]) which can be applied to find all those solutions of an algebraic ODE which belong to class W . Now, combining these two features allows one to find in closed form all the particular solutions of (7) which are meromorphic.

Let us first recall the classical definition of the *elliptic order* of an elliptic function: this is the common number of poles or zeros, counting multiplicity, inside a fundamental parallelogram. The method is based on the following well known theorem of Briot and Bouquet on first order Briot-Bouquet differential equations [4, 19]:

Theorem 3. *Any elliptic function obeys a first order algebraic differential equation of the form*

$$F(u, u') \equiv \sum_{k=0}^m \sum_{j=0}^{2m-2k} a_{j,k} u^j u'^k = 0, \quad a_{0,m} \neq 0, \quad (15)$$

where m is the elliptic order of u .

In the case of (7), if the solution u is elliptic, it has either one double pole (then $m = 2$, and only one Laurent series contributes to this solution) or two double poles (then $m = 4$, and the two Laurent series contribute to this solution) in a fundamental parallelogram. Therefore the subequation method must be applied with, successively, $m = 2$ and $m = 4$.

If u is a degenerate elliptic function, the above theorem does not apply, but any Laurent series expansion around a pole still has the polar order two and there exist at most two such expansions, therefore u must still obey an ODE of the form (15), with $m = 2$ or 4. Now let u be a meromorphic solution of (7). From section 2, we know that we can determine uniquely and recursively the coefficients of the two possible Laurent series of u in a neighborhood of a movable pole $z = z_0$. After putting these Laurent series into (15), the equation (15) should vanish identically. This generates a countably infinite (hence overdetermined and easy to solve) system of linear equations in the finitely many unknowns $a_{j,k}$'s. By solving it for the $a_{j,k}$'s, the first order ODEs $F(u, u') = 0$ are known explicitly and all admit (7) as a differential consequence. This is why we call the obtained first order ODEs $F(u, u') = 0$ a *subequation* of (7). For details of the implementation of the algorithm, see [29].

3.1 Subequations obeyed by one Laurent series

These subequations have degree $m = 2$. For a given Laurent series (13), the method yields one subequation, at the price of one constraint among the fixed coefficients K, b, c, s_1, s_2, s_3 of (7),

$$\begin{cases} F_2 \equiv cu'^2 - 4(u - s_1 - bc)^3 + 20(s_1^2 - s_2 - b^2c^2)(u - s_1 - bc) \\ \quad + 10(s_3 - 3s_1s_2 + s_1^3 + 2bcs_2 - 2bcs_1^2 + 4b^3c^3) = 0, \\ c^2K = 10(-128b^4c^4 - 88b^2c^2(s_2 - s_1^2) + 18bc(-2s_1^3 + 3s_2s_1 - s_3)). \end{cases} \quad (16)$$

The solution for the other Laurent series is obtained by changing c to $-c$. This first order ODE (16) is nothing else than the canonical equation of Weierstrass, up to some translation and rescaling.

This defines the codimension-one³ solution of (7)

$$\begin{cases} u = s_1 + bc + c\wp(z - z_0, g_2, g_3), \\ g_2 = 20(s_1^2 - s_2 - b^2c^2)/c^2, \\ g_3 = 10(s_3 - 3s_1s_2 + s_1^3 + 2bcs_2 - 2bcs_1^2 + 4b^3c^3)/c^3. \end{cases} \quad (17)$$

This solution is not new, and has already been obtained [28] by assuming u (which has movable double poles) to be a polynomial in (\wp, \wp') having double poles, i.e. reducing to an affine function of \wp .

When the genus of the curve $F_2(u, u') = 0$ is zero, i.e. when $g_2^3 - 27g_3^2 = 0$, this solution becomes a rational function of one exponential, obtained by the degeneracy formula

$$\forall x, d: \wp(x, 3d^2, -d^3) = -d + \frac{3d}{2} \coth^2 \sqrt{\frac{3d}{2}} x. \quad (18)$$

³ Following a frequent terminology, we define the *codimension* of a solution of a differential equation as the number of constraints required among the parameters of the equation.

This codimension-two genus-zero solution is

$$\begin{cases} u = s_1 + bc + c \left(k^2 \tanh^2 kz - \frac{2}{3}k^2 \right), \\ k^4 = 15(s_1^2 - s_2 - (bc)^2)/c^2, \\ s_3 = \frac{1}{135}(3s_1 + 3bc + ck^2)(45s_1^2 - 15s_1c(3b + k^2) - 4c^2k^4 + 30bc^2k^2 - 90b^2c^2), \\ K = -\frac{2}{9}c^2(3b + k^2)^2(45b^2 - 30bk^2 + k^4), \end{cases} \quad (19)$$

in which the arbitrary origin of z has been omitted.

3.2 Subequations obeyed by two Laurent series

The subequation then has degree $m = 4$. One thus finds two subequations and only two. One of them, as expected, is factorizable into the product of the two second degree subequations (16) (one for c , the other for $-c$) with the additional condition that the K in (16) be invariant under parity on c , i.e. $2s_1^3 - 3s_2s_1 + s_3 = 0$.

The other subequation is irreducible and has genus one,

$$\begin{cases} F_4 \equiv c^2 \left(u'^2 + 12b((u - s_1)^2 + 2\sigma_2) \right)^2 - 16 \left((u - s_1)^2 + 2\sigma_2 \right)^3 = 0, \\ \sigma_2 := \frac{1}{3}(5(s_2 - s_1^2) + 8b^2c^2), \quad 2s_1^3 - 3s_2s_1 + s_3 = 0, \\ c^2K = -\frac{16}{3} (8b^2c^2 + 5(s_2 - s_1^2))^2 (11b^2c^2 - 10(s_2 - s_1^2)). \end{cases} \quad (20)$$

The condition on (s_1, s_2, s_3) expresses that the three fixed points of (3) are equispaced, like $-1, 0, 1$ in (2).

In order to integrate this genus one equation $F_4(u, u') = 0$, one must first establish a birational transformation $(u, u') \leftrightarrow (\wp, \wp')$ between $F_4 = 0$ and the canonical equation of Weierstrass $\wp'^2 = 4\wp^2 - g_2\wp - g_3$, i.e. two pairs of rational transformations

$$u = R_1(\wp, \wp'), \quad u' = R_2(\wp, \wp'), \quad \wp = R_3(u, u'), \quad \wp' = R_4(u, u'), \quad (21)$$

where R_i , $i = 1, 2, 3, 4$ are rational in their arguments. The algorithm consists of mapping (*via* a birational transformation) the given algebraic curve in (u, u') to an algebraic curve in (v, v') in which v has a lower elliptic order than u , until elliptic order 2 (that of \wp) has been reached. Outlined by Briot and Bouquet [4, §250–251 p. 395] for the so-called “binomial” and “trinomial” equations, it has been implemented in the Maple package `algcurves` [20] for any genus one first order equation. Then a final scaling yields the desired solution.

This codimension-two elliptic solution of (7) can be written in terms of either an

even or an odd elliptic function (just like $\sin(z - z_0) = -\cos(z - z_1)$, $z_1 - z_0 = \pi/2$),

$$\left\{ \begin{array}{l} u = s_1 - c(e_2 - e_3) \frac{(\wp_0 + b)^2 + g_2/4 - 3b^2}{\wp_0^2 - b\wp_0 - g_2/4 + b^2} = s_1 - \frac{d}{dz} \left(\frac{c_1 c}{\wp_1 - e_0} \right), \\ s_3 \text{ and } K \text{ as in (20),} \\ g_2 = 2(e_1^2 + e_2^2 + e_3^2) = 10(s_1^2 - s_2 + 2b^2c^2)/(3c^2), \\ g_3 = 4e_1e_2e_3 = 2(5(s_1^2 - s_2) + 4b^2c^2)b/(3c^2), \\ (e_2 - e_3)^2 = (11b^2c^2 + 10(s_1^2 - s_2)), \\ c_1^2 = 4(e_0 + b)^2(2e_0 - b), \quad \sigma_2 = 2c^2(e_0 + b)^2, \\ \Delta = g_2^3 - 27g_3^2 = \frac{4}{27c^6} (11b^2c^2 + 10(s_1^2 - s_2)) (8b^2c^2 - 5(s_1^2 - s_2))^2, \end{array} \right. \quad (22)$$

in which \wp_j is short for $\wp(z - z_j, g_2, g_3)$, $j = 0, 1$, and z_0, z_1 are constants of integration. The equivalence between the conditions $K = 0$ and $\Delta = 0$ is a consequence of the suitable definition of K in (7).

The degeneracy $\Delta = 0$ splits into two cases. For $s_2 = s_1^2 - (8/5)(bc)^2$, the curve $F_4 = 0$ is reducible, this is a particular case of subequation $F_2 = 0$. For $s_2 = s_1^2 + (11/10)(bc)^2$, the curve $F_4 = 0$ has genus zero, therefore its solution is a rational function of one exponential, readily obtained from the odd expression of u in (22) by the degeneracy formula (18), the final result is a solution of (7) rational in hyperbolic tangent,

$$\left\{ \begin{array}{l} u = s_1 - \frac{2cc_1}{3b} \frac{d}{dz} (1 + \tanh^2 kz)^{-1} = s_1 - \frac{4cc_1k}{3b} \frac{(1 - \tanh^2 kz) \tanh kz}{(1 + \tanh^2 kz)^2}, \\ k^2 = \frac{3b}{2}, \quad c_1^2 = -54b^3, \\ K = 0, \quad s_2 = s_1^2 + \frac{11}{10}(bc)^2, \quad s_3 = s_1 \left(s_1^2 + \frac{33}{10}(bc)^2 \right), \end{array} \right. \quad (23)$$

in which the arbitrary origin of z has been omitted.

The solutions (17), (19), (22) and (23) are thus all the meromorphic solutions of (7), and the last two ones, to the best of our knowledge, seem to be new⁴. These exact solutions, in particular the new ones, can be of an important practical use to check the validity of numerical simulations.

4 Comparison with known results

Let us now examine how the complex solutions of (3) found in section 3 compare with some existing results on the real-valued smooth solutions of (2). First we summarize the results obtained so far on (2) for real Z, U, a .

When $a > 0$,

- [33] for almost all $a \in [0, 2\sqrt{2}]$, Eq. (2) admits a homoclinic solution in $H^2(\mathbb{R})$;

⁴After submission, we were informed of similar results on (3) by Kudryashov and Sinelshchikov [23] in which these authors apply another method.

- [32] for all $a \in [0, a_0]$, where $a_0 = 1.228 \dots$ (see footnote on page 2).

Eq. (2) admits a homoclinic solution U such that $U - 1 \in H^2(\mathbb{R})$.

When $a \leq 0$, any bounded smooth solution of (2) has the following properties [1]:

- $a \leq 0 \Rightarrow |U| \leq \sqrt{2}$;
- $a \leq -2\sqrt{2} \Rightarrow |U| \leq 1$.

Let us reduce (3) to (2) by assuming both independent variables Z and z to be equal and real ($\mu = 1$ in (6)). Let us restrict the domain of the four solutions of (2) associated to (17), (19), (22) and (23) to the real line and examine whether those resulting solutions possess one or more of the following properties: real-valuedness, boundedness, homoclinic topology.

Let us first examine the two genus-one solutions of (2). Easily deduced from (17) and (22), they are respectively

$$U = \pm\sqrt{-120} \left(\wp(Z, g_2, g_3) + \frac{a}{60} \right), \quad g_2 = -\frac{a^2 + 10}{180}, \quad g_3 = \frac{a(a^2 + 5)}{5400}, \quad (24)$$

and

$$\left\{ \begin{array}{l} U = \pm\sqrt{\frac{100 - 11a^2}{90} \frac{(a^2 + 25 - 360a\wp(Z, g_2, g_3) - 10800\wp(Z, g_2, g_3)^2)}{2a^2 - 25 + 180a\wp(Z, g_2, g_3) - 10800\wp(Z, g_2, g_3)^2}}, \\ g_2 = \frac{a^2 - 5}{540}, \quad g_3 = \frac{2a^3 - 25a}{162000}, \\ e_1 = -\frac{a}{60}, \quad e_2 = \frac{3a + \sqrt{3(11a^2 - 100)}}{360}, \quad e_3 = \frac{3a - \sqrt{3(11a^2 - 100)}}{360}, \end{array} \right. \quad (25)$$

where the arbitrary origins of Z have been omitted. For the above two solutions, if a is real, then g_2 and g_3 are also real. By Theorem 3.16.2 in [22], \wp (and therefore \wp') is then real-valued for all $Z \in \mathbb{R}$ and \wp has a real period. This implies that the elliptic solution (24) is not real-valued and that the solution (25) is real-valued if and only if $a^2 < 100/11$.

Let us now prove that, when it is real, the solution (25) is bounded on \mathbb{R} . The necessary conditions for U to be extremal are either $\wp(Z) = \infty$ or $\wp'(Z) = 0$ (which implies $\wp(Z) = -a/60$ since e_2, e_3 are not real and $\wp'^2 = 4(\wp - e_1)(\wp - e_2)(\wp - e_3)$) or

$$270a\wp^2(Z) + (a^2 - 50)\wp(Z) + \frac{a(a^2 - 5)}{24} = 0. \quad (26)$$

The last event (26) can only occur for $0 < a < \sqrt{100/11}$. It is because from $\wp'^2 = 4(\wp - e_1)(\wp - e_2)(\wp - e_3)$ and $\wp - e_2 = \overline{(\wp - e_3)}$, we know that $\wp - e_1 \geq 0$ as \wp' is real on \mathbb{R} . Therefore \wp attains the global minimum value $e_1 = -a/60$ when it is restricted on the real line. Now if $-\sqrt{100/11} < a \leq 0$, then it can be checked easily

that the two zeros of the equation $270ap^2 + (a^2 - 50)p + \frac{a(a^2 - 5)}{24} = 0$ are both less than e_1 and this implies that \wp cannot satisfy (26) on the real line if $-\sqrt{100/11} < a \leq 0$.

$$\left\{ \begin{array}{l} U_{1,-} = \mp \sqrt{\frac{10}{9} - \frac{11a^2}{90}} \quad \text{when } \wp(Z) = -\frac{a}{60}, \\ U_{1,+} = \pm \sqrt{\frac{10}{9} - \frac{11a^2}{90}} \quad \text{when } \wp(Z) = \infty, \\ U_{2,\pm} = \pm \sqrt{\frac{10}{9} + \frac{8a^2}{45}} \quad \text{when } \wp(Z) = \frac{50 - a^2 \pm \sqrt{(100 - 11a^2)(4a^2 + 25)}}{540a}. \end{array} \right. \quad (27)$$

Therefore, whenever it is real-valued, the solution (25) is bounded on \mathbb{R} , with the following bounds for U^2 (because of parity, it is sufficient to consider U^2),

$$\left\{ \begin{array}{l} -10/\sqrt{11} < a \leq 0 : 0 \leq U^2 \leq \frac{10}{9} - \frac{11a^2}{90}, \\ 0 \leq a < +10/\sqrt{11} : 0 \leq U^2 \leq \frac{10}{9} + \frac{8a^2}{45}. \end{array} \right. \quad (28)$$

Within one real period, the graph of $U(Z)$ (see Figure 2) displays either two opposite extrema (case $-10/\sqrt{11} < a \leq 0$) or six extrema (case $0 \leq a < 10/\sqrt{11}$) made of two extrema $U_{1,\pm}$ and four extrema $U_{2,\pm}$.

All these results agree with those in [1] mentioned at the beginning of this section.

Let us now examine whether the solutions (19) and (23) of (2) are real and can represent a bounded or homoclinic solution. The first genus zero solution (19) when converted to a solution $U(Z)$ of (2) implies a constraint on a , namely,

$$(4a^2 + 25)(8a^4 + 55a^2 + 200) = 0, \quad (29)$$

which admits no real solution for a , so this solution must be discarded.

The meromorphic solution $U(Z)$ of (2) defined by the second genus zero solution (23) is,

$$U = \pm \frac{4\sqrt{30}(1 - \tanh^2 kZ) \tanh kZ}{\sqrt{11}(1 + \tanh^2 kZ)^2}, \quad k^4 = \frac{1}{176}, \quad a = 40k^2, \quad (30)$$

where the arbitrary origin has been omitted. When restricted to the real Z line, this solution is real-valued if and only if $k^2 = 1/(4\sqrt{11})$ and $a = 10/\sqrt{11}$. It is then bounded on \mathbb{R} and homoclinic, and displays one minimum and one maximum, $U_{\text{extrema}} = \pm\sqrt{30/11} \approx 1.65$, reached for $\tanh kZ = \varepsilon(\sqrt{2} - 1)$, where $\varepsilon^2 = 1$, in agreement with the limit $a \rightarrow 10/\sqrt{11}$ in (27).

To summarize, the exact solution (30) with $k^2 = 1/(4\sqrt{11})$ and $a = 10/\sqrt{11}$ is the unique meromorphic homoclinic solution (up to a translation of origin) mentioned in the introduction. See Figure 1.

The bound of the solution (25) is an increasing function of a . It would be interesting to know whether the same trend will be observed for the class of all bounded smooth solutions of (2), and whether the bound $\sqrt{30/11}$ is sharp for any bounded smooth solution whenever $0 < a < 10/\sqrt{11}$.




Figure 1:
Trigonometric solution $U(Z)$ of (2) defined by (30) when $a = 10/\sqrt{11}$, $k = 1/(2 \times 11^{1/4})$.




Figure 2:
Elliptic solution $U(Z)$ of (2) defined by (25), from left to right:
(i) $a = -1.5$, $b = a/60$, $g_2 \approx 0.0134$, $g_3 \approx 2.731 \times 10^{-4}$, $\Delta \approx 3.6499 \times 10^{-7}$;
(ii) $a = 1.5$, $b = a/60$, $g_2 \approx 0.0134$, $g_3 \approx 2.731 \times 10^{-4}$, $\Delta \approx 3.6499 \times 10^{-7}$.

Remark 4. We could not obtain a codimension zero singlevalued solution of (7). If it exists, this solution, which is **not** meromorphic and therefore not elliptic, is locally represented by the two Laurent series (13), which excludes by construction the contribution of the irrational Fuchs indices $(7 \pm \sqrt{-71})/2$. Such series have been shown to be convergent by Chazy [5]. Although there is no analytic evidence of the existence of this nonmeromorphic solution, a numerical study by Padé approximants in the similar situation of the traveling wave reduction of the Kuramoto-Sivashinsky equation [35, 9] does not yield any evidence of its nonexistence either. It could be possible that this singlevalued, nonmeromorphic solution displays a movable natural boundary.

References

- [1] J.B. van den Berg, Uniqueness of solutions for the extended Fisher-Kolmogorov equation, C. R. Acad. Sci. Paris **326**:447–452 (1998).

- [2] J.B. van den Berg, L.A. Peletier and W.C. Troy, Global branches of multi bump periodic solutions of the Swift-Hohenberg equation, *Arch. Rational Mech. Anal.* **158**:91–153 (2001).
- [3] J.B. van den Berg, *Dynamics and equilibria of fourth order differential equations*, Thesis, Leiden University, 2003.
<http://www.math.vu.nl/~janbouwe/pub/dyn.pdf>
- [4] C. Briot et J.-C. Bouquet, *Théorie des fonctions elliptiques*, 1ère édition (Mallet-Bachelier, Paris, 1859); 2ième édition (Gauthier-Villars, Paris, 1875).
- [5] J. Chazy, Sur les équations différentielles du troisième ordre et d'ordre supérieur dont l'intégrale générale a ses points critiques fixes, *Acta Math.* **34**):317–385 (1911).
- [6] Chiang Yik-man and R. Halburd, On the meromorphic solutions of an equation of Hayman, *J. Math. Anal. Appl.* **281**:663–677 (2003).
- [7] R. Conte, The Painlevé approach to nonlinear ordinary differential equations, *The Painlevé property, one century later*, pp. 77–180, ed. R. Conte, CRM series in mathematical physics, Springer, New York, 1999.
<http://arXiv.org/abs/solv-int/9710020>
- [8] R. Conte and M. Musette, Elliptic general analytic solutions, *Stud. Appl. Math.* **123**:63–81 (2009). <http://arxiv.org/abs/0903.2009>
- [9] R. Conte and M. Musette, *The Painlevé handbook*, Springer, Berlin, 2008.
- [10] R. Conte and T.W. Ng, Meromorphic solutions of a third order nonlinear differential equation, *J. Math. Phys.* **51**:033518(2010).
- [11] V. Croquette, M. Mory and F. Schosseler, Rayleigh-Bénard convective structures in a cylindrical container, *J. Physique* **44**:293–301 (1983).
- [12] M. Cross and H. Greenside, *Pattern Formation and Dynamics in Nonequilibrium Systems*, Cambridge University Press, 2009.
- [13] M.C. Cross and P.C. Hohenberg, Pattern formation outside of equilibrium, *Rev. Modern Phys.* **65**:851–1112 (1993).
- [14] A.E. Eremenko, Meromorphic solutions of equations of Briot-Bouquet type, *Teor. Funktsii, Funktsional'nyi Analiz i vyk Prilozhen., Vyp.* **16**:48–56 (1982). [English : *Amer. Math. Soc. Transl. Ser. 2* **133**:15–23 (1986)].
- [15] A.E. Eremenko, Meromorphic traveling wave solutions of the Kuramoto-Sivashinsky equation, *J. Math. Phys., Anal. Geom.* **2**:278–286 (2006).
- [16] A.E. Eremenko, L.W. Liao and T.W. Ng, Meromorphic solutions of higher order Briot-Bouquet differential equations, *Math. Proc. Cambridge Philos. Soc.* **146**:197–206 (2009).

- [17] S. Greenside and W.M. Coughran, Jr., Nonlinear pattern formation near the onset of Rayleigh-Bénard convection, *Phys. Rev. A* **30**:398–428 (1984).
- [18] W.K. Hayman, *Meromorphic functions*, Clarendon Press, Oxford, 1964.
- [19] E. Hille, *Ordinary differential equations in the complex domain*, J. Wiley and sons, New York, 1976.
- [20] Mark van Hoeij, package “algcures”, Maple V (1997).
<http://www.math.fsu.edu/~hoeij/algcures.html>
- [21] R. Hoyle, *Pattern formation: an introduction to methods*, Cambridge University Press, Cambridge, 2006.
- [22] G. A. Jones, David Singerman, *Complex functions: an algebraic and geometric viewpoint*, Cambridge University Press, Cambridge, 1987.
- [23] N.A. Kudryashov and D.I. Sinelshchikov, Exact solutions of the Swift-Hohenberg equation with dispersion, *Commun.Nonlinear Sci . Numer. Simulat.* **17** 26–34 (2012).
- [24] J. Kwapisz, Uniqueness of the stationary wave for the extended Fisher-Kolmogorov equation, *J. Diff. Eq.* **165**:235–253 (2000).
- [25] I. Laine, *Nevanlinna theory and complex differential equations*, de Gruyter, Berlin and New York, 1992.
- [26] J. Lega, J.V. Moloney and A.C. Newell, Swift-Hohenberg equation for lasers, *Phys. Rev. Lett.* **73**:2978–2981 (1994).
- [27] S. Longhi and A. Geraci, Swift-Hohenberg equation for optical parametric oscillators, *Phys. Rev. A* **54**:4581–4584 (1996).
- [28] K. Maruno, A. Ankiewicz and N.N. Akhmediev, Exact soliton solutions of the one-dimensional complex Swift-Hohenberg equation, *Phys. D* **176**:44–66 (2003).
- [29] M. Musette and R. Conte, Analytic solitary waves of nonintegrable equations, *Phys. D* **181**:70–79 (2003).
- [30] L.A. Peletier and V. Rottschäfer, Pattern selection of solutions of the Swift-Hohenberg equation, *Phys. D* **194**:95–126 (2004).
- [31] L.A. Peletier and W.C. Troy, *Spatial patterns: higher order models in physics and mechanics*, Springer, 2001.
- [32] S. Santra and J. Wei, Homoclinic solutions for fourth order traveling wave equations, *SIAM J. Math. Anal.* **41**:2038–2056 (2009).
- [33] D. Smets and J. B. van den Berg, Homoclinic solutions for Swift-Hohenberg and suspension bridge type equations, *J. Diff. Eq* **184**:78–96 (2002).

- [34] J. Swift and P.C. Hohenberg, Hydrodynamic fluctuations at the convective instability, *Phys. Rev. A* **15**:319–328 (1977).
- [35] Yee T.-l., R. Conte, and M. Musette, Sur la “solution analytique générale” d’une équation différentielle chaotique du troisième ordre, 195–212, *From combinatorics to dynamical systems*, eds. F. Fauvet and C. Mitschi, IRMA Lectures in mathematics and theoretical physics **3**, de Gruyter, Berlin, 2003. <http://arXiv.org/abs/nlin.PS/0302056>