



Title	Emergence of NDM-1-producing Enterobacteriaceae in China
Author(s)	Ho, PL; Li, Z; Lai, EL; Chiu, SS; Cheng, VCC
Citation	Journal of Antimicrobial Chemotherapy, 2012, v. 67 n. 6, p. 1553-1555
Issued Date	2012
URL	http://hdl.handle.net/10722/157694
Rights	This is a pre-copy-editing, author-produced PDF of an article accepted for publication in Journal of Antimicrobial Chemotherapy following peer review. The definitive publisher-authenticated version Journal of Antimicrobial Chemotherapy, 2012, v. 67 n. 6, p. 1553-1555 is available online at: http://jac.oxfordjournals.org/content/67/6/1553

1 Emergence of NDM-1-producing Enterobacteriaceae in China

2 Pak-Leung Ho^{1*}, Zhen Li¹, Eileen L. Lai¹, Susan S. Chiu², Vincent CC Cheng¹

3

4 ¹*Department of Microbiology and Carol Yu Centre for Infection, The University of Hong*

5 *Kong, Queen Mary hospital, Hong Kong*

6 ²*Department of Paediatrics and Adolescent Medicine, The University of Hong Kong, Queen*

7 *Mary hospital, Hong Kong*

8

9 Running title: New Delhi Metallo-beta-lactamases

10 Keywords: carbapenem resistance; carbapenemases; cefotaximases; *Escherichia coli*

11

12 *Corresponding author. Mailing address: Division of Infectious Diseases, Department of

13 Microbiology, The University of Hong Kong, Queen Mary hospital, Pokfulam Road,

14 Pokfulam, Hong Kong SAR, CHINA. Tel: +852-2855 4897; Fax: +852-2855 1241; E-mail:

15 plho@hkucc.hku.hk

16 Word count = 785 (excluding references)

17

18 Sir,

19

20 The worldwide dissemination of bacteria producing New Delhi metallo- β -lactamase (NDM)-
21 is a major public health issue.¹ Since NDM producers are primarily found among individuals
22 with history of hospitalization or travel to the India subcontinent, many hospitals have
23 implemented microbiological screening of patients with such an epidemiological history.^{1,2}

24 In mid-2011, the stool sample of a one year-old infant was found to have CRE upon
25 admission screening. The infant was admitted because of cough and intermittent fever in the
26 preceding two weeks. The family had travelled to and stayed in Hunan province, China in the
27 preceding month. Following onset of the symptoms, the infant had been admitted to a
28 hospital in Hunan for 3 days. Patient was given a diagnosis of broncholitis and had been
29 treated with a course of intravenous cefoperazone. At presentation to our hospital, patient had
30 fever of 38 °C. Chest examination was unremarkable. Patient was treated conservatively and
31 the fever resolved without further antibiotic treatment. Patient was discharged 2 days later. In
32 accordance with the screening policy, stool samples were obtained for surveillance culture.

33 In brief, a red-bean size faecal pellet was suspended in saline. A 10 μ L aliquot of the
34 suspension was then removed and plated directly onto one ChromID ESBL plate. In addition,
35 a broth enrichment step was performed by inoculating another 10 μ L aliquot of the faecal
36 suspension into nutrient broth with 1 mg/L meropenem, incubated at 37 °C overnight, and
37 then subcultured onto MacConkey plate with 1 mg/L meropenem. All plates were incubated
38 at 37 °C in air for 20 h. All distinct colony types recovered from either the chromogenic or
39 the MacConkey media were investigated for evidence of carbapenemase activity using the
40 combined disc method and boronic acid or EDTA as inhibitor.³ All isolates were identified
41 using VITEK 2 and antimicrobial susceptibility was determined by the CLSI disc diffusion
42 method.⁴

43 After the patient's stool samples were found to carry CRE. Stool samples from the
44 infant's parents and other family members were also cultured using the same methodology. A
45 total of four CRE isolates were recovered from the faecal samples of the child and her mother
46 (Table 1). Cultures of the faecal samples from the other household members (father,
47 grandfather, grandmother and aunt) were negative. All four isolates (two *Escherichia coli*,
48 one *Klebsiella pneumoniae* and one *Enterobacter aerogenes*) exhibited synergy with EDTA
49 in the combined disc testing. No synergy with boronic acid was observed. PCR and
50 sequencing, using previously described methods confirmed presence of NDM-1 in the four
51 CRE isolates.² Additional β -lactamases including other metallo- β -lactamases (IMP, VIM,
52 GIM, SPM, SIM), CTX-M and OXA-48-like genes were also sought by PCR and
53 sequencing.⁵ This allowed identification of an additional extended-spectrum β -lactamase
54 (ESBL) gene in the *K. pneumoniae* (CTX-M-65) and the two *E. coli* (CTX-M-57) isolates.
55 Next, we studied the relationship between the two *E. coli* isolates by PFGE after digestion of
56 their genomic DNAs with *Xba*I. The two isolates shared the same PFGE banding pattern,
57 indicating that the two isolates were clonally related. Multilocus sequence typing showed that
58 the two *E. coli* strains and the *K. pneumoniae* strain belonged to ST744 and ST483,
59 respectively. S1-PFGE demonstrated that the strains had one to three plasmids with sizes of
60 50-90 kb. Hybridization demonstrated that *bla*_{NDM-1} gene was harboured on the 50 kb plasmid
61 in all the strains. In conjugation experiments,² the 50 kb *bla*_{NDM-1} harbouring plasmid could
62 be transferred to J53 *E. coli* recipient at high frequencies (up to 10⁻¹ per donor cell). In the
63 transconjugants, there was no co-transfer of resistance to the non- β -lactam antibiotics
64 (amikacin, ciprofloxacin, chloramphenicol, fosfomycin, nalidixic acid, sulphonamides,
65 tetracycline and trimethoprim). Transconjugants with the *bla*_{NDM-1} harbouring plasmid as the
66 only plasmid was investigated by PCR-based replicon typing and *bla*_{CTX-M} PCR. The finding

67 showed that the 50 kb plasmid belonged to an untypeable replicon type and was *bla*_{CTX-M}
68 negative.

69 Members of this family had no history of travel to the Indian subcontinent. The index
70 patient is the only one with a history of hospitalization. Therefore, the infant has mostly
71 likely acquired the *bla*_{NDM-1} gene during the hospitalization in Hunan province. This indicates
72 that some hospitals in mainland China could be reservoirs of the *bla*_{NDM-1} gene,⁶ which may
73 or may not initially have reached there from the Indian subcontinent. While both the infant
74 and her mother shared the same *E. coli* strain, it is impossible to tell if there was intra-familial
75 transmission as opposed to the mother and the infant both becoming colonized while in
76 hospital. In China, the burden of CTX-M-producing Enterobacteriaceae is tremendous.
77 Therefore, accumulation of NDM-1 in multiple CTX-M-producing Enterobacteriaceae
78 species is worrying.¹ In conclusion, this report shows the spread of NDM-1 among persons
79 with no established links to the Indian subcontinent and demonstrates the usefulness of
80 admission screening for early identification of NDM-1 for patients who have been treated
81 aboard.

82

83 **Acknowledgements**

84 This study is supported by a block grant from the Research Fund for the Control of Infectious
85 Diseases (RFCID) of the Health and Food Bureau of the Hong Kong SAR Government. We
86 thank the patient for giving written consent to publication.

87

88 **Transparency declaration**

89 None to declare.

90

91

92 Table 1. Characteristics of four carbapenem-resistant Enterobacteriaceae in this study

	Strain (bacterial species)			
	CRE379 (EA)	CRE380 (KP)	CRE396 (EC)	CRE397 (EC)
Specimen source	index	index	index	index's mother
β -lactamase gene content ^a	NDM-1	NDM-1, CTX-M-65	NDM-1, CTX-M-57	NDM-1, CTX-M-57
Plasmid content (size in kb) ^b	<u>50</u>	<u>50</u>	<u>50</u> , 90	<u>50</u> , 80, 90
Inhibition zone diameter (mm, with/without EDTA)				
Ertapenem	24/9	20/6	16/13	24/6
Imipenem	23/12	25/15	25/15	27/11
Meropenem	24/11	24/10	25/13	27/8
Co-resistance pattern				
Amikacin	S	S	S	S
Chloramphenicol	S	R	R	R
Ciprofloxacin	S	R	R	R
Cotrimoxazole	S	R	R	R
Fosfomycin	M	S	S	S
Gentamicin	S	R	S	S
Nitrofurantoin	R	R	S	S
Tigecycline ^c	M	R	S	S

93 EA, *Enterobacter aerogenes*; EC, *Escherichia coli*; KP, *Klebsiella pneumoniae*

94 ^a β -lactamase group-specific PCR was used to assay for presence of NDM, IMP, VIM, KPC,
95 GES and OXA-48-like genes.

96 ^bThe plasmid showed to harbour the *bla*_{NDM-1} gene by probe hybridization was underlined.

97 ^cThe FDA disc breakpoints was used to interpret tigecycline susceptibility results: sensitive
98 ≥ 19 mm, intermediate 15-18 mm and resistant ≤ 14 mm.

99

References

100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123

1. Nordmann P, Poirel L, Toleman MA *et al*. Does broad-spectrum beta-lactam resistance due to NDM-1 herald the end of the antibiotic era for treatment of infections caused by Gram-negative bacteria? *J Antimicrob Chemother* 2011; **66**, 689-92.
2. Ho PL, Lo WU, Yeung MK *et al*. Complete sequencing of pNDM-HK encoding NDM-1 carbapenemase from a multidrug-resistant *Escherichia coli* strain isolated in Hong Kong. *PLoS One* 2011; **6**, e17989.
3. Giske CG, Gezelius L, Samuelsen O *et al*. A sensitive and specific phenotypic assay for detection of metallo-beta-lactamases and KPC in *Klebsiella pneumoniae* with the use of meropenem disks supplemented with aminophenylboronic acid, dipicolinic acid and cloxacillin. *Clin Microbiol Infect* 2011; **17**, 552-6.
4. Clinical and Laboratory Standards Institute. *Performance standards for antimicrobial susceptibility testing: twenty-first informational supplement M100-S21*. CLSI, Wayne, PA, USA, 2011.
5. Woodford N. Rapid characterization of beta-lactamases by multiplex PCR. *Methods Mol Biol* 2010; **642**, 181-92.
6. Chen Y, Zhou Z, Jiang Y *et al*. Emergence of NDM-1-producing *Acinetobacter baumannii* in China. *J Antimicrob Chemother* 2011; **66**, 1255-9.