



Title	Prevalence and diversity of Synergistetes taxa in periodontal health and disease
Author(s)	You, M; Mo, S; Watt, RM; Leung, WK
Citation	Journal of Periodontal Research, 2013, v. 48 n. 2, p. 159-168
Issued Date	2013
URL	http://hdl.handle.net/10722/163713
Rights	The definitive version is available at www3.interscience.wiley.com

Journal of

PERIODONTAL RESEARCH

**Prevalence and diversity of Synergistetes taxa in
periodontal health and disease**

Journal:	<i>Journal of Periodontal Research</i>
Manuscript ID:	JRE-05-12-1842.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	You, Meng; The University of Hong Kong, Faculty of Dentistry Mo, Sisu; The University of Hong Kong, Faculty of Dentistry Watt, Rory; The University of Hong Kong, Faculty of Dentistry Leung, W. Keung; The University of Hong Kong, Faculty of Dentistry;
Keywords:	Biofilms, Dental plaque, Periodontitis, Phylogeny, RNA, Ribosomal, 16S, Synergistetes

SCHOLARONE™
Manuscripts

proof

1
2
3 **Prevalence and diversity of *Synergistetes* taxa in periodontal**
4
5
6 **health and disease**
7
8

9
10 **Meng You¹, Sisu Mo², Rory M. Watt^{2*} and W. Keung Leung^{1*}**
11
12

13
14
15 **Running title: Human subgingival *Synergistetes***
16
17

18
19 ¹Oral Diagnosis and Polyclinics and ²Oral Biosciences, Faculty of Dentistry, The University
20 of Hong Kong, Prince Philip Dental Hospital, 34 Hospital Road, Sai Ying Pun, Hong Kong
21 of Hong Kong, Prince Philip Dental Hospital, 34 Hospital Road, Sai Ying Pun, Hong Kong
22 SAR, China
23
24

25
26
27
28 *Corresponding addresses:
29

30 Rory M. Watt: Oral Biosciences, Faculty of Dentistry, The University of Hong Kong, Prince
31 Philip Dental Hospital, 34 Hospital Road, Sai Ying Pun, Hong Kong, Tel: (+852) 2859-0482,
32 Fax: (+852) 2547-6133 E-mail: rmwatt@hku.hk
33
34

35
36
37 W. Keung Leung: Oral Diagnosis and Polyclinics, Faculty of Dentistry, The University of
38 Hong Kong, Prince Philip Dental Hospital, 34 Hospital Road, Sai Ying Pun, Hong Kong, Tel:
39 (+852) 2859-0417, Fax: (+852) 2858-2532, E-mail: ewkleung@hkucc.hku.hk
40
41
42
43

44
45
46 Key words: Biofilms; Dental plaque; Periodontitis; Phylogeny; RNA, Ribosomal, 16S;
47
48 *Synergistetes*
49

50
51
52 Date of resubmission: June 18, 2012
53
54
55
56
57
58
59
60

1
2
3 You M, Mo S, Watt RM, Leung WK. Prevalence and diversity of *Synergistetes* taxa in
4
5 periodontal health and disease. *J Periodont Res*
6
7
8

9 10 **Abstract**

11
12 *Background and Objective:* Members of the phylum *Synergistetes* have previously been
13
14 identified within periodontitis subgingival plaque and are considered putative
15
16 periodontopathogens. This study compared the diversity of subgingival *Synergistetes* in a
17
18 cohort of subjects with periodontitis (n=10) versus periodontitis-free controls (n=10).
19
20

21
22
23 *Materials and Methods:* Pooled subgingival plaque samples from all deep periodontal
24
25 pockets or all sulci were collected from the periodontitis and periodontitis-free subjects,
26
27 respectively. Bacterial 16S rRNA genes were PCR-amplified from purified subgingival
28
29 plaque DNA using a *Synergistetes* 'selective' primer set. PCR products were cloned and
30
31 sequenced to analyze the prevalence and diversity of *Synergistetes* operational taxonomic
32
33 units (OTUs) present in plaque samples of both subject groups.
34
35
36

37
38
39 *Results:* A total of 1030 non-chimeric 16S rRNA clones were obtained, of which 162
40
41 corresponded to members of the phylum *Synergistetes*. A significantly larger number of
42
43 *Synergistetes* clones were obtained from periodontitis subgingival plaque than from
44
45 periodontitis-free controls (25.4% vs. 5.9%, $p<0.001$). All *Synergistetes* clones corresponded
46
47 to cluster A oral *Synergistetes*, and fell into 31 OTUs (99% sequence identity cut-off). 29
48
49 *Synergistetes* OTUs were detected in the periodontitis group while 8 were detected in the
50
51 periodontitis-free group ($p<0.001$). Five *Synergistetes* OTUs; including one OTU
52
53 corresponding to the recently-characterized species *Fretibacterium fastidiosum*, were more
54
55 prevalent in the periodontitis subjects ($p<0.05$).
56
57
58
59
60

1
2
3
4
5 *Conclusion:* OTUs belonging to oral *Synergistetes* cluster A were more readily detectable and
6
7 were more diverse in subgingival plaque from periodontitis subjects compared with
8
9 periodontitis-free controls. Specific *Synergistetes* OTUs appear to be associated with
10
11 periodontitis.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Manuscript proof

1
2
3 Periodontitis is a chronic polymicrobial infectious inflammatory disease affecting the
4
5 periodontium; which causes the progressive destruction of the soft and hard tooth supporting
6
7 tissues (1, 2). It is generally considered to result from microbial dysbiosis, i.e. a 'pathogenic
8
9 shift' in the composition of the subgingival microbia that promotes and potentiates
10
11 pathogenic processes (3, 4). Of the 13 bacterial phyla known to inhabit the oral cavity (5), 9
12
13 have been detected in the subgingival plaque from both periodontally healthy (6) and
14
15 diseased subjects (7). The predominant phyla found in the periodontal niches are *Firmicutes*,
16
17 *Proteobacteria*, *Bacteroidetes* and *Actinobacteria*; which have been estimated to account for
18
19 more than 90% of the total bacteria present (6). In surveys of the oral microbiome that have
20
21 utilized 'universal' 16S rRNA molecular detection protocols, *Synergistetes* was defined as a
22
23 'rare' phylum, as it had a relatively low detection frequency of ca. 0.1-2% (6, 8-10). However,
24
25 other estimations based on microscopic fluorescence *in situ* hybridization (FISH) studies
26
27 have indicated that *Synergistetes* taxa may account for as much as 11% of the microbiota
28
29 within deep pockets (11).
30
31
32
33

34 Members of the *Synergistetes* inhabit diverse ecological environments (12, 13). They
35
36 are fastidious, gram-negative anaerobic straight or curved bacilli, some of which are motile
37
38 (14, 15). The *Synergistetes* phylum was only recently described, having 9 component genera
39
40 (16). A significant number of taxa, most of which were previously placed in the phyla
41
42 *Deferribacteres* (7, 10, 17-19) and *Firmicutes* (20, 21), were incorporated into this new
43
44 phylum. In previous culture-independent surveys of oral bacterial diversity, 16S rRNA gene
45
46 sequences corresponding to members of the phylum *Synergistetes* have been detected in a
47
48 variety of oral niches; especially within diseased periodontal pockets, endodontic infections
49
50 and carious dentine (6-9, 22, 23). The oral *Synergistetes* have been divided into two principal
51
52 clusters, based on their phylogenetic origins (11, 14). Cluster A is the larger group
53
54 comprising mainly uncultivated phylotypes, with one member *Fretibacterium fastidiosum*
55
56
57
58
59
60

1
2
3 (also referred to as SGP1) isolated and characterized recently (24, 25). Cluster B is
4
5 considerably smaller, containing two cultivated species: *Pyramidobacter piscolens* (26) and
6
7 *Jonquetella anthropi* (27).
8

9
10 The *Synergistetes* population appears to be more diverse in periodontitis niches than
11
12 in dental caries or endodontic infections (14). The presence or (clonal) abundance of certain
13
14 *Synergistetes* taxa have previously been shown to be associated with periodontal disease, and
15
16 consequently certain members of this phylum may be considered putative periodontopathogens
17
18 (7, 11, 28). In a recent UK-based study, Vartoukian *et al.* compared the diversity and
19
20 detection frequency of *Synergistetes* OTUs within several oral sites in a cohort of subjects
21
22 with periodontitis (n=5) versus subjects with good periodontal health (n=5) (11). The authors
23
24 found that the richness of *Synergistetes* OTUs, and the prevalence and abundance of one
25
26 taxon in particular (OTU 4.2), were associated with periodontitis. Notwithstanding these
27
28 results however, we still have a very limited understanding of the genetic diversity,
29
30 distributions and disease-associations of oral *Synergistetes* taxa in global populations.
31
32

33
34 To directly address these issues, we employed a 16S rRNA clone library-based
35
36 approach to investigate the diversity and prevalence of members of the *Synergistetes* phylum
37
38 in the subgingival plaque of Chinese subjects with periodontitis, versus periodontitis-free
39
40 controls. We found that *Synergistetes* clones were significantly more prevalent and more
41
42 diverse in periodontitis subgingival plaque compared with periodontitis-free controls. The
43
44 presence of certain *Synergistetes* OTUs (phylotypes) was associated with periodontitis. Our
45
46 results add to the growing body of evidence supporting the putative role of members of this
47
48 phylum in the pathogenesis of periodontal disease.
49
50

51 52 53 54 55 56 **Materials and methods** 57 58 59 60

Subject recruitment, clinical examination and sampling methods

Ten periodontitis patients and ten periodontitis-free controls were recruited for this study with informed consent. Ethical approval was granted by the Institutional Review Board of the University of Hong Kong/Hospital Authority, Hong Kong West Cluster (UW 11-154). All subjects were non-smokers or former smokers who had quit smoking for more than 6 months. All subjects were ethnically Chinese, who lived in Hong Kong. Exclusion criteria included a history of periodontal or antimicrobial therapy within the past 6 months, any underlying systemic medical conditions; or less than 20 standing teeth. The inclusion criteria for the periodontitis group were four or more posterior teeth with periodontal pockets of ≥ 6 mm and clinical attachment level (CAL) of ≥ 4 mm, plus radiographic evidence of alveolar bone loss. The periodontitis-free subjects had no probing depths of ≥ 3 mm, no detectable clinical attachment loss or radiographic bone loss. Full mouth bleeding on probing (BOP) scores and probing pocket depths (PPD) were recorded to characterize the periodontal status. After scrupulous removal of supragingival plaque, subgingival plaque samples were collected from as many periodontal pockets (sites with PPD ≥ 5 mm in periodontitis subjects) or sulci (periodontitis-free subjects) using sterile Gracey or universal curettes, and were pooled to generate a single subgingival plaque sample for each subject ($n = 20$). The collected plaque samples were suspended in 1.0 ml phosphate buffered saline (PBS, pH 8.0), placed on ice, and processed in the laboratory within 30 minutes after collection.

DNA extraction, amplification, cloning and sequencing

Subgingival plaque samples were washed twice (PBS, pH 7.4, 1 ml), and DNA was extracted (Wizard Genomic DNA Purification Kit, Promega) following the manufacturer's protocol. DNA pellets were dissolved in 100 μ l of rehydration buffer. The universal forward primer TPU1 (5'-AGAGTTTGATCMTGGCTCAG-3') (29) and the C90 reverse primer (5'-

1
2
3 GTTACGACTTCACCCTCCT-3') (5, 19) were used to PCR-amplify ca. 1500bp 16S rRNA
4
5 gene fragments, using the purified DNA from subgingival plaque samples as templates. PCR
6
7 mixtures (50 µl) contained: (5×) PCR buffer (10 µl), template DNA (2 µl, ca. 50 ng), MgCl₂
8
9 (2.5 mM), dNTPs (0.2 mM each), TPU1 and C90 primers (0.4 µM each), LongAmp DNA
10
11 polymerase (1.5 U, New England Biolabs). PCR reactions were performed using a
12
13 'touchdown' protocol on a GeneAmp PCR System 9700 (Applied Biosystems): initial
14
15 denaturation (94°C, 480s); then 6 cycles of: denaturation (94°C, 30s), annealing (52°C, 30s,
16
17 decreasing 0.5°C every cycle), extension (65°C, 150s); followed 23 cycles of denaturation
18
19 (94°C, 20s), annealing (49°C, 20s), extension (65°C, 120s); final extension (65°C, 600s).
20
21 PCR products of ca. 1500bp were gel purified (QIAquick Gel Extraction Kit, Qiagen).
22
23 Purified PCR amplicons from each subject were individually TOPO-cloned into pCR2.1
24
25 vectors (TOPO TA Cloning Kit, Invitrogen). Ligation mixtures (n=20) were electroporated
26
27 into *Escherichia coli* DH10B, plated onto Luria-Bertani (LB) 1% agar plates supplemented
28
29 with kanamycin (50µg/ml) and X-gal (5-bromo-4-chloro-indolyl-β-D-galactopyranoside,
30
31 20µg/ml), then incubated overnight at 37°C. Plasmid DNA was purified from 45-60 (white)
32
33 colonies from each transformant plate (QIAprep Spin Miniprep Kits, QIAgen). 45-60
34
35 plasmids from each subject (were subsequently sequenced in both directions using M13
36
37 forward and reverse primers [Beijing Genome Institute (BGI-Hong Kong Co. Ltd), Tai Po,
38
39 Hong Kong].
40
41
42
43
44
45
46

47 **Primer and 16S rRNA gene sequence analysis**

48
49 The selectivity and coverage of the C90 reverse primer (5'-GTTACGACTTCACCCTCCT-3')
50
51 for amplifying 16S rRNA gene sequences from members of the *Synergistetes* phylum was
52
53 estimated *in silico*, using the Probe Match programme of the Ribosomal Database Project
54
55 (RDP) 10 online resource (<http://rdp.cme.msu.edu/>) (30). The 16S rRNA gene sequence data
56
57
58
59
60

1
2
3 were assembled and trimmed using CODONCODE ALIGNER 3.7.1 (Codon Code
4 Corporation, Dedham, MA). Sequences were aligned using the 16S ribosomal SILVA
5 bacteria dataset (31) as a template, and were manually corrected before further analysis.
6
7 Chimeric sequences were identified using Chimera Slayer and 24 suspicious sequences were
8 removed. RDP Classifier was used to taxonomically-classify valid sequences to at least the
9 level of phylum (32).
10
11
12
13
14
15
16
17

18 **Taxonomic assignment Estimates of microbial diversity**

19
20 A distance matrix was generated and sequences were assigned to Operational Taxonomic
21 Units (OTUs) at various sequence identity cut-off levels (99%, 97%, 95%) using the furthest
22 neighbor algorithm in the Mothur software package (32). This algorithm requires that the
23 distance between every sequence within an OTU lie within the specified identity threshold
24 (33). Representative sequences for each *Synergistetes* OTU found in this study were
25 deposited in the NCBI GenBank (accession numbers JX010869 to JX010946, Supplementary
26 Table S2). Rarefaction curves for OTU were calculated using Mothur.
27
28
29
30
31
32
33
34
35
36
37

38 **Community comparisons**

39
40 *Synergistetes* OTU populations in each subject were compared and visualized as
41 dendrograms according to the Yue and Clayton theta structural diversity measure (34). The
42 structure of *Synergistetes* communities present in the periodontitis and **periodontitis-free**
43 groups were compared using β -libshuff (35, 36), unweighted and weighted UniFrac analysis
44 (37, 38) and Parsimony p -tests (39). The p -value threshold was set to 0.05.
45
46
47
48
49
50
51
52
53

54 **Phylogenetic relationships**

1
2
3 An alignment comprising representative 16S rRNA sequences from each identified non-
4 *Synergistetes* OTU were used as an input to examine phylogenetic relationships. NJ trees
5 were constructed following evolutionary distance analysis of aligned sequences, corrected by
6 the Jukes-Cantor substitution model using MEGA 5 (40), and were supported by 100
7 bootstrap replicates. The absolute clonal abundance for each OTU was visualized as a heat-
8 map using ggplot2 from the R statistical computing package (<http://www.r-project.org/>) (41).
9
10
11
12
13
14
15
16
17

18 **Statistical analysis**

19
20 Standard descriptive statistical analysis was conducted to synthesize the clinical data.
21 Assumption tests for normality and equality of variance were performed prior to all statistical
22 comparative analyses of datasets obtained from the two cohorts. Non-parametric Mann-
23 Whitney *U* tests were used for all statistical tests between the two cohorts. The *p*-value
24 threshold was set to 0.05.
25
26
27
28
29
30
31
32
33
34
35
36

37 **Results**

38 **Clinical evaluation of periodontal health status in the two subject groups**

39
40 Subjects with periodontitis who sought treatment at the Prince Philip Dental Hospital and
41 subjects with periodontitis-free attending regular check-up were recruited for this study, with
42 informed consent and ethical approval. There was no difference in gender distribution
43 between periodontitis and periodontitis-free control groups; but the average age of the
44 periodontitis-free group was slightly higher than that of the periodontitis group (Table 1). The
45 periodontal health status of all subjects was as summarized in Table 1. The mean percentage
46 of BOP sites for the periodontitis group was significantly higher than that for the
47 periodontitis-free group. None of the periodontitis-free group had probing pocket depths ≥ 4
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 mm, radiographic evidence of bone loss or furcation involvement in multi-rooted teeth (data
4 not shown), confirming that they had good periodontal status. All subgingival sites in the
5 periodontitis-free group were sampled to maximize the total amount of plaque collected
6 (Table 1). Whilst it was possible to collect subgingival plaque from every diseased
7 periodontal pocket sampled in the periodontitis group, it was not possible to obtain detectable
8 levels of plaque from every sulcal site in the periodontitis-free group. As may be seen in
9 Table 1, the sampled sites from the periodontitis subjects had a significantly higher average
10 probing depth and percentage of BOP.
11
12
13
14
15
16
17
18
19

20 21 22 ***In silico* evaluation of PCR primer amplification specificity**

23
24 In combination with a ‘universal’ forward PCR primer [e.g. TPU1 (29)], the C90 reverse
25 primer (5, 19) has previously been shown to predominantly amplify near-full length 16S
26 rRNA (*rrs*) gene fragments (corresponding to positions 8-1503 of the *Escherichia coli* 16S
27 rRNA gene) from members of the *Spirochaetes* (7, 19, 42, 43) and *Synergistetes* phyla
28 (previously referred to as *Synergistes/Flexistipes/Deferribacteres* taxa) (19). To
29 systematically evaluate the hybridization specificity of the C90 primer, we used the Probe
30 match software tool to analyze its sequence homology with 16S rRNA sequences from
31 representative *Synergistetes* OTUs. Results indicated that the C90 primer should anneal with
32 cluster B oral *Synergistetes* sequences without mismatches (Supplementary Table S1). The
33 C90 primer exactly matched some cluster A oral *Synergistetes* sequences, whilst there was a
34 single G/A nucleotide mismatch with other sequences at position 1485 (*E. coli* numbering).
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 52 **Construction and analysis of 16S rRNA clone libraries**

53
54 Using the TPU1 plus C90 primer set, ca. 1500bp 16S rRNA gene fragments were
55 successfully amplified from DNA template purified from the pooled subgingival plaque
56
57
58
59
60

1
2
3 samples from each of the 20 subjects. Plasmid libraries of 'TOPO-cloned' 16S rRNA genes
4
5 were separately constructed for each subject, and the PCR amplicons within ca. 45-60
6
7 plasmid clones from each subject were sequenced. After poor-quality reads and chimeras
8
9 were filtered out, a total of 1,030 high-quality 16S rRNA sequences were obtained. There
10
11 was no significant difference in the mean number of plasmid clones obtained from the
12
13 periodontitis (52 ± 8.0) and periodontitis-free control (51 ± 8.4) groups. 16S rRNA sequences
14
15 were assigned a taxonomy using RDP classifier (32), with results summarized in Table 2. In
16
17 terms of clonal abundance (i.e. number of plasmid clones obtained), the predominant phyla
18
19 detected were *Spirochaetes* (59.7%), *Synergistetes* (15.7%), *Actinobacteria* (16.2%),
20
21 *Fusobacteria* (3.6%), *Proteobacteria* and *Firmicutes* (2.4%) and *Proteobacteria* (2.3%). This
22
23 indicated that whilst the TPU1 plus C90 primer set predominantly amplified 16S rRNA genes
24
25 from *Spirochaetes*, they exhibited a reasonably-good selectivity for *Synergistetes*. The
26
27 composition of the *Spirochaetes* clones obtained will be discussed elsewhere (You *et al.*,
28
29 manuscript in preparation).
30
31
32
33

34 There were significantly higher numbers of *Synergistetes* clones amplified from
35
36 periodontitis subjects compared with periodontitis-free controls (Table 2). Of the 162
37
38 *Synergistetes* clones detected, 132 (>80%) clones were obtained from the periodontitis
39
40 subject group. The mean number of *Synergistetes* clones obtained per subject was 13.2 (range:
41
42 8-21) and 3.0 (range: 0-7) from the periodontitis and periodontitis-free groups, respectively.
43
44

45 Compared with the periodontitis group, a relatively large proportion of clones
46
47 corresponding to members of the *Actinobacteria*, and a sizable number of clones belonging to
48
49 *Firmicutes*, *Fusobacteria* and *Proteobacteria* phyla were detected in the periodontitis-free
50
51 group (See Table 2). However, as the TPU1 and C90 primers do not have good coverage for
52
53 members of these phyla (data not shown), these findings are probably not representative of
54
55 the actual populations of these bacteria present; and as such, correlations with clinical health
56
57
58
59
60

1
2
3 status was not performed. As has been previously proposed (7, 11, 19), this may possibly
4
5 reflect the low abundance of stringently-matched taxa (e.g. *Spirochaetes*, *Synergistetes*),
6
7 resulting in the amplification of more abundant and less-stringently matched 16S rRNA
8
9 sequences (especially *Actinobacteria* spp.).
10

11 12 13 14 **Prevalence of *Synergistetes* taxa within the periodontitis and periodontitis-free subject** 15 16 **groups**

17
18 The *Synergistetes* clones were divided into distinct taxonomic groups, i.e. OTUs, based on
19
20 the levels of shared 16S rRNA gene sequence identity. The prevalence of OTUs in both
21
22 subject groups were analyzed using three different sequence identity cut-off values: 99%, 97%
23
24 and 95% (Table 3). Rarefaction curves were plotted using this data to estimate the OTU
25
26 richness within the two subject groups at each of these levels (see Supplementary Figure S1).
27
28 Graphs were plotted based on four different parameters: the numbers of subjects sampled
29
30 (Panel A); as well as the total number of clones sampled from both groups (Panel B), the
31
32 periodontitis-free group only (Panel C) or the periodontitis group only (Panel D). In all four
33
34 panels it may be seen that all the rarefaction curves plotted using OTU composition based on
35
36 a 97% or 95% sequence identity cut-off level plateaued-out at low OTU richness values,
37
38 whilst curves plotted used a 99% sequence identity cut-off did not plateau over the range
39
40 shown. When rarefaction curves were respectively plotted for the number of OTUs observed
41
42 versus number of clones sequenced for the two cohorts using a 99% cut-off value, only the
43
44 curve for the periodontitis-free group appeared to reach an asymptote. This result was
45
46 consistent with the Chao 1 diversity estimates described below.
47
48
49
50

51
52 Using a 99% 16S rRNA sequence identity cut-off, a total of 31 *Synergistetes* OTUs
53
54 were present in the overall dataset (Table 3). These OTUs were given alphanumeric names
55
56 reflecting their origin (1P-10P for the 10 periodontitis subjects, 1H-10H for the periodontitis-
57
58
59
60

1
2
3 free subjects). It should be noted that some 'P' OTUs were detectable in both periodontitis
4 and **periodontitis-free** subjects; with the name indicating the subject group from which the
5 representative clones were first obtained. Substantially more OTUs were detected in the
6 periodontitis group (n=29), compared with the periodontitis-free controls (n=8). Twenty three
7 OTUs were detected exclusively in the periodontitis group, six OTUs were common to both
8 groups (5H23, 1P28, 7P1, 4P26, 8H6, 4P12), whilst only two OTUs (3H20, 5H10) was found
9 exclusively in the periodontitis-free group (also see Figure 1). The mean number of
10 *Synergistetes* OTUs detected per subject was 6.7 (SD \pm 3.0; range: 3-13) in the periodontitis
11 group and 1.7 (SD \pm 1.4; range: 0-4) in the periodontitis-free group (p <0.001, Mann-Whitney
12 U test). The total estimated number of *Synergistetes* OTUs (as calculated using a Chao 1
13 index) was 53.8; with a projected 47.5 and 8.5 OTUs present in the periodontitis and
14 periodontitis-free groups, respectively.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The prevalence of the 31 *Synergistetes* OTUs within the subjects, and their clonal abundance (i.e. absolute numbers of plasmids encoding 16S rRNA genes that correspond to a specific OTU) were represented using a grey-scale heatmap (Figure 1). The three most widely distributed OTUs were 4P26 (detected in 10 subjects), 7P1 (9 subjects) and 2P9 (7 subjects). Sixteen OTUs were present in two or more subjects, whilst 15 OTUs were only found in one subject. However, it should be noted that individual OTUs defined using a 99% sequence identity cut-off may comprise several closely-related *Synergistetes* 16S rRNA sequences. Non-parametric Mann-Whitney U tests revealed that five *Synergistetes* OTUs were detected with a significantly higher frequency in the periodontitis group, compared with the periodontitis-free group. These were OTUs 2P9 (p <0.01; analogous to *Synergistetes* OTU 2 clone 7.1 (25)); 4P12 (p <0.01; analogous to *F. fastidiosim* SGIP^T (1435/1440 nt, 99.7%; previously known as *Synergistetes* SGP1) (24, 25) and the '*Deferribacteres*' oral clone W090 clone (1434/1440 nt, 99.6% identity) (17)); 7P1 (p <0.01, analogous to '*Deferribacteres*' oral

1
2
3 clone BH007 (1433/1434 nt, 99.9% identity) (7)); 7P22 ($p<0.05$) and 6P18 ($p<0.05$); also
4
5 discussed below.
6
7
8

9 10 **Phylogenetic relationships between *Synergistetes* OTUs**

11
12 Phylogenetic relationships between the 31 *Synergistetes* OTUs identified in the clinical
13
14 samples were analyzed using a Neighbor-joining (NJ) approach. The overall topology of the
15
16 NJ phylogenetic tree generated was generally well-supported with high bootstrap values
17
18 (Figure 1). It may be seen that the OTUs belonged to various well-separated genetic lineages
19
20 that diverged at an early point in their evolutionary history. Several OTUs appeared to be
21
22 quite genetically distinct (e.g. 6P18, 2P9, 3P46), whilst others were part of ‘branches’
23
24 containing two or three OTUs that shared reasonably high levels (ca. 98.5-99.5%) of 16S
25
26 rRNA gene sequence similarity (e.g. 4P36, 4P12, 3P2).
27
28

29
30 Figure 2 shows the phylogenetic relationships between the thirty-one OTUs detected
31
32 in the subjects analyzed here and the three named *Synergistetes* species previously isolated
33
34 from the oral cavity, as well as the corresponding *Synergistetes* Human Oral Taxa (HOT)
35
36 defined in the Human Oral Microbiome Database (HOMD) (5) Twenty-three of the thirty-one
37
38 OTUs identified here could be assigned to an existing *Synergistetes* HOT (using a 98.5%
39
40 sequence identity cut-off). Eight OTUs (6P48, 4P10, 5P11, 8P65, 4P40, 3H20, 4P23, 7P51)
41
42 did not correspond to an existing *Synergistetes* HOT. We did not detect any clones or isolates
43
44 corresponding to oral cluster B *Synergistetes* within either of our subject groups.
45
46
47
48

49 50 **Community-based analysis of *Synergistetes* OTUs detected in subjects**

51
52 A hierarchical OTU-based comparison analysis was performed to determine whether the two
53
54 subject groups contained similar or disparate communities of *Synergistetes* OTUs (99% cut-
55
56 off). As may be seen in the dendrogram shown in Figure 3, based on their resident
57
58
59
60

1
2
3 populations of identified OTUs, the subjects formed two quite distinct clusters that correlated
4
5 closely to their periodontal health status. The only notable outlier was periodontitis subject P1,
6
7 whose resident *Synergistetes* clones were similar to those present in healthy subjects H7 and
8
9 H4. J-libshuff, unweighted and weighted UniFrac and Parsimony *p*-tests were all used to
10
11 systematically compare the differences in communities of *Synergistetes* OTUs present in the
12
13 periodontitis and periodontitis-free subject sets. All these hypothesis tests (that employ
14
15 different algorithms) indicated that the populations within the two groups were significantly
16
17 different, with overall *p*-values of less than 0.001 obtained.
18
19
20
21
22
23
24

25 Discussion

26
27
28 In this study, we used a 16S rRNA gene-based approach to systematically analyze and
29
30 compare members of the *Synergistetes* phylum that were present in pooled subgingival
31
32 plaque sampled from periodontal pockets of 10 periodontitis patients, versus pooled
33
34 subgingival plaque collected from the sulci of an equal number of periodontitis-free controls.
35
36 Of the 162 *Synergistetes* clones identified, significantly more clones were obtained from the
37
38 periodontitis group (n=132, 81.5%) than the periodontitis-free group (n=30, 18.5%; *p*<0.001),
39
40 consistent with the hypothesis that certain members are 'periodontopathogens'.
41
42

43
44 Subgingival niches in subjects with periodontal or endodontic infections are populated
45
46 with anaerobic microbial biofilm communities that are able to produce peptides and free
47
48 amino acids through the proteolytic degradation of host tissues (44). Such niches appear to be
49
50 ideal environments for members of the *Synergistetes* taxa, which are proposed to acquire
51
52 nutrients predominantly via amino acid metabolism (14). Correspondingly, a significant
53
54 number of *Synergistetes* taxa have been detected or isolated from diseased periodontal sites
55
56 (7, 10, 17, 20, 45, 46) and endodontic infections (19, 21, 47-51); as well as dental caries
57
58
59
60

1
2
3 lesions (52). Using a similar 16S rRNA clone-based approach, it has previously been shown
4
5 that the *Synergistetes* levels detected in subgingival plaque was associated with periodontal
6
7 health status; constituting ca. 1.5% of all clones obtained from periodontal pockets, but only
8
9 0.1% of clones from periodontally-healthy sites (10). Although *Synergistetes* have been
10
11 frequently identified in periodontitis subjects, several studies targeting periodontal healthy
12
13 subjects have failed to detect taxa belonging to the *Synergistetes* phylum (53, 54). In our
14
15 cohort, *Synergistetes* could be detected, albeit at low frequencies, in 8/10 periodontitis-free
16
17 subjects, and in all periodontitis subjects. In agreement with previous reports (6), our results
18
19 support the hypothesis that *Synergistetes* are typically present in 'healthy' human oral
20
21 microbiota at low levels.
22
23

24
25 Taxa within the *Synergistetes* phylum share less than 75% 16S rRNA sequence
26
27 identity with other phyla. *Synergistetes* taxa that have previously been (co)-detected within
28
29 specific niches; such as in animal hosts, soil samples, or human subgingival plaque generally
30
31 share only ca. 80% 16S rRNA sequence similarity (12, 16). *Synergistetes* thus far identified
32
33 within the oral cavity form two major clusters: A and B, with the majority of taxa belonging
34
35 to cluster A (14, 25). To date, only one species of cluster A oral *Synergistetes* has been
36
37 isolated and characterized: *Fretibacterium fastidiosum* (24), which was previously referred to
38
39 as the SGP1 clone (25). The smaller cluster B includes 2 cultivated species: *Pyramidobacter*
40
41 *piscolens*, with several strains isolated from odontogenic abscesses, inflamed gingival
42
43 crevices or periodontal pockets; and *Jonquetella anthropi*, with strain E3_33 isolated from an
44
45 endodontic infection (14, 26, 27).
46
47
48

49
50 *Synergistetes* populations within periodontitis subgingival plaque were previously
51
52 reported to be more diverse than those in endodontic infections or dental caries, and certain
53
54 taxa were identified exclusively in periodontal sites (14). Using a '*Synergistetes* specific'
55
56 primer set that amplified a ca. 800bp fragment of the 16S rRNA gene, Vartoukian *et al.* (11)
57
58
59
60

1
2
3 detected 12 *Synergistetes* OTUs (using a 99% cut-off value) in plaque sampled from the
4
5 periodontal environment in 5 healthy and 5 periodontitis subjects. Here, we used the
6
7 ‘*Synergistetes* selective’ TPU1+C90 primers to PCR-amplify near full-length 16S rRNA
8
9 fragments (ca. 1500bp), enabling us to identify 31 *Synergistetes* OTUs. 12 of the
10
11 *Synergistetes* OTUs in our dataset corresponded to known cluster A members, and 19
12
13 corresponded to previously undescribed OTUs. Analogous to Vartoukian *et al.* (11), we
14
15 defined the *Synergistetes* OTUs using a 99% sequence identity cut-off. Whilst this OTU
16
17 definition gives higher levels of taxonomic resolution, it may over-estimate the numbers of
18
19 ‘species-level’ phylotypes present. We detected 16S rRNA sequences belonging to OTUs that
20
21 corresponded to all 15 of the previously-identified oral *Synergistetes* cluster A taxa listed in
22
23 Supplementary Table S1, even though many of these putatively encode a G/A single
24
25 nucleotide mismatch with the C90 reverse primer. This indicated that we were able to sample
26
27 a broad range of oral *Synergistetes* taxa using our approach. It should be noted however, that
28
29 the recently-reported M98 *Spirochaetes* and *Synergistetes* selective reverse primer (5)
30
31 contains a degenerate Y (C/T) residue in place of a T residue which is present in the C90
32
33 primer, and should theoretically have a better range of coverage.
34
35
36
37

38
39 Consistent with previous reports (11), we did not detect any oral cluster B taxa in our
40
41 subject groups. Other molecular analyses of oral (especially periodontitis) microbiota have
42
43 similarly noted that the vast majority of *Synergistetes* taxa (previously referred to as
44
45 *Deferibacteres/Flexistipes/Synergistes*) corresponded to cluster A, with relatively few
46
47 cluster B taxa detected (7, 10, 17, 45). Sequence analysis indicated that the C90 reverse
48
49 primer can perfectly complement all previously-identified cluster B OTUs, indicating that
50
51 they should be detected if present at levels comparable to those of the cluster A taxa (see
52
53 Supplementary Table S1). Consequently, it is not possible to state whether oral cluster B
54
55 *Synergistetes* are absent, or are present at very low levels within our subject set.
56
57
58
59
60

1
2
3 The communities of *Synergistetes* were significantly more OTU-rich in the
4 periodontitis subjects (mean: 6.7 OTUs, range: 3-13) compared with the periodontitis-free
5 subjects (mean: 1.7, range: 0-4). This is in good agreement with the results of Vartoukian *et*
6 *al.* (11), who noted that periodontitis cases (mean: 7.8 OTUs, range: 7-10) also had
7 significantly higher OTU-richness than controls (mean, 3.4, range 0-6). These authors also
8 noted that certain individuals could be 'dominated' with a specific *Synergistetes* OTU (up to
9 64% relative clonal abundance), whilst others contained a greater OTU diversity with a fairly
10 even prevalence. Similarly, in our data, control subject H7 contained 7 clones corresponding
11 to a single OTU (1P28), whilst periodontitis subject P6 harbored 13 different OTUs, which
12 were each present on 1-6 plasmid clones.
13
14
15
16
17
18
19
20
21
22
23
24

25 In their landmark (US-based) study published in 2001, Paster and co-workers
26 identified *Synergistetes* clones BH017, D084, W028 and W090 as putative periodontal
27 pathogens on the basis of their prevalence in diseased subjects, and absence in healthy
28 subjects (7). Using primer sets specific for clones BH017/D084 and clone W090, Kumar and
29 co-workers subsequently found that the BH017 and D084 clones, but not W090, were
30 associated with chronic periodontitis in a US cohort (17). Based on results from a 16S rRNA
31 clone-based investigation, Kumar *et al.* later found that the occurrence and prevalence of
32 *Synergistetes* clones BH007 and W090, but not BH017, were strongly associated with
33 periodontal disease (10). In a UK-based study, Vartoukian *et al.* (11) subsequently found that
34 a previously undescribed *Synergistetes* taxon (OTU 4.2) was significantly associated with
35 periodontitis; whilst other phylotypes including: BH007, W090 and W028 had no significant
36 correlation with disease. **Notably, we did not detect a sequence corresponding to OTU 4.2**
37 **within our cohort.** Therefore, despite consistent results linking members of the *Synergistetes*
38 phylum with (chronic) periodontitis, the association between specific *Synergistetes* clones
39 and periodontal disease remains somewhat inconclusive.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Here, we found that the increased prevalence of 5 *Synergistetes* OTUs were
4 associated with periodontitis: 7P1, 2P9 and 4P12 ($p<0.01$); and 7P22 and 6P18 ($p<0.05$).
5
6 Whilst OTUs 7P22 and 6P18 are novel; OTU 7P1 corresponds to clone BH007 (10); 2P9
7 corresponds to clone 7.1 (25). OTU 4P12 shares >99.6% sequence identity with clone W090
8 (7, 10) and *F. fastidiosum* SGIP^T (24, 25), which may both be designated as Human Oral
9 Taxon 363 (*Synergistetes* [G-3] sp.) according to the Human Oral Microbiome Database
10 (HOMD) classification scheme (5). Although our understanding of taxonomy in the
11 *Synergistetes* phylum is rudimentary, it seems probable that both the 4P12 and W090 clones
12 correspond to *F. fastidiosum* strains. Our results suggest that certain *Synergistetes* taxa
13 belonging to cluster A may be widely distributed in global populations, as they have now
14 been detected in the oral cavities of subjects residing in the UK, US and China. Whilst our
15 finding that 4P12 and 7P1 OTUs are strongly associated with periodontitis is consistent with
16 two previous studies performed in the US (7, 10), they are not consistent with results from
17 two other studies performed in the UK (11) and US (17). This highlights the fact that the
18 phylogeny, global distributions and disease-associations of oral *Synergistetes* taxa require
19 further detailed investigation.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 In brief conclusion, there was a greater diversity and clonal abundance of OTUs
39 belonging to cluster A oral *Synergistetes* detected in subgingival plaque from periodontitis
40 patients, compared to periodontitis-free controls. Five *Synergistetes* OTUs were found
41 predominantly or exclusively in the periodontitis group. Our results are consistent with the
42 hypothesis that members of the phylum *Synergistetes* play a role in the occurrence or
43 pathogenesis of periodontal disease.
44
45
46
47
48
49
50
51

52 53 54 55 56 **Acknowledgements** 57 58 59 60

1
2
3 RMW acknowledges financial support from the Infection and Immunology Strategic
4
5 Research Theme of the University of Hong Kong, and the Research Grants Council of Hong
6
7 Kong, via a General Research Fund (GRF) grant [#781911]. WKL acknowledges support
8
9 from the University of Hong Kong: Research and Conference Grant, Small Project Fund
10
11 [#10206659] and the University Research Committee Fund [#10205584].
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Manuscript proof

References

1. Kornman KS. Mapping the pathogenesis of periodontitis: a new look. *J Periodontol* 2008; **79**: 1560-1568.
2. Page RC, Offenbacher S, Schroeder HE, Seymour GJ, Kornman KS. Advances in the pathogenesis of periodontitis: summary of developments, clinical implications and future directions. *Periodontol 2000* 1997; **14**: 216-248.
3. Schaudinn C, Gorur A, Keller D, Sedghizadeh PP, Costerton JW. Periodontitis: an archetypical biofilm disease. *J Am Dent Assoc* 2009; **140**: 978-986.
4. Zarco MF, Vess TJ, Ginsburg GS. The oral microbiome in health and disease and the potential impact on personalized dental medicine. *Oral Dis* 2012; **18**: 109-120.
5. Dewhirst FE, Chen T, Izard J, *et al.* The human oral microbiome. *J Bacteriol* 2010; **192**: 5002-5017.
6. Bik EM, Long CD, Armitage GC, *et al.* Bacterial diversity in the oral cavity of 10 healthy individuals. *Isme J* 2010; **4**: 962-974.
7. Paster BJ, Boches SK, Galvin JL, *et al.* Bacterial diversity in human subgingival plaque. *J Bacteriol* 2001; **183**: 3770-3783.
8. Zaura E, Keijser BJ, Huse SM, Crielaard W. Defining the healthy "core microbiome" of oral microbial communities. *BMC Microbiol* 2009; **9**: 259.
9. Keijser BJ, Zaura E, Huse SM, *et al.* Pyrosequencing analysis of the oral microflora of healthy adults. *J Dent Res* 2008; **87**: 1016-1020.
10. Kumar PS, Griffen AL, Moeschberger ML, Leys EJ. Identification of candidate periodontal pathogens and beneficial species by quantitative 16S clonal analysis. *J Clin Microbiol* 2005; **43**: 3944-3955.

- 1
2
3 11. Vartoukian SR, Palmer RM, Wade WG. Diversity and morphology of members of the
4
5 phylum "*Synergistetes*" in periodontal health and disease. *Appl Environ Microbiol*
6
7 2009; **75**: 3777-3786.
- 8
9 12. Horz HP, Citron DM, Warren YA, Goldstein EJ, Conrads G. *Synergistes* group
10
11 organisms of human origin. *J Clin Microbiol* 2006; **44**: 2914-2920.
- 12
13 13. Godon JJ, Moriniere J, Moletta M, Gaillac M, Bru V, Delgenes JP. Rarity associated
14
15 with specific ecological niches in the bacterial world: the '*Synergistes*' example.
16
17 *Environ Microbiol* 2005; **7**: 213-224.
- 18
19 14. Vartoukian SR, Palmer RM, Wade WG. The division "*Synergistes*". *Anaerobe* 2007;
20
21 **13**: 99-106.
- 22
23 15. Marchandin H, Damay A, Roudiere L, *et al.* Phylogeny, diversity and host
24
25 specialization in the phylum *Synergistetes* with emphasis on strains and clones of
26
27 human origin. *Res Microbiol* 2010; **161**: 91-100.
- 28
29 16. Jumas-Bilak E, Roudiere L, Marchandin H. Description of '*Synergistetes*' phyl. nov.
30
31 and emended description of the phylum '*Deferribacteres*' and of the family
32
33 *Syntrophomonadaceae*, phylum '*Firmicutes*'. *Int J Syst Evol Microbiol* 2009; **59**:
34
35 1028-1035.
- 36
37 17. Kumar PS, Griffen AL, Barton JA, Paster BJ, Moeschberger ML, Leys EJ. New
38
39 bacterial species associated with chronic periodontitis. *J +Dent Res* 2003; **82**: 338-
40
41 344.
- 42
43 18. Kumar PS, Leys EJ, Bryk JM, Martinez FJ, Moeschberger ML, Griffen AL. Changes
44
45 in periodontal health status are associated with bacterial community shifts as assessed
46
47 by quantitative 16S cloning and sequencing. *J Clin Microbiol* 2006; **44**: 3665-3673.
- 48
49 19. Siqueira JF, Jr., Rocas IN, Cunha CD, Rosado AS. Novel bacterial phylotypes in
50
51 endodontic infections. *J Dent Res* 2005; **84**: 565-569.
- 52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
20. Hutter G, Schlagenhauf U, Valenza G, *et al.* Molecular analysis of bacteria in periodontitis: evaluation of clone libraries, novel phylotypes and putative pathogens. *Microbiology* 2003; **149**: 67-75.
 21. Munson MA, Pitt-Ford T, Chong B, Weightman A, Wade WG. Molecular and cultural analysis of the microflora associated with endodontic infections. *J Dent Res* 2002; **81**: 761-766.
 22. Kumar PS, Mason MR, Brooker MR, O'Brien K. Pyrosequencing reveals unique microbial signatures associated with healthy and failing dental implants. *J Clin Periodontol* 2012; **39**: 425-433.
 23. Crielaard W, Zaura E, Schuller AA, Huse SM, Montijn RC, Keijser BJ. Exploring the oral microbiota of children at various developmental stages of their dentition in the relation to their oral health. *BMC Med Genomics* 2011; **4**: 22.
 24. Vartoukian S, Downes J, Palmer RM, Wade WG. *Fretibacterium fastidiosum* gen. nov., sp. nov., isolated from the human oral cavity. *Int J Syst Evol Microbiol* 2012, doi: 10.1099/ijss.0.041038-0.
 25. Vartoukian SR, Palmer RM, Wade WG. Cultivation of a *Synergistetes* strain representing a previously uncultivated lineage. *Environ Microbiol* 2010; **12**: 916-928.
 26. Downes J, Vartoukian SR, Dewhirst FE, *et al.* *Pyramidobacter piscolens* gen. nov., sp. nov., a member of the phylum 'Synergistetes' isolated from the human oral cavity. *Int J Syst Evol Microbiol* 2009; **59**: 972-980.
 27. Jumas-Bilak E, Carlier JP, Jean-Pierre H, *et al.* *Jonquetella anthropi* gen. nov., sp. nov., the first member of the candidate phylum 'Synergistetes' isolated from man. *Int J Syst Evol Microbiol* 2007; **57**: 2743-2748.

- 1
2
3 28. Griffen AL, Beall CJ, Campbell JH, et al. Distinct and complex bacterial profiles in
4 human periodontitis and health revealed by 16S pyrosequencing. *ISMEJ* 2012; **6**:
5 1176-1185.
6
7
8
9
10 29. Lane DJ. 16S/23S rRNA sequencing, *In*: E S, M G, eds. *Nucleic Acid Techniques in*
11 *Bacterial Systematics*. Chichester, England: John Wiley and Sons Ltd, 1991: 115–173.
12
13 30. Cole JR, Wang Q, Cardenas E, et al. The Ribosomal Database Project: improved
14 alignments and new tools for rRNA analysis. *Nucleic Acids Res* 2009; **37**: D141-145.
15
16
17 31. Pruesse E, Quast C, Knittel K, et al. SILVA: a comprehensive online resource for
18 quality checked and aligned ribosomal RNA sequence data compatible with ARB.
19 *Nucleic Acids Res* 2007; **35**: 7188-7196.
20
21
22
23 32. Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid
24 assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ*
25 *Microbiol* 2007; **73**: 5261-5267.
26
27
28
29 33. Schloss PD, Westcott SL. Assessing and improving methods used in operational
30 taxonomic unit-based approaches for 16S rRNA gene sequence analysis. *Appl*
31 *Environ Microbiol* 2011; **77**: 3219-3226.
32
33
34 34. Schloss PD, Westcott SL, Ryabin T, et al. Introducing mothur: open-source, platform-
35 independent, community-supported software for describing and comparing microbial
36 communities. *Appl Environ Microbiol* 2009; **75**: 7537-7541.
37
38
39 35. Schloss PD, Larget BR, Handelsman J. Integration of microbial ecology and statistics:
40 a test to compare gene libraries. *Appl Environ Microbiol* 2004; **70**: 5485-5492.
41
42
43 36. Singleton DR, Furlong MA, Rathbun SL, Whitman WB. Quantitative comparisons of
44 16S rRNA gene sequence libraries from environmental samples. *Appl Environ*
45 *Microbiol* 2001; **67**: 4374-4376.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 37. Lozupone C, Knight R. UniFrac: a new phylogenetic method for comparing microbial
4 communities. *Appl Environ Microbiol* 2005; **71**: 8228-8235.
5
6
7 38. Lozupone C, Hamady M, Knight R. UniFrac--an online tool for comparing microbial
8 community diversity in a phylogenetic context. *BMC Bioinformatics* 2006; **7**: 371.
9
10 39. Schloss PD, Handelsman J. Introducing TreeClimber, a test to compare microbial
11 community structures. *Appl Environ Microbiol* 2006; **72**: 2379-2384.
12
13 40. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: molecular
14 evolutionary genetics analysis using maximum likelihood, evolutionary distance, and
15 maximum parsimony methods. *Mol Biol Evol* 2011; **28**: 2731-2739.
16
17 41. Wickham H. *ggplot2: elegant graphics for data analysis*: Springer New York, 2009.
18
19 42. Choi BK, Paster BJ, Dewhirst FE, Gobel UB. Diversity of cultivable and uncultivable
20 oral spirochetes from a patient with severe destructive periodontitis. *Infect Immun*
21 1994; **62**: 1889-1895.
22
23 43. Dewhirst FE, Tamer MA, Ericson RE, *et al*. The diversity of periodontal spirochetes
24 by 16S rRNA analysis. *Oral Microbiol Immunol* 2000; **15**: 196-202.
25
26 44. Socransky S, Haffajee A. Periodontal microbial ecology. *Periodontol 2000* 2005; **38**:
27 135-187.
28
29 45. Paster BJ, Olsen I, Aas JA, Dewhirst FE. The breadth of bacterial diversity in the
30 human periodontal pocket and other oral sites. *Periodontol 2000* 2006; **42**: 80-87.
31
32 46. de Lillo A, Ashley FP, Palmer RM, *et al*. Novel subgingival bacterial phylotypes
33 detected using multiple universal polymerase chain reaction primer sets. *Oral*
34 *Microbiol Immunol* 2006; **21**: 61-68.
35
36 47. Siqueira JF, Jr., Rocas IN. Uncultivated phylotypes and newly named species
37 associated with primary and persistent endodontic infections. *J Clin Microbiol* 2005;
38 **43**: 3314-3319.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 48. Machado de Oliveira JC, Gama TG, Siqueira JF, Jr., Rocas IN, Peixoto RS, Rosado
4 AS. On the use of denaturing gradient gel electrophoresis approach for bacterial
5 identification in endodontic infections. *Clin Oral Investig* 2007; **11**: 127-132.
6
7
8
9
10 49. Rocas IN, Baumgartner JC, Xia T, Siqueira JF, Jr. Prevalence of selected bacterial
11 named species and uncultivated phylotypes in endodontic abscesses from two
12 geographic locations. *J Endod* 2006; **32**: 1135-1138.
13
14
15
16 50. Saito D, Leonardo Rde T, Rodrigues JL, Tsai SM, Hofling JF, Goncalves RB.
17 Identification of bacteria in endodontic infections by sequence analysis of 16S rDNA
18 clone libraries. *J Med Microbiol* 2006; **55**: 101-107.
19
20
21
22
23 51. Sakamoto M, Rocas IN, Siqueira JF, Jr., Benno Y. Molecular analysis of bacteria in
24 asymptomatic and symptomatic endodontic infections. *Oral Microbiol Immunol* 2006;
25 **21**: 112-122.
26
27
28
29 52. Munson MA, Banerjee A, Watson TF, Wade WG. Molecular analysis of the
30 microflora associated with dental caries. *J Clin Microbiol* 2004; **42**: 3023-3029.
31
32
33
34 53. Aas JA, Paster BJ, Stokes LN, Olsen I, Dewhirst FE. Defining the normal bacterial
35 flora of the oral cavity. *J Clin Microbiol* 2005; **43**: 5721-5732.
36
37
38
39 54. Xie G, Chain PS, Lo CC, *et al.* Community and gene composition of a human dental
40 plaque microbiota obtained by metagenomic sequencing. *Mol Oral Microbiol* 2010;
41 **25**: 391-405.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Legends

Fig. 1. Phylogenetic relationships and clonal abundance of *Synergistetes* OTUs identified in this study. Left hand side: NJ phylogenetic tree composed of the *Synergistetes* OTUs identified in this study. Bootstrap values $\geq 50\%$ are shown beside branch points. Right hand side: The absolute clonal abundance of the *Synergistetes* OTUs detected in each subject is displayed using a heat map with grey-scale boxes; with corresponding values indicated in the key shown near the base of the figure. The names of previously reported oral *Synergistetes* clones or isolates that correspond to the OTUs defined here are indicated. The statistical significance of the differences in clonal abundances of specific OTUs between the periodontitis (P) and periodontitis-free (H) subject groups are indicated (* $p < 0.05$, ** $p < 0.01$, Mann-Whitney U test).

Fig. 2. Phylogenetic relationships between *Synergistetes* OTUs detected here and previously identified species of oral *Synergistetes* species and Human Oral Taxon (HOT) numbers listed in the Human Oral Microbiome Database (HOMD). The phylogenetic tree was constructed from corresponding 16S rRNA gene sequences using a Neighbour-joining method (Jukes-Cantor model), and was rooted against *Treponema socranskii* subsp. *socranskii* ATCC 35536. Bootstrap values greater than 50% are shown beside the branch points. *Synergistetes* OTUs were assigned to HOTs using a 98.5% sequence identity cut-off value. The eight OTUs that do not correspond to existing *Synergistetes* HOTs are indicated in bold text.

Fig. 3. Similarities in the composition of *Synergistetes* OTUs detected in the 20 subjects (H1-H10, P1-P10) included in the study. The dendrogram indicates the similarity of *Synergistetes* communities detected in each subject according to the structure-based Yue and Clayton theta

1
2
3 coefficient, using a 99% 16S rRNA sequence identity cut-off for defining OTUs. P:
4
5 Periodontitis, H: Periodontitis-free (control) subject groups.
6
7
8

9
10 *Fig. S1.* Rarefaction curves for *Synergistetes* OTUs defined using 99%, 97% and 95% 16S
11 rRNA sequence similarity cut-off values. Rarefaction curves for the number of *Synergistetes*
12 OTUs were plotted using four different parameters for the x-axis, shown in Panels A-D. The
13 rarefaction curves for the three defined OTU cut-off values were plotted on each panel: 99%
14 (black), 97% (dark grey) and 95% (light grey). Panel A: rarefaction curves for the numbers of
15 *Synergistetes* OTUs, constructed by sub-sampling the pool of subjects analyzed. Panel B:
16 rarefaction curves for OTU numbers based on the total number of *Synergistetes* 16S rRNA
17 clones analyzed. Panel C: rarefaction curves for OTU numbers based on the *Synergistetes*
18 16S rRNA clones analyzed from the periodontitis subjects only. Panel D: rarefaction curves
19 for OTU numbers based on the *Synergistetes* 16S rRNA clones analyzed from the
20 periodontitis-free subjects only.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Demography and clinical profile of study subjects

Characteristics	Categories	Periodontal status		Statistics ^a	p-value
		Periodontitis (n = 10)	Periodontitis- free (n = 10)		
Age (Mean ±SD, year)		40.7 ± 10.6	54.0 ± 10.6	U	0.014
Gender	Male	3	3	χ^2	NS
	Female	7	7		
<i>Full mouth periodontal profile</i>					
Standing teeth		28.0 ± 2.2	27.5 ± 1.1	U	NS
% BOP		88.6 ± 10.3	30.4 ± 12.7	U	<0.001
% PPD = 4-5 mm		22.0 ± 8.8	0	NA	
% PPD ≥ 6 mm		15.8 ± 15.0	0	NA	
<i>Profile of sample sites</i>					
No. of sites sampled	mean	42.7 ± 31.2	165.0 ± 6.5	U	<0.001
	range	8 - 81	156 - 180		
% BOP	mean	99.8 ± 0.5	30.4 ± 12.7	U	<0.001
	range	98.3 – 100.0	11.0 – 48.2		
Mean PPD (mm)	mean	6.2 ± 0.5	1.9 ± 0.6	U	<0.001
	range	5.21 – 6.63	1.3 – 2.4		

Results are Number unless otherwise indicated; BOP = bleeding on probing; NS = not significant; NA: not applicable; PPD: probing pocket depth

^aU: Mann-Whitney U test; χ^2 : chi square test.

Table 2. Absolute and relative clonal abundance at the phylum level


Phylum	Periodontitis-			<i>p</i> -value ^a
	Periodontitis	free	Total	
<i>Spirochaetes</i>	365 (70.2%)	250 (49.0%)	615 (59.7%)	<0.05
<i>Synergistetes</i>	132 (25.4%)	30 (5.9%)	162 (15.7%)	<0.001
<i>Actinobacteria</i>	20 (3.8%)	147 (28.8%)	167 (16.2%)	<0.05
<i>Firmicutes</i>	NA	25 (4.9%)	25 (2.4%)	<0.05
<i>Fusobacteria</i>	3 (0.6%)	34 (6.7%)	37 (3.6%)	NS
<i>Proteobacteria</i>	NA	24 (4.7%)	24 (2.3%)	<0.05
Total	520 (100%)	510 (100%)	1030 (100%)	NS

NA: not applicable; NS: not significant

^aMann-Whitney U test


Table 3. The number of observed *Synergistetes* OTUs

Level	Periodontitis	Periodontitis-free	Total
Unique sequences	119	24	143
OTU (99%)	29	8	31
OTU (97%)	8	3	8
OTU (95%)	5	2	5




192x159mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



199x170mm (300 x 300 DPI)

proof



0.1

92x104mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

Manuscript proof

