



Title	Study on the interaction between DNA-binding domain (DBD) of human androgen receptor (AR) and SWIRM domain of lysine-specific demethylase 1 (LSD1)
Author(s)	Lo, KC; Zhu, G; Sze, KH
Citation	The 2011 Croucher Advanced Study Institute Conference on Structure-Based Screening and Design of Ligands for Protein Targets, Hong Kong, 12-15 December 2011
Issued Date	2011
URL	http://hdl.handle.net/10722/169360
Rights	Creative Commons: Attribution 3.0 Hong Kong License

STUDY ON THE INTERACTION BETWEEN DNA-BINDING DOMAIN (DBD) OF HUMAN ANDROGEN RECEPTOR (AR) AND SWIRM DOMAIN OF LYSINE-SPECIFIC DEMETHYLASE 1 (LSD1)

Lo KC¹, Zhu G², and Sze KH¹

¹Chemistry Department and Open Laboratory of Chemical Biology of the Institute of Molecular Technology for Drug Discovery and Synthesis, The University of Hong Kong, Pokfulam Road, Hong Kong

²Department of Biochemistry, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong, SAR, PR China

h0415676@hku.hk

Gene regulations of prokaryotes and eukaryotes are different. Prokaryotic gene regulation requires simply binding of regulatory proteins to help with or avoid forming transcription complex. For eukaryotes like humans, however, their regulation needs “chromatin remodeling” with the association of regulatory proteins involving the opening up of DNA-histone protein complex chromatin and unwinding DNA. Without remodeling, RNA polymerases responsible of the transcription process cannot get access and perform transcription.

Lysine-specific demethylase 1 is one of the chromatin remodeling enzymes. It can demethylate specifically the N-tail mono- or di-methylated K4 residue on histone H3 by oxidation (Y. Shi et al., 2004). LSD1 has three known domains: FAD-binding domain, demethylase domain and SWIRM domain. Till now, the exact function of SWIRM domain of LSD1 is still unclear, although its solution structure has been reported.

In 2005, Metzger’s group found that the LSD1 SWIRM domain could bind the N-terminus, the DNA-binding domain (DBD) and the ligand-binding domain (LBD) of androgen receptor (AR) by performing a GST-pull down assay (Metzger et al., 2005); and shows that SWIRM domain has the strongest interaction with DBD among the domains of AR. Therefore, my study is to investigate the interaction between AR-DBD and SWIRM-LSD1 domains by NMR spectroscopy.

Acknowledgements. This work was supported by grants from the Hong Kong University Research Grant and the Research Grants Council of Hong Kong for KH Sze (GRF 7755/08M, GRF 7765/09M).