



Title	The prognostic and therapeutic potential of AMP-activated protein kinase in ovarian cancer
Author(s)	Chan, DW; Ngan, HYS
Citation	The ASGO 2nd International Workshop on Gynecologic Oncology, Kyoto, Japan, 28 September 2012. In Conference Program, 2012, p. 32
Issued Date	2012
URL	http://hdl.handle.net/10722/169438
Rights	Creative Commons: Attribution 3.0 Hong Kong License

The Prognostic and Therapeutic Potential of AMP-activated Protein Kinase in Ovarian Cancer

David W Chan, Hextan YS Ngan

Department of Obstetrics and Gynecology, LKS Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, P.R.China

Ovarian cancer is one of the leading causes of cancer-associated death in women. The high mortality is due to its poor prognosis as most cases are found in late stages. Therefore, searching reliable tumor markers is urgently needed for clinical management of this disease. Altered cellular metabolism is a crucial phenomenon for the development and progression of ovarian cancer. AMP-activated protein kinase (AMPK) acts as a key intracellular energy sensor and regulator for governing energy balance homeostasis. It also closely links with cancer cell metabolism. Others and we have reported that the activation of AMPK by pharmacological agents shows cytotoxicity to cancer cells, indicating that targeting AMPK could be a promising therapeutic approach. On the other hand, our study also demonstrated that the AMPK activity had an inversely correlation between tumor stage and/or high grade ovarian cancer. Importantly, the reduced AMPK activity was associated with the increased ERK/FOXO1 signaling cascade in advanced ovarian cancers. The ERK/FOXO1 signaling pathway has been shown to possess oncogenic effects on promoting oncogenesis in numerous human cancers and showed a progressive increased pattern from low- to high-grade tumor. In fact, using ovarian cancer cell models, we showed that the activation of AMPK by pharmaceutical or natural AMPK activators could inhibit not only ERK/FOXO1 activity but also the tumor growth. The suppressive functions were equivalent to ERK or FOXO1 inhibitors in ovarian cancer cells or other gynecological carcinoma cell lines. Taken together, these findings suggest that the decreased AMPK activity may act as a prognostic marker in ovarian cancer because it is significantly correlated with the increased ERK/FOXO1 signaling cascade and the advanced stage ovarian cancers. Moreover, targeting AMPK by either pharmaceutical or natural AMPK activators may be explored as an alternative therapeutic regimen in the treatment of this disease.

Curriculum Vitae

Academic Qualifications: BSc. (Hons) (HKBU) 1992, MPhil (HKU) 1996, PhD (Monash) 2002

Present Academic Position: Research Assistant Professor, Honary Assistant Professor
Department of O & G, HKU, 2007-present

Previous relevant research work:

2006-2007	Postdoctoral Fellow and Hon. Assistant Professor, Dept. of O&G, HKU.
2003-2006	Postdoctoral Fellow, Department of Pathology, HKU.
2001-2003	Research Officer, Peter MacCallum Cancer Institute, Uni. of Melbourne.
2000-2001	Research Officer, MIRD, Monash University.
1996-1997	Technician, Department of Medicine, HKU.