



| | |
|--------------------|---|
| Title | Erlotinib-induced autophagy in epidermal growth factor receptor mutated non-small cell lung cancer |
| Author(s) | Li, Y; Lam, SK; Mak, JCW; Zheng, C; Ho, JCM |
| Citation | Lung cancer, 2013, v. 81 n. 3, p. 354-361 |
| Issued Date | 2013 |
| URL | http://hdl.handle.net/10722/184573 |
| Rights | NOTICE: this is the author's version of a work that was accepted for publication in Lung cancer. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Lung cancer, 2013, v. 81 n. 3, p. 354-361. DOI: 10.1016/j.lungcan.2013.05.012 |

1 **Erlotinib-induced autophagy in epidermal growth factor receptor mutated non-small cell**
2 **lung cancer**

3 Yuan-yuan Li MPhil, Sze-kwan Lam PhD, Judith Choi-wo Mak PhD, Chun-yan Zheng MPhil,
4 James Chung-man Ho MD

5 Division of Respiratory Medicine, Department of Medicine

6 The University of Hong Kong, Queen Mary Hospital, Hong Kong SAR

7

8 **Address for correspondence:**

9 Dr. James C. Ho M.D. FRCP

10 Department of Medicine, The University of Hong Kong

11 Queen Mary Hospital, Pokfulam, Hong Kong SAR, China

12 Tel: (852) 2255 4999 Fax: (852) 2872 5828 Email: jhocm@hku.hk

13 **Word count:2981**

14 **Keywords:** erlotinib, autophagy, epidermal growth factor receptor, resistance, lung cancer

15

16 **Abbreviations:** AMPK, 5' adenosine monophosphate-activated protein kinase; Atg, autophagy-
17 related protein; LC3, microtubule-associated protein light chain 3; mTOR, mammalian target of
18 rapamycin; NFκB, nuclear factor kappa beta; PARP, poly (ADP-ribose) polymerase.

1 **Abstract**

2 **Purpose:** Erlotinib is a commonly used tyrosine kinase inhibitor (TKI) in non-small cell lung
3 cancer (NSCLC). Autophagy is a catabolic process in response to stress and deprivation of
4 nutrients. This study aims to investigate whether autophagy confers acquired resistance to
5 erlotinib treatment in NSCLC.

6 **Methods:** Four NSCLC cell lines (HCC827, HCC4006, H358 and H1975) with different
7 epidermal growth factor receptor (EGFR) mutation status (exon 19 deletion, exon 19 deletion,
8 wild-type and L858R/T790M respectively) were selected. MTT assay, crystal violet staining and
9 Annexin-V assay were performed to determine cell viability and apoptosis. Autophagic proteins
10 were detected by Western blot. Acidic vesicular organelle (AVO) formation was determined by
11 acridine orange staining. Autophagy inhibitor (chloroquine) and RNA interference were used to
12 demonstrate the biological effect of erlotinib-induced autophagy.

13 **Results:** In line with EGFR mutation status, it was shown that both HCC827 and HCC4006 cells
14 were sensitive to erlotinib, while H358 and H1975 cell lines were resistant. Erlotinib treatment at
15 clinically relevant concentrations induced autophagy (increased LC3II expression, Atg-5/Atg12
16 conjugation, formation of AVO and p62 degradation) in sensitive NSCLC cell lines, via p53
17 nuclear translocation, AMPK activation and mTOR suppression. Addition of chloroquine, as an
18 autophagy inhibitor, enhanced erlotinib sensitivity in sensitive cells. Similarly, silencing of Atg5
19 or Beclin-1 significantly increased sensitivity to erlotinib in both sensitive cell lines. In contrast,
20 there was no induction of autophagy in resistant H358 and H1975 cell lines upon erlotinib
21 exposure.

1 **Conclusions:** Erlotinib can induce both apoptosis and autophagy in sensitive NSCLC cell lines
2 with activating EGFR mutation (exon 19 del). Inhibition of autophagy can further enhance
3 sensitivity to erlotinib in EGFR-mutated NSCLC, suggesting that autophagy may serve as a
4 protective mechanism.

5 (Word count of abstract: 274)

6

1 **Introduction**

2 Non-small cell lung cancer (NSCLC) is one of the leading causes of cancer deaths worldwide [1].
3 The clinical benefit of platinum-based chemotherapy in advanced NSCLC has reached a plateau
4 [2, 3]. Epidermal growth factor receptor (EGFR) is frequently deregulated [4, 5] or
5 overexpressed [6] in NSCLC. The EGFR downstream signaling pathways are pivotal in the
6 initiation and progression of cancers, serving as one of the key therapeutic targets in NSCLC [7].
7 Erlotinib, as one of the prototypical EGFR tyrosine kinase inhibitors (TKIs), has been approved
8 by U.S. Food and Drug Administration (FDA) for treatment of NSCLC [8-10], however,
9 development of acquired resistance (mostly secondary EGFR mutation (T790M in exon 20) [11,
10 12] and MET amplification [13]) has almost been the rule during treatment. Better understanding
11 of additional resistance mechanisms, with corresponding therapeutic interventions, could
12 potentially overcome or delay the development of acquired resistance to EGFR TKI treatment.

13
14 Autophagy is characterized by the formation of double-membrane vacuoles (autophagosomes) in
15 the cytoplasm. The key steps of autophagic process include the induction and nucleation of
16 autophagic vesicles, followed by fusion with lysosomes and subsequent degradation of their
17 contents [14, 15]. During autophagy, Beclin-1 complex recruits type III PI3K VPS34 to generate
18 phosphatidylinositol 3-phosphate. Autophagy-related protein 12 (Atg12) and microtubule-
19 associated protein light chain 3 (LC3) are involved in the expansion of autophagosome
20 membranes. The E1-like Atg7 and E2-like Atg10 covalently link Atg12 with Atg5, which
21 together bind Atg16 to form pre-autophagosomal structures. LC3 is cleaved by Atg 4 and
22 lipidated to form LC3II, which is commonly used to monitor autophagy [16]. The catabolic
23 cellular self-degradation serves to produce building blocks for molecular synthesis and maintain
24 energy homeostasis during periods of nutrient deprivation [16].

1

2 It is still controversial whether autophagy causes cancer cell death or sustains cancer survival
3 under stressful conditions. Abrogation of autophagy, with pharmacological inhibitors (e.g.
4 chloroquine and bafilomycin A1) or shRNA knockdown of autophagic proteins, has been shown
5 to re-sensitize cancer cells to chemotherapy or radiation [17, 18]. However, others suggested
6 synergistic or additive effect in combining chemotherapy with autophagy inducer like rapamycin
7 in a lung cancer cell line model [19]. The effect of autophagy on EGFR TKI treatment in
8 NSCLC is less well-known. In a recent study, autophagy was induced in a relatively resistant
9 NSCLC cell line model upon exposure to EGFR TKI at high concentrations [20].

10

11 Therapeutic exposure of cancer cells to an anti-cancer agent would be considered as a form of
12 stress only if they are prone to the growth inhibitory or cytotoxic effects. Since EGFR TKI
13 preferentially works on NSCLC with activating EGFR mutations, we postulate that autophagy
14 would only be induced among the sensitive EGFR-mutated NSCLC treated with clinically
15 relevant concentrations of EGFR TKI, which could serve as a cancer survival mechanism. In this
16 study, we investigated the effect of erlotinib treatment on autophagy in 4 NSCLC cell lines with
17 different EGFR mutation status, in order to understand the possible mechanisms involved and
18 therapeutic implications.

19

20 **Materials and Methods**

21 *Cell lines and cultures*

22 Four human NSCLC cell lines, HCC827 (EGFR exon 19 del), HCC4006 (EGFR exon 19 del),
23 H358 (EGFR WT) and H1975 (EGFR exon 21 L858R/exon 20 T790M) were obtained from

1 American Type Culture Collection (Manassas, VA, USA). All cells were maintained in RPMI-
2 1640 medium (Gibco®, Life Technologies, Carlsbad, California, USA) with 10% fetal bovine
3 serum (FBS) (Gibco®, Life Technologies) and cultured in a humidified atmosphere of 5% CO₂
4 at 37°C.

5

6 *Drugs and reagents*

7 Erlotinib (Selleck, Houston, USA) and bafilomycin A1 (Sigma-Aldrich, St. Louis, Missouri,
8 USA) were diluted in dimethyl sulfoxide (DMSO) (Sigma-Aldrich). Chloroquine (Sigma-
9 Aldrich) was diluted in phosphate buffered saline (PBS). Acridine orange hydrochloride was
10 purchased from Sigma-Aldrich. Anti-LC3B and anti-beta-actin were obtained from Sigma-
11 Aldrich. Anti-Beclin-1, Atg5/12, p62, p53, p-AMPK(Thr172), AMPK, p-NFκB (s536), NFκB, p-
12 mTOR(ser2448), and mTOR were obtained from Cell Signaling Technology (Danvers,
13 Massachusetts, USA). siRNA targeting Beclin-1 and Atg5 were purchased from Ambion Life
14 Technologies (Carlsbad, California, USA). Control siRNA was purchased from Santa Cruz
15 (California, USA).

16

17 *Cell viability assay*

18 Cells were cultured in 96-well plates (5,000 cells in 0.2 ml culture medium/well). After drug
19 treatment, 20 μl 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (0.5
20 mg/ml) (Sigma-Aldrich) was added for 2 h, and followed by lysis with DMSO. The absorbance
21 at 570 nm was measured using a microplate reader Fluo Star Optima (Bmg Labtec GmbH,
22 Ortenberg, Germany). For experiments using chloroquine, cells were fixed with 4%
23 formaldehyde (Sigma-Aldrich) for 10 min and then stained with 0.05% crystal violet (Sigma-

1 Aldrich) in 30% ethanol for 10 min at room temperature. Plates were washed four times with tap
2 water. Cells were lysed with 1% SDS solution (Sigma-Aldrich), and dye uptake was measured at
3 550 nm using a microplate reader Fluo Star Optima. The absorbance in treatment group was
4 normalized to those of the untreated cells (control). All experiments were done in triplicates.

5

6 ***Western blot analysis***

7 Cells were lysed on ice with RIPA lysis buffer [10 mM Tris, 150 mM NaCl, 1 mM
8 methylenediaminetetra-acetic acid, 1% Triton X-100, 0.5% NP40, freshly added 0.2 mM PMSF
9 in isopropanol, 1:50 phosphatase inhibitor cocktail 2 (Sigma-Aldrich), 1:50 protease inhibitor
10 cocktail (Sigma-Aldrich)] for 1 h. The supernatants were collected after centrifugation. Nuclear
11 and cytoplasmic proteins were extracted using NE-PER nuclear and cytoplasmic reagents from
12 Thermo Scientific according to instructions (Rockford, IL, USA). The protein concentration in
13 the supernatants was quantified with Bradford Protein Assay (Bio-Rad, Berkeley, California,
14 USA). For each supernatant, 30-50 µg protein was loaded on 7.5-15% of sodium dodecyl sulfate-
15 polyacrylamide gel electrophoresis and then transferred onto polyvinylidene fluoride membranes
16 (GE Healthcare, Buckinghamshire, UK). The membranes were incubated with primary
17 antibodies overnight at 4°C. Further incubation with the corresponding secondary antibody (Cell
18 Signaling technology) for 90 min at 4°C was then carried out. Detection was performed using an
19 enhanced chemiluminescence (ECL) kit (GE Healthcare). Quantification was done with Image J
20 (National Institute of Health, USA). Results were obtained from at least 3 independent
21 experiments.

22

23 ***Small interfering RNA (siRNA) transfection***

1 Knockdown of specific RNA was performed, including *beclin1* (siRNA ID: s16539; sense
2 sequence: CAGAUACUCUUUUAGACCATT; antisense sequence:
3 UGGUCUAAAAGAGUAUCGTG), *ATG5* (siRNA ID: s18160; sense sequence:
4 GCUAUAUCAGGAUGAGAUATT; antisense sequence: UAUCUCAUCCUGAUUAUAGCGT)
5 and a scrambled siRNA (sc-37007). Briefly, a mixture of siRNA and transfection reagent (Santa
6 Cruz) in culture medium was incubated with cells for 6 h. Then 5% FBS in culture medium was
7 added and incubated for 48 h. Corresponding protein downregulation was confirmed with
8 Western blot. Drug treatment experiments were conducted for 48 h, cell viability and apoptotic
9 events were then measured.

10

11 ***Apoptotic assay***

12 Cells were resuspended in binding buffer at $1-5 \times 10^6$ /mL. Five microliter of fluorochrome-
13 conjugated Annexin-V and 5 μ l of 7-AAD Viability Staining Solution (BD biosciences, New
14 Jersey, USA) were added to 100 μ l of the cell suspension, and incubated 15 min at room
15 temperature. Flow cytometry (Beckman Coulter, Inc., USA) was performed to detect cells
16 undergoing apoptosis.

17

18 ***Acridine orange staining***

19 Autophagy is the process of sequestering cytoplasmic proteins into the lytic compartment, being
20 characterized by the formation of acidophilic vesicular organelles (AVO) [21]. Cells were
21 stained by 1 μ g/ml AO in PBS at 37°C for 15 min, then washed and visualized with fluorescence
22 microscope (Carl Zeiss, Axioskop 2 plus, New York, USA) equipped with a mercury 100-W

1 lamp, 490-nm band-pass blue excitation filters, a 500 nm dichroic mirror and a 515 nm long
2 pass-barrier filter.

3

4 ***Statistical analysis***

5 Data from triplicate experiments were presented in mean \pm standard deviation (SD). Comparison
6 between groups was performed using Student's two-tailed t-test by Prism (GraphPad Software,
7 La Jolla, Southern California, USA). A p-value < 0.05 was taken as statistically significant.

8

9

1 **Results**

2

3 *Autophagy induced in sensitive NSCLC cells after erlotinib treatment*

4 Four NSCLC cell lines with different EGFR mutation status were chosen: HCC827 (exon 19 del),
5 HCC4006 (exon 19 del), H358 (wild-type) and H1975 (exon 21 L858R, exon 20 T790M). Using
6 MTT assay, HCC827 and HCC4006 cells were relatively sensitive to erlotinib with IC₅₀ values
7 22 ± 5.1 and 46.2 ± 6.4 nM respectively after 72 h treatment, while H358 and H1975 cells were
8 relatively resistant with IC₅₀ values 7.4 ± 2.9 and 18.6 ± 8.8 μ M respectively. Upon treatment
9 with erlotinib, autophagy was induced only in the 2 sensitive NSCLC cell lines. Accordingly, the
10 conversion of LC3/Atg8 from the cytoplasmic form (LC3-I (18 kDa)) to the autophagosomic
11 form (LC3-II (16 kDa)) after 24 h exposure to low concentration of erlotinib (0.1 and 0.2 μ M)
12 was only evident in HCC827 and HCC4006 cells (sensitive) (Fig. 1A), but not in H358 and
13 H1975 cells (resistant) (Fig. 1B). Similarly, autophagic flux was observed in sensitive NSCLC
14 cells after adding bafilomycin A1 (BFA), suggesting induction of autophagy (Fig. 1C). BFA
15 further increased LC3-II expression compared with erlotinib treatment alone, which supported
16 that the increase of LC3-II with erlotinib treatment was due to increased production rather than
17 decreased degradation. After 48 h of erlotinib treatment, acidic vesicular organelles (AVOs), as
18 markers for late stage of autophagy, were detected with acridine orange staining. Fluorescence
19 microscopy demonstrated that cells stained positive for AVOs were more frequently seen with
20 erlotinib treatment compared to control in HCC827 and HCC4006 cells, but not in H358 and
21 H1975 cells (Fig. 1D, E, F, G). The conjugation of Atg5/Atg12 increased in a time-dependent
22 manner when HCC827 and HCC4006 cells were treated with erlotinib (Fig. 1H). Finally, p62
23 was degraded in a time-dependent manner after erlotinib exposure in HCC827 and HCC4006

1 cells (Fig. 1I). Taken together, these findings provided strong evidence that erlotinib induced
2 autophagy in sensitive NSCLC cells.

3

4 ***Pathways involved in autophagy induction in sensitive NSCLC cells***

5 Previous studies have shown that p53/AMPK/mTOR axis plays an important role in the
6 regulation of autophagy. The appearance of autophagic p62 degradation (Fig. 1I) was
7 accompanied by the p53 translocation from cytosol to nucleus (Fig. 2A and 2B), AMPK
8 activation through up-regulation of p-AMPK (Thr172) (Fig. 2C), p-NFkB activation (Fig. 2D)
9 and down-regulation of p-mTOR (ser2448) (Fig. 2E) after erlotinib treatment (0.2 μ M) in both
10 HCC827 and HCC4006 cells.

11

12 ***Inhibition of autophagy enhanced pro-apoptotic effects of erlotinib in sensitive NSCLC cells***

13 To investigate whether inhibition of autophagy would affect the cytotoxicity of erlotinib,
14 NSCLC cells were treated with erlotinib in the presence of 10 μ M chloroquine (autophagy
15 inhibitor). The concentration of chloroquine was chosen that did not cause significant cell death
16 (data not shown), but was sufficient to inhibit autophagy as shown by accumulation of LC3-II
17 (Fig. 3A). Based on cell viability assay using crystal violet staining, the combination of
18 chloroquine and erlotinib significantly decreased cell viability compared to erlotinib alone in
19 sensitive NSCLC cells, but not in the resistant cell lines H358 and H1975 (Fig. 3B). Similarly,
20 addition of chloroquine to erlotinib increased the percentage of apoptotic cells in sensitive
21 NSCLC cells, but not in resistant cells (Fig. 3C and Table 1). Furthermore, the cleavage of PARP
22 was also enhanced in the presence of chloroquine in the sensitive cells (Fig. 3D).

23

1 ***Knockdown of ATG5 and Beclin-1 increased sensitivity to erlotinib***

2 To further confirm the effect of autophagy inhibition on erlotinib-induced apoptosis, a genetic
3 approach through RNA interference to target autophagy-related genes, Atg 5 and Beclin-1, was
4 used. The transfection efficiency of siRNA against Atg5 and Beclin-1 was demonstrated with
5 Western blot showing significant suppression of Atg5 and Beclin-1 protein expression (Fig. 4A),
6 without significant loss of cell viability (data not shown). Treatment with siRNA against Atg5 (6
7 nM) or Beclin-1 (4 nM) did not result in significant cell death. The cell viability in erlotinib
8 combined with ATG-silencing (HCC827: $37.5 \pm 10.5\%$; HCC4006: $46 \pm 3.6\%$) or beclin-
9 silencing (HCC827: $37.7 \pm 3.2\%$; HCC4006: $25.4 \pm 13.0\%$) was significantly lower than
10 erlotinib combined with control-silencing (HCC827: $62.9 \pm 8.9\%$; HCC4006: $63.6 \pm 4.1\%$,
11 $p < 0.01$) (Figure 4B). Apoptotic cell death in erlotinib combined with ATG-silencing ($51.8 \pm$
12 1.9%) or beclin-silencing ($53.4 \pm 2.0\%$) was increased comparing with erlotinib combined with
13 control-silencing ($26.7 \pm 2.9\%$, $p < 0.01$) in HCC827 cells. Similarly, in HCC4006, apoptotic cell
14 death was significantly enhanced in erlotinib combined with ATG-silencing ($53.1 \pm 3.4\%$) or
15 beclin-silencing ($61.9 \pm 5.1\%$), comparing with erlotinib combined with control-silencing (31.2
16 $\pm 10.6\%$, $p < 0.01$) (Figure 4C).

17

18 **Discussion**

19 We have reported *in vitro* evidence of induction of autophagy with erlotinib treatment in NSCLC
20 cell lines carrying activating EGFR mutations. More importantly, our findings supported the role
21 of autophagy in enhancing NSCLC survival in the presence of life-threatening insult from EGFR
22 TKI, serving as a potentially novel mechanism of acquired resistance. Our original hypothesis
23 that this phenomenon would only occur in sensitive EGFR-mutated NSCLC cell lines exposed to

1 EGFR TKI (erlotinib), as demonstrated in this study, is in line with autophagy as a secondary
2 reaction to the pro-apoptotic effects of erlotinib.

3

4 In autophagy, lysosomes fuse with autophagosomes to form autophagolysosomes, responsible
5 for breakdown of membrane-bound cytoplasmic contents. LC3 exists in two forms (LC3-I and its
6 lipidated derivative LC3-II), which is located in autophagosomal membranes prior to their fusion
7 with lysosomes [22]. Autophagy can be detected as early as 24 h by the conversion of LC3-I to
8 LC3-II, which is further supported by demonstration of autophagic flux [23]. In brief, LC3-II
9 protein accumulation can be due to either increase in LC3-II production (i.e. enhanced autophagy)
10 or decrease in its autophagolysosomal breakdown (i.e. suppressed autophagy). Further increase
11 in LC3-II protein expression after treatment with an inhibitor of late-stage autophagy like BFA
12 would support increased LC3-II production. During early induction of autophagy, ubiquitin-
13 mediated association of Atg5 and Atg12 is required to recruit other proteins to the
14 autophagosomal membrane and form the autophagic vacuole [24]. At a later stage of autophagy,
15 acidic vesicular organelles (AVO) would be formed, serving as another marker of autophagy
16 [25]. p62/sequestosome 1 is an ubiquitin-binding scaffold protein that can be specifically
17 degraded by autophagy [26]. Thus, degradation of p62, as a cargo of autophagosome, can also
18 provide indirect evidence of autophagy. Therefore, based on increased LC3-II protein expression
19 (with autophagic flux), Atg 5/Atg 12 and AVO, as well as decreased p62 protein expression, we
20 have demonstrated ample evidence that autophagy was induced with erlotinib treatment in
21 sensitive (EGFR-mutated), but not in resistant NSCLC cells. Furthermore, using both
22 pharmacological blockade (chloroquine, a specific lysosome inhibitor blocking the fusion of
23 autophagosomes and lysosomes [27]) and gene silencing approach (siRNA knockdown of Atg5

1 and Beclin-1), inhibition of autophagy would enhance EGFR TKI (erlotinib) sensitivity among
2 EGFR-mutated NSCLC cell lines.

3

4 Many conventional cytotoxic drugs have been shown to induce autophagy, however, the role of
5 autophagy in chemotherapy and targeted cancer therapy remains controversial. The anti-cancer
6 drugs that can induce autophagy include cetuximab (anti-EGFR antibody) [25], imatinib (BCR-
7 ABL tyrosine kinase inhibitor) [28], TNF-related apoptosis inducing ligand (TRAIL) [29],
8 among many others. Some studies have suggested combining autophagy inducers with anti-
9 cancer therapy, including combination of TKI with autophagy inducing drugs (rapamycin and
10 imatinib) for treating NSCLC [19]. The combination effect was evaluated in EGFR wild-type
11 NSCLC cell lines, suggesting potential benefit of combining TKI with rapamycin or imatinib
12 [19]. On the other hand, other studies have shown beneficial effect of autophagy inhibition in
13 enhancing tumor cell death. Inhibition of autophagy in pre-clinical models improved response to
14 TRAIL-mediated apoptosis in leukemic and colon cancer cell lines [9], and enhanced apoptosis
15 caused by anti-EGFR antibody cetuximab [30]. Recently, autophagy was shown to be induced
16 with EGFR TKI (gefitinib and erlotinib) treatment in a resistant (but not in sensitive) NSCLC
17 cell line model (A549 and NCI-H1299 cells), which was suggested as a possible resistance
18 mechanism to erlotinib [20]. The apparent discrepancy from our findings could be due to several
19 reasons. The concentrations of EGFR TKI that led to significant features of autophagy were
20 relatively high (25 μ M erlotinib or gefitinib), in contrast to the use of clinically relevant
21 concentrations in our study ($\leq 0.2 \mu$ M erlotinib). Though erlotinib is relatively specific for EGFR,
22 there could still be possible off-target effects especially at excessive concentrations. Notably,
23 EGFR knockdown with specific siRNA did not abrogate erlotinib-induced autophagy in that

1 study of resistant cell line model, suggesting possible EGFR-independent actions of erlotinib.
2 Therefore, we believe that our EGFR-mutated NSCLC cell line model with appropriate
3 concentrations of erlotinib should be more clinically relevant.

4

5 The mammalian target of rapamycin (mTOR) acts as a major anti-autophagy protein which is
6 regulated by upstream signaling pathway PtdIns3K/Akt, p53-AMP-activated protein kinase and
7 several other proteins [31]. Regarding the mechanisms involved in autophagy induction, recent
8 studies have indicated that both inhibition of PI3K/Akt/mTOR signaling and translocation of p53
9 into the nucleus could trigger autophagy in cancer cell [32]. It has been reported that many drugs
10 could induce autophagy through inhibition of PI3K/Akt/mTOR pathway [19, 20, 33]. Recently,
11 more attention has been paid on the autophagy modulation function of the tumor suppressor p53
12 protein. Nuclear p53 can transcriptionally activate autophagy inducers, while cytoplasmic p53
13 inhibits autophagy [34-36]. Following EGFR inhibition, its downstream signals including Akt
14 and mTOR would be effectively abolished resulting in both apoptosis and autophagy. In
15 addition, our findings also confirmed p53 nuclear translocation and APMK activation upon
16 erlotinib treatment in EGFR-mutated NSCLC cell lines. These concerted efforts could eventually
17 lead to effective blockade of mTOR, and release of its negative control on autophagy.

18

19 There are several limitations of our study. Due to limited number of NSCLC cell lines in our
20 model, our findings could be cell line-specific biological behavior rather than a generalizable
21 phenomenon. The measurement of autophagy was limited to within 72 h in our system, which
22 might not reflect the situation of prolonged exposure to erlotinib as in clinical application. Also,

1 the influence of tumor microenvironment on autophagic response to erlotinib could not be
2 assessed in our model. Future studies of autophagy in acquired resistant cell lines derived from
3 parental lines (HCC827 and HCC4006) or clinical tumor samples obtained from lung cancer
4 patients with acquired resistance to EGFR TKI will help to further elucidate the clinical
5 significance of autophagy as a resistance mechanism to EGFR TKI.

6

7 In conclusion, autophagy can be induced upon exposure to clinically relevant concentrations of
8 erlotinib in a NSCLC cell line model with activating EGFR mutations. Strategies to inhibit
9 autophagy may enhance sensitivity to EGFR TKI and potentially overcome acquired resistance.

10

1

2 **CONFLICT OF INTEREST STATEMENT**

3 All authors in this manuscript have no financial and personal relationships with other people or
4 organizations that could inappropriately influence our work. None of the authors have conflict of
5 interest to declare.

6

7 **ACKNOWLEDGEMENTS**

8 There is no funding source to declare. No writing assistance was sought in the preparation of this
9 report.

10

11

1 **References**

- 2 1. Sun S, Schiller JH, Gazdar AF. Lung cancer in never smokers--a different disease. *Nat Rev Cancer*
3 2007;7: 778-790.
- 4 2. Spiro SG, Silvestri GA. One hundred years of lung cancer. *Am J Respir Crit Care Med* 2005;172:
5 523-529.
- 6 3. Spiro SG, Tanner NT, Silvestri GA, Janes SM, Lim E, Vansteenkiste JF, Pirker R. Lung cancer:
7 progress in diagnosis, staging and therapy. *Respirology* 2010;15: 44-50.
- 8 4. Ray M, Salgia R, Vokes EE. The role of EGFR inhibition in the treatment of non-small cell lung
9 cancer. *Oncologist* 2009;14: 1116-1130.
- 10 5. Mendelsohn J, Baselga J. The EGF receptor family as targets for cancer therapy. *Oncogene*
11 2000;19: 6550-6565.
- 12 6. Hirsch FR, Varella-Garcia M, Bunn PA, Di Maria MV, Veve R, Bremnes RM, Baron AE, Zeng C,
13 Franklin WA. Epidermal growth factor receptor in non-small-cell lung carcinomas: Correlation
14 between gene copy number and protein expression and impact on prognosis. *Journal of Clinical*
15 *Oncology* 2003;21: 3798-3807.
- 16 7. Henson ES, Gibson SB. Surviving cell death through epidermal growth factor (EGF) signal
17 transduction pathways: Implications for cancer therapy. *Cellular Signalling* 2006;18: 2089-2097.
- 18 8. Herbst RS, Bunn PA. Targeting the epidermal growth factor receptor in non-small cell lung
19 cancer. *Clinical Cancer Research* 2003;9: 5813-5824.
- 20 9. Bunn PA, Franklin W. Epidermal growth factor receptor expression, signal pathway, and
21 inhibitors in non-small cell lung cancer. *Seminars in Oncology* 2002;29: 38-44.
- 22 10. Ji HB, Zhao XJ, Yuza Y, Shimamura T, Li DN, Protopopov A, Jung BL, McNamara K, Xia HL, Glatt KA,
23 Thomas RK, Sasaki H, Horner JW, Eck M, Mitchell A, Sun YP, Al-Hashem R, Bronson RT,
24 Rabindran SK, Discafani CM, Maher E, Shapiro GI, Meyerson M, Wong KK. Epidermal growth

- 1 factor receptor variant III mutations in lung tumorigenesis and sensitivity to tyrosine kinase
2 inhibitors. Proceedings of the National Academy of Sciences of the United States of America
3 2006;103: 7817-7822.
- 4 11. Kosaka T, Yatabe Y, Endoh H, Yoshida K, Hida T, Tsuboi M, Tada H, Kuwano H, Mitsudomi T.
5 Analysis of epidermal growth factor receptor gene mutation in patients with non-small cell lung
6 cancer and acquired resistance to gefitinib. Clinical Cancer Research 2006;12: 5764-5769.
- 7 12. Kwak EL, Sordella R, Bell DW, Godin-Heymann N, Okimoto RA, Brannigan BW, Harris PL, Driscoll
8 DR, Fidias P, Lynch TJ, Rabindran SK, McGinnis JP, Wissner A, Sharma SV, Isselbacher KJ,
9 Settleman J, Haber DA. Irreversible inhibitors of the EGF receptor may circumvent acquired
10 resistance to gefitinib. Proceedings of the National Academy of Sciences of the United States of
11 America 2005;102: 7665-7670.
- 12 13. Pao W, Miller VA, Politi KA, Riely GJ, Somwar R, Zakowski MF, Kris MG, Varmus H. Acquired
13 resistance of lung adenocarcinomas to gefitinib or erlotinib is associated with a second mutation
14 in the EGFR kinase domain. Plos Medicine 2005;2: 225-235.
- 15 14. Klionsky DJ, Emr SD. Cell biology - Autophagy as a regulated pathway of cellular degradation.
16 Science 2000;290: 1717-1721.
- 17 15. Baehrecke EH. Autophagy: Dual roles in life and death? Nature Reviews Molecular Cell Biology
18 2005;6: 505-510.
- 19 16. Rabinowitz JD, White E. Autophagy and Metabolism. Science 2010;330: 1344-1348.
- 20 17. Apel A, Herr I, Schwarz H, Rodemann HP, Mayer A. Blocked autophagy sensitizes resistant
21 carcinoma cells to radiation therapy. Cancer Research 2008;68: 1485-1494.
- 22 18. Bellodi C, Lidonnici MR, Hamilton A, Helgason GV, Soliera AR, Ronchetti M, Galavotti S, Young
23 KW, Selmi T, Yacobi R, Van Etten RA, Donato N, Hunter A, Dinsdale D, Tirro E, Vigneri P, Nicotera
24 P, Dyer MJ, Holyoake T, Salomoni P, Calabretta B. Targeting autophagy potentiates tyrosine

- 1 kinase inhibitor-induced cell death in Philadelphia chromosome-positive cells, including primary
2 CML stem cells. *Journal of Clinical Investigation* 2009;119: 1109-1123.
- 3 19. Gorzalczany Y, Gilad Y, Amihai D, Hammel I, Sagi-Eisenberg R, Merimsky O. Combining an EGFR
4 directed tyrosine kinase inhibitor with autophagy-inducing drugs: A beneficial strategy to
5 combat non-small cell lung cancer. *Cancer Letters* 2011;310: 207-215.
- 6 20. Han WD, Pan HM, Chen Y, Sun J, Wang YS, Li J, Ge WT, Feng LF, Lin XY, Wang XJ, Wang X, Jin HC.
7 EGFR Tyrosine Kinase Inhibitors Activate Autophagy as a Cytoprotective Response in Human
8 Lung Cancer Cells. *Plos One* 2011;6.
- 9 21. Kanzawa T, Kondo Y, Ito H, Kondo S, Germano I. Induction of autophagic cell death in malignant
10 glioma cells by arsenic trioxide. *Cancer Research* 2003;63: 2103-2108.
- 11 22. Wang AL, Boulton ME, Dunn WA, Rao HV, Cai J, Lukas TJ, Neufeld AH. Using LC3 to monitor
12 autophagy flux in the retinal pigment epithelium. *Autophagy* 2009;5: 1190-1193.
- 13 23. Mizushima N, Yoshimori T. How to interpret LC3 immunoblotting. *Autophagy* 2007;3: 542-545.
- 14 24. Hanada T, Noda NN, Satomi Y, Ichimura Y, Fujioka Y, Takao T, Inagaki F, Ohsumi Y. The Atg12-
15 Atg5 conjugate has a novel E3-like activity for protein lipidation in autophagy. *Journal of*
16 *Biological Chemistry* 2007;282: 37298-37302.
- 17 25. Li X, Fan Z. The epidermal growth factor receptor antibody cetuximab induces autophagy in
18 cancer cells by downregulating HIF-1 α and Bcl-2 and activating the beclin 1/hVps34 complex.
19 *Cancer Research* 2010;70: 5942-5952.
- 20 26. Bjorkoy G, Lamark T, Pankiv S, Overvatn A, Brech A, Johansen T. Monitoring autophagic
21 degradation of p62/SQSTM1. *Methods Enzymol* 2009;452: 181-197.
- 22 27. Huang B, Meng N, Zhao B, Zhao J, Zhang Y, Zhang S, Miao J. Protective effects of a synthesized
23 butyrolactone derivative against chloroquine-induced autophagic vesicle accumulation and the

- 1 disturbance of mitochondrial membrane potential and Na⁺,K⁺-ATPase activity in vascular
2 endothelial cells. *Chem Res Toxicol* 2009;22: 471-475.
- 3 28. Ertmer A, Huber V, Gilch S, Yoshimori T, Erfle V, Duyster J, Elsasser HP, Schatzl HM. The
4 anticancer drug imatinib induces cellular autophagy. *Leukemia* 2007;21: 936-942.
- 5 29. Han J, Hou W, Goldstein LA, Lu C, Stolz DB, Yin XM, Rabinowich H. Involvement of protective
6 autophagy in TRAIL resistance of apoptosis-defective tumor cells. *Journal of Biological Chemistry*
7 2008;283: 19665-19677.
- 8 30. Li X, Lu Y, Pan T, Fan Z. Roles of autophagy in cetuximab-mediated cancer therapy against EGFR.
9 *Autophagy* 2010;6: 1066-1077.
- 10 31. Wullschleger S, Loewith R, Hall MN. TOR signaling in growth and metabolism. *Cell* 2006;124:
11 471-484.
- 12 32. Wang N, Pan W, Zhu M, Zhang M, Hao X, Liang G, Feng Y. Fangchinoline induces autophagic cell
13 death via p53/sestrin2/AMPK signalling in human hepatocellular carcinoma cells. *Br J Pharmacol*
14 2011;164: 731-742.
- 15 33. Viola G, Bortolozzi R, Hamel E, Moro S, Brun P, Castagliuolo I, Ferlin MG, Basso G. MG-2477, a
16 new tubulin inhibitor, induces autophagy through inhibition of the Akt/mTOR pathway and
17 delayed apoptosis in A549 cells. *Biochem Pharmacol* 2012;83: 16-26.
- 18 34. Maiuri MC, Malik SA, Morselli E, Kepp O, Criollo A, Mouchel PL, Carnuccio R, Kroemer G.
19 Stimulation of autophagy by the p53 target gene Sestrin2. *Cell Cycle* 2009;8: 1571-1576.
- 20 35. Ghavami S, Mutawe MM, Sharma P, Yeganeh B, McNeill KD, Klonisch T, Unruh H, Kashani HH,
21 Schaafsma D, Los M, Halayko AJ. Mevalonate cascade regulation of airway mesenchymal cell
22 autophagy and apoptosis: a dual role for p53. *Plos One* 2011;6: e16523.


1 36. Morselli E, Shen S, Ruckstuhl C, Bauer MA, Marino G, Galluzzi L, Criollo A, Michaud M, Maiuri
2 MC, Chano T, Madeo F, Kroemer G. p53 inhibits autophagy by interacting with the human
3 ortholog of yeast Atg17, RB1CC1/FIP200. *Cell Cycle* 2011;10: 2763-2769.

4


5

6

1 Legends of figures







1 **Figure 1.** Autophagy induction in HCC827, HCC4006, H358 and H1975 cells after erlotinib
2 treatment. (A) LC3-II expression was increased in HCC827 and HCC4006 cells after 24h
3 erlotinib (0.1 μ M, 0.2 μ M) treatment. (B) LC3-II expression was unchanged in H358 and H1975
4 cells after 24h erlotinib (up to 10 μ M) treatment. (C) Autophagic flux with bafilomycin A1
5 treatment was shown in both HCC827 and HCC4006 cells. Acidic vesicular organelle (AVO)
6 formation detected by acridine orange staining in (D) HCC827 and (E) HCC4006 cells, but not
7 in (F) H358 and (G) H1975 cells after 48h erlotinib treatment. Chloroquine (CQ) treatment
8 served as a positive control. Arrows indicate the AVO positive cells. (H) Atg5/12 conjugation
9 and (I) p62 degradation with erlotinib treatment (0.2 μ M) in HCC827 and HCC4006 cells. Data
10 are shown as mean \pm SD (error bars) of 3 independent experiments, depicted with representative
11 immunoblots. Statistical comparison was made between treatment vs control groups, or different
12 time points vs baseline (0h), unless otherwise indicated. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$;
13 ns, not significant.




1

2 **Figure 2.** Pathways involved in autophagy induction with erlotinib treatment (0.2 μ M). Nuclear
 3 translocation of p53 was detected in (A) HCC827 and (B) HCC4006 cells. (C) AMP-activated
 4 protein kinase and (D) p-NF κ B pathways were activated. (E) mTOR pathway was inhibited.
 5 Data are shown as mean \pm SD (error bars) of 3 independent experiments, depicted with
 6 representative immunoblots. Statistical comparison was made between different time points vs
 7 baseline (0h). *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

8



1 **Figure 3.** Inhibition of autophagy with chloroquine (CQ) enhanced the effect of erlotinib
2 treatment in EGFR-mutated, but not in wild-type, NSCLC cells. (A) LC3II accumulated after
3 24h of chloroquine exposure in HCC827, HCC4006, H358 and H1975 cells. (B) Cell viability by
4 crystal violet staining in HCC827, HCC4006, H358 and H1975 cells after erlotinib ±
5 chloroquine treatment. (C) Apoptotic cell death detected by Annexin V/7-AAD flow cytometry
6 in HCC827, HCC4006, H358 and H1975 cells after erlotinib ± chloroquine treatment. (D)
7 Cleaved PARP detected by Western blot after erlotinib, chloroquine, or combined treatment in
8 sensitive NSCLC cells. Data are shown as mean ± SD (error bars) of 3 independent experiments,
9 depicted with representative immunoblots. Statistical comparison was made between treatment
10 vs control groups, unless otherwise indicated. *, p< 0.05; **, p< 0.01; ***, p < 0.001; ns, not
11 significant.



1

2 **Figure 4.** Combination of erlotinib with siRNA targeting autophagy related genes Atg5 and

3 Beclin-1 in sensitive NSCLC cells. (A) Atg 5 and Beclin-1 were efficiently downregulated in

1 HCC827 and HCC4006 cells, with representative immunoblots shown. GAPDH served as the
2 loading control. (B) Cell viability determined by MTT. (C) Apoptotic cell death determined by
3 flow cytometry. Data are shown as mean \pm SD (error bars) of 3 independent experiments.
4 Statistical comparison was made between treatment groups as indicated. *, $p < 0.05$; **, $p < 0.01$;
5 ***, $p < 0.001$; ns, not significant.

6 |
7

Table 1. The percentage of apoptotic cells (Annexin-V/7-AAD assay) in different cell lines treated with erlotinib, chloroquine or their combination.

| Cell line | Erlotinib | Chloroquine | Chloroquine + Erlotinib |
|------------------|------------------|--------------------|--------------------------------|
| HCC827 | 21.2 ± 4.2% | 13.0 ± 2.5% | 52.3 ± 2.8% |
| HCC4006 | 9.4 ± 2.4% | 9.0 ± 1.8% | 49.3 ± 4.8% |
| H358 | 17.4 ± 2.8% | 7.5 ± 0.7% | 17.7 ± 1.7% |
| H1975 | 6.9 ± 1.8% | 5.5 ± 2.0% | 8.7 ± 2.8% |