



Title	Assessment of the neurocentral synchondrosis in pediatric spines and the "developmental" etiology of Schmorl's nodes
Author(s)	Luk, KDK; Law, TK; Khong, PL; Anthony, M; Samartzis, D
Citation	The 40th Annual Meeting of the International Society for the Study of the Lumbar Spine (ISSLS 2013), Scottsdale, AZ., 13-17 May 2013. In Oral and Special Posters, 2013, p. 64, abstract SP13
Issued Date	2013
URL	http://hdl.handle.net/10722/189810
Rights	Creative Commons: Attribution 3.0 Hong Kong License

ASSESSMENT OF THE NEUROCENTRAL SYNCHONDROSIS IN PEDIATRIC SPINES AND THE "DEVELOPMENTAL" ETIOLOGY OF SCHMORL'S NODES

(1) Keith Luk, (2) Travis Law, (2) Pek-Lan Khong, (2) Marina Portia-Anthony, (1) Dino Samartzis

1) Department of Orthopaedics and Traumatology, University of Hong Kong, Hong Kong, SAR China 2) Department of Diagnostic Radiology, University of Hong Kong, Hong Kong, SAR China

INTRODUCTION: Schmorl's nodes of the thoraco-lumbar region have been associated with the presence and severity of disc degeneration in adults. The etiology of Schmorl's nodes remains precarious; however a unique multilevel phenotype of "kissing" nodes seems to suggest a developmental origin. The neurocentral synchondrosis (NCS) are cartilaginous growth plates near the neural arch ossification centres in a growing vertebrae that may play a direct role in the development of endplate abnormalities, such as Schmorl's nodes. This study assessed the NCS in pediatric spines to raise discussion of a developmental component of Schmorl's nodes.

METHODS: A retrospective imaging study of pediatric patients at a single institute assessed over a 5 year period (age range: 0 to 10 years) was performed. Patients with spinal disorders (e.g. scoliosis) were excluded. 102 patients (57 males, 45 females) and a total of 113 sets of MRI images were reviewed. The thoraco-lumbar regions (T12-S1) were evaluated. Sagittal T1- and T2-weighted images were assessed for the presence or absence of the NCS, defined as a hypointense vertical line located between the vertebral body anteriorly and the posterior arch. The presence of Schmorl's nodes was also noted.

RESULTS: NCS was noted in 46% of the MRIs. No statistically significant difference in the disappearance of the NCS between different age groups ($p=0.063$) or gender ($p=0.706$) was found. The NCS was noted to be completely fused at the midpoint of the vertebrae. Indentation of the vertebral endplates resembling Schmorl's node at many of the rostral and caudal ends of the unfused NCS were observed (Figure 1).

CONCLUSIONS: The significance of further characterizing the nature of NCS closure may lie in potential associations with failure of complete closure with endplate abnormalities, such as Schmorl's nodes. Our MRI study provides a foundation that a developmental etiology of Schmorl's nodes exists.