The HKU Scholars Hub The University of Hong Kong 香港大學學術庫



Title	Segregation of the polyphyletic genus Polyalthia (Annonaceae)
Author(s)	Xue, B; Su, CFY; Thomas, DC; Saunders, RMK
Citation	The 9th International Flora Malesiana Symposium, Bogor, Indonesia, 27-31 August 2013. In Programme Book, 2013, p. 23
Issued Date	2013
URL	http://hdl.handle.net/10722/190646
Rights	Creative Commons: Attribution 3.0 Hong Kong License

## Segregation of the polyphyletic genus Polyalthia (Annonaceae)

Xue, B.\*1, Su, C.F.Y.1,2, Thomas, D.C.1,3 & Saunders, R.M.K.11 School of Biological Sciences, The University of Hong Kong, P. R China 2Duke-NUS Graduate Medical School Singapore 3Netherlands Centre for Biodiversity Naturalis, section NHN, Leiden University \*e-mail: Xue Bine xuebine@gmail.com

The circumscription of the species-rich genus *Polyalthia* (Annonaceae, with ca. 155 species) has long been recognised to be highly problematic: as previously circumscribed, the genus was a morphologically heterogeneous assemblage lacking conspicuous synapomorphies, and was one of the most significant impediments to revisionary and floristic taxonomic research in the family. Recent molecular phylogenetic analyses have confirmed that *Polyalthia* is polyphyletic. A broader taxonomic sampling of *Polyalthia* species and associated taxa is included in the present phylogenetic analyses to enable the recognition of monophyletic genera with clearly defined diagnostic morphological characters. Bayesian, maximum parsimony and maximum likelihood analyses of chloroplast *matK*, *rbcL* and *trnL-F* sequences consistently gave congruent topologies, with *Polyalthia* species distributed in four well-supported clades. Based on molecular and morphological data, existing *Polyalthia* species are classified into the following four genera: (1) Hubera: a newly established genus consists of 27 Polyalthia species; (2) Marsypopetalum: six species, following the transfer of five *Polyalthia* species; (3) *Monoon*: a revived generic name with 56 species, following the transfer of 37 Polyalthia species, 18 Enicosanthum species, and Woodiellantha sympetala; and (4) Polyalthia s.str.: ca. 86 species, following the transfer of 11 Haplostichanthus species.