



Title	MicroRNA-143 is downregulated in breast cancer and regulates DNA methyltransferases 3A in breast cancer cells
Author(s)	Ng, EKO; Li, R; Shin, VY; Siu, JMT; Ma, ESK; Kwong, A
Citation	Tumor Biology, 2014, v. 35 n. 3, p. 2591-2598
Issued Date	2014
URL	http://hdl.handle.net/10722/193223
Rights	The original publication is available at www.springerlink.com

MicroRNA-143 is downregulated in breast cancer and regulates DNA methyltransferases 3A in breast cancer cells

Enders K. O. Ng · Rufina Li · Vivian Y. Shin · Jennifer M. Siu · Edmond S. K. Ma · Ava Kwong

Received: 23 May 2013 / Accepted: 16 October 2013
© International Society of Oncology and BioMarkers (ISOBM) 2013

Abstract MicroRNAs (miRNAs) are small non-protein-coding RNAs that regulate expression of a wide variety of genes including those involved in cancer development. Here, we investigate the role of miR-143 in breast cancer. In this study, we showed that miR-143 was frequently downregulated in 80 % of breast carcinoma tissues compared to their adjacent noncancerous tissues. Ectopic expression of miR-143 inhibited proliferation and soft agar colony formation of breast cancer cells and also downregulated DNA methyltransferase 3A (DNMT3A) expression on both mRNA and protein levels. Restoration of miR-143 expression in breast cancer cells reduces PTEN hypermethylation and increases TNFRSF10C methylation. DNMT3A was demonstrated to be a direct target of miR-143 by luciferase reporter assay. Furthermore, miR-143 expression was observed to be inversely correlated with DNMT3A mRNA and protein expression in breast cancer tissues. Our findings suggest that miR-143 regulates DNMT3A in breast cancer cells. These findings elucidated a tumor-suppressive role of miR-143 in epigenetic aberration of breast cancer, providing a potential development of miRNA-based treatment for breast cancer.

Keywords miR-143 · DNMT3A · Breast cancer · Tumor suppressor · PTEN 36
37

Abbreviations		38 Q3
miRNA	microRNA	40
DNMT	DNA methyltransferase	43
qRT-PCR	Quantitative reverse transcription–polymerase chain reaction	44 46
3'UTR	3' Untranslated region	48
MTT	3-[4,5-Dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide	50 51 52

Introduction 53 54

Breast cancer is one of the three most commonly diagnosed cancers among women, accounting for about 30 % of patients [1]. In the past decades, despite the dedication of research and resources to the development of biomarkers for diagnosis and prognosis, unpredictable response and development of resistance to adjuvant therapy remain major challenges in breast cancer management. Although mammography diagnosis for breast cancer is the currently used screening tool, the cost incurred and expertise required for mammogram has hampered wide application of this procedure. On the other hand, alternative methods such as ultrasound screening has very operator-dependent sensitivity, and tumor markers such as CA15.3 and carcinoembryonic antigen (CEA) are also nonspecific and has limited sensitivity and specificity [2]. Thus, there is still a pressing need to elucidate novel mechanism of breast cancer development so as to develop a cost-effective and accurate screening method for this cancer. 54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71

Recently, the emergence of small non-protein-coding RNAs, microRNAs (miRNAs), playing important roles in 72
73

E. K. O. Ng · R. Li · V. Y. Shin · J. M. Siu · A. Kwong
Department of Surgery, The University of Hong Kong,
Hong Kong SAR, Hong Kong

E. K. O. Ng · E. S. K. Ma
Department of Molecular Pathology, Hong Kong Sanatorium
and Hospital, Hong Kong SAR, Hong Kong

A. Kwong
The Hong Kong Hereditary Breast Cancer Family Registry,
Hong Kong, Hong Kong

A. Kwong (✉)
Chief of Breast Surgery Division, The University of Hong Kong,
Hong Kong SAR, Hong Kong
e-mail: akwong@asiabreastregistry.com

74 oncogenesis, has opened new opportunities for early cancer
 75 diagnosis [3, 4]. Evidence suggests that miRNA expression
 76 profiles can cluster similar tumor types together more
 77 accurately than the expression profiles of protein-coding
 78 mRNA genes [5]. Furthermore, miRNA expression signatures
 79 have been used to predict prognosis [6, 7]. Importantly,
 80 expression of some miRNAs correlated with the molecular
 81 subtypes and with two major features of breast cancer (grade
 82 and ER status) [8]. Therefore, miRNA has a great potential to
 83 be a novel biomarker for breast cancer and holds promising
 84 potential for individualizing patients' treatment regimens [9],
 85 although, as yet, there is still limited knowledge on the exact
 86 mRNA target of the deregulated miRNA in breast cancer.
 87 Research shows that each miRNA could target up to 200
 88 mRNA transcripts, and a single mRNA could have multiple
 89 miRNA binding sites [10]. This finding indicates that there is
 90 a great demand to further investigate on the mRNA targets and
 91 understand the functional role of these differentially expressed
 92 miRNAs, so as to elucidate their potential as therapeutic
 93 agents or targets.

94 In this study, we investigated the functional role of miR-
 95 143 in breast cancer. MiR-143, located on chromosome 5q33,
 96 is a miRNA found to be deregulated in colon cancer [11] and
 97 bladder cancer [12]. It is previously demonstrated that miR-
 98 143 targets on DNA methyltransferase 3A (DNMT3A)
 99 mRNA [11]. DNMT3A is the member of the
 100 methyltransferase family. DNMT3A and 3B are responsible
 101 for de novo methylation in the genome [13], while DNMT1 is
 102 responsible for maintaining methylation in the genome. The
 103 expression level of DNMT3A is high in early embryonic stage
 104 and downregulated in differentiated cells; maintaining high
 105 expression of DNMT3A in embryonic cells will inhibit cell
 106 differentiation [13].

107 Until now, the role of DNMT3A in cancer is less studied
 108 than DNMT3B. There are reports showing that DNMT3A
 109 deficiency promotes tumor growth and progression [13].
 Q4 110 The downregulation of miR-143 in tumor can lead to the
 111 overexpression of DNMT3A, which in turn causes
 112 hypermethylation and silencing of the tumor suppressor genes
 113 and contributes to tumorigenesis.

114 One of the most researched tumor suppressor genes is the
 115 phosphatase and tensin homolog (PTEN) which acts as a
 Q5 116 negative regulator of PI3K/AKT signaling pathway [14]. A
 117 high proportion of human cancers have a mutated form of
 118 PTEN or abnormal PTEN expression, and this attributed to
 119 40 % of breast cancer [15]. Mutation and inactivation of
 120 PTEN gene lead to hyperactivation of PI3K/AKT pathway,
 121 which causes cell cycle deregulation and suppression of
 122 apoptosis [16]. Evidences showed that breast cancer patients
 123 with defective PTEN have poor prognosis and high grade
 124 tumor [15].

125 We aimed to show that miR-143 and DNMT3A are both
 126 deregulated in breast cancer and prove that overexpression of

DNMT3A has caused a change in methylation status of PTEN 127
 and TNFRSF10C which contributed to tumorigenesis. These 128
 results help to understand the molecular mechanism of how 129
 miR-143 promotes cancer progression. 130

Materials and methods 131

Cell lines and tissue samples 132

Five human breast cancer cell lines including MCF-7, MD- 133
 MB-231, MD-MB-468, T47D, and SK-BR-3 and two colon 134
 cancer cell lines HT-29 and SW480 (American Type Culture 135
 Collection, Manassas, VA) were cultured at 37 °C in 10 % 136
 CO₂ atmosphere and maintained routinely in Dulbecco's 137
 modified Eagle's Medium (DMEM) supplemented with 138
 10 % fetal bovine serum and 2 mM of L-glutamine 139
 (Invitrogen, Carlsbad, CA). A total of 20 pairs of primary 140
 breast tumors and noncancerous tissue counterparts were 141
 collected. All samples were collected from patients who 142
 underwent surgical resection of tumors. Informed consent 143
 has been obtained from each patient. This project was 144
 approved by the Institutional Review Board of the University 145
 of Hong Kong. 146

Real-time quantitative PCR 147

Total RNA containing small RNA was extracted from tissues 148
 and cell lines by TRIzol reagent (Invitrogen) according to the 149
 instructions of the manufacturer. SYBR Green real-time 150
 qPCR assay for miRNA expression was used as previously 151
 described [11, 17]. In brief, 100 ng of total RNA containing 152
 miRNA was polyadenylated and reverse-transcribed to cDNA 153
 by using miScript Reverse Transcription Kit (Qiagen) 154
 according to the manufacturer's instructions. Real-time qPCR 155
 was performed using miScript SYBR Green PCR Kit 156
 (Qiagen) in ABI PRISM 7900 HT System (Applied 157
 Biosystems, Foster City, CA). The miR-143-specific forward 158
 primer sequence was 5'-TGAGATGAAGCACTGTAGCTC- 159
 3' and was designed based on the miRNA sequences obtained 160
 from the miRBase database. Human U6 snRNA was used for 161
 normalization. For DNMT3A mRNA qPCR, total RNA was 162
 reverse-transcribed to cDNA by using miScript Reverse 163
 Transcription Kit (Qiagen) according to the manufacturer's 164
 instructions. Gene-specific primers for DNMT3A gene were 165
 used as previously described [17]. The mRNA expression was 166
 normalized to β-actin. ΔCt was calculated by subtracting the 167
 Ct values of U6 or β-actin from the Ct values of the gene of 168
 interest. ΔΔCt was then calculated by subtracting the ΔCt of 169
 the control from the ΔCt of cancer sample. Fold change of 170
 gene was calculated by the equation $2^{-\Delta\Delta Ct}$. 171

172	Ectopic miR-143 expression	Luciferase activity assay	215
173	Ectopic expression of miR-143 in breast cancer cells (MD-	DNMT3A 3'UTR containing an intact miR-143 recognition	216
174	MB-231 and MCF7) was achieved by transfection with	sequence was amplified, and the PCR product (199 bp) was	217
175	mature miR-143 mimic (Qiagen). Cells were plated in culture	subcloned into pGL3 basic vector (Promega, Madison, WI)	218
176	dishes or 6/96-well plates for 24 h and transfected with 1 nM	immediately downstream of luciferase gene, as described	219
177	of mimic with HiPerFect Transfection Reagent (Qiagen) for	previously [11]. A pGL3 construct containing DNMT3A 3'	220
178	24 h. Precursor control (Ambion, Austin, TX) was used as	UTR with point mutations in seed sequence was also	221
179	negative control. Cells were then subjected to further assays or	synthesized using Site-Directed Mutagenesis Kit (Stratagene,	222
180	for RNA/protein extraction.	La Jolla, CA) according to the manufacturer's instructions.	223
		Cells were co-transfected with 800 ng of pGL3 constructs	224
		with or without miR-143 precursor for 24 h. Each sample	225
181	Cell proliferation assay	was co-transfected with 0.05 µg pRL-CMV plasmid	226
		expressing Renilla luciferase to monitor the transfection	227
182	Cell proliferation was measured by 3-(4,5-dimethylthiazol-2-	efficiency (Promega, Madison, WI). Luciferase activity assay	228
183	yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-	was performed 24 h after transfection using Dual-Luciferase	229
184	tetrazolium (MTT) assay (Promega Corporation, Madison,	Reporter Assay System (Promega). Relative luciferase	230
185	WI). MB-231 (2×10^6) and MCF-7 (5×10^6) cells were	activity was normalized with Renilla luciferase activity.	231
186	seeded in a 96-well plate for 24 h, transfected with 1 nM		
187	miR-143 mimic (Qiagen) and HiPerFect Transfection	Methylation-sensitive PCR	232
188	Reagent (Qiagen) for 24 h and further grown in normal		
189	medium for 3 days. Thereafter, cells were incubated in	Genomic DNA from cell lines with or without miR-143-mimic	233
190	0.1 mg/ml MTT at 37 °C for 3 h and lysed in dimethyl	transfection used for methylation analysis was extracted by	234
191	sulfoxide (DMSO) at room temperature for 30 min. The	DNeasy Mini Kit (Qiagen) according to the user manual.	235
192	absorbance in each well was measured at 580 nm by a	Methyl-Profilier DNA Methylation qPCR Primer Assays (SA	236
193	microplate reader.	Biosciences) was used to determine the methylation status of	237
		the promoter in different genes. In brief, 250 µg of genomic	238
		DNA was used for enzyme digestion by using a Methyl-	239
194	Anchorage-independent colony formation assay	Profilier Enzyme Kit. For each sample, mock digestion (Mo),	240
		methylation-sensitive digestion (Ms), methylation-dependent	241
195	Soft agar plates were prepared in 24-well plates with a bottom	digestion (Md), and double digestion (Msd) was performed by	242
196	layer of 0.6 % Noble agar in serum-free DMEM. Cells were	adding different combinations of enzyme according to the	243
197	trypsinized, and 500 cells were seeded onto the bottom layer	manufacturer's protocol and was placed in 37 °C heating block	244
198	after being mixed with 0.3 % Noble agar in DMEM	for 6 h, followed by heat inactivation at 65 °C for 20 min.	245
199	supplemented with 10 % fetal calf serum. Plates were	SYBR Green-based qPCR was performed with a panel of 26	246
200	incubated at a 37 °C incubator for 3 weeks. The number of	breast cancer methylated gene promoters (MeAH-011C, SA	247
201	colonies was counted after stained with 0.05 % crystal violet	Biosciences), on PRISM 7900 HT. Ct was obtained after	248
202	for 1 h and washed extensively with phosphate-buffered saline	qPCR, and the relative amount of methylation was calculated	249
203	(PBS).	by first determining the relative amount of DNA resistance to	250
		enzyme digestion (Cr): $2^{-\Delta Ct(Msd-Mo)}$. Then the degree of	251
		methylation of each gene promoter can be calculated as follows:	252
204	Western blot analysis	(1) amount of hypermethylation (C_{HM}): $(2^{-\Delta Ct(Ms-Mo)} - Cr) / (1 -$	253
		$Cr)$; (2) amount of hypomethylation (C_{UM}): $(2^{-\Delta Ct(Md-Mo)} - Cr) /$	254
205	Cells were lysed in Lammeli's lysis buffer, resolved in SDS-	$(1 - Cr)$; and (3) amount of intermediately methylated DNA: $1 -$	255
206	PAGE minigel, and transferred onto Immobilon-P membrane	$C_{HM} - C_{UM}$.	256
207	(Millipore, Billerica, MA). Membranes were probed with 1:1,		
208	000 diluted primary antibodies against DNMT3A (Cell	Statistical analysis	257
209	Signaling) at room temperature for 2 h, washed extensively		
210	with 0.1 % Tween-20 in PBS, and incubated with secondary	Paired <i>t</i> test was used in the expression comparison of miR-	258
211	antibodies conjugated with horse-radish peroxidase (1:10,000	143 between paired breast tumor and adjacent noncancerous	259
212	dilution). The signals were visualized with enhanced	tissues. Two-sided Student's <i>t</i> test was used to analyze MTT	260
213	chemiluminescence (Amersham Life Science Inc.,	assay, anchorage-independent soft agar assay, and luciferase	261
214	Buckinghamshire, UK).	reporter assay. Data are expressed as the mean ± SD from at	262

263 least three independent experiments. All *P* values are two-
 264 sided, and a value of less than 0.05 was considered statistically
 265 significant. All statistical calculations were performed by the
 266 SPSS software (version 13.0, Chicago, IL, USA).

267 **Results**

268 MiR-143 is downregulated in breast tumor and human breast
 269 cancer cell lines

270 MiR-143 has been reported to be downregulated in other
 271 cancers like bladder cancer and colon cancer. To examine the
 272 expression levels of miR-143 in breast cancer, 20 pairs of breast
 273 tumor with adjacent normal tissue and five breast cancer cell
 274 lines were quantified by real-time PCR. The expression level of
 275 miR-143 in all the five breast cancer cell lines (MCF7, MB-
 276 231, MB-468, T47D, and SK-BR-3) was lower than that of
 277 noncancerous breast tissue (Fig. 1a). For patient samples, low
 278 expression of miR-143 was found in tumor compared with the
 279 adjacent normal tissues (*P*<0.05, Wilcoxon test; Fig. 1b).

280 The effect of miR-143 on cell growth and DNMT3A
 281 expression

282 Low expression levels of miR-143 in breast cancer cells
 283 suggest that miR-143 has a role in breast cancer
 284 carcinogenesis. To prove this, enforced expression of miR-
 285 143 on cell growth in MB-231 and T47D breast cancer cells
 286 was tested by MTT assay and colony formation assay. After
 287 transfection with miR-143, both MB-231 and T47D showed a
 288 significant decrease in growth rate (22 % decrease for MB-
 289 231 and 30 % decrease for T47D; Fig. 2a, b). Colony
 290 formation assay was performed to determine the degree of
 291 invasiveness in different cell lines after miR-143 mimic

transfection. Figure 2c showed the overexpression of miR-
 143 after transfection with miR-143 mimic when compared to
 control the precursor. The increased miR-143 expression
 significantly reduced anchorage-dependent growth in both
 cell lines as shown in Fig. 2c (all *P*<0.05; Mann–Whitney
 test), confirming that miR-143 also affects the malignant
 transformation phenotypes. These results suggested that
 miR-143 has a role in suppressing tumor cell growth. We then
 examined the correlation between miR-143 and DNMT3A.
 Our results indicated that restored expression of miR-143
 leads to decreased expressions of DNMT3A mRNA
 (Fig. 2d) and protein (Fig. 2e) in both cell lines, which suggest
 a potential regulatory role of miR-143 on DNMT3A.

Direct interaction between DNMT3A and miR-143

To confirm that DNMT3A is the direct target of miR-143,
 luciferase assay was performed. In short, wild-type (WT) or
 mutated (MUT) 3'UTR of DNMT3A (11) was subcloned into
 downstream of the firefly luciferase reporter and co-
 transfected with miR-143 precursor or precursor control into
 both MB-231 and T47D breast cancer cell lines. In the
 presence of miR-143, the relative luciferase activity of breast
 cancer cell lines with WT construct was significantly reduced
 (Fig. 3b; *P*<0.05 for MB-231, *P*<0.01 for T47D; Mann–
 Whitney test). While no significant suppressive effect by
 miR-143 was found in cells transfected with the MUT
 construct, this suggested a direct and specific interaction of
 miR-143 on DNMT3A 3'UTR in breast cancer cells.

Expression relationship between miR-143 and DNMT3A
 in breast tumor tissue

To confirm the relationship between miR-143 and DNMT3A,
 we assessed the expression of miR-143 and DNMT3A protein
 in breast tumor tissues from ten patients. As shown in Fig. 4a,
 there is no correlation between miR-143 and DNMT3A
 mRNA expression. However, DNMT3A protein levels were
 inversely correlated with miR-143 (*r*=−0.61, *P*<0.05;
 Spearman's correlation; Fig. 4b).

Methylation status of PTEN

To examine the effect of DNMT3A downregulation on
 methylation status of PTEN gene, we performed the
 methylation-sensitive PCR. As shown in Fig. 5, ectopic
 expression of miR-143 in MB-231 reduced hypermethylated
 DNA on PTEN gene promoter from 50 to 2.3 %, while that of
 unmethylated DNA raised from 50 to 97.7 %. In addition,
 hypermethylated TNFRSF10C reduced from 50 to 25 %,
 whereas unmethylated TNFRSF10C reduced from 50 to
 0.8 %, and that of intermediate methylated DNA increased
 from 0 to 73.8 %.

Q7 **Fig. 1** Downregulated miR-143 expression in both primary breast tumor tissues and breast cancer cell lines. **a** Relative miR-143 expression in breast cancer cell lines was much lower than the noncancerous breast tissue (NI). **b** Relative miR-143 expression between tumor and their paired adjacent nontumor tissues from 20 patients by real-time qPCR. Expression of miR-143 (Log₁₀ scale at Y-axis) was normalized to U6. Statistical difference was analyzed by Wilcoxon test, *P*<0.05





Fig. 2 Functional effect of ectopic miR-143 expression in MB-231 and T47D cells. Ectopic miR-143 expression reduced growth rate of both **a** MB-231 and **b** T47D cell lines assessed by MTT (Mann–Whitney test, $*P < 0.05$, $*P < 0.001$). **c** Anchorage-independent growth of cancer cells, examined by soft agar colony formation assay, was reduced. Cells were plated in 0.3 % Noble agar for 3 weeks. The number of colonies was counted after being stained with 0.05 % crystal violet (Mann–Whitney test, $*P < 0.05$). Overexpression of miR-143 reduced both **d** mRNA and **e** protein expression of DNMT3A. Cells were transfected with miR-143 precursor or control precursor for 24 h and then lysed for RNA or protein extraction. DNMT3A mRNA was detected by real-time qPCR (Mann–Whitney test, $*P < 0.05$), and the protein expression was detected by Western blotting with anti-DNMT3A antibody. β -actin was used as a loading control



339 **Discussion**

340 Since the discovery of miRNAs, the differential expression
 341 pattern of miRNAs in various cancers has been reported;
 342 however, the functional roles of individual miRNAs towards
 343 cellular transformation and tumorigenesis continue to be
 344 actively studied. Increasing evidence showed that miRNAs

might be involved in tumorigenesis by regulating oncogenes 345
 or tumor suppressor genes. A recent report showed that 346
 miRNA and epigenetic methylations are interconnected and 347
 contributed to tumorigenesis [18, 19]. In this study, we 348
 showed that miRNA can affect methylation through altering 349
 methyltransferase synthesis, which in turn affects tumor 350
 malignancy. 351




Fig. 3 DNMT3A is the direct target of miRNA-143. **a** The wild-type (WT) and mutant (MUT) DNMT3A 3'UTR, with or without point mutations in the seed sequence. **b** Ectopic miR-143 expression inhibited WT, but not MUT DNMT3A 3'UTR reporter activity in MB-231 and T47D cells. Cells were co-transfected with miR-143 precursor and either WT or MUT DNMT3A 3'UTR reporter construct. Luciferase activity assay was performed at 24 h post-transfection (Mann-Whitney test, * $P < 0.05$, ** $P < 0.01$)

352 It has been reported previously that miR-143 is deregulated
 353 in colorectal cancer [20], prostate cancer [21], B cell
 354 lymphoma [22], etc. In this study, we demonstrated that
 355 miR-143 is downregulated not only in breast cancer cell lines
 356 but also in primary breast tumors. The frequent
 357 downregulation of miR-143 suggests a tumor-suppressive role
 358 in breast cancer. We verified this by the enforced expression of




Fig. 4 Expression level of miR-143 and DNMT3A were tested in ten tumor samples. **a** Expression relationship between miR-143 and DNMT3A mRNA. **b** Scatter plot of the fold changes of miR-143 and DNMT3A protein (Log₁₀ scale at both X- and Y-axis) in (Spearman correlation, $r = -0.61$, $P < 0.05$)




Fig. 5 Effect of ectopic miR-143 expression on PTEN and TNFRSF10C promoter methylation status in MB-231 cells. HM, hypermethylated; UM, unmethylated; IM, intermediate methylated

miR-143 in breast cancer cells, resulting in a suppression of
 malignant transformation.

To further understand the tumor-suppressive role of miR-143, in silico target prediction (PicTar and TargetScan 5) is used for target prediction. Despite a large number of predicted potential targets for miR-143, only a limited amount was verified. There were reports showing that miR-143 acted on extracellular signal-regulated kinase 5 (Erk5) which in turn affects the mitogen-activated protein kinase (MAPK) pathways [21]. MAPK is an important pathway for oncogenesis, as it involves in cell proliferation, differentiation, and migration [23]. Apart from Erk5, DNMT3A is also a predicted target of miR-143.

DNMT3A together with DNMT1 and DNMT3B are catalytically active DNMTs responsible for genome methylation [24]. DNMT1 is a maintenance DNA methyltransferase for retaining methylation pattern, with inefficient de novo methylation ability. DNMT3A and DNMT3B are de novo methyltransferase with different targets [25]. Increasing evidence showed that these DNMTs work together to maintain a normal methylation pattern, and deregulation of either one could promote malignancies [26].

In this study, we showed that there is a correlation between miR-143 and DNMT3A in breast cancer. Enforced expression of miR-143 suppressed tumor transformation and DNMT3A mRNA and protein. Site-directed mutation on the 3'UTR of DNMT3A revealed the presence of specific binding site of miR-143. An inverse correlation of miR-143 and DNMT3A expression in human breast samples further consolidated miR-143 negatively regulated DNMT3A.

Genome-wide hypomethylation is common in cancer genomes which causes genome instability [27]; whereas site-specific hypermethylation in the promoter region of the tumor suppressor gene causing gene silencing is often observed [28]. Deregulation of methyltransferase could be

394 due to mutations in methyltransferase gene or imbalanced
 395 methyltransferase biogenesis. MiRNAs modulate
 396 posttranscription repression and maintain the balance of gene
 397 expression level in the cells [29]. We demonstrated that miR-
 398 143 targeted on DNMT3A gene and caused transcriptional
 399 repression. Low miR-143 expression increased the expression
 400 of DNMT3A enzyme which caused hypermethylation in other
 401 tumor-suppressing genes.

402 PTEN has long been known for its tumor-suppressive
 403 property; inactivation of PTEN could lead to various cancers
 404 [30]. Homozygous mutation is often found in familial and
 405 sporadic cancer. In breast cancer, reduction or complete
 406 absence of PTEN protein is found in about 40 % of the cases,
 407 mostly due to loss of heterozygosity (LOH), rarely somatic
 408 mutation [31]. The low mutation rate and high LOH suggested
 409 that epigenetic modification is responsible for the lost or
 410 reduced expression of PTEN protein. Many reports showed
 411 that promoter CpG hypermethylation is the reason for PTEN
 412 expression silencing. Methylation of PTEN was also shown to
 413 correlate with estrogen and progesterone receptor level which
 414 is highly related to the invasiveness of breast cancer, or even
 415 drug resistance [16, 32]. Recent research showed that PTEN
 416 expression is methylation-dependent and is preferentially
 417 methylated by DNMT3A [33]. By depleting DNMT3A,
 418 PTEN expression could be resumed due to demethylation of
 419 the CpG islands in the promoter region. Therefore, depletion
 420 of DNMT3A could exhibit antiproliferative effects.

421 Being the preferential targets of DNMT3A [33], changes in
 422 methylation level of PTEN genes depend on the DNMT3A
 423 level inside the cells. Our results showed for the first time that in
 424 restoration of miR-143, the methylation status of PTEN has
 425 been changed. With lowered expression of DNMT3A after
 426 miR-143 transfection, percentage of hypermethylated DNA in
 427 PTEN promoter drastically reduced, while that of unmethylated
 428 DNA increased. This clearly showed that miR-143 indirectly
 429 control the PTEN expression level through DNMT3A.

430 The tumor necrosis factor receptor superfamily member
 431 10C (TNFRSF10C) located on 8p22-p21 encodes a protein
 432 in the TNF receptor superfamily [34]. It has an extracellular
 433 TNF-related apoptosis-inducing ligand (TRAIL)-binding
 434 domain and a transmembrane domain, but lacks cytoplasmic
 435 death domain. This antagonistic receptor protects cell from
 436 TRAIL-induced apoptosis. Deletion of TNFRSF10C locus
 437 has been reported in lung cancer [35] and prostate cancer
 438 [36], while methylation of TNFRSF10C has been reported
 439 in lung cancer, pancreatic cancer, and breast cancer [37]. A
 440 recent study reported that the higher frequency of
 441 TNFRSF10C methylation resulted in tumor cell growth,
 442 suggesting a tumor-suppressive role in carcinogenesis [38].
 443 In our study, we are the first to find a decrease in
 444 hypermethylation of TNFRSF10C after being transfected
 445 with miR-143 mimic. This may suggest a potential role of
 446 DNMT3A in the methylation of TNFRSF10C expression.

In conclusion, we found that miR-143 was frequently
 downregulated in breast cancer, which might be a potential
 tumor suppressor. The direct targeting of miR-143 on
 DNMT3A suggested for the first time that miR-143 took part
 in the regulation of DNA methylation and caused PTEN and
 TNFRSF10C methylation. These novel findings provided a
 new insight into the relationship of miRNA and methylation,
 which may provide a new direction for the development of
 miRNA-based target treatment.

Conflicts of interest None

References

1. Parkin DM, Bray F, Ferlay J, Pisani P. Global cancer statistics, 2002. *CA Cancer J Clin.* 2005;55:74–108. 460
 2. Duffy MJ. Ca 15-3 and related mucins as circulating markers in breast cancer. *Ann Clin Biochem.* 1999;36:579–86. 461
 3. Calin GA, Croce CM. MicroRNA signatures in human cancers. *Nat Rev Cancer.* 2006;6:857–66. 462
 4. He L, He X, Lim LP, de Stanchina E, Xuan Z, Liang Y, et al. A microRNA component of the p53 tumour suppressor network. *Nature.* 2007;447:1130–4. 463
 5. Lu J, Getz G, Miska EA, Alvarez-Saavedra E, Lamb J, Peck D, et al. MicroRNA expression profiles classify human cancers. *Nature.* 2005;435:834–8. 464
 6. Calin GA, Ferracin M, Cimmino A, Di Leva G, Shimizu M, Wojcik SE, et al. A microRNA signature associated with prognosis and progression in chronic lymphocytic leukemia. *N Engl J Med.* 2005;353:1793–801. 465
 7. Schetter AJ, Leung SY, Sohn JJ, Zanetti KA, Bowman ED, Yanaihara N, et al. MicroRNA expression profiles associated with prognosis and therapeutic outcome in colon adenocarcinoma. *JAMA.* 2008;299:425–36. 466
 8. Blenkiron C, Goldstein LD, Thorne NP, Spiteri I, Chin SF, Dunning MJ, et al. MicroRNA expression profiling of human breast cancer identifies new markers of tumor subtype. *Genome Biol.* 2007;8:R214. 467
 9. van't Veer LJ, Dai HY, van de Vijver MJ, He YD, Hart AA, Mao M, et al. Gene expression profiling predicts clinical outcome of breast cancer. *Nature.* 2002;415:530–6. 468
 10. Lewis BP, Burge CB, Bartel DP. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell.* 2005;120:15–20. 469
 11. Ng EK, Tsang WP, Ng SS, Jin HC, Yu J, Li JJ, et al. MicroRNA-143 targets DNA methyltransferases 3a in colorectal cancer. *Br J Cancer.* 2009;101:699–706. 470
 12. Lin TX, Dong W, Huang J, Pan Q, Fan X, Zhang C, et al. MicroRNA-143 as a tumor suppressor for bladder cancer. *J Urol.* 2009;181:1372–80. 471
 13. Gao Q, Steine EJ, Barrasa MI, Hockemeyer D, Pawlak M, Fu D, et al. Deletion of the de novo DNA methyltransferase Dnmt3a promotes lung tumor progression. *Proc Natl Acad Sci U S A.* 2011;108:18061–6. 472
 14. Maehama T, Dixon JE. PTEN: a tumour suppressor that functions as a phospholipid phosphatase. *Trends Cell Biol.* 1999;9:125–8. 473
 15. Bose S, Crane A, Hibshoosh H, Mansukhani M, Sandweis L, Parsons R. Reduced expression of PTEN correlates with breast cancer progression. *Hum Pathol.* 2002;33:405–9. 474
 16. Phuong NT, Kim SK, Lim SC, Kim HS, Kim TH, Lee KY, et al. Role of PTEN promoter methylation in tamoxifen-resistant breast cancer cells. *Breast Cancer Res Treat.* 2011;130:73–83. 475

507 17. Ng EKO, Chong WWS, Jin H, Lam EK, Shin VY, Yu J, et al. 539
 508 Differential expression of microRNAs in plasma of patients with 540
 509 colorectal cancer: a potential marker for colorectal cancer screening. 541
 510 Gut. 2009;58:1375–81. 542
 511 18. Lujambio A, Calin GA, Villanueva A, Ropero S, Sánchez-Céspedes M, 543
 512 Blanco D, et al. A microRNA DNA methylation signature for human 544
 513 cancer metastasis. Proc Natl Acad Sci U S A. 2008;105:13556–61. 545
 514 19. Sharma S, Kelly TK, Jones PA. Epigenetics in cancer. 546
 515 Carcinogenesis. 2010;31:27–36. 547
 516 20. Chen X, Guo X, Zhang H, Xiang Y, Chen J, Yin Y, et al. Role of miR- 548
 517 143 targeting KRAS in colorectal tumorigenesis. Oncogene. 549
 518 2009;28:1385–92. 550
 519 21. Clape C, Fritz V, Henriquet C, Apparailly F, Fernandez PL, Iborra F, 551
 520 et al. MiR-143 interferes with ERK5 signaling, and abrogates 552
 521 prostate cancer progression in mice. PLoS One. 2009;4:e7542. 553
 522 22. Sandhu SK, Croce CM, Garzon R. Micro-RNA expression and function 554
 523 in lymphomas. Adv Hematol. 2011. doi:10.1155/2011/347137. 555
 524 23. Dhillon AS, Hagan S, Rath O, Kolch W. MAP kinase signalling 556
 525 pathways in cancer. Oncogene. 2007;26:3279–90. 557
 526 24. Robertson KD, Uzvolgyi E, Liang G, Talmadge C, Sumegi J, 558
 527 Gonzales FA, et al. The human DNA methyltransferases (DNMTs) 559
 528 1, 3a and 3b: coordinate mRNA expression in normal tissues and 560
 529 overexpression in tumors. Nucleic Acids Res. 1999;27:2291–8. 561
 530 25. Okano M, Bell DW, Haber DA, Li E. DNA methyltransferases 562
 531 Dnmt3a and Dnmt3b are essential for de novo methylation and 563
 532 mammalian development. Cell. 1999;99:247–57. 564
 533 26. Liang G, Chan MF, Tomigahara Y, Tsai YC, Gonzales FA, Li E, et al. 565
 534 Cooperativity between DNA methyltransferases in the maintenance 566
 535 methylation of repetitive elements. Mol Cell Biol. 2002;22:480–91. 567
 536 27. Chen RZ, Pettersson U, Beard C, Jackson-Grusby L, Jaenisch R. 568
 537 DNA hypomethylation leads to elevated mutation rates. Nature. 569
 538 1998;395:89–93. 570
 571 28. Baylin SB. DNA methylation and gene silencing in cancer. Nat Clin 539
 Pract Oncol. 2005;2 Suppl 1:S4–11. 540
 29. Baranwal S, Alahari SK. MiRNA control of tumor cell invasion and 541
 metastasis. Int J Cancer. 2010;126:1283–90. 542
 30. Chalhoub N, Baker SJ. PTEN and the PI3-kinase pathway in cancer. 543
 Annu Rev Pathol. 2009;4:127–50. 544
 31. Rhei E, Kang L, Bogomolnii F, Federici MG, Borgen PI, Boyd J. 545
 Mutation analysis of the putative tumor suppressor gene PTEN/ 546
 MMAC1 in primary breast carcinomas. Cancer Res. 1997;57: 547
 3657–9. 548
 32. Khan S, Kumagai T, Vora J, Bose N, Sehgal I, Koeffler PH, et al. 549
 PTEN promoter is methylated in a proportion of invasive breast 550
 cancers. Int J Cancer. 2004;112:407–10. 551
 33. Chik F, Szyf M. Effects of specific DNMT gene depletion on cancer 552
 cell transformation and breast cancer cell invasion; toward selective 553
 DNMT inhibitors. Carcinogenesis. 2011;32:224–32. 554
 34. Ashkenazi A. Targeting death and decoy receptors of the tumour- 555
 necrosis factor superfamily. Nat Rev Cancer. 2002;2:420–30. 556
 35. Tessema M, Yu YY, Stidley CA, Machida EO, Schuebel KE, Baylin 557
 SB, et al. Concomitant promoter methylation of multiple genes in 558
 lung adenocarcinomas from current, former and never smokers. 559
 Carcinogenesis. 2009;30:1132–8. 560
 36. Cheng Y, Kim JW, Liu W, Dunn TA, Luo J, Loza MJ, et al. Genetic 561
 and epigenetic inactivation of TNFRSF10C in human prostate 562
 cancer. Prostate. 2009;69:327–35. 563
 37. Shivapurkar N, Toyooka S, Toyooka KO, Reddy J, Miyajima K, 564
 Suzuki M, et al. Aberrant methylation of trail decoy receptor genes 565
 is frequent in multiple tumor types. Int J Cancer. 2004;109:786– 566
 92. 567
 38. Cai HH, Sun YM, Miao Y, Gao WT, Peng Q, Yao J, et al. Aberrant 568
 methylation frequency of tnfrsf10c promoter in pancreatic cancer cell 569
 lines. Hepato-Biliary-Pancreat Dis Int. 2011;10:95–100. 570