



Title	Purpurin Triggers Caspase-Independent Apoptosis in <i>Candida dubliniensis</i> Biofilms
Author(s)	Tsang, WK; Wong, PK; Yang, H; Li, NF
Citation	PLoS One, 2013, v. 8 n. 12, p. article no. e86032
Issued Date	2013
URL	http://hdl.handle.net/10722/193576
Rights	Creative Commons: Attribution 3.0 Hong Kong License

Purpurin Triggers Caspase-Independent Apoptosis in *Candida dubliniensis* Biofilms

Paul Wai-Kei Tsang*, Alan Pak-Kin Wong, Hai-Ping Yang, Ngai-For Li

Oral BioSciences, Faculty of Dentistry, the University of Hong Kong, HKSAR, China

Abstract

Candida dubliniensis is an important human fungal pathogen that causes oral infections in patients with AIDS and diabetes mellitus. However, *C. Dubliniensis* has been frequently reported in bloodstream infections in clinical settings. Like its phylogenetically related virulent species *C. albicans*, *C. Dubliniensis* is able to grow and switch between yeast form and filamentous form (hyphae) and develops biofilms on both abiotic and biotic surfaces. Biofilms are recalcitrant to antifungal therapies and *C. Dubliniensis* readily turns drug resistant upon repeated exposure. More than 80% of infections are associated with biofilms. Suppression of fungal biofilms may therefore represent a viable antifungal strategy with clinical relevance. Here, we report that *C. dubliniensis* biofilms were inhibited by purpurin, a natural anthraquinone pigment isolated from madder root. Purpurin inhibited *C. dubliniensis* biofilm formation in a concentration-dependent manner; while mature biofilms were less susceptible to purpurin. Scanning electron microscopy (SEM) analysis revealed scanty structure consisting of yeast cells in purpurin-treated *C. dubliniensis* biofilms. We sought to delineate the mechanisms of the anti-biofilm activity of purpurin on *C. Dubliniensis*. Intracellular ROS levels were significantly elevated in fungal biofilms and depolarization of MMP was evident upon purpurin treatment in a concentration-dependent manner. DNA degradation was evident. However, no activated metacaspase could be detected. Together, purpurin triggered metacaspase-independent apoptosis in *C. dubliniensis* biofilms.

Citation: Tsang PW-K, Wong AP-K, Yang H-P, Li N-F (2013) Purpurin Triggers Caspase-Independent Apoptosis in *Candida dubliniensis* Biofilms. PLoS ONE 8(12): e86032. doi:10.1371/journal.pone.0086032

Editor: Joy Sturtevant, Louisiana State University, United States of America

Received: October 24, 2013; **Accepted:** December 9, 2013; **Published:** December 23, 2013

Copyright: © 2013 Tsang et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by Research Fund for the Control of Infectious Diseases, the Food and Health Bureau of the Government of the HKSAR (Project No. 11100992 to PWKT); Health and Medical Research Fund, the Food and Health Bureau of the Government of the HKSAR (Project No. 12111292 to PWKT); Seed Funding Programme for Basic Research from The University of Hong Kong (Project No. 201211159011 to PWKT). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

* E-mail: pwktsang@hku.hk

Introduction

Invasive fungal infections have long been one of the most important medical problems in humans [1-3]. *Candida* fungi, among others, are prevalent human fungal pathogens that cause both superficial and systemic diseases (candidiasis) in patients with impaired immunity. In severe cases, the mortality and morbidity range from 40-60% [4,5]. In fact, candidiasis is the fourth leading type of hospital-acquired infections in clinical settings [6,7]. Although *C. albicans* is the major causative agent of candidiasis, recent epidemiological and clinical studies have shown an escalating number of bloodstream infections caused by non-*albicans Candida* species, which accounted for 36-63% of candidemia [8-11].

C. dubliniensis is now firmly recognized as an emerging and medically-relevant opportunistic human fungal pathogen, especially in the oral cavity of patients with AIDS and diabetes mellitus. Epidemiological studies indicated a worldwide spread of *C. dubliniensis*-related infections [12,13]. *C. dubliniensis* has

been isolated from other body sites including respiratory tract and blood [14,15], with up to 7% of candidemia caused by this pathogenic fungus [16]. In addition, azole-resistant *C. dubliniensis* isolates have been frequently reported in antifungal interventions, in particular to repeated and lengthy treatments [17,18], suggesting a dire need for novel approaches and strategies in treating this notorious human fungal pathogen.

In the course of our continuing efforts to characterize small molecules with novel antifungal activity, we have demonstrated the potent *in vitro* antifungal activity of purpurin, a natural red anthraquinone pigment in madder roots (*Rubia tinctorum* L.), against a panel of six pathogenic *Candida* species [19]. In particular, purpurin was found inhibitory to *C. albicans* biofilm development by downregulation of the expression of hypha-specific genes and the central morphogenetic regulator Ras1p [20]. *C. dubliniensis* is a close relative to *C. albicans* as they share >80% identity in genome sequence [21]. *C. dubliniensis* and *C. albicans* are the only *Candida* species that can form

true hyphae [22], and *C. dubliniensis* biofilms have been documented in bloodstream infections and on the surfaces of different biomaterials in the presence of saliva and serum [23-25]. *Candida* biofilms are heterogeneous sessile communities of yeast and hyphal cells in extracellular matrix, and are highly resistant to antifungal chemotherapy [26,27]. It has been estimated that 80% of microbial infections are biofilm-associated [28]. Thus, suppression of biofilms could be an effective measure to tackle with *Candida* virulence and pathobiology. The present study was designed to investigate the *in vitro* effect of purpurin on *C. dubliniensis* biofilms, with special attention on its correlation to cell demise.

Results and Discussion

The main etiopathologic role of biofilms in pathogenic *Candida* fungi is its ability to maintain a community of invasive fungal population that serves as a reservoir for recurrent host infection and disease dissemination [29]. Similar to its phylogenetically closely related species *C. albicans*, *C. dubliniensis* forms true hyphae and biofilms on both abiotic and biotic surfaces [24]. The increasing clinical prevalence of *C. dubliniensis* as an emerging human fungal pathogen is reflected by an astonishing number of reports in detecting bloodstream isolates and medical devices such as catheters and implanted materials [17,18,30].

Our laboratory recently deciphered the mechanisms of action of purpurin against *Candida* fungi through perturbation of mitochondrial homeostasis and initiation of apoptosis [19]. More experiments indicated that purpurin also interferes with *C. albicans* biofilm development at sub-MIC levels (3 $\mu\text{g/ml}$) [20]. In the present study, we explored the antifungal activity of purpurin on *C. dubliniensis* biofilms and evaluated several biochemical hallmarks of cell demise (apoptosis), an area that is still relatively unexplored in this *Candida* species. The antifungal activity of purpurin on *C. dubliniensis* biofilms was evaluated semi-quantitatively by using 3-bis(2-methoxy-4-nitro-5-sulfo-phenyl)-2H-tetrazolium-5-carboxanilide (XTT) reduction assay. The inhibitory effect was concentration-dependent, as revealed by a progressive reduction in cell viability with increasing concentrations of purpurin (Figure 1A). The metabolic activity of *C. dubliniensis* during biofilm formation was reduced by ~45% at 1 $\mu\text{g/ml}$ and by ~65% at 3 $\mu\text{g/ml}$. Pre-formed (mature) *C. dubliniensis* biofilms were less susceptible to purpurin; the cell viability was reduced by ~17% at 1 $\mu\text{g/ml}$ and by ~40% at 3 $\mu\text{g/ml}$ (Figure 1B). Scanning electron microscopy (SEM) images showed that purpurin inhibited biofilm development in *C. dubliniensis*, a scanty architecture with reduced hyphal growth was evident in purpurin-treated biofilms (Figure 2A-D). Considering the differences in phenotypic properties of *C. dubliniensis* to its sibling *C. albicans*, our data demonstrated that *C. dubliniensis* is more vulnerable as lower concentration of purpurin (1 $\mu\text{g/ml}$) was sufficient to abrogate biofilm development. As in other biofilm-forming microbes, the structurally well-organized mature *C. dubliniensis* biofilms were more resistant to purpurin, presumably due to poor penetration.

The *in vitro* effect of purpurin on *C. dubliniensis* biofilms prompted us to evaluate the presence of apoptotic features, namely reactive oxygen species (ROS), mitochondrial membrane potential (MMP), and DNA degradation, upon exposure to purpurin. Apoptosis is a highly sophisticated cellular process that leads to cell death in multicellular organisms and is crucial for normal growth, development, and cell maintenance. *Candida* fungi exhibit biochemical hallmarks typical to mammalian apoptosis in response to diverse environmental cues [31-36], in which mitochondria play a pivotal role in controlling both caspase-dependent and caspase-independent cell death pathways. Mitochondrial dysfunction is regarded as the onset of apoptosis [37-40]. Moreover, it has been shown that activation of *C. albicans* metacaspase is associated with elevation of intracellular ROS levels [41].

Therefore, we sought to evaluate the effect of purpurin on mitochondrial homeostasis. Purpurin increased the intracellular ROS levels in *C. dubliniensis* biofilms in a concentration-dependent manner, as revealed by a progressive elevation of fluorescence intensity. At 1 $\mu\text{g/ml}$, the endogenous ROS levels were increased by ~11%, and by ~27% at higher concentration (10 $\mu\text{g/ml}$) (Figure 3). Immense elevation of intracellular ROS levels is regarded as an onset of apoptosis, which is followed by depolarization of MMP [42]. To this end, we used the lipophilic fluorescent dye 5,5',6,6'-tetrachloro-1,1',3,3'-tetraethyl-imidacarbocyanine iodide (JC-1) to measure MMP in purpurin-treated *C. dubliniensis* biofilms. Intact MMP allows the cationic dye to enter the mitochondrial matrix and form J-aggregates (red fluorescence) beyond critical concentration. On the contrary, cells with depolarized MMP fluoresce green due to monomeric JC-1 [19]. The ratio of red fluorescence (FL-2) to green fluorescence (FL-1) represents the change of MMP in cell population. As expected, purpurin elicited a concentration-dependent depolarization of MMP in *C. dubliniensis* biofilms. At 1 $\mu\text{g/ml}$, MMP was decreased by ~12% and by 57% at 10 $\mu\text{g/ml}$ (Figure 4).

Collapse of MMP triggers a series of downstream cellular events that lead to disorganization of DNA and therefore cell death [43,44]. We assessed the effect of purpurin on DNA integrity in *C. dubliniensis* biofilms by using the TUNEL assay. DNA degradation results in a multitude of 3'-OH termini of DNA ends that can be labelled by fluorescent-tagged deoxyuridine triphosphate nucleotides using terminal deoxynucleotidyl transferase-mediated dUTP nick end labelling (TUNEL) assay [45]. Untreated fungal cells did not emit green fluorescence; while TUNEL-positive cells (green fluorescence) were detected upon purpurin treatment in *C. dubliniensis* biofilms (Figure 5A-D).

Activation of caspase is a distinct biochemical marker of apoptosis. Although no *bona fide* caspase is found in fungi, a distant relative called metacaspase has been characterized in *Saccharomyces cerevisiae* and *C. albicans* [41,46]. We therefore examined the presence of activated metacaspases in purpurin-treated *C. dubliniensis* biofilms by using FITC-VAD-FMK. FITC-VAD-FMK is a cell permeable caspase inhibitor that specifically binds to activated caspases [47]. However, no green fluorescence could be observed in purpurin-treated *C.*




Figure 1. Effect of purpurin on *Candida dubliniensis* biofilms. A) Biofilm formation of *C. dubliniensis* MYA-646. B) Mature biofilms. In both cases, biofilms were incubated with the indicated concentration of purpurin for 24 h at 37°C. The metabolic activity of the biofilms was assessed semi-quantitatively using XTT reduction assay. The activity of samples without purpurin treatment (i.e. DMSO only) (0 µg/ml) was taken as 100%. Results shown were the average of three independent experiments \pm SD. * p <0.05 when compared with the untreated controls.

doi: 10.1371/journal.pone.0086032.g001




Figure 2. Scanning electron microscopy images of *Candida dubliniensis* biofilms. Representative SEM images showing the topography of *C. dubliniensis* biofilms. Fungal biofilms were formed on plastic coverslips, processed, and coated with gold before viewing (magnification 500 \times). A) Biofilm formation in the absence of purpurin. B) Biofilm formation in the presence of purpurin (1 μ g/ml). C) Pre-formed biofilms in the absence of purpurin. D) Pre-formed biofilms in the presence of purpurin (1 μ g/ml).

doi: 10.1371/journal.pone.0086032.g002

dubliniensis biofilms (data not shown), suggesting an absence of activated metacaspases. Nevertheless, a putative metacaspase sequence (Cd36_85170) has been annotated in *C. dubliniensis* genome (<http://www.candidagenome.org/>), and we have detected the presence of activated metacaspase upon exposure of planktonic cells of *C. dubliniensis* to apoptotic levels of amphotericin B or acetic acid (P. W. K. Tsang, unpublished data). Functional significance of this uncharacterized open reading frame in apoptosis is in progress in our laboratory.

C. dubliniensis poses a serious health threat to humans and puts heavy economic burden on our society. Strategies that work by switching on endogenous cell death mechanisms specific to *C. dubliniensis* could be a novel and tangible antifungal approach against this emerging human fungal pathogen. Our findings clearly demonstrated that purpurin

triggers mitochondrial-mediated apoptotic pathway in *C. dubliniensis* biofilms without the involvement of activated metacaspases, suggesting that purpurin-mediated cell death in *C. dubliniensis* does not require the proteolytic activity of metacaspases. This is an entirely new antifungal action mechanism which is different from the standard antifungal agents in clinical settings: flucytosine inhibits DNA synthesis; polyenes cause membrane damage; azoles affect sterol metabolism; and echinocandins inhibit cell wall synthesis. Nevertheless, the usefulness of these antifungal agents has been hampered by severe side effects, poor therapeutic index, and the emergence of multidrug resistance [48,49]. Therefore, the unique antifungal action of purpurin on *C. dubliniensis* could be of particular clinical relevance in the design of more selective and effective therapeutic treatments. One aspect could be killing of *C. dubliniensis* biofilms via induction of the




Figure 3. Effect of purpurin on intracellular ROS levels. *C. dubliniensis* biofilms were treated with different concentrations of purpurin (from 1 µg/ml to 10 µg/ml), and intracellular ROS levels were determined using DCFDA. Fluorescence intensity was measured at 485 nm excitation and 535 nm emission. Results shown were the average of three independent experiments ± SD. * $p < 0.05$ when compared with the untreated controls.

doi: 10.1371/journal.pone.0086032.g003

endogenous mitochondrial-mediated apoptotic pathway. Another aspect could be a combined use of purpurin with the conventional antifungal agents to reduce potential side effects and the likelihood of acquired multidrug resistance.

In conclusion, the findings of the present study provide solid evidence that purpurin triggered apoptosis-like features in *C. dubliniensis* biofilms. As *C. dubliniensis* and *C. albicans* belong to the CTG clade, it is not surprising that they share common core pathways in metabolism. Nonetheless, a comparative analysis of calcineurin signalling pathways between *C. dubliniensis* and *C. albicans* indicated differential scenarios in pH homeostasis [50]. Another study revealed a rewiring of iron assimilation gene expression in *C. dubliniensis* [51]. Thus, comparing metabolic pathways between these two species not only helps gain further insights into their evolutionary divergence, but also alludes to a functional characterization of core machineries responsible for their pathogenicity. For instance, a better understanding of the cell death mechanisms can be beneficial to the design of innovative antifungal strategies that target the core components specifically.

Materials and Methods

Strains, cultivation and chemicals

The reference laboratory strain *C. dubliniensis* MYA-646 was used throughout the study, and was routinely cultured in YPD agar (10 g/l yeast extract, 20 g/l peptone, 20 g/l dextrose, 20 g/l agar) at 30°C. To prepare a standard cell suspension, a single colony was inoculated into YNB medium (6.7 g/l yeast nitrogen base w/o amino acids, 20 g/l dextrose) and incubated for 18 h at 30°C with agitation (200 rpm). Fungal cells were harvested by centrifugation, washed twice with phosphate-buffered saline (PBS) (pH 7.2), and resuspended at 1×10^7 cells/ml. Purpurin (purity >99%) was purchased from TimTec Inc. (Newark, DE, USA). Stock solution of purpurin was prepared in distilled dimethyl sulphoxide (DMSO) and kept at -20°C until use. The final concentration of DMSO was 1% in all assays. Other chemicals were obtained from commercial suppliers with the highest grade available.

Effect of Purpurin on *C. dubliniensis* Biofilm Formation and Pre-Formed (Mature) Biofilms

Fungal biofilms were prepared as described [52] on commercially available, pre-sterilized, flat-bottomed 96-well




Figure 4. Depolarization of MMP in purpurin-treated *Candida dubliniensis* biofilms. Purpurin-treated *C. dubliniensis* biofilms were incubated with JC-1. MMP was measured by flow cytometer at FL-1 (525 nm) and FL-2 (595 nm), and expressed as ratio of FL-2/FL-1. Results shown were the average of three independent experiments \pm SD. * $p < 0.05$ when compared with the untreated controls.

doi: 10.1371/journal.pone.0086032.g004

polystyrene microtitre plates (Iwaki). Standard cell suspension of *C. dubliniensis* (100 μ l) was transferred into the wells and incubated for 1.5 h at 37°C with agitation (80 rpm). After the adhesion phase, the liquid was aspirated and the wells were washed twice with PBS to remove non-adherent cells. Fresh YNB medium (200 μ l) containing purpurin (1 μ g/ml or 3 μ g/ml) was added to each well and the plates were incubated for an additional 24 h at 37°C.

To investigate the effect of purpurin on pre-formed biofilms, *C. dubliniensis* biofilms were prepared for 24 h at 37°C as described above. The wells were washed twice with PBS and fresh YNB medium (200 μ l) containing purpurin (1 μ g/ml or 3 μ g/ml) was added and the plates were incubated for an additional 24 h at 37°C. YNB medium with 1% DMSO was included in control wells. The metabolic activity of the *C. dubliniensis* biofilms, representing viability of the fungal cells, was determined by using a standard XTT reduction assay that measures mitochondrial dehydrogenase activity [52]. XTT solution (1 g/l; 40 μ l) was mixed with freshly prepared menadione solution (0.4 mM; 2 μ l) at 20:1 (v/v) immediately prior to the assay. Thereafter, PBS (158 μ l) was mixed with XTT-menadione solution (42 μ l) and transferred to each well containing pre-washed biofilms, and incubated in the dark for 3

h at 37°C. Colour changes were measured at 492 nm with a microtitre plate reader (SpectraMax 340 tunable microplate reader; Molecular Devices).

Measurement of endogenous ROS levels

The effect of purpurin on intracellular ROS levels in *C. dubliniensis* biofilms was examined by using the fluorescent dye 2',7'-dichlorofluorescein diacetate (DCFDA) (Molecular Probes, CA, USA) [45]. Briefly, after treatment with different concentrations of purpurin (from 1 μ g/ml to 10 μ g/ml), the biofilms were washed twice with PBS, added with DCFDA (final concentration: 20 μ M) and incubated for 1 h at 37°C. Fluorescence intensity was measured in a fluorescence plate reader (TECAN Polarion, Tecan UK Ltd, Theale, UK) with a 485 nm excitation and 535 nm emission.

Measurement of MMP

The effect of purpurin on MMP of *C. dubliniensis* biofilms was analyzed as described by using JC-1 [19]. Briefly, after treatment with different concentrations of purpurin (from 1 μ g/ml to 10 μ g/ml), the biofilms were washed thrice with PBS, followed by incubation with JC-1 (0.25 μ M) (Molecular Probes, CA) for 15 min at 35°C. Fluorescence intensities at FL-1




Figure 5. Effect of purpurin on DNA integrity (TUNEL assay). Representative fluorescent images showing the occurrence of DNA degradation in purpurin-treated *C. dubliniensis* biofilms captured by confocal microscope. A) Biofilms in the absence of purpurin (Bright Field). B) Biofilms in the absence of purpurin (FITC). C) Biofilms in the presence of purpurin (1 µg/ml) (Bright Field). D) Biofilms in the presence of purpurin (1 µg/ml) (FITC). Cells with damaged DNA emitted intense green fluorescence. The bar represents 50 µm.

doi: 10.1371/journal.pone.0086032.g005

(unhealthy cells emit green fluorescence, 525 nm) and FL-2 (healthy cells emit red fluorescence, 595 nm) were recorded by

a Beckman-Coulter flow cytometer, and the results were expressed as a ratio of the mean values at FL-2 and FL-1.

Detection of activated metacaspases

The presence of activated metacaspases in *C. dubliniensis* biofilms was evaluated by using CaspACE™ FITC-VAD-FMK In Situ Marker (Promega, Madison, WI, USA). Briefly, purpurin-treated *C. dubliniensis* biofilms were washed thrice with PBS and incubated with CaspACE™ FITC-VAD-FMK In Situ Marker (10 µM) for 20 min at 37°C in the dark. The cells were washed twice and resuspended in PBS. Aliquots of fungal cells were added to poly-L-lysine-coated glass slides and analyzed by fluorescence microscopy using a confocal laser scanning microscope (Fluoview FV 1000, Olympus, Tokyo, Japan). Fungal cells containing activated metacaspases fluoresced green.

TUNEL assay

DNA fragmentation in *C. dubliniensis* biofilms after treating with purpurin was detected by using the *In Situ* Cell Death Detection Kit (fluorescein) (Roche, Penzberg, Germany) [31]. Briefly, after incubation with purpurin, the biofilms were washed thrice with PBS and fixed with 2% paraformaldehyde for 1 h at 20°C. The biofilms were rinsed thrice with PBS, and then incubated with permeabilization solution (0.1% Triton X-100, 0.1% sodium citrate) on ice for 2 min. The biofilms were rinsed twice with PBS and labelled with 50 µl TUNEL reaction mixture for 1 h at 37°C in a humidified incubator in the dark. The cells were mounted under glass coverslip and analyzed by fluorescence microscopy using a confocal laser scanning microscope.

SEM analysis

C. dubliniensis biofilms were prepared on custom-made, tissue culture-treated, polystyrene coverslips as described

[20,52]. Thereafter, the coverslips were washed twice with PBS and placed in 1% osmium tetroxide for 1 h. Samples were subsequently washed with distilled water, dehydrated in a series of ethanol solutions (70% for 10 min, 95% for 10 min and 100% for 20 min), and air-dried overnight in a desiccator prior to sputter coating with gold (JFC1 100; JEOL). The surface topographies of the *C. dubliniensis* biofilms were viewed with a scanning electron microscope (Philip XL30CP).

Statistical analysis

All experiments were performed in triplicate in three different occasions and all data were expressed as the mean values with the corresponding standard deviation (SD). Statistical significance between treated and control groups was assessed by Mann-Whitney U test. A *p*-value of <0.05 was considered statistically significant.

Acknowledgements

We thank Mr. Raymond Tong, Mr. Geoffrey Ng and Ms. Becky Cheung for their excellent technical assistance.

Author Contributions

Conceived and designed the experiments: PWKT. Performed the experiments: PWKT APKW HPY NFL. Analyzed the data: PWKT APKW. Contributed reagents/materials/analysis tools: PWKT. Wrote the manuscript: PWKT.

References

- Parize P, Rammaert B, Lortholary O (2012) Emerging invasive fungal diseases in transplantation. *Curr Infect Dis Rep* 14: 668-675. doi: 10.1007/s11908-012-0296-y. PubMed: 23065419.
- Garcia-Vidal C, Viasus D, Carratalà J (2013) Pathogenesis of invasive fungal infections. *Curr Opin Infect Dis* 26: 270-276. doi:10.1097/QCO.0b013e32835fb920. PubMed: 23449139.
- Hundalani S, Pammi M (2013) Invasive fungal infections in newborns and current management strategies. *Expert Rev Anti Infect Ther* 11: 709-721. doi:10.1586/14787210.2013.811925. PubMed: 23829639.
- Corner BE, Magee PT (1997) *Candida* pathogenicity: unravelling the threads of infection. *Curr Biol* 7: R691-R694. doi:10.1016/S0960-9822(06)00357-5. PubMed: 9382797.
- Wenzel RP, Gennings C (2005) Bloodstream infections due to *Candida* species in the intensive care unit: identifying especially high-risk patients to determine prevention strategies. *Clin Infect Dis* 41: S389-S393. doi:10.1086/430923. PubMed: 16108005.
- Bustamante CI (2005) Treatment of *Candida* infection: a view from the trenches! *Curr Opin Infect Dis* 18: 490-495. doi:10.1097/01.qco.0000191516.43792.61. PubMed: 16258321.
- Miceli MH, Díaz JA, Lee SA (2011) Emerging opportunistic yeast infections. *Lancet Infect Dis* 11: 142-151. doi:10.1016/S1473-3099(10)70218-8. PubMed: 21272794.
- Bassetti M, Righi E, Costa A, Fasce R, Molinari MP et al. (2006) Epidemiological trends in nosocomial candidemia in intensive care. *BMC Infect Dis* 6: 21. doi:10.1186/1471-2334-6-21. PubMed: 16472387.
- Colombo AL, Nucci M, Salomão R, Branchini ML, Richtmann R et al. (1999) High rate of non-albicans candidemia in Brazilian tertiary care hospitals. *Diagn Microbiol Infect Dis* 34: 281-286. doi:10.1016/S0732-8893(99)00042-5. PubMed: 10459478.
- Dimopoulos G, Ntziora F, Rachiotis G, Armaganidis A, Falagas ME (2008) *Candida albicans* versus non-*albicans* intensive care unit-acquired bloodstream infections: differences in risk factors and outcome. *Anesth Analg* 106: 523-529. doi:10.1213/ane.0b013e3181607262. PubMed: 18227310.
- Chow JK, Golan Y, Ruthazer R, Karchmer AW, Carmeli Y et al. (2008) Factors associated with candidemia caused by non-*albicans Candida* species versus *Candida albicans* in the intensive care unit. *Clin Infect Dis* 46: 1206-1213. doi:10.1086/529435. PubMed: 18444857.
- Sullivan DJ, Moran GP, Coleman DC (2005) *Candida dubliniensis*: ten years on. *FEMS Microbiol Lett* 253: 9-17. doi:10.1016/j.femsle.2005.09.015. PubMed: 16213674.
- Willis AM, Coulter WA, Sullivan DJ, Coleman DC, Hayes JR et al. (2000) Isolation of *Candida dubliniensis* from insulin-using diabetes mellitus patients. *J Oral Pathol Med* 29: 86-90. doi:10.1034/j.1600-0714.2000.290206.x. PubMed: 10718404.
- Polacheck I, Strahilevitz J, Sullivan D, Donnelly S, Salkin IF et al. (2000) Recovery of *Candida dubliniensis* from non-human immunodeficiency virus-infected patients in Israel. *J Clin Microbiol* 38: 170-174. PubMed: 10618082.
- Jabra-Rizk MA, Johnson JK, Forrest G, Mankes K, Meiller TF et al. (2005) Prevalence of *Candida dubliniensis* fungemia at a large teaching hospital. *Clin Infect Dis* 41: 1064-1067. doi:10.1086/432943. PubMed: 16142677.
- Sullivan DJ, Moran GP, Pinjon E, Al-Mosaid A, Stokes C et al. (2004) Comparison of the epidemiology, drug resistance mechanisms, and virulence of *Candida dubliniensis* and *Candida albicans*. *FEMS Yeast Res* 4: 369-376. doi:10.1016/S1567-1356(03)00240-X. PubMed: 14734017.
- Moran GP, Sullivan DJ, Henman MC, McCreary CE, Harrington BJ et al. (1997) Antifungal drug susceptibilities of oral *Candida dubliniensis*

- isolates from human immunodeficiency virus (HIV)-infected and non-infected subjects and generation of stable fluconazole-resistant derivatives *in vitro*. *Antimicrob Agents Chemother* 41: 617-623. PubMed: 9056003.
18. Fanci R (2009) Breakthrough *Candida dubliniensis* fungemia in an acute myeloid leukemia patient during voriconazole therapy successfully treated with caspofungin. *J Chemother* 21: 105-107. PubMed: 19297283.
 19. Kang K, Fong WP, Tsang PWK (2010) Novel antifungal activity of purpurin against *Candida* species *in vitro*. *Med Mycol* 48: 904-911. doi: 10.3109/13693781003739351. PubMed: 20392152.
 20. Tsang PWK, Bandara HMHN, Fong WP (2012) Purpurin suppresses *Candida albicans* biofilm formation and hyphal development. *PLOS ONE* 7: e50866. doi:10.1371/journal.pone.0050866. PubMed: 23226409.
 21. Jackson AP, Gamble JA, Yeomans T, Moran GP, Saunders D et al. (2009) Comparative genomics of the fungal pathogens *Candida dubliniensis* and *Candida albicans*. *Genome Res* 19: 2231-2244. doi: 10.1101/gr.097501.109. PubMed: 19745113.
 22. Moran GP, Coleman DC, Sullivan DJ (2012) *Candida albicans* versus *Candida dubliniensis*: why is *C. albicans* more pathogenic? *Int J Microbiol* 2012: 205921
 23. Pannanusorn S, Fernandez V, Römling U (2013) Prevalence of biofilm formation in clinical isolates of *Candida* species causing bloodstream infection. *Mycoses* 56: 264-272. doi:10.1111/myc.12014. PubMed: 23113805.
 24. Ramage G, Vande Walle K, Wickes BL, López-Ribot JL (2001) Biofilm formation by *Candida dubliniensis*. *J Clin Microbiol* 39: 3234-3240. doi: 10.1128/JCM.39.9.3234-3240.2001. PubMed: 11526156.
 25. Ásmundsdóttir LR, Erlendsdóttir H, Agnarsson BA, Gottfredsson M (2009) The important of strain variation in virulence of *Candida dubliniensis* and *Candida albicans*: results of a blinded histopathological study of invasive candidiasis. *Clin Microbiol Infect* 15: 576-585. doi:10.1111/j.1469-0691.2009.02840.x. PubMed: 19604278.
 26. Lorenz MC, Bender JA, Fink GR (2004) Transcriptional response of *Candida albicans* upon internalization by macrophages. *Eukaryot Cell* 3: 1076-1087. doi:10.1128/EC.3.5.1076-1087.2004. PubMed: 15470236.
 27. ten Cate JM, Klis FM, Pereira-Cenci T, Crielaard W, de Groot PW (2009) Molecular and cellular mechanisms that lead to *Candida* biofilm formation. *J Dent Res* 88: 105-115. doi:10.1177/0022034508329273. PubMed: 19278980.
 28. Dongari-Bagtzoglou A (2008) Pathogenesis of mucosal biofilms infections: challenges and progress. *Expert Rev Anti Infect Ther* 6: 201-208. doi:10.1586/14787210.6.2.201. PubMed: 18380602.
 29. Ruhnke M (2006) Epidemiology of *Candida albicans* infections and role of non-*Candida albicans* yeasts. *Curr Drug Targets* 7: 495-504. doi: 10.2174/138945006776359421. PubMed: 16611037.
 30. Khan Z, Ahmad S, Joseph L, Chandy R (2012) *Candida dubliniensis*: an appraisal of its clinical significance as a bloodstream pathogen. *PLOS ONE* 7: e32952. doi:10.1371/journal.pone.0032952. PubMed: 22396802.
 31. Shirtliff ME, Krom BP, Meijering RA, Peters BM, Zhu J et al. (2009) Farnesol-induced apoptosis in *Candida albicans*. *Antimicrob Agents Chemother* 53: 2392-2401. doi:10.1128/AAC.01551-08. PubMed: 19364863.
 32. Phillips AJ, Sudbery I, Ramsdale M (2003) Apoptosis induced by environmental stresses and amphotericin B in *Candida albicans*. *Proc Natl Acad Sci U S A* 100: 14327-14332. doi:10.1073/pnas.2332326100. PubMed: 14623979.
 33. Andrés MT, Viejo-Díaz M, Fierro JF (2008) Human lactoferrin induces apoptosis-like cell death in *Candida albicans*: critical role of K⁺-channel-mediated K⁺ efflux. *Antimicrob Agents Chemother* 52: 4081-4088. doi: 10.1128/AAC.01597-07. PubMed: 18710913.
 34. Al-Dhaheri RS, Douglas LJ (2010) Apoptosis in *Candida* biofilms exposed to amphotericin B. *J Med Microbiol* 59: 149-157. doi:10.1099/jmm.0.015784-0. PubMed: 19892857.
 35. Keyhani E, Khavari-Nejad S, Keyhani J, Attar F (2009) Acriflavine-mediated apoptosis and necrosis in yeast *Candida utilis*. *Ann N Y Acad Sci* 1171: 284-291. doi:10.1111/j.1749-6632.2009.04682.x. PubMed: 19723067.
 36. Kang K, Wong KS, Fong WP, Tsang PWK (2011) Metergoline-induced cell death in *Candida krusei*. *Fungal Biol* 115: 302-309. doi:10.1016/j.funbio.2011.01.001. PubMed: 21354537.
 37. Pradelli LA, Bénétteau M, Ricci JE (2010) Mitochondrial control of caspase-dependent and -independent cell death. *Cell Mol Life Sci* 67: 1589-1597. doi:10.1007/s00018-010-0285-y. PubMed: 20151314.
 38. Estaquier J, Vallette F, Vayssiere JL, Mignotte B (2012) The mitochondrial pathways of apoptosis. *Adv Exp Med Biol* 942: 157-183. doi:10.1007/978-94-007-2869-1_7. PubMed: 22399422.
 39. Borutaite V (2010) Mitochondria as decision-makers in cell death. *Environ Mol Mutagen* 51: 406-416. PubMed: 20209625.
 40. Karbowski M (2010) Mitochondria on guard: role of mitochondrial fusion and fission in the regulation of apoptosis. *Adv Exp Med Biol* 687: 131-142. doi:10.1007/978-1-4419-6706-0_8. PubMed: 20919642.
 41. Cao YY, Huang S, Dai B, Zhu Z, Lu H et al. (2009) *Candida albicans* cells lacking CaMCA1-encoded metacaspase show resistance to oxidative stress-induced death and change in energy metabolism. *Fungal Genet Biol* 46: 183-189. doi:10.1016/j.fgb.2008.11.001. PubMed: 19049890.
 42. Pozniakovskiy AI, Knorre DA, Markova OV, Hyman AA, Skulachev VP et al. (2005) Role of mitochondria in the pheromone- and amiodarone-induced programmed cell death of yeast. *J Cell Biol* 168: 257-267. doi: 10.1083/jcb.200408145. PubMed: 15657396.
 43. Madeo F, Herker E, Wissing S, Jungwirth H et al. (2004) Apoptosis in yeast. *Curr Opin Microbiol* 7: 655-660. doi:10.1016/j.mib.2004.10.012. PubMed: 15556039.
 44. Ramsdale M (2008) Programmed cell death in pathogenic fungi. *Biochim Biophys Acta* 1783: 1369-1380. doi:10.1016/j.bbamcr.2008.01.021. PubMed: 18294459.
 45. Kang K, Fong WP, Tsang PWK (2010) Antifungal activity of baicalein against *Candida krusei* does not involve apoptosis. *Mycopathologia* 170: 391-396. doi:10.1007/s11046-010-9341-2. PubMed: 20614252.
 46. Madeo F, Herker E, Maldener C, Wissing S, Lächelt S et al. (2002) A caspase-related protease regulates apoptosis in yeast. *Mol Cell* 9: 911-917. doi:10.1016/S1097-2765(02)00501-4. PubMed: 11983181.
 47. Barroso G, Taylor S, Morshedi M, Manzur F, Gaviño F et al. (2006) Mitochondrial membrane potential integrity and plasma membrane translocation of phosphatidylserine as early apoptotic markers: a comparison of two different sperm subpopulations. *Fertil Steril* 85: 149-154. doi:10.1016/j.fertnstert.2005.06.046. PubMed: 16412746.
 48. Odds FC, Brown AJ, Gow NA (2003) Antifungal agents: mechanism of action. *Trends Microbiol* 11: 272-279. doi:10.1016/S0966-842X(03)00117-3. PubMed: 12823944.
 49. Ghannoum MA, Rice LB (1999) Antifungal agents: mode of action, mechanisms of resistance, and correlation of these mechanisms with bacterial resistance. *Clin Microbiol Rev* 12: 501-517. PubMed: 10515900.
 50. Zhang J, Heitman J, Chen YL (2012) Comparative analysis of calcineurin signaling between *Candida dubliniensis* and *Candida albicans*. *Commun Integr Biol* 5: 122-126. doi:10.4161/cib.18833. PubMed: 22808313.
 51. Moran GP (2012) Transcript profiling reveals rewiring of iron assimilation gene expression in *Candida albicans* and *Candida dubliniensis*. *FEMS Yeast Res* 12: 918-923. doi:10.1111/j.1567-1364.2012.00841.x. PubMed: 22888912.
 52. Bandara HMHN, Lam OL, Watt RM, Jin LJ, Samaranyake LP (2010) Bacterial lipopolysaccharides variably modulate *in vitro* biofilm formation of *Candida* species. *J Med Microbiol* 59: 1225-1234. doi: 10.1099/jmm.0.021832-0. PubMed: 20576747.