



Title	Paradoxical Impact of Two Folate Receptors, FR and RFC, in Ovarian Cancer: Effect on Cell Proliferation, Invasion and Clinical Outcome
Author(s)	Siu, MKY; Kong, DSH; Chan, HY; Wong, ESY; Ip, PPC; Jiang, L; Ngan, HYS; Le, X-F; Cheung, ANY
Citation	PLoS ONE, 2012, v. 7 n. 11
Issued Date	2012
URL	http://hdl.handle.net/10722/196720
Rights	Creative Commons: Attribution 3.0 Hong Kong License

Paradoxical Impact of Two Folate Receptors, FR α and RFC, in Ovarian Cancer: Effect on Cell Proliferation, Invasion and Clinical Outcome

Michelle K. Y. Siu^{1*}, Daniel S. H. Kong¹, Hoi Yan Chan¹, Esther S. Y. Wong¹, Philip P. C. Ip¹, LiLi Jiang¹, Hextan Y. S. Ngan², Xiao-Feng Le³, Annie N. Y. Cheung^{1*}

1 Department of Pathology, University of Hong Kong, Hong Kong, Special Administrative Region of China, **2** Department of Obstetrics and Gynaecology, University of Hong Kong, Hong Kong, Special Administrative Region of China, **3** Department of Experimental Therapeutics, Division of Cancer Medicine, University of Texas M. D. Anderson Cancer Center, Houston, Texas, United States of America

Abstract

Despite being an essential vitamin, folate has been implicated to enhance tumor growth, as evidenced by reports on overexpression of folate receptor alpha (FR α) in carcinomas. The role of another folate transporter, reduced folate carrier (RFC), is largely unknown. This study investigated the roles of folate, FR α and RFC in ovarian cancers. We demonstrated FR α mRNA and protein overexpression and reduced RFC expression in association with FR α gene amplification and RFC promoter hypermethylation, respectively. FR α overexpression was associated with tumor progression while RFC expression incurred a favorable clinical outcome. Such reciprocal expression pattern was also observed in ovarian cancer cell lines. Folate was shown to promote cancer cell proliferation, migration and invasion *in vitro*, and down-regulate E-cadherin expression. This effect was blocked after either stable knockdown of FR α or ectopic overexpression of RFC. This hitherto unreported phenomenon suggests that, RFC can serve as a balancing partner of FR α and confer a protective effect in patients with high FR α -expressing ovarian carcinomas, as evidenced by their prolonged overall and disease-free survivals. In conclusion, we report on the paradoxical impact of FR α (putative oncogenic) and RFC (putative tumor suppressive) in human malignancies. FR α and RFC may potentially be explored as therapeutic target or prognostic marker respectively. We recommend caution and additional research on folate supplements in cancer patients.

Citation: Siu MKY, Kong DSH, Chan HY, Wong ESY, Ip PPC, et al. (2012) Paradoxical Impact of Two Folate Receptors, FR α and RFC, in Ovarian Cancer: Effect on Cell Proliferation, Invasion and Clinical Outcome. PLoS ONE 7(11): e47201. doi:10.1371/journal.pone.0047201

Editor: Ming Tat Ling, Queensland University of Technology, Australia

Received: June 1, 2012; **Accepted:** September 10, 2012; **Published:** November 7, 2012

Copyright: © 2012 Siu et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was supported by funding from the American Institute for Cancer Research Post-doctoral Award and the University of Hong Kong Seed Funding, and the Conference and Research Grant from the University of Hong Kong. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: anycheun@hkucc.hku.hk (ANYC); mkysiu@hkucc.hku.hk (MKYS)

Introduction

Ovarian carcinomas account for the highest mortality amongst all gynecological cancers in the world [1,2]. While the incidence of ovarian carcinomas varies between different ethnic groups, its incidence in Asian countries is on a rising trend [3,4]. The reasons for this remain largely unknown or controversial. Several lifestyle risk factors have been implicated. These include diet, obesity, fertility and parity statuses. On the other hand, it has become a general belief that high intake of micronutrients such as folate, vitamin C, vitamin E may protect against cancers [5]. As such, better understanding of the effects of nutritional elements on carcinogenesis is important to improve the strategies for cancer prevention and management.

Folate is a water-soluble B vitamin found in most vegetables. A high dietary folate intake has been reported to associate with a lower risk of developing ovarian cancers, in particular, those who consume alcohol [6,7,8,9]. It is closely related to its function on DNA synthesis and its involvement in the related methionine metabolic pathway essential for DNA methylation. Folate deficiency would therefore leads to DNA hypomethylation, altered gene expression and the misincorporation of uracil into DNA,

leading to chromosome damage, all of which, are key factors for carcinogenesis [10,11]. It would appear that folate is an important vitamin essential in normal functioning of cells, and to prevent the initiation of cancer. However, there is increasing evidence to show that folate may in fact enhance cancer progression in established carcinomas of colon and rectum, breast and prostate [12,13,14].

Folate uptake involves several transporters, such as folate receptors, and reduced folate carrier (RFC) [15,16]. Folate receptor alpha (FR α), a single chain glycosyl-phosphatidylinositol-anchored membrane protein, enhances folate uptake through endocytosis. Its overexpression has been reported in ovarian cancers, implying that it may promote tumor growth [17,18,19,20]. RFC is an ubiquitously expressed transporter for natural folates and classical antifolates, and can control folate uptake in a bi-directional manner [16]. Loss of RFC with subsequent effects of folate deficiency was found to promote cancer progression in colorectal cancer [16,21]. It would therefore be logical to assume that these two folate transporters, FR α and RFC, exert different effects in cancer progression.

Although overexpression of FR α in ovarian cancers has been established, the expression status and functional roles of RFC

remain largely unknown. In this study, we investigated the expression, genetic and epigenetic profiles of FR α and RFC in normal ovarian epithelium and ovarian cancer, and correlated with clinicopathological parameters. Their functional roles and possible downstream targets on cell proliferation, migration and invasion in relation to folate in ovarian cancer were also assessed. We endeavoured to better understand the roles of folate and its transporters in ovarian carcinogenesis, and explore the possible effects of folate intake in cancer patients.

Materials and Methods

Clinical Samples and cell lines

One hundred and fifty three formalin-fixed paraffin embedded samples of ovarian tumors, including 11 inclusion cysts/benign cystadenomas (22–63 years; mean age, 50 years), 19 borderline tumors (20–46 years; mean age, 30 years), 83 carcinomas (34 to 83 years; mean 51 years) of different histological subtypes and 44 corresponding metastatic foci (Table 1), were collected from the Department of Pathology, Queen Mary Hospital, the University of Hong Kong. All patients underwent surgery and 67 patients with ovarian cancers were also treated with chemotherapy including platinum/paclitaxel. The follow-up period ranged from five to 209 months (mean 63 months). Thirty three randomly selected clinical samples of ovarian tumors and their corresponding normal counterparts, including fallopian tubes and/or contralateral ovaries, with available frozen blocks were also retrieved. Informed consent was obtained by all patients and the use of these clinical samples was approved by Institutional Review Board of the University of Hong Kong/Hospital Authority Hong Kong West Cluster (HKU/HA HKW IRB)(Institutional Review Board number: UW10-129). Haematoxylin Eosin stained sections of the frozen blocks of each sample were reviewed by two of us (A.N.Y.C. and P.P.C.I.) to confirm the diagnosis and to ensure that more than 80% tumor cells were present in the tumor blocks.

Two immortalized ovarian epithelial cell lines, HOSE 6-3 and HOSE 17-1, and nine ovarian cancer cell lines, SKOV-3, OVCAR-3, OVCA 420, OVCA433, OC316, Dov13, ES-2, TOV21G, SW626 (ATCC; Manassas, VA) were cultured as previously described [22,23].

Real-time PCR (qPCR)

Total RNA from frozen clinical samples and cancer cell lines was extracted using Trizol reagent (Invitrogen). Genomic DNA contamination was removed by treating with DNase I (Invitrogen) treatment. 2.5 μ g total RNA was reverse transcribed by SuperScript Reverse Transcriptase (Invitrogen, San Diego, CA). Genomic DNA was extracted using phenol/chloroform (Invitrogen). qPCR was performed with ABI Prism 7700 sequence detection system (Applied Biosystems, Foster City, CA) as described [22,23,24]. Primer sequences for evaluating mRNA expression of FR α , RFC and GAPDH (as internal control) were as follows: FR α sense, 5'- AAGTGCAGCTGGGAGCT -3', and antisense, 5'- CATTGCACAGAACAGTGGGTG -3'; RFC sense, 5'- CGAAACCTCGGCTTCGGAGC -3', and antisense, 5'- GCACGTAGTAGACCACAGG -3'; GAPDH sense, 5'- TCCATGACAACCTTTGGTATCGTG -3', and antisense, 5'- ACAGTCTTCTGGGTGGCAGTG -3'. The PCR purity was confirmed by gel electrophoresis. Primer sequences (sense, 5'- GTATGCATGGCTTCCTGCAGG -3', and antisense, 5'- ACTTGTTAAACCCTGTAGAGAGG -3') for evaluating FR α gene copy number were designed based on the genomic sequence of intron 3 and intron 4 of FR α (Ensemble database). TRAT1 was used as the reference gene [25]. Ovarian cancer samples having \geq

two-fold increase from their corresponding normal counterparts were considered as positive for FR α gene amplification.

Immunoblotting

Cells were harvested with lysis buffer (0.125 M Tris, pH 6.8 at 22°C containing 1% NP-40 (v/v), 2 mM EDTA, 2 mM N-ethylmaleimide, 2 mM PMSF, 1 mM sodium orthovanadate and 0.1 μ M sodium okadaate), and cleared by centrifugation at 4°C. Protein concentration was determined by DC (detergent compatible) protein assay (Bio-Rad Laboratories, Hercules, CA). 20 μ g protein was resolved by SDS-PAGE, transferred to polyvinylidene difluoride membrane, and hybridized with antibodies specific to FR α (1:1000; Alexis Biochemical, San Diego, CA; ALX-804-439), RFC (1:1000; Affinity BioReagents; Golden, CO; PA1-9553), E-cadherin (1:5000; BD Biosciences; Palo Alto, CA; 610182), and actin (1:1000; Sigma, St. Louis, MO; A5060) and appropriate secondary antibodies (Santa Cruz Biotechnology, Santa Cruz, CA). The blots were developed by Enhanced Chemiluminescence (ECL) Plus detection system (Amersham, Arlington Heights, UK), and visualized with X-ray film (Galen Medical Group, Chattanooga, TN) [22,23,24].

Immunohistochemistry

Immunohistochemical staining was performed as described in earlier reports [22,23,24]. For FR α immunohistochemistry, paraffin sections were treated with goat anti-folate receptor antibody conjugated with horse radish peroxidase (1:200; Abcam; Cambridge, MA; ab20572). Since the anti-FR α antibody from Alexis Biochemical used for immunoblotting failed to have satisfactory result on immunostaining using paraffin sections, another antibody from Abcam was used. Although this antibody may recognize other isoforms of FR, the beta and gamma isoforms of FR were reported to be predominantly expressed in placenta and hematopoietic cells but not in other tissues [26,27]. This antibody can therefore be used to detect FR α immunoreactivity in ovarian cancers. For RFC immunohistochemistry, chicken anti-RFC antibody (1:200; Affinity BioReagents) was applied, followed by biotin-rabbit anti-chicken IgG (H+L). 3-diaminobenzidine-hydrogen peroxide was used as chromogen. Microwave antigen recovery using citrate buffer (pH 6.0) was performed. Omission or substitution of the primary antibody with preimmune IgG serum was used as a negative control. Intensity in stained epithelial cells was scored as 0 (negative), 1 (faint), 2 (moderate), and 3 (strong). The percentage of stained cells was rated as 0 (<5%), 1 (5%–25%), 2 (26%–50%), 3 (51%–75%) and 4 (>75%). Immunoreactivity was assessed by multiplying the staining intensity by the percentage of stained cells to give a composite a composite “HistoScore” [22,23]. High and low levels of FR α and RFC were defined by “HistoScores” cut off at mean.

Demethylation treatment

SKOV-3 and OVCA420 cells were treated with 0, 5 or 10 μ M 5-Aza-2'-deoxycytidine (5-aza-dc, a DNA methylation inhibitor) for 72 hours [28]. Control cells were treated with equal volume of dimethyl sulfoxide (DMSO). Total RNA was extracted from cells. The transcription activity of RFC was determined by qPCR.

DNA preparation, bisulfite treatment and methylation-specific PCR (MSP) analysis

Genomic DNA from frozen clinical samples was extracted using phenol/chloroform. Bisulfite treatment was performed as described [28]. Primers specific to the methylated (sense, 5'- TTCGTCTAGTTTGCGAATG -3', and antisense, 5'- CAA-CACGTACCTAAACGCGA -3') and unmethylated (sense, 5'-

Table 1. Correlation of FR α and RFC immunoreactivities with different diagnostic categories and clinicopathological parameters in ovarian cancer.

Characteristics	Case (n)	FR α		RFC	
		Mean \pm SD	<i>P</i> -value	Mean \pm SD	<i>P</i> -value
Diagnostic categories					
Cysts/benign	11	1.75 \pm 1.30		5.36 \pm 3.11	
Borderline	19	3.96 \pm 2.47		4.50 \pm 3.06	
Carcinomas	83	7.10 \pm 3.25	<u><0.001*</u>	2.94 \pm 2.33	<u><0.008*</u>
Carcinomas \ddagger	21	8.14 \pm 3.06		2.45 \pm 2.40	
Metastatic foci \ddagger	44	7.60 \pm 3.18	0.522 \dagger	3.11 \pm 1.80	0.128 \dagger
Stage (FIGO)					
I	28	5.94 \pm 2.83		2.45 \pm 1.90	
II	12	8.35 \pm 3.36		3.38 \pm 1.98	
III	25	7.72 \pm 3.22		3.60 \pm 2.52	
IV	14	7.14 \pm 3.56	0.092*	2.57 \pm 2.87	0.188*
I	28	5.94 \pm 2.83		2.45 \pm 1.90	
II–IV	51	7.71 \pm 3.31	<u>0.022\dagger</u>	3.27 \pm 2.50	0.113 \dagger
Histological grade					
1	18	5.68 \pm 1.94		3.67 \pm 2.46	
2	38	7.21 \pm 3.39		2.61 \pm 2.16	
3	25	8.18 \pm 3.47	<u>0.042*</u>	2.80 \pm 2.30	0.262*
Low (1)	18	5.68 \pm 1.94		3.67 \pm 2.46	
High (2–3)	63	7.60 \pm 3.43	<u>0.022\dagger</u>	2.68 \pm 2.20	0.123 \dagger
Histology					
Serous	30	8.53 \pm 3.30		3.23 \pm 2.85	
Clear Cell	20	5.13 \pm 2.74		2.50 \pm 2.43	
Endometrioid	26	7.60 \pm 2.83		3.02 \pm 1.73	
Mucinous	7	4.71 \pm 1.90	<u>0.001*</u>	2.64 \pm 1.55	0.780*
Non-mucinous	76	7.32 \pm 3.27		2.97 \pm 2.39	
Mucinous	7	4.71 \pm 1.90	<u>0.027\dagger</u>	2.64 \pm 1.55	0.947 \dagger
Chemoresensitivity\S					
Sensitive	56	7.38 \pm 3.29		2.99 \pm 2.30	
Resistant	15	6.34 \pm 3.03	0.259 \dagger	2.50 \pm 2.21	0.564 \dagger

Intensity values are expressed as “Histoscores” as specified in Methods.

*Kruskal–Wallis rank test;

\dagger Mann–Whitney test;

\ddagger Randomly selected primary carcinomas with matched metastatic foci.

\S Chemoresensitive-patients remained disease free more than 6 months after completion of first-line chemotherapy.

Those with significant *P*-values are underlined.

doi:10.1371/journal.pone.0047201.t001

TTTGTGTTAGTTTGTGAATGG -3', and antisense, 5'-ACAACACATACCTAAACACAA -3') RFC promoter A were reported previously [29]. The annealing temperature was 52°C and 56°C for methylated and unmethylated promoter A respectively. MSP products were detected by electrophoresis on 2% agarose gel with ethidium bromide staining. Normal lymphocyte DNA methylated with Sssl methyltransferase was used as positive control. Untreated genomic DNA and water blanks without DNA were used as negative controls.

Stable knockdown of FR α , ectopic overexpression of RFC and folate treatment in SKOV-3

SKOV-3, an ovarian cancer cell line with relatively high FR α and low RFC expression, was used. To stable knockdown FR α ,

cells were transfected with a set of shRNA constructs against human FR α , pRS-sh FR α (Origene, Rockville, MD), selected with puromycin (1.5 μ g/ml) [22,23,24]. The pRS vector was used as controls. To transient overexpress RFC, pcDNA3-RFC plasmid (kindly provided by Prof L Matherly, Michigan Cancer Foundation) and the empty pcDNA3 vector (control) was transfected into control SKOV-3 cells using Lipofectamine 2000 (Invitrogen) [22,23,24]. Cells were cultured in Medium 199 (Invitrogen)/MCDB 105 (Sigma) medium containing 22.7 nM folic acid and supplemented with 10% fetal bovine serum (FBS) (JRH Biosciences, Lenexa, KS) [22,23]. shFR α cells and RFC overexpressing cells (2 days after transfection) were pretreated with folate-free RPMI 1640 medium (Invitrogen) supplemented with 10% dialyzed FBS containing 0.6 nM folic acid (Invitrogen) for 2 days, trypsinized, counted, plated for functional assays and then treated




Figure 1. Overexpression of FR α in ovarian cancer. (A) qPCR analysis of FR α mRNA in ovarian tumors and the corresponding non-tumor counterparts. (B) mRNA (upper panel) and protein (lower panel) expression of FR α in two immortalized ovarian epithelial cell lines, HOSE 6-3 and HOSE-17-1, and nine ovarian cancer cell lines, OVCAR-3, SKOV-3, OVCA 420, OVCA433, OC316, Dov13, ES-2, TOV21G, SW626, as assessed by qPCR and immunoblotting respectively. (C) Immunoreactivity of FR α in serous (a) and mucinous (e) benign ovarian cystadenomas, serous (b) and mucinous (f) borderline ovarian tumors and serous (c), mucinous (g), clear cell (d) and endometrioid (h) ovarian carcinomas. Scale bar = 100 μ m. (D) Kaplan-Meier overall (left panel) and disease-free (right panel) survival curves for ovarian cancer patients with high and low levels of FR α (cut off at mean). doi:10.1371/journal.pone.0047201.g001

with different doses of folic acid, a synthetic folate, including 0, 6, 12 and 60 nM. The folic acid concentrations used are based on the physiological range in plasma, which ranges from <7 nM in individuals with a negative folate balance to >50 nM in

individuals with >400 μ g/d of folate consumption [30], which is the estimated folate intake by supplement nonusers in North America [13]. The folic acid deficiency concentration selected (12 nM) was based on the observation that such concentration is

the lowest requirement for cell growth [31]. Protein was extracted 2 days after treatment.

MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl tetrazolium bromide) assay

Cell proliferation was determined by MTT assay (Sigma) as described [22,24]. Cells were seeded in 96-well plates with 2000 cells/well. At specific time points, 10 μ l MTT was added to each well. Plates were incubated at 37°C for 4 h, followed by addition of 100 μ l DMSO to each well for dye extraction. Cell proliferation was determined by measuring the absorbance of samples at 570 nm with 630 nm as the reference wavelength.

In vitro migration and invasion assays

In vitro migration and invasion assays were performed as described [22,23,24]. 1.25×10^5 cells were plated on the upper side of a Transwell insert and allowed to migrate through an 8- μ m pore size membrane (migration assays) or invade through a Matrigel-coated membrane (invasion assays). Cells at the upper side of the membrane were removed and the migrated or invaded cells were fixed with methanol, stained with 0.5% crystal violet, and counted under a light microscope in 5 random fields after 24 h or 48 h respectively.

Statistical Analysis

Statistical analysis was performed using SPSS 15.0 for Windows (SPSS Inc., Chicago, IL). Mann-Whitney test was used for comparison between two groups whereas Kruskal–Wallis rank test was used for comparison among multiple groups. Survival analysis was performed by Kaplan–Meier analysis and log-rank test. Cox regression analysis was used for multivariate survival analysis. P values < 0.05 were considered as statistically significant.

Results

Overexpression of FR α was associated with ovarian tumor progression

By qPCR, significantly higher FR α mRNA was found in cancer samples when compared with the corresponding non-tumor counterparts after normalization with GAPDH ($P = 0.015$) (Figure 1A). By immunohistochemistry, strong FR α immunoreactivity was observed in ovarian cancers in contrast to moderate staining of FR α in borderline tumors and weak or absence of staining in benign cystadenomas/inclusion cysts (Figure 1C). Indeed, significantly higher FR α immunoreactivity was detected in ovarian cancers and borderline tumors than in benign cystadenomas/inclusion cysts (all $P < 0.05$, Table 1). At cell lines level, six out of nine ovarian cancer cell lines also showed up-regulation of FR α mRNA and protein expression with SKOV-3, OVCAR-3 and SW626 showing strong expression while OVCA 420, Dov13 and

Table 2. Correlation between clinicopathological parameters and mRNA expression of FR α and RFC in ovarian cancers.

Characteristics	FR α			RFC		
	mRNA expression			mRNA expression		
	Normal (no. of cases)	Increased (no. of cases)	P -value*	Normal (no. of cases)	Decreased (no. of cases)	P -value*
Stage (FIGO)						
I	11	5		4	12	
II	1	2		1	2	
III	2	9		2	9	
IV	2	1	0.062	1	2	0.920
Early (I)	11	5		4	12	
Late (II–IV)	5	12	<u>0.024</u>	4	13	1.000
Histological grade						
1	1	1		0	2	
2	11	5		5	11	
3	4	11	0.064	3	12	0.545
Low (1–2)	12	6		5	13	
High (3)	4	11	<u>0.037</u>	3	12	0.699
Histology						
Serous	3	9		2	10	
Endometrioid	5	6		4	7	
Clear Cell	4	2		1	5	
Mucinous	4	0	0.052	1	3	0.695
Non-mucinous	12	17		7	22	
Mucinous	4	0	<u>0.044</u>	1	3	1.000
Serous	3	9		2	10	
Non-serous	13	8	0.071	6	15	0.443

*Fisher's exact test. Those with significant P -values are underlined.

doi:10.1371/journal.pone.0047201.t002

TOV21G showed weak expression when compared with two normal ovarian epithelium cell lines in which no FR α mRNA and protein expression was detected (Figure 1B).

In clinical samples, high FR α mRNA expression and immunoreactivity were found to be significantly associated with advanced stages of disease and poor histological grade, factors associated with tumor aggressiveness (Tables 1 and 2). However, no significant difference of FR α immunoreactivity was found between the chemosensitive and chemoresistant cases (Table 1). Kaplan-Meier-survival analyses also did not reveal association between high FR α expression and overall or disease-free survival (Figure 1D).

RFC was down-regulated in ovarian cancers and correlated with good prognosis of patients

In contrast to FR α , ovarian cancer samples displayed significantly lower RFC mRNA when compared with the corresponding non-tumor counterparts as assessed by qPCR ($P=0.001$) (Figure 2A). Immunohistochemical analysis also revealed strong RFC immunoreactivity in inclusion cysts/benign cystadenomas and weak expression in ovarian cancers (Figure 2C). Six out of nine ovarian cancer cell lines also displayed down-regulation of RFC mRNA and protein expression when compared with two normal ovarian epithelium cell lines (Figure 2B).

RFC mRNA and immunoreactivity did not correlate with stages of disease, histological grade and histological subtypes (Tables 1 and 2). Interestingly, there was a significant association between low expression of RFC, and shorter overall ($P=0.034$) and disease-free ($P=0.011$) survival (Figure 2D). Moreover, among ovarian cancers with high FR α expression, the overall ($P=0.007$) and disease-free ($P=0.008$) survival was significantly longer in those with high RFC expression (Figure 2E).

FR α gene amplification and RFC promoter methylation contributed to dysregulated gene expression in ovarian cancers

By qPCR, 11 out of 33 (33.3%) cancer samples displayed FR α gene (FOLR1, chromosome 11q13.3) amplification when compared with the corresponding non-tumor counterparts. All amplified cases showed elevated mRNA expression. FR α amplification was correlated with its mRNA expression ($P<0.001$, Fisher's exact test) (Table 3).

On the other hand, reduced expression of RFC in ovarian cancers was related to hypermethylation. After treatment of SKOV-3 and OVCA420 ovarian cancer cells by 5-aza-dc, a DNA methylation inhibitor, two-fold and 2.5-fold increase of RFC gene expression was detected respectively (Figure 3A). Furthermore, promoter hypermethylation of RFC gene (chromosome 21q22.2) was found in 14 out of 33 (42.4%) ovarian cancer samples by MSP. Representative examples of MSP were shown in Figure 3B. In contrast, only 3 out of 33 (9%) of non-tumor samples showed hypermethylation. Unmethylated alleles were detected in all tumor and non-tumor samples. Promoter hypermethylation of RFC significantly inversely correlated with its mRNA expression ($P=0.005$, Fisher's exact test) (Table 3). By MSP, we also detected RFC promoter hypermethylation in five ovarian cancer cell lines SKOV-3, OVCA 420, OVCA433, TOV21G and SW626 (Figure 3B), all of them showed down-regulated RFC mRNA and protein expression (Figure 2B). In contrast, no methylated alleles were detected (Figure 3B) in the normal ovarian epithelium cell line HOSE 6-3 and two cancer cell lines OVCAR-3 and OC316, which displayed RFC mRNA and protein expression (Figure 2B),




Figure 2. Down-regulation of RFC in ovarian cancers and correlation with prognosis of patients. (A) qPCR analysis of RFC mRNA in ovarian tumors and the corresponding non-tumor counterparts. (B) mRNA (upper panel) and protein (lower panel) expression of RFC in two immortalized ovarian epithelial cell lines, HOSE 6-3 and HOSE-17-1, and nine ovarian cancer cell lines, SKOV-3, OVCAR-3, OVCA 420, OVCA433, OC316, Dov13, ES-2, TOV21G, SW626, as assessed by qPCR and immunoblotting respectively. (C) Immunoreactivity of RFC in inclusion cyst (a) and serous ovarian carcinomas (b). Scale bar = 100 μ m. (D) Kaplan-Meier overall (left panel) and disease-free (right panel) survival curves for ovarian cancer patients with high and low levels of RFC (cut off at mean). (E) Kaplan-Meier overall (left panel) and disease-free (right panel) survival curves for high FR α expressed ovarian cancer patients with high and low levels of RFC (cut off at mean). doi:10.1371/journal.pone.0047201.g002

Table 3. Correlation of FR α amplification and RFC promoter methylation with their mRNA expression in ovarian cancers.

FR α		Gene amplification		P-value (Fisher's exact test)
		Non-amplified (no. of cases)	Amplified (no. of cases)	
mRNA expression	Normal (no. of cases)	16	0	<0.001
	Increased (no. of cases)	6	11	
RFC		Promoter DNA methylation		P-value (Fisher's exact test)
		Unmethylated (no. of cases)	Methylated (no. of cases)	
mRNA expression	Normal (no. of cases)	8	0	0.005
	Decreased (no. of cases)	11	14	

doi:10.1371/journal.pone.0047201.t003

Knockdown of FR α altered folate-mediated cell proliferation in SKOV-3 cells

After confirming the specific knockdown of FR α mRNA and protein expression in SKOV-3 cells (Figure 4A), we first determined the effects of FR α on folate-mediated cell proliferation by MTT assay. On days 2 and 4, no significant change of cell proliferation was found in control and shFR α SKOV-3 cells after folate treatment. By day 6, control cells showed proliferation in 12 and 60 nM folate treatment. On Day 8, 6, 12 and 60 nM folate-treated control cells showed dose-dependent proliferation. In contrast, knockdown of FR α blocked folate-mediated cell proliferation (Figure 4B).

Folate through FR α induced SKOV-3 cell migration and invasion and down-regulated E-cadherin

Next, we tested the effect of folate and FR α on SKOV-3 cell migration and invasion. Based on the effects of folate on cell proliferation, 12 and 60 nM doses were chosen for treating control and shFR α SKOV-3 cells. Transwell migration and invasion assays showed that 12 and 60 nM folate significantly induced cell migration and invasion in control cells whereas knockdown of FR α blocked folate-mediated cell migration and invasion (Figure 4C). We then determined the possible downstream target for folate mediated effect on cell migration and invasion. The expression of E-cadherin, an important cell-cell adhesion molecule essential for regulating cell motility, was found to be reduced dose-dependently after folate treatment (Figure 4D). Such down-regulation of E-




Figure 3. Promoter hypermethylation of RFC gene in ovarian cancers. (A) The relative mRNA expression of RFC in SKOV-3 and OVCA 420 after 5-aza-dc treatment with indicated concentrations for 72 hours. Each experiment was performed in triplicate. Bars, means of fold change \pm SD. *, $P < 0.05$. (B) Representative ovarian cancers (CA) (upper panel) and ovarian cell lines (lower panel) of MSP on RFC methylation status. M, DNA marker; P, positive control; N, negative control; M, methylated alleles; U, unmethylated alleles.

doi:10.1371/journal.pone.0047201.g003




Figure 4. Folate induced SKOV-3 cell proliferation, migration and invasion and down-regulated E-cadherin through FR α . (A) Stable knockdown of FR α mRNA and protein in SKOV-3 as detected by qPCR (left panel) and immunoblotting (right panel) respectively. **, $P < 0.005$. (B) Cell proliferation rate of control and shFR α SKOV-3 cells treated with 6, 12 and 60 nM folate at 2, 4, 6 and 8 days displayed as fold change relative to

control without folate treatment (0 nM). $n=3$; *, $P<0.05$. (C) *In vitro* migration (left panel) and invasion assays (right panel) in control and shFR α SKOV-3 cells treated with 0, 12 and 60 nM folate using Transwell membrane without or with Matrigel coating respectively. Upper panels: representative images of migrating or invading SKOV-3 cells. Lower panels: Cell migration or invasion from SKOV-3 presented as percentage of control treated with 0 nM folate; $n=3$; *, $P<0.05$; **, $P<0.005$. (D) Immunoblotting on FR α and E-cadherin using protein lysates prepared from control and shFR α SKOV-3 (left panel). Relative E-cadherin protein level as analyzed by ImageJ software (US National Institutes of Health); $n=3$; *, $P<0.05$; **, $P<0.005$ (right panel).
doi:10.1371/journal.pone.0047201.g004

cadherin after folate treatment was also abrogated after knock-down of FR α .

Ectopic overexpression of RFC in high FR α -expressing SKOV-3 counteracted folate-mediated cell proliferation, migration and invasion and restored E-cadherin expression

We have demonstrated overexpression of FR α and reduced expression of RFC in ovarian cancers, suggesting that they may

exert opposite roles in the progression of ovarian cancer. More importantly, in patients with high FR α , the overall and disease-free survival was significantly longer in those with high RFC expression, implicating the protective role of RFC in high FR α cancers. To elucidate such protective role, *in vitro* functional studies were performed on FR α -positive SKOV-3 cells with ectopically expressed RFC after folate treatment. RFC was found to counteract folate-mediated cell proliferation (Figure 5A), migration and invasion (Figure 5B). Moreover, down-regulation of E-




Figure 5. Ectopic overexpression of RFC in FR α -positive SKOV-3 counteracted folate-mediated cell proliferation, migration and invasion, and down-regulation of E-cadherin. (A) Cell proliferation rate of SKOV-3 cells with ectopically expressed RFC or control vector (control) treated with 60 nM folate after 5 days displayed as fold change compared to control without folate treatment (0 nM); $n=3$; *, $P<0.05$. (B) *In vitro* migration (left panel) and invasion assays (right panel) in SKOV-3 cells with ectopically expressed RFC or control vector treated with 0 and 60 nM folate displayed as percentage of control treated with 0 nM folate; $n=3$; *, $P<0.05$; **, $P<0.005$. (C) Immunoblotting of RFC and E-cadherin using protein lysates prepared from SKOV-3 cells with ectopically expressed RFC or control vector (left panel). Relative E-cadherin protein level as analyzed by ImageJ software (US National Institutes of Health); $n=3$; *, $P<0.05$; **, $P<0.005$ (right panel).
doi:10.1371/journal.pone.0047201.g005

cadherin in cells after folate treatment was also abrogated after overexpressing RFC (Figure 5C).

Discussion

In this study, we demonstrated the progressive increase in FR α mRNA and protein expression from non-tumor tissues, benign and borderline tumors to carcinomas. In addition, FR α gene amplification as a possible mechanism of its overexpression was also demonstrated for the first time in ovarian cancers. Overexpression of FR α in ovarian cancers [19,20] as well as in cancers of kidney, lung and breast have been previously reported [32]. Our results also indicate that a high FR α expression correlates with poor histological grade and advanced stages of disease, suggesting its roles in ovarian tumor progression.

In contrast to FR α , a lower RFC mRNA and protein expression in ovarian cancers was found when comparing with normal tissues or benign tumors. RFC is ubiquitously expressed in normal tissue and is the major folate transport system for transporting natural folates, such as 5-methyl or 5-formyl tetrahydrofolate (THF), and antifolates such as methotrexate (MTX) and pemetrexed [16]. 5-methyl THF is a cofactor essential for DNA methylation which normally leads to the suppression of oncogenes [33]. Thus, loss of RFC has been described to contribute to colonic carcinogenesis [21]. In this study, we found that reduced expression of RFC was significantly associated with shorter overall and disease-free survivals, suggesting that RFC may be considered as a marker for good prognosis in ovarian cancer patients. Moreover, among patients with high FR α expressing ovarian cancers, the overall and disease-free survival was significantly better in those with high RFC expression than those without, implicating the protective role of RFC for patients with these tumors.

We also demonstrated that FR α amplification and RFC promoter methylation correlated with mRNA expression in ovarian cancers. In earlier reports, RFC promoter methylation has been found in breast cancer cells [34] and primary lymphomas [35]. Our findings suggested that up-regulation of FR α (a putative oncogenic folate transporter) and down-regulation of RFC (a putative tumor suppressor type folate transporter) were controlled genetically and epigenetically respectively during ovarian cancer development.

As noted in the introduction above, folate is essential for DNA synthesis [12,13,14], thereby indirectly exerts its effect on cell proliferation. Overexpression of the FR α in NIH/3T3 cells has been reported to induce increased cell growth in vitro and in vivo [36]. Using folate at various dosages ranging from 12 nM (considered as dietary deficient in North America) to 60 nM (considered normal for supplement non-users), we were able to demonstrate cell proliferation in FR α -positive SKOV-3 cells [13]. Conversely, when the FR α was knockdown by shRNA approach, this folate-mediated cell proliferation in SKOV-3 cells was lost, confirming the fact that folate indeed transports through FR α during the process of ovarian cancer cell proliferation. Similarly, intracellular expression of anti-FR antibodies in ovarian cancer cells has been reported to exert growth inhibitory effects as shown by reduced colony formation in soft agar [37].

Besides its effects on cell proliferation, we also demonstrated for the first time that folate increased SKOV-3 cell migration and invasion, possibly through the downregulation of cell-cell adhesion molecule E-cadherin. This effect was abrogated after the

knockdown of FR α . In ovarian cancer, reduced E-cadherin expression has been described in the metastases but not in the corresponding primary ovarian tumors [38]. Patients with such loss of E-cadherin expression were found to have significantly shorter survival [39]. Moreover, simultaneous expression of caveolin-1 and E-cadherin in ovarian cancer cells stabilized adherens junctions through inhibition of src-related kinases [40] whereas loss of E-cadherin enhanced ovarian cancer metastasis through up-regulation of α 5-integrin [41]. In our study, we were able to show E-cadherin down-regulation in folated-treated SKOV-3 cells, suggesting that the folate-mediated enhancing effect on ovarian cancer cell migration and invasion probably acts through FR α via down-regulation of E-cadherin expression.

Interestingly, our study also demonstrates folate-mediated cell proliferation, migration, invasion and E-cadherin reduction in FR α -positive SKOV-3. Such effect is abrogated with ectopically expressed RFC, supporting its tumour suppressive effect in FR α -expressed cells. This in vitro finding further explains the in vivo finding that in patients with high FR α expressing ovarian cancers, the overall and disease-free survival was significantly longer in those with concomitant high RFC expression.

It has been suggested that vitamin supplements, including folate, is beneficial to health. Folate is usually taken as folic acid, its synthetic form, which is fortified in many food products. Supplementation is often believed to be of value to those who suffer from long-term illnesses. Nevertheless, the beneficial effect of folate supplement among cancer patients is controversial [12,13,14]. Earlier studies have shown that folate plays a dual role in colorectal, breast and prostate cancers [12,13,14]. Our results also indicated that folate may potentially enhance the progression and growth of ovarian cancer cells, in particular, those with high FR α and low RFC expressions. Although folate may prevent cancer initiation, once the neoplasia is established, it appears to enhance cancer progression. Should folate supplements, generally considered by the public as a healthy option, therefore be taken more cautiously? Additional studies to further explore the benefits or harmful effects of folate supplement in cancer patients are necessary.

In conclusion, we demonstrated that folate and FR α contribute to the progression and growth of ovarian cancer cells through the regulation of cell proliferation, migration and invasion. In contrast, RFC can serve as a balancing partner of FR α and seems to exert a protective role in ovarian cancer patients, conferring longer survival among patients with cancers that showed a high FR α expression status. We also demonstrated a mechanistic link between folate, FR α , RFC and E-cadherin. The potential of FR α and RFC as alternative molecular therapeutic target or prognostic marker in ovarian cancers should be further explored, respectively.

Acknowledgments

We thank Prof L Matherly for providing the RFC plasmid.

Author Contributions

Conceived and designed the experiments: MKYS ANYC. Performed the experiments: MKYS DSHK HYC ESYW. Analyzed the data: MKYS ANYC. Contributed reagents/materials/analysis tools: LLJ HYSN XFL. Wrote the paper: MKYS ANYC. Critically reviewed the paper: PPCI.

References

- Jemal A, Siegel R, Ward E, Murray T, Xu J, et al. (2007) Cancer statistics, 2007. *CA Cancer J Clin* 57: 43–66.
- Goodman MT, Howe HL, Tung KH, Hotes J, Miller BA, et al. (2003) Incidence of ovarian cancer by race and ethnicity in the United States, 1992–1997. *Cancer* 97: 2676–2685.

3. Marugame T, Hirabayashi Y (2007) Comparison of time trends in ovary cancer incidence (1973–1997) in East Asia, Europe, and the USA, from Cancer Incidence in Five Continents Vols IV VIII. *Jpn J Clin Oncol* 37: 802–803.
4. The Hong Kong Cancer Registry (2007) Department of Health Annual Report. Hong Kong, Department of Health, Hospital Authority.
5. Woodside JV, McCall D, McGartland C, Young IS (2005) Micronutrients: dietary intake v. supplement use. *Proc Nutr Soc* 64: 543–553.
6. Larsson SC, Giovannucci E, Wolk A (2004) Dietary Folate Intake and Incidence of Ovarian Cancer: The Swedish Mammography Cohort. *J Natl Cancer Inst* 96: 396–402.
7. Kelemen LE, Sellers TA, Vierkant RA, Harnack L, Cerhan JR (2004) Association of folate and alcohol with risk of ovarian cancer in a prospective study of postmenopausal women. *Cancer Causes Control* 15: 1085–1093.
8. Kelemen LE, Sellers TA, Keeney GL, Lingle WL (2005) Multivitamin and alcohol intake and folate receptor alpha expression in ovarian cancer. *Cancer Epidemiol Biomarkers Prev* 14: 2168–2172.
9. Pelucchi C, Mereghetti M, Talamini R, Negri E, Montella M, et al. (2005) Dietary folate, alcohol consumption, and risk of ovarian cancer in an Italian case-control study. *Cancer Epidemiol Biomarkers Prev* 14: 2056–2058.
10. Blount BC, Mack MM, Wehr CM, MacGregor JT, Hiatt RA, et al. (1997) Folate deficiency causes uracil misincorporation into human DNA and chromosome breakage: implications for cancer and neuronal damage. *Proc Natl Acad Sci* 94: 3290–3295.
11. Duthie SJ (1999) Folic acid deficiency and cancer: mechanisms of DNA instability. *Br Med Bull* 55: 578–592.
12. Smith AD, Kim YI, Refsum H (2008) Is folic acid good for everyone? *Am J Clin Nutr* 87: 517–533.
13. Kim YI (2008) Folic acid supplementation and cancer risk: point. *Cancer Epidemiol Biomarkers Prev* 17: 2220–2225.
14. Ulrich CM (2008) Folate and cancer prevention—where to next? Counterpoint. *Cancer Epidemiol Biomarkers Prev* 17: 2226–2230.
15. Matherly LH, Goldman DI (2003) Membrane transport of folates. *Vitam Horm* 66: 403–456.
16. Matherly LH, Hou Z, Deng Y (2007) Human reduced folate carrier: translation of basic biology to cancer etiology and therapy. *Cancer Metastasis Rev* 26: 111–128.
17. Ross JF, Chaudhuri PK, Ratnam M (1994) Differential regulation of folate receptor isoforms in normal and malignant tissues in vivo and in established cell lines. Physiologic and clinical implications. *Cancer* 73: 2432–2443.
18. Campbell IG, Jones TA, Foulkes WD, Trowsdale J (1991) Folate-binding protein is a marker for ovarian cancer. *Cancer Res* 51: 5329–5338.
19. Toffoli G, Cernigoi C, Russo A, Gallo A, Bagnoli M, et al. (1997) Overexpression of folate binding protein in ovarian cancers. *Int J Cancer* 74: 193–198.
20. Kalli KR, Oberg AL, Keeney GL, Christianson TJ, Low PS, et al. (2008) Folate receptor alpha as a tumor target in epithelial ovarian cancer. *Gynecol Oncol* 108: 619–626.
21. Ma DW, Finnell RH, Davidson LA, Callaway ES, Spiegelstein O, et al. (2005) Folate transport gene inactivation in mice increases sensitivity to colon carcinogenesis. *Cancer Res* 65: 887–897.
22. Siu MK, Wong ES, Chan HY, Kong DS, Woo NW, et al. (2010) Differential expression and phosphorylation of Pak1 and Pak2 in ovarian cancer: effects on prognosis and cell invasion. *Int J Cancer* 127: 21–31.
23. Siu MK, Chan HY, Kong DS, Wong ES, Wong OG, et al. (2010) p21-activated kinase 4 regulates ovarian cancer cell proliferation, migration, and invasion and contributes to poor prognosis in patients. *Proc Natl Acad Sci U S A* 107: 18622–18627.
24. Siu MK, Wong ES, Chan HY, Ngan HY, Chan KY, et al. (2008) Overexpression of NANOG in gestational trophoblastic diseases: effect on apoptosis, cell invasion, and clinical outcome. *Am J Pathol* 173: 1165–1172.
25. Woenckhaus J, Steger K, Sturm K, Munstedt K, Franke FE, et al. (2007) Prognostic value of PIK3CA and phosphorylated AKT expression in ovarian cancer. *Virchows Arch* 450: 387–395.
26. Nakashima-Matsushita N, Homma T, Yu S, Matsuda T, Sunahara N, et al. (1999) Selective expression of folate receptor beta and its possible role in methotrexate transport in synovial macrophages from patients with rheumatoid arthritis. *Arthritis Rheum* 42: 1609–1616.
27. Kelemen LE (2006) The role of folate receptor alpha in cancer development, progression and treatment: cause, consequence or innocent bystander? *Int J Cancer* 119: 243–250.
28. Shen DH, Chan KY, Khoo US, Ngan HY, Xue WC, et al. (2005) Epigenetic and genetic alterations of p33ING1b in ovarian cancer. *Carcinogenesis* 26: 855–863.
29. Yang R, Li WW, Hoang BH, Kim H, Banerjee D, et al. (2008) Quantitative correlation between promoter methylation and messenger RNA levels of the reduced folate carrier. *BMC Cancer* 8: 124.
30. Kimura M, Umegaki K, Higuchi M, Thomas P, Fenech M (2004) Methylene tetrahydrofolate reductase C677T polymorphism, folic acid and riboflavin are important determinants of genome stability in cultured human lymphocytes. *J Nutr* 134: 48–56.
31. Beetsra S, Salisbury C, Turner J, Altree M, McKinnon R, et al. (2005) Lymphocytes of BRCA1 and BRCA2 germ-line mutation carriers, with or without breast cancer, are not abnormally sensitive to the chromosome damaging effect of moderate folate deficiency. *Carcinogenesis*: [Epub ahead of print].
32. Parker N, Turk MJ, Westrick E, Lewis JD, Low PS, et al. (2005) Folate receptor expression in carcinomas and normal tissues determined by a quantitative radioligand binding assay. *Anal Biochem* 338: 284–293.
33. Choi SW, Mason JB (2000) Folate and carcinogenesis: an integrated scheme. *J Nutr* 130: 129–132.
34. Worm J, Kirkin AF, Dzhandzhugazyan KN, Guldberg P (2001) Methylation-dependent silencing of the reduced folate carrier gene in inherently methotrexate-resistant human breast cancer cells. *J Biol Chem* 276: 39990–40000.
35. Ferreri AJ, Dell’Oro S, Capello D, Ponzoni M, Iuzzolino P, et al. (2004) Aberrant methylation in the promoter region of the reduced folate carrier gene is a potential mechanism of resistance to methotrexate in primary central nervous system lymphomas. *Br J Haematol* 126: 657–664.
36. Bottero F, Tomassetti A, Canevari S, Miotti S, Menard S, et al. (1993) Gene transfection and expression of the ovarian carcinoma marker folate binding protein on NIH/3T3 cells increases cell growth in vitro and in vivo. *Cancer Res* 53: 5791–5796.
37. Figini M, Ferri R, Mezzanzanica D, Bagnoli M, Luison E, et al. (2003) Reversion of transformed phenotype in ovarian cancer cells by intracellular expression of anti folate receptor antibodies. *Gene Ther* 10: 1018–1025.
38. Veatch AL, Carson LF, Ramakrishnan S (1994) Differential expression of the cell-cell adhesion molecule E-cadherin in ascites and solid human ovarian tumor cells. *Int J Cancer* 58: 393–399.
39. Darai E, Scoazec JY, Walker-Combrouze F, Mlika-Cabanne N, Feldmann G, et al. (1997) Expression of cadherins in benign, borderline, and malignant ovarian epithelial tumors: a clinicopathologic study of 60 cases. *Hum Pathol* 28: 922–928.
40. Miotti S, Tomassetti A, Facetti I, Sanna E, Berno V, et al. (2005) Simultaneous expression of caveolin-1 and E-cadherin in ovarian carcinoma cells stabilizes adherens junctions through inhibition of src-related kinases. *Am J Pathol* 167: 1411–1427.
41. Sawada K, Mitra AK, Radjabi AR, Bhaskar V, Kistner EO, et al. (2008) Loss of E-cadherin promotes ovarian cancer metastasis via alpha 5-integrin, which is a therapeutic target. *Cancer Res* 68: 2329–2339.